

Immunochemical and Mass Spectrometric Analysis of N()-(Carboxymethyl)lysine Content of AGE-BSA Systems Prepared with and without Selected Antiglycation Agents.

Connolly, L. (2010). Immunochemical and Mass Spectrometric Analysis of N()-(Carboxymethyl)lysine Content of AGE-BSA Systems Prepared with and without Selected Antiglycation Agents. *Journal of Agricultural and Food Chemistry*, 58(22)(22), 11955-11961. DOI: 10.1021/jf102543g

Published in:
Journal of Agricultural and Food Chemistry

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Immunochemical and Mass Spectrometric Analysis of *N*^ε-(Carboxymethyl)lysine Content of AGE–BSA Systems Prepared with and without Selected Antiglycation Agents

CHOU SREY,^{*,†} SIMON A. HAUGHEY,[†] LISA CONNOLLY,[†]
 MARIA DOLORES DEL CASTILLO,[§] JENNIFER M. AMES,[#] AND CHRISTOPHER T. ELLIOTT[†]

[†]Institute of Agri-Food and Land Use, School of Biological Sciences, Queen's University,
 Belfast, Northern Ireland BT7 5AG, [§]Department of Food Analysis and Bioactivity,
 Institute of Food Science Research (CSIC-UAM), C/Nicolás Cabrera 9, 28049 Madrid, Spain, and
[#]School of Life Sciences, Northumbria University at Newcastle, England NE1 8ST

The present study was designed to compare surface plasmon resonance (SPR) biosensor, enzyme-linked immunosorbent assay (ELISA), and ultraperformance liquid chromatography–tandem mass spectrometry (UPLC-MS/MS) methods for the analysis of *N*^ε-(carboxymethyl)lysine (CML) in glucose–bovine serum albumin (BSA) model systems and to investigate the possible inhibitory effect of selected compounds (α -tocopherol, ferulic acid, rutin, thiamin, thiamin monophosphate, and thiamin pyrophosphate) on CML formation. The reported levels of CML detected were dependent upon the method of analysis employed. The highest reported concentrations were obtained with the SPR biosensor, whereas the lowest were found by ELISA. However, a high correlation was observed between these two immunochemical procedures. CML concentrations were dependent upon the type and concentration of the candidate CML inhibitor. All inhibitory compounds investigated, with the exception of α -tocopherol, decreased the level of CML formation in the glucose–BSA system.

KEYWORDS: *N*^ε-(Carboxymethyl)lysine (CML); advanced glycation endproducts (AGEs); CML inhibitor; enzyme-linked immunosorbent assay (ELISA); surface plasmon resonance (SPR) biosensor; ultraperformance liquid chromatography–tandem mass spectrometry (UPLC-MS/MS)

INTRODUCTION

Advanced glycation endproducts (AGEs) are a class of Maillard reaction (MR) products (MRPs). Chemically, the MR involves a reaction between a free amino group, for example, the ϵ -amino groups of lysine residues within protein, with the carbonyl group of a reducing sugar, such as glucose (1). *N*^ε-(Carboxymethyl)lysine (CML), one of the best known AGEs, can be formed through a number of different pathways. The condensation reaction between glucose and the ϵ -amino group of lysine forms fructose-lysine (the Amadori rearrangement product, ARP, of the reaction), which is subsequently oxidized to form CML. Glyoxal can be formed from the oxidation of glucose, and it can also react directly with the ϵ -amino group of lysine to form CML (2). CML has been associated with aging and diseases such as renal failure and diabetes (3, 4). Furthermore, the accumulation of CML in the hearts of patients with diabetes may contribute to the increased risk of heart failure associated with hyperglycemia (5). Therefore, the search for inhibitors of CML formation is of significant medical interest. The effectiveness of various potential glycation inhibitors has been tested in model systems (6, 7), in foods (8), and in vivo (9). Mechanisms include reactive

carbonyl trapping (6, 10), antioxidant activity (7, 8), sugar autoxidation inhibition (6), and amino group binding inhibition/competition (9, 11).

Several analytical methods have been reported for the detection and quantification of CML, including LC-MS/MS (12), GC-MS (13), and ELISA (14). Recently, new analytical approaches such as UPLC-MS/MS (15) and surface plasmon resonance imaging (SPRI) biosensor assays (16) have been employed for the quantitative analysis of CML. Only a small number of reports on inter-/intralaboratory comparisons of CML data obtained via different analytical approaches have been published to date in the scientific literature (13, 16). Biosensor assays have proved to be versatile, robust, and capable of producing rapid and reliable data for the analysis of a wide range of components in complex food matrices with minimal sample preparation (17–20). The main difference between ELISA and SPR biosensor assays is that the SPR biosensor approach is a label-free technique which relies for detection upon an increase in molecular mass due to antibody–analyte interactions on a chip surface.

Research has been conducted on AGE inhibitors (8, 21), but a small number of reports have been focused on CML. No comparison of levels of CML determined by SPR, ELISA, and UPLC-MS/MS has been reported. Therefore, the present study aimed to employ three analytical methods in the detection of CML and to investigate the effect of three antioxidants

*Address correspondence to this author at the Institute of Agri-Food and Land Use, Queen's University, David Keir Building, Stranmillis Road, Belfast BT9 5AG, Northern Ireland [phone + 44 (0)28 9097 6543; e-mail chousrey@yahoo.fr or csrey01@qub.ac.uk].

71 (α -tocopherol, ferulic acid, and rutin hydrate) and competitors
72 (thiamin, thiamin monophosphate, and thiamin pyrophosphate)
73 on the formation of CML, in AGE-bovine serum albumin
74 (BSA) model systems.

75 MATERIALS AND METHODS

76 **Reagents and Apparatus.** All of the reagents used in this study were of
77 analytical grade. α -Tocopherol, bovine serum albumin (BSA, fraction V),
78 ferulic acid, glucose, glyoxylic acid, keyhole limpet hemocyanin (KLH),
79 *N*-hydroxysuccinimide (NHS), *N*-(3-dimethylaminopropyl)-*N'*-ethylcar-
80 bodiimide hydrochloride (EDC), polyethylene glycol sorbitan monolaurate
81 (Tween 20), rutin hydrate, sodium cyanoborohydride, sodium phosphate
82 monobasic, sodium phosphate dibasic, sodium pyruvate, thiamin hydro-
83 chloride, thiamin monophosphate, thiamin pyrophosphate, and other
84 chemicals were purchased from Sigma-Aldrich (Gillingham, U.K.).
85 Dimethyl sulfoxide (DMSO), Hybridoma Feeder Supplement (Doma-
86 Drive), Dulbecco's modified eagle's medium (DMEM), heat-inactivated
87 fetal calf serum (HI-FCS), hypoxanthine aminopterin thymidine (HAT)
88 medium, penicillin streptomycin (pen strep), and polyethylene glycol
89 (PEG) were from Invitrogen (Paisley, U.K.). Gelatin and horseradish
90 peroxidase-linked anti-mouse immunoglobulin were obtained from
91 DAKO (Cambridge, U.K.). 3,3',5,5'-Tetramethylbenzidine (TMB)
92 solution was from Chemicon International (Temecula, CA). The
93 optical surface plasmon resonance (SPR) biosensor system (Biacore
94 Q), Biosensor chip (CM5), ethanolamine hydrochloride (1 M), and
95 HBS-EP buffer (0.01 M HEPES, 0.15 M NaCl, 3 mM EDTA, 0.005%
96 polysorbate 20 (v/v), pH 7.4) were supplied by GE Healthcare
97 (Uppsala, Sweden). Nunc-Immuno 96 microwell plates (NUNC
98 Brand Products) were from Thermo-Scientific, Denmark. The Tecan
99 Safire plate reader was from Vector Scientific, Ireland. The Genevac
100 evaporator (EZ-2) was from Ipswich, U.K. A Waters (Manchester, U.
101 K.) Acquity UPLC triple-quadrupole MS/MS was used for mass
102 spectrometric analyses.

103 **Sample Preparation.** *Glycated Bovine Serum Albumin (Glycated*
104 *BSA).* BSA (10 mg, equivalent to 8.85 mM lysine), glucose (90 mg, 0.5 M),
105 and inhibitors (8.85 and 88.5 mM α -tocopherol, ferulic acid, rutin hydrate,
106 thiamin hydrochloride, thiamin monophosphate, thiamin pyrophosphate)
107 were mixed in sodium phosphate buffer (0.2 M, pH 7.2, 1 mL) to bring the
108 molecular ratio of lysine/inhibitor to 1:1 and 1:10. The mixed samples were
109 incubated, in a 5 mL glass bottle with a screw-tight lid, at 50 °C for 10 days
110 and vortexed once per day. Glycated BSA, without addition of an
111 inhibitor, was prepared by mixing BSA (10 mg) with glucose (90 mg) in
112 phosphate buffer (0.2 M, pH 7.2, 1 mL) and used as a positive control to
113 calculate the percentage inhibition of CML formation caused by the
114 trialed compounds. All of the samples were prepared in triplicate and were
115 stored at -20 °C prior to analysis.

116 *N^ε-(Carboxymethyl)lysine-Bovine Serum Albumin/Keyhole Limpet*
117 *Hemocyanin Protein (CML-BSA/KLH).* CML-BSA was prepared by
118 incubating BSA (10 mg, equivalent to 8.85 mM lysine) with glyoxylic acid
119 (8.85 mM) in phosphate buffer (pH 7.5, 0.5 M) for 1 h at 37 °C, and then
120 sodium cyanoborohydride (17.70 mM) was added and the incubation
121 continued for a further 23 h. The pH was adjusted to 7.4 with NaOH
122 (0.1 M), if required. CML-modified KLH was prepared by incubating
123 KLH (10 mg) with glyoxylic acid (260 mM) in phosphate buffer (pH 7.5,
124 0.5 M) for 1 h at 37 °C, and then sodium cyanoborohydride (520 mM) was
125 added and the incubation continued for a further 23 h. Blank samples were
126 also prepared as previously stated, but with the omission of glyoxylic acid.
127 The incubated solutions were dialyzed against phosphate buffer (0.05 M,
128 pH 7.2) containing 0.15 M NaCl and stored at -20 °C. The conjugates
129 were used to prepare immunogens to raise antibodies and for use in the
130 ELISA and SPR assays.

131 *N^ε-(Carboxyethyl)lysine-Modified Bovine Serum Albumin (CEL-*
132 *BSA).* CEL-BSA was prepared by incubating BSA (20 mg/mL), sodium
133 pyruvate (17.14 mM, 17.14 μ L), and sodium cyanoborohydride
134 (25.71 mM, 25.7 μ L) in phosphate buffer (0.2 M, pH 7.4, 1 mL). The solu-
135 tion was incubated at 37 °C for 24 h. A control was also prepared using the
136 same conditions but with the omission of sodium pyruvate.

137 **Preparation of a Monoclonal Antibody to CML.** The immuniza-
138 tion and fusion procedure described previously by Stewart et al. (17) was

used to prepare anti-CML monoclonal antibody. Briefly, three BALB/c
mice were immunized at 3 week intervals with CML-KLH immunogen
(20 μ g of protein). Primary and secondary booster immunizations were
administered using Quil A adjuvant by subcutaneous injection. Third
(20 μ g protein) and fourth (80 μ g protein) boosters were administered by
intraperitoneal injection with Freund's complete adjuvant. Tail bleeds
taken from mice 10 days after each booster were tested using ELISA and
SPR assays. The most responsive mouse, as determined by antibody titer,
was selected and, 4 days prior to the fusion being performed, received a
final booster intraperitoneally of the immunogen (100 μ g of protein) in
phosphate-buffered saline (pH 7.2). The fusion was performed according
to a modification of the method of Kohler and Milstein (22). A single cell
suspension was collected from the spleen of the immunized mouse and
fused with SP2/O-Ag14 myeloma cells using polyethylene glycol. After
10–14 days of fusion, the resulting hybrid cells (hybridomas) were
screened using ELISA and SPR assays. Serum from the final heart bleed
of the fusion mouse was used as a positive control, and cell culture me-
dium buffer was used as a negative control, in the screening assays.
The hybridomas that produced antibodies specific for CML but did not
bind to CEL were selected for further investigation. All cell lines that gave
a strong binding to CML-coated biosensor chips (as detected by SPR)
were cloned twice and selected for scale-up antibody production, and the
products were stored in liquid nitrogen.

Development of an Enzyme-Linked Immunosorbent Assay (ELISA).
Ninety-six-well Nunc Maxisorp plates were coated with CML-BSA (1 μ g/
mL, 100 μ L) and blocked with PBS/gelatin (1%) blocking buffer (0.1 M,
pH 7.2, 200 μ L) overnight at room temperature. After the blocking buffer
was discarded, 50 μ L of glycated BSA (with or without inhibitors, 300 μ g/
mL) and 50 μ L of anti-CML antibody, 1:15000-fold dilution, were added
to the wells and incubated by shaking at 37 °C for 90 min. The supernatant
was discarded and the plate washed three times with wash buffer (1%
Tween 20 and 0.9% NaCl). The secondary goat anti-mouse HRP antibody
was added (2000 dilution) and incubated at 37 °C for 60 min. The
supernatant was again discarded, and the plate was washed three times
with wash buffer. TMB (100 μ L) was added to each well and developed in
darkness for 5 min. The substrate reaction was stopped using sulfuric acid
(2.5 M, 25 μ L/well). Absorbance was read at 450 nm using
a microplate reader. A range of CML-BSA standards (50%, 0–100 μ g
of BSA/mL, equivalent to 0–3500 μ g of CML/g BSA) were also added to
a number of wells and used to generate a calibration curve. Prism
5 software was used to calculate CML concentration in the samples.
Buffer (no antibody) was used as a negative control. Unheated glycated
BSA (prepared with and without an inhibitor) was tested for background
effects.

Development of the SPR Biosensor Assay. *CML-BSA Immobilization onto a Biosensor Chip (CM5).* The CM5 sensor chip is composed
of a glass slide coated with a thin layer of gold to which a carboxymethyl-
ated dextran matrix is covalently attached. The conditions for the
immobilization of the CML-BSA were optimized (e.g., concentration of
CML-BSA, speed of injection, and contact time) to ensure a high surface
coverage of the protein. The immobilization procedure was performed
within the biosensor unit using Biacore control software. Briefly, the con-
centration, injection speed, and contact time of each solution were carried
out as follows: EDC (0.4 M, 50%) and NHS (0.1 M, 50%) were mixed and
injected for 7 min (10 μ L/min) onto the CM5 chip (flow cell 1), to activate
carboxyl groups on the chip. Jeffamine diluted (1:5) in sodium borate
buffer (pH 8.5, 63 mM) was injected (5 μ L/min, 7 min) to cover the surface
of the flow cell with amino groups. The flow cell was then deactivated with
ethanolamine-HCl (10 μ L/min, 3 min). After deactivation, CML-BSA
solutions were coated for 10 min (1 mg/mL). The coated chip was ready for
use immediately or could be stored at 4 °C for several months under dry
conditions. After use, the chip was washed with distilled water and dried
under a gentle stream of nitrogen.

Samples Analysis Using the SPR Biosensor Assay. Glycated BSA
samples (10 mg of BSA/mL, 2 μ L) and HBS-EP buffer (pH 7.4, 38 μ L)
were pipetted and mixed in a 96-well plate to bring the final concentration
to 0.5 mg of BSA/mL. The CML-specific monoclonal antibodies were
diluted in HBS-EP buffer (60-fold dilution). The antibody was mixed
automatically with glycated BSA (50:50) in wells of a 96-microtiter plate
and then injected over the CML-BSA coated chip at a flow rate of 10 μ L/
min and a contact time 2 min. The chip surface was regenerated




Figure 1. CML concentration in glycated BSA as determined by SPR biosensor analysis. No inhibitor, AGE-BSA without addition of any inhibitor; α -tocopherol, AGE-BSA with addition of α -tocopherol; ferulic acid, AGE-BSA with addition of ferulic acid; rutin, AGE-BSA with addition of rutin; thiamin hydrochloride, AGE-BSA with addition of thiamin hydrochloride; thiamin monophosphate, AGE-BSA with addition of thiamin monophosphate; thiamin pyrophosphate, AGE-BSA with addition of thiamin pyrophosphate. Data are expressed as the mean \pm SD ($n = 3$).

210 with 50 mM NaOH (flow rate = 25 μ L/min, 1 min contact time).
 211 The concentration of CML in glycated samples was calculated against a
 212 CML-BSA calibration curve (50%, 0–100 μ g of BSA/mL, equivalent to
 213 0–3500 μ g of CML/g BSA, $R^2 = 0.997$) and using Prism 5 software.

214 **UPLC-MS/MS Analysis.** The CML content of glycated BSA was
 215 determined by UPLC-MS/MS (15). Samples were prepared for analysis by
 216 sodium borohydride reduction, protein isolation using TCA precipitation,
 217 protein hydrolysis with 6 M HCl at 110 $^{\circ}$ C for 24 h, and solid phase
 218 extraction using a C_{18} cartridge. Protein hydrolysates (equivalent to 7.5 μ g
 219 of protein, 7.5 μ L) were injected into a BEH C_{18} UPLC column (Waters,
 220 2.1 \times 50 mm, 1.7 μ m) housed in a column oven at 50 $^{\circ}$ C in gradient
 221 elution mode. Solvent A was nonafluoropentanoic acid (NFPA, 5 mM),
 222 and solvent B was acetonitrile. The injection time was 7.5 min. The analysis
 223 was performed using a Waters Acquity UPLC (Manchester, U.K.) coupled
 224 to a Waters Premier triple-quadrupole MS operating in multiple
 225 reaction monitoring (MRM) mode. The flow rate was 0.2 mL/min. The
 226 MS was operated in electrospray ionization (ESI) positive mode using
 227 MRM mode. The CML data were analyzed using MassLynx software.
 228 CML concentrations in the samples were quantified by means of reference
 229 to the internal standard. Data were reported as the mean \pm SD.

230 **Statistical Analysis.** Statistical analysis (ANOVA) was performed to
 231 determine differences between three groups of means ($P < 0.01$). Limits of
 232 detection (LOD) and limits of quantification (LOQ) of CML concentration
 233 analysis by immunochemical methods (SPR and ELISA) were determined
 234 from three independent runs of 20 unheated AGE-BSA samples. The
 235 LOD and LOQ of CML concentration, analysis by UPLC-MS/MS
 236 method, were 1.62 and 5.41 μ g of CML/g of BSA, respectively (15).

237 RESULTS

238 **Anti-CML Monoclonal Antibody.** Fourteen days after fusion,
 239 447 hybridoma supernatants were screened by two methods 447
 240 hybridoma supernatants gave 23 positive reactors to CML-
 241 BSA, whereas screening by SPR biosensor revealed 5 positive
 242 reactors. These 5 positive reactors (shown by SPR) were also
 243 found to be positive binders by ELISA and were considered
 244 to be true positive binders. After further testing using CML-
 245 BSA and CEL-BSA in assay inhibition studies, 1 of the 5
 246 positive reactors was selected for full assay development and
 247 designated 2C1. This monoclonal antibody was chosen be-
 248 cause it exhibited the highest specificity of the five positive
 249 reactors for CML-BSA but not CEL-BSA as observed in
 250 inhibition binding ELISA. 2C1 was applied to ELISA and
 251 SPR assays for the determination of CML in glycated BSA.

252 **SPR Analysis of CML in Glycated BSA.** The SPR-measured
 253 level of CML in glycated BSA model samples is shown in Figure 1.
 254 CML levels in the samples were (45 \pm 4.76)–(1603 \pm 156) μ g of

255 CML/g of BSA. The highest level of CML was in BSA glycated
 256 either in the absence of any inhibitor (1543 \pm 119 μ g of CML/g of
 257 BSA) or in the presence of the antioxidant α -tocopherol (1603 \pm
 258 156 μ g of CML/g of BSA). The lowest concentrations of CML
 259 were found in BSA glycated in the presence of rutin or thiamin
 260 pyrophosphate (127 \pm 5.20 and 45 \pm 4.76 μ g of CML/g of BSA,
 261 respectively). BSA glycated in the presence of thiamin or thiamin
 262 monophosphate gave similar concentrations of CML (223 \pm 20.1
 263 and 233 \pm 20.2 μ g of CML/g of BSA), about 5-fold higher than in
 264 the presence of thiamin pyrophosphate.

265 **ELISA Analysis of CML in Glycated BSA.** CML levels in
 266 glycated BSA measured by ELISA are shown in Figure 2. Con-
 267 centrations were (6 \pm 0.80)–(900 \pm 99.0) μ g of CML/g of BSA.
 268 Again, the highest levels were in BSA glycated without an
 269 inhibitor and in BSA glycated in the presence of α -tocopherol
 270 (852 \pm 27.82 and 900 \pm 99.0 μ g of CML/g of BSA, respectively).
 271 Once more, BSA glycated in the presence of rutin or thia-
 272 min pyrophosphate provided the lowest concentration of CML
 273 (8 \pm 0.55 and 6 \pm 0.37 μ g of CML/g of BSA, respectively),
 274 whereas samples containing thiamin and thiamin monophos-
 275 phate gave very similar levels of CML formation (41 \pm 2.20 and
 276 66 \pm 3.43 μ g of CML/g of BSA, respectively), which were about
 277 7-fold higher than those obtained in the presence of thiamin
 278 pyrophosphate.

279 **UPLC-MS/MS Analysis of CML in Glycated BSA.** CML levels
 280 of glycated BSA measured by UPLC-MS/MS are shown in
 281 Figure 3. The concentrations of CML in samples varied between
 282 194 \pm 19.7 and 1075 \pm 68.6 μ g/g of BSA. The highest concentra-
 283 tions of CML were determined in glycated BSA in the absence of
 284 an inhibitor and samples containing α -tocopherol (960 \pm 14.4
 285 and 1075 \pm 68.6 μ g of CML/g of BSA, respectively), whereas the
 286 lowest concentrations of CML were found in glycated samples
 287 containing rutin and thiamin pyrophosphate (389 \pm 17.7 and 194
 288 \pm 19.7 μ g of CML/g of BSA, respectively). The level of CML in
 289 glycated BSA in the presence of thiamin or thiamin monophos-
 290 phate gave similar data (467 \pm 25.4 and 433 \pm 39.7 μ g of CML/g
 291 of BSA, respectively), which were 2.4-fold higher than those of
 292 thiamin pyrophosphate.

293 **Comparison of CML Data Determined by SPR, ELISA, and**
 294 **UPLC-MS/MS.** SPR, ELISA, and UPLC-MS/MS methods for
 295 CML analysis were validated and the results of a range of samples
 296 compared by linear regression. The three methods showed similar
 297 trends with regard to analysis of samples of BSA glycated in the
 298 presence of different inhibitors. CML concentrations obtained by




Figure 2. CML concentration in glycated BSA determined by ELISA assay. No inhibitor, AGE–BSA without addition of any inhibitor; α -tocopherol, AGE–BSA with addition of α -tocopherol; ferulic acid, AGE–BSA with addition of ferulic acid; rutin, AGE–BSA with addition of rutin; thiamin hydrochloride, AGE–BSA with addition of thiamin hydrochloride; thiamin monophosphate, AGE–BSA with addition of thiamin monophosphate; thiamin pyrophosphate, AGE–BSA with addition of thiamin pyrophosphate. Data are expressed as the mean \pm SD ($n = 3$).




Figure 3. CML concentration in glycated BSA determined by UPLC-MS/MS analysis. No inhibitor, AGE–BSA without addition of any inhibitor; α -tocopherol, AGE–BSA with addition of α -tocopherol; ferulic acid, AGE–BSA with addition of ferulic acid; rutin, AGE–BSA with addition of rutin; thiamin hydrochloride, AGE–BSA with addition of thiamin hydrochloride; thiamin monophosphate, AGE–BSA with addition of thiamin monophosphate; thiamin pyrophosphate, AGE–BSA with addition of thiamin pyrophosphate. Data are expressed as the mean \pm SD ($n = 3$).

F4 299 SPR, ELISA, and UPLC-MS/MS methods are shown in **Figure 4**.
 300 Correlations between CML levels determined by SPR versus
 301 ELISA, UPLC-MS/MS versus SPR, and UPLC-MS/MS versus
 302 ELISA methods of analysis are shown in panels **a**, **b**, and **c**,
 F5 303 respectively, of **Figure 5**.

304 The SPR biosensor analysis of CML gave the highest LOD and
 305 LOQ (1.7 and 17 μg of CML/g of BSA, respectively), whereas
 306 ELISA analysis gave the lowest (1.51 and 5.20 μg of CML/g of
 307 BSA, respectively). The LOD and LOQ for UPLC-MS/MS were
 308 1.62 and 5.41 μg of CML/g of BSA, respectively, very similar
 309 to the values for the ELISA. The intra-assay % CV for samples
 310 analyzed by each method was found to be $< 10\%$. The inter-
 311 assay % CV was also $< 10\%$ for the SPR and UPLC-MS/MS
 312 methods, but the variation for the ELISA was up to 24% for
 313 some samples. CML concentrations detected by ELISA [(6 \pm
 314 0.80)–(900 \pm 99.0) μg /g of BSA] were significantly lower
 315 ($P < 0.01$) compared to SPR biosensor [(45 \pm 4.76)–(1603 \pm
 316 156) μg /g of BSA] and UPLC-MS/MS [(194 \pm 19.7)–(1075 \pm
 317 68.6) μg /g of BSA] methods (**Figure 4**). CML levels detected in
 318 glycated BSA using the SPR biosensor were 1.6-fold higher than
 319 those obtained by ELISA, but there was good linearity between
 320 the methods ($R^2 = 0.977$) (**Figure 5a**). The concentration of CML
 321 obtained by UPLC-MS/MS was 0.79-fold lower than by SPR
 322 analysis (**Figure 5b**), but 1.35-fold higher than by ELISA assay

(**Figure 5c**). The coefficients of determination of a linear correlation
 CML values obtained between UPLC-MS/MS versus SPR and
 UPLC-MS/MS versus ELISA were fairly good ($R^2 = 0.85$ and
 0.75, respectively).

DISCUSSION

327
 328 An earlier study compared a SPRI biosensor method and an
 329 ELISA for the analysis of AGEs in serum from Zucker diabetic
 330 fatty (ZDF) rats and Zucker lean (ZL) rats (16). The authors
 331 reported that the two methods gave similar results, although the
 332 data from the two assays were expressed in different units. It is not
 333 possible to compare those data sets with CML concentrations
 334 determined by the three methods used in the current study. The
 335 differences in values obtained by the two methods may be due to
 336 the different equipment used. In the present study a SPR
 337 biosensor assay was employed, whereas Kim et al. (16) used
 338 a surface plasmon resonance imaging (SPRI) biosensor. A pub-
 339 lisher study (23) reported the detection levels of progesterone in
 340 bovine milk to be higher when analyzed by a biosensor compared
 341 to an ELISA, but, in agreement with our study, a reasonable
 342 correlation ($R^2 = 0.75$) was observed between both methods of
 343 analysis. In contrast, Yman et al. (19), measuring tropomyosin in
 344 crabmeat (surimi), illustrated that biosensor analysis reported
 345 lower levels than an ELISA. The discrepancies observed between




Figure 4. Comparison of CML concentration detected by three methods of analysis (SPR, ELISA, and UPLC-MS/MS): (a) amount of AGE inhibitor used was equimolar with respect to the lysine content of BSA; (b) amount of AGE inhibitor used was 10-fold greater than the lysine content of BSA. No inhibitor, AGE-BSA without addition of any inhibitor; α -tocopherol, AGE-BSA with addition of α -tocopherol; ferulic acid, AGE-BSA with addition of ferulic acid; rutin, AGE-BSA with addition of rutin; thiamin hydrochloride, AGE-BSA with addition of thiamin hydrochloride; thiamin monophosphate, AGE-BSA with addition of thiamin monophosphate; thiamin pyrophosphate, AGE-BSA with addition of thiamin pyrophosphate. Data are expressed as the mean \pm SD ($n = 3$).

346 the methods applied in these different papers may be due to
347 different instrumentations in the case of biosensor-based assay or
348 interfering factors that are present in the samples (24).

349 Compared to UPLC-MS/MS, immunochemical (ELISA and
350 SPR) analysis has advantages such as simpler sample prepara-
351 tion, speed, and cost. However, the advantages of UPLC-MS/MS
352 include the smaller volumes of sample (7.5 μ L, 7.5 μ g of protein)
353 required compared to ELISA and SPR methods. The correlation
354 of the CML levels found in glycated samples, with or without
355 inhibitors, analyzed by immunochemical methods and UPLC-
356 MS/MS were also well correlated. Charissou et al. (13) reported
357 a good correlation between ELISA and GC-MS data for CML
358 levels in model milk (slope = 1.18) and powdered formulas, but
359 satisfactory linear or nonlinear fitting in liquid formula was not
360 observed. These authors also reported that data for CML in
361 liquid milk were almost 10-fold higher when analyzed by ELISA
362 compared to GC-MS (13). This is in contrast to the present study
363 in which measured concentrations of CML in glycated BSA
364 obtained by the ELISA were lower compared to those obtained
365 by UPLC-MS/MS.

366 In the current study, the concentration of CML in glycated
367 BSA was dependent on the concentration and the nature of the
368 inhibitor used. α -Tocopherol did not affect CML formation,
369 whatever the concentration of α -tocopherol applied. In contrast,
370 ferulic acid, rutin, thiamin hydrochloride, and thiamin derivatives

(thiamin monophosphate and thiamin pyrophosphate) inhibited
371 CML formation, and the inhibitory effect increased with con-
372 centration. In good agreement with the current study, Yin
373 and Chan (25) published that α -tocopherol did not inhibit
374 CML and pentosidine formation in the glycated BSA model
375 system. This is possibly due to α -tocopherol insolubility in the
376 aqueous media (phosphate buffer, 0.2 M, pH 7.2) used in both
377 investigations.

378 Ferulic acid is a free radical scavenger (26, 27), and in the
379 current study its inhibitory effect on CML formation was con-
380 centration dependent. The strong free radical scavenging
381 (hydroxyl or superoxide radical) activity of ferulic acid is
382 due to its phenolic nucleus and extended side-chain conjuga-
383 tion, which allow it to form a resonance-stabilized phenoxy
384 radical (8, 26, 27). The anti-CML effect of ferulic acid is
385 attributed to the second phase of the glycation reaction, that
386 is, glyoxal production from sugar or Amadori product
387 oxidation. The data presented here agree with an earlier
388 study (8) suggesting that ferulic acid (0.25 mg/mL) inhibits
389 AGE formation as a result of its free radical scavenging
390 capacities. However, much lower concentrations of ferulic
391 acid appear not to inhibit CML formation (7).

392 Rutin, a powerful antioxidant and antiglycation agent, inhib-
393 ited CML formation in the current study due to its free radical
394 scavenging capacity (21), which mainly inhibits glyoxal (10)
395




Figure 5. Correlation between CML concentrations obtained by SPR, ELISA, and UPLC-MS/MS analysis: (a) correlation of CML levels obtained by SPR and ELISA analysis; (b) correlation of CML levels obtained by UPLC-MS/MS and SPR analysis; (c) correlation of CML levels obtained by UPLC-MS/MS and ELISA analysis.

396 formation. The data agree with earlier studies (6, 10, 21) suggest-
 397 ing that rutin inhibits all stages of protein glycation formation,
 398 that is, autoxidation of glucose, glyoxal formation, retroaldo
 399 condensation of Schiff base, and oxidative degradation of Ama-
 400 dori products to CML. Furthermore, dietary rutin has been
 401 proven to reduce glycation in tissue protein of streptozotocin-
 402 induced diabetic rats (28). Data from previous studies (6, 10, 21, 28)
 403 and our current investigation would suggest that rutin is a power-
 404 ful antioxidant which inhibits CML formation both in vitro
 405 and in vivo.

406 Thiamin and its derivatives, thiamin monophosphate and
 407 thiamin pyrophosphate, are not antioxidants, but do inhibit
 408 CML formation, dependent on their concentration investigated,
 409 in the current study. The inhibitory effect may be through
 410 a competitive mechanism between the amino group of thiamin/
 411 thiamin derivatives with the amino group of lysine residues
 412 within protein during glycation, as well as α -oxoaldehyde forma-
 413 tion (29). The data agree with CML plasma levels of diabetic rats,
 414 which were reduced by thiamin administered orally (9). Besides
 415 this evidence, Booth et al. (11) also report a similar effect of

thiamin pyrophosphate on AGE formation (98%), even at low 416
 concentrations. However, the authors (11) did not observe any 417
 antiglycation capacity for thiamin and thiamin monophosphate. 418
 In this investigation we found thiamin pyrophosphate to exhibit 419
 a greater inhibitory effect on CML formation than thiamin and 420
 thiamin monophosphate. Each has a similar chemical structure, 421
 and all contain a functional amino group. The proposed mechanism 422
 may be due to the diphosphate group on thiamin pyrophosphate 423
 interfering in the reaction rate between the amino group of 424
 thiamin pyrophosphate and the carbonyl group of a reducing 425
 sugar during the glycation process, thus inhibiting the formation 426
 of CML through a competitive reaction with the amino group on 427
 the protein. 428

In conclusion, the concentration of CML formed was dependen- 429
 on the types and concentrations of AGE inhibitor, and the 430
 reported level of CML was found to be dependent on the methods 431
 of analysis (SPR biosensor, ELISA, and UPLC-MS/MS). How- 432
 ever, good correlations were observed between those three 433
 analysis methods of analysis. The concentrations of CML 434
 detected by SPR and UPLC-MS/MS were closer compared to 435

436 an ELISA analysis. The SPR biosensor has a number of advan-
437 tages over ELISA such as the need for smaller volumes of
438 reagents, no need for a labeled compound, higher repeatability,
439 high automation, and higher precision between runs. However,
440 the biosensor assay required a higher concentration of the anti-
441 body. Immunochemical analysis of CML, in the AGE-BSA
442 model system, has advantages over the UPLC-MS/MS method
443 such as reduced sample preparation, reduced analysis time,
444 increased speed, and lower costs. α -Tocopherol had no measur-
445 able effect on CML formation in AGE-BSA model systems. In
446 contrast, ferulic acid, rutin, thiamin, and thiamin metabolites,
447 thiamin monophosphate and thiamin pyrophosphate, showed
448 various degrees of antiglycation capacity on CML formation.
449 These compounds may be used for health therapy.

450 ABBREVIATIONS USED

451 CML, N^{ϵ} -(carboxymethyl)lysine; CEL, N^{ϵ} -(carboxyethyl)lysine;
452 AGEs, advanced glycation endproducts; BSA, bovine serum
453 albumin; KLH, keyhole limpet hemocyanin; CML-BSA,
454 N^{ϵ} -(carboxymethyl)lysine-modified bovine serum albumin;
455 BSA, N^{ϵ} -(carboxyethyl)lysine-modified bovine serum albumin;
456 CML-KLH, N^{ϵ} -(carboxymethyl)lysine-modified keyhole limpet
457 hemocyanin; ELISA, enzyme-linked immunosorbent assay; SPR,
458 surface plasmon resonance; UPLC-MS/MS, ultraperformance
459 liquid chromatography-tandem mass spectrometry.

460 ACKNOWLEDGMENT

461 We thank Julie Meneely, Dr. Linda Stewart, and Dr. Katrina
462 Campbell for providing technical support and guidance.

463 LITERATURE CITED

- 464 (1) Lima, M.; Assar, S. H.; Ames, J. M. Formation of N^{ϵ} -(carb-
465 oxymethyl)lysine and loss of lysine in casein glucose-fatty acid model
466 systems. *J. Agric. Food Chem.* **2010**, *58*, 1954-1958.
- 467 (2) Ames, J. M. Determination of N^{ϵ} -(carboxymethyl)lysine in foods
468 and related systems. *Ann. N.Y. Acad. Sci.* **2008**, *1126*, 20-24.
- 469 (3) Semba, R. D.; Fink, J. C.; Sun, K.; Windham, B. G.; Ferrucci, L.
470 Serum carboxymethyl-lysine, a dominant advanced glycation end
471 product, is associated with chronic kidney disease: the Baltimore
472 longitudinal study of aging. *J. Renal Nutr.* **2010**, *20*, 74-81.
- 473 (4) Matsumoto, T.; Ozawa, Y.; Taguchi, K.; Kobayashi, T.; Kamata, K.
474 Diabetes-associated changes and role of N^{ϵ} -(carboxymethyl)lysine in
475 big ET-1-induced coronary vasoconstriction. *Peptides* **2010**, *31*,
476 346-353.
- 477 (5) Schalkwijk, C. G.; Baidoshvili, A.; Stehouwer, C. D.; van Hinsbergh,
478 V. W.; Niessen, H. W. Increased accumulation of the glycoxidation
479 product N^{ϵ} -(carboxymethyl)lysine in hearts of diabetic patients:
480 generation and characterization of a monoclonal anti-CML
481 antibody. *Biochim. Biophys. Acta* **2004**, *1636*, 82-89.
- 482 (6) Cervantes-Laurean, D.; Schramm, D. D.; Jacobson, E. L.;
483 Halaweish, I.; Bruckner, G. G.; Boissonneault, G. A. Inhibition of
484 advanced glycation end product formation on collagen by rutin and
485 its metabolites. *J. Nutr. Biochem.* **2006**, *17*, 531-540.
- 486 (7) Wu, J. W.; Hsieh, C. L.; Wang, H. Y.; Chen, H. Y. Inhibitory effects
487 of guava (*Psidium guajava* L.) leaf extracts and its active compounds
488 on the glycation process of protein. *Food Chem.* **2009**, *113*, 78-84.
- 489 (8) Wang, J.; Sun, B.; Cao, Y.; Tian, Y. Protein glycation inhibitory
490 activity of wheat bran feruloyl oligosaccharides. *Food Chem.* **2009**,
491 *112*, 350-353.
- 492 (9) Karachalias, N.; Babaei-Jadidi, R.; Kupich, C.; Ahmed, N.;
493 Thornalley, P. J. High-dose thiamine therapy counters dyslipidemia
494 and advanced glycation of plasma protein in streptozotocin-induced
495 diabetic rats. *Ann. N.Y. Acad. Sci.* **2005**, *1043*, 777-783.
- 496 (10) Pashikanti, S.; de-Alba, D. R.; Boissonneault, G. A.; Cervantes-
497 Laurean, D. Rutin metabolites: Novel inhibitors of nonoxidative
498 advanced glycation end products. *Free Radical Biol. Med.* **2010**, *48*,
499 656-663.

- (11) Booth, A. A.; Khalifah, R. G.; Hudson, B. G. Thiamine pyrophos-
500 phate and pyridoxamine inhibit the formation of antigenic advanced
501 glycation end-products: comparison with amino guanidine. *Biochem.*
502 *Biophys. Res. Commun.* **1996**, *220*, 113-119. 503
- (12) Hegele, J.; Buetler, T.; Delatour, T. Comparative LC-MS/MS
504 profiling of free and protein-bound early and advanced glycation-
505 induced lysine modifications in dairy products. *Anal. Chim. Acta*
506 **2008**, *617*, 85-96. 507
- (13) Charissou, A.; Ait-Ameur, L.; Birlouez-Aragon, I. Evaluation of
508 a gas chromatography/mass spectrometry method for the quantifi-
509 cation of carboxymethyllysine in food samples. *J. Chromatogr., A*
510 **2007**, *1140*, 189-194. 511
- (14) Kioto, W.; Araki, T.; Horiuchi, S.; Nagai, R. Conventional antibody
512 against N^{ϵ} -(carboxymethyl)lysine (CML) shows cross-reaction to
513 N^{ϵ} -(carboxyethyl)lysine (CEL): immunochemical quantification of
514 CML with a specific antibody. *J. Biochem.* **2004**, *136*, 331-337. 515
- (15) Assar, S. H.; Moloney, C.; Lima, M.; Magee, R.; Ames, J. M. M.
516 Determination of N^{ϵ} -(carboxymethyl)lysine in food systems by
517 ultraperformance liquid chromatography-mass spectrometry.
518 *Amino Acids* **2009**, *36*, 317-326. 519
- (16) Kim, Y. S.; Yi, S. Y.; Kim, J.; Kim, M.; Kim, C. S.; Chung, B. H.; Kim,
520 J. S. Novel application of surface plasmon resonance biosensor chips for
521 measurement of advanced glycation end products in serum of Zucker
522 diabetic fatty rats. *Biosens. Bioelectron.* **2009**, *25*, 248-252. 523
- (17) Stewart, L. D.; Elliott, C. T.; Walker, A. D.; Curran, R. M.;
524 Connolly, L. Development of a monoclonal antibody binding
525 okadaic acid and dinophysistoxins-1, -2 in proportion to their
526 toxicityequivalence factors. *Toxicol.* **2009**, *54*, 491-498. 527
- (18) Campbell, K.; Huet, A. C.; Charlier, C.; Higgins, C.; Delahaut, P.;
528 Elliott, C. T. Comparison of ELISA and SPR biosensor technology
529 for the detection of paralytic shellfish poisoning toxins. *J. Chroma-
530 togr., B* **2009**, *877*, 4079-4089. 531
- (19) Yman, I. M.; Eriksson, A.; Johansson, M. A.; Hellenäs, K. E. Food
532 allergen detection with biosensor immunoassays. *J. AOAC Int.* **2006**,
533 *89*, 856-861. 534
- (20) Dupont, D.; Muller-Renaud, S. Quantification of proteins in dairy
535 products using an optical biosensor. *J. AOAC Int.* **2006**, *89*, 843-848. 536
- (21) Wu, C. H.; Yen, G. C. Inhibitory effect of naturally occurring
537 flavonoids on the formation of advanced glycation endproducts
538 *J. Agric. Food Chem.* **2005**, *53*, 3167-3173. 539
- (22) Köhler, G.; Milstein, C. Continuous cultures of fused cells secreting
540 antibody of predefined specificity. *Nature* **1975**, *256*, 495-497. 541
- (23) Gillis, E. H.; Gosling, J. P.; Sreenan, J. M.; Kane, M. Development
542 and validation of a biosensor-based immunoassay for progesterone
543 in bovine milk. *J. Immunol. Methods* **2002**, *267*, 131-138. 544
- (24) Dorrian, C. A.; Cathcart, S.; Clausen, J.; Shapiro, D.; Dominiczak,
545 M. H. Factors in human serum interfere with the measurement of
546 advanced glycation endproducts. *Cell Mol Biol.* **1998**, *44*, 1069-1079. 547
- (25) Yin, M. C.; Chen, K. C. Nonenzymatic antioxidative and antigly-
548 cative effects of oleanolic acid and ursolic acid. *J. Agric. Food Chem.*
549 **2007**, *55*, 7177-7181. 550
- (26) Srinivasan, M.; Sudheer, A. R.; Menon, V. P. Ferulic acid: ther-
551 apeutic potential its antioxidant property. *J. Clin. Biochem. Nutr.*
552 **2007**, *40*, 92-100. 553
- (27) Kanski, J.; Aksenova, M.; Stoyanova, A.; Butterfield, D. A. Ferulic
554 acid antioxidant protection against hydroxyl and peroxy radical
555 oxidation in synaptosomal and neuronal cell culture systems in vitro:
556 structure-activity studies. *J. Nutr. Biochem.* **2002**, *13*, 273-281. 557
- (28) Nagasawa, T.; Tabata, N.; Ito, Y.; Aiba, Y.; Nishizawa, N.; Kitts,
558 D. D. Dietary G-rutin suppresses glycation in tissue proteins of
559 streptozotocin-induced diabetic rats. *Mol. Cell. Biochem.* **2003**, *252*,
560 141-147. 561
- (29) Babaei-Jadidi, R.; Karachalias, N.; Ahmed, N.; Battah, S.;
562 Thornalley, P. J. Prevention of incipient diabetic nephropathy by
563 high-dose thiamine and benfotiamine. *Diabetes* **2003**, *52*, 2110-2120. 564

Received for review July 1, 2010. Revised manuscript received October 565
7, 2010. Accepted October 11, 2010. We acknowledge The Department 566
of Agriculture and Rural Development for Northern Ireland for a 567
studentship (to C.S.). 568