



Animal (2012), 6:7, pp 1040–1048 © The Animal Consortium 2012
doi:10.1017/S1751731111002655

Merging and characterising phenotypic data on conventional and rare traits from dairy cattle experimental resources in three countries

G. Banos^{1,2†}, M. P. Coffey², R. F. Veerkamp³, D. P. Berry⁴ and E. Wall²

¹Department of Animal Production, Faculty of Veterinary Medicine, Aristotle University of Thessaloniki, GR-54124 Thessaloniki, Greece; ²Sustainable Livestock Systems Group, Scottish Agricultural College, Roslin Institute Building, Easter Bush, Midlothian EH25 9RG, Scotland, UK; ³Animal Breeding and Genomics Centre, Wageningen UR Livestock Research, 8200 AB Lelystad, The Netherlands; ⁴Animal and Bioscience Research Department, Animal & Grassland Research and Innovation Centre, Teagasc, Moorepark, Co. Cork, Republic of Ireland

(Received 3 June 2011; Accepted 12 October 2011; First published online 4 January 2012)

This study set out to demonstrate the feasibility of merging data from different experimental resource dairy populations for joint genetic analyses. Data from four experimental herds located in three different countries (Scotland, Ireland and the Netherlands) were used for this purpose. Animals were first lactation Holstein cows that participated in ongoing or previously completed selection and feeding experiments. Data included a total of 60 058 weekly records from 1630 cows across the four herds; number of cows per herd ranged from 90 to 563. Weekly records were extracted from the individual herd databases and included seven traits: milk, fat and protein yield, milk somatic cell count, liveweight, dry matter intake and energy intake. Missing records were predicted with the use of random regression models, so that at the end there were 44 weekly records, corresponding to the typical 305-day lactation, for each cow. A total of 23 different lactation traits were derived from these records: total milk, fat and protein yield, average fat and protein percentage, average fat-to-protein ratio, total dry matter and energy intake and average dry matter intake-to-milk yield ratio in lactation weeks 1 to 44 and 1 to 15; average milk somatic cell count in lactation weeks 1 to 15 and 16 to 44; average liveweight in lactation weeks 1 to 44; and average energy balance in lactation weeks 1 to 44 and 1 to 15. Data were subsequently merged across the four herds into a single dataset, which was analysed with mixed linear models. Genetic variance and heritability estimates were greater ($P < 0.05$) than zero for all traits except for average milk somatic cell count in weeks 16 to 44. Proportion of total phenotypic variance due to genotype-by-environment (sire-by-herd) interaction was not different ($P > 0.05$) from zero. When estimable, the genetic correlation between herds ranged from 0.85 to 0.99. Results suggested that merging experimental herd data into a single dataset is both feasible and sensible, despite potential differences in management and recording of the animals in the four herds. Merging experimental data will increase power of detection in a genetic analysis and augment the potential reference population in genome-wide association studies, especially of difficult-to-record traits.

Keywords: merging phenotypes, dairy genetics, experimental resources, dry matter intake

Implications

The feasibility of merging data from different experimental herds paves the way to pooling valuable records of rare and difficult-to-record traits, such as those associated with feed intake and efficiency, and body energy balance, from resource populations located in various sites and different countries. Use of the enhanced database would increase the experimental power of effect detection, facilitate genomic selection and enable other genetic analyses that would otherwise be difficult, if not impossible, to conduct separately in each country/herd.

Introduction

Undertaking genetic studies of animal phenotypes presupposes accurate recording on sufficient numbers of animals to help dissect the genetics of the trait. The development of elaborate national monitoring schemes has facilitated routine population-wide on-farm accurate recording of several conventional traits, mostly associated with production (International Committee for Animal Recording, 2011). Certain indicators of functional traits, such as milk somatic cell count in dairy cattle, are included in these programmes. Nevertheless, several increasingly important traits associated with health, fitness and efficiency are currently not possible to routinely record in the commercial

[†] E-mail: banos@vet.auth.gr

population. Understanding the genetic background of such traits depends then on data from experimental resource populations where animals are raised in controlled, closely monitored environments. Such populations, however, are usually of limited size. Combining data from different experimental herds would provide an expanded dataset that would allow a more rigorous genetic analysis of difficult- and expensive-to-record traits.

Furthermore, the advent of high-density single nucleotide polymorphism (SNP) arrays has encouraged the application of this technology to the analysis of economically important traits, especially rare and expensive phenotypes such as feed intake, health and fertility. The amount of data required to find significant associations between SNP markers and phenotypes has led to a number of institutions and countries exploring the possibility of sharing phenotypes and genotypes to enable genomic analyses to be undertaken. This has raised a number of analytical issues regarding the use of ostensibly similar data collected at different sites and subjected to different experimental treatments.

The objectives of this study were to (i) demonstrate the feasibility of merging phenotypic data from different experimental resources and (ii) characterise the merged database and assess its suitability for a joint genetic analysis as a single dataset.

Material and methods

Animals and experimental herds

Data used in this study were collected from first lactation Holstein cows raised in four distinct experimental resource herds in three different countries (one herd in each of Scotland and Ireland, and two in the Netherlands). In all cases, cows were used in various ongoing or previously completed experiments conducted at different time periods, as described below.

Scotland – Crichton herd

Data originated from the Scottish Agricultural College Dairy Research Centre based at Crichton Royal Farm, Scotland. These cows had previously comprised the Langhill herd, Edinburgh (Veerkamp *et al.*, 1995; Pryce *et al.*, 1999) and were transferred to Crichton Royal Farm in September 2001. The herd normally consists of ~200 milking cows divided evenly between two genetic groups (control *v.* selection) established in 1992 as part of a still ongoing selection experiment. Cows in each genetic group are further split randomly into two diet groups (high-concentrates *v.* high-forage) for the purposes of a feeding experiment, which is also in progress. The two genetic groups on a particular diet are managed together.

Control and selection group cows are allocated to the same group as their dams and remain there throughout their productive life. Cow sires in the selection group are picked on the basis of their genetic merit for milk fat and protein yield; available sires with the highest genetic evaluation for fat plus protein kg are chosen at the time of artificial insemination. Sires of control group cows are selected to

have the average genetic merit for fat plus protein per kg of UK animals at the time of breeding. In both groups, matings are arranged such that the inbreeding coefficient remains less than 6%.

Furthermore, animals are randomly allocated to either the high-concentrate or the high-forage (low-concentrate) group at first calving in such a way as to keep the groups balanced for number of cows and sires. The high-forage system consists entirely of home-grown feeds, including maize and other whole-crop cereals, and the cows are grazed on grass during the summer months. The winter ration consists of grass silage, maize silage and alkalage at a ratio of 60:20:20 on a dry matter basis, plus a protein supplement. The ration is fed as a total mixed ration. At least 75% of the dry matter of the ration is designed to come from forages. The target metabolisable energy content is 11.5 MJ/kg dry matter with a target CP content of 180 g/kg dry matter. Forages are supplemented with a range of energy and protein sources to meet the targets shown above. The high-concentrate system cows are housed all year, with access to an exercise area during the summer months. Their ration also contains the three forages mentioned above, in the same dry matter ratio to each other, with a supplement blend of energy and protein ingredients. The target ration metabolisable energy content is 12.3 MJ/kg dry matter with a target CP content of 185 g/kg dry matter.

All cows are kept together and treated the same at all times, except where the production systems require management differences. Cows are milked three times per day.

For the purposes of the present study, data pertained to 563 cows equally distributed across the four experimental groups that had calved between 1992 and 2009. Individual records available for each cow were: daily milk yield (sum of three milkings) and liveweight, weekly milk fat and protein yield and milk somatic cell count, and three times weekly dry matter intake.

Ireland – Moorepark herd

This experimental resource is located at the Teagasc Moorepark Research Farm, Republic of Ireland. Data were collated from several studies that had been previously conducted (Buckley *et al.*, 2000; Kennedy *et al.*, 2003; O'Donovan and Delaby, 2005; Horan *et al.*, 2006; Kennedy *et al.*, 2006; McCarthy *et al.*, 2007; McEvoy *et al.*, 2007). In brief, these studies compared either alternative genotypes of Holstein–Friesian cows raised on different production systems or alternative grazing strategies or grass varieties. Different strains of Holstein–Friesians were evaluated on contrasting grass-based production systems (Buckley *et al.*, 2000; Kennedy *et al.*, 2003; Horan *et al.*, 2006; McCarthy *et al.*, 2007). Animals within strain were randomly assigned, at the start of lactation, to feed systems differing in stocking rate and/or concentrate input. Perennial ryegrass (*Lolium perenne*) was the predominant pasture species in these studies and a rotational grazing system was operated. Annual concentrate feeding level across studies varied from 325 to 1452 kg per cow. All cows calved in the spring and were milked twice daily.

Table 1 Data description

Herd (country)	No. of records	No. of cows	No. of sires of cows	Calving years
Crichton (Scotland)	22 426	563	93	1992 to 2009
NBZ (the Netherlands)	1346	90	49	2003 to 2004
Moorepark (Ireland)	18 612	449	80	1998 to 2008
TGEN (the Netherlands)	18 312	549	94	1990 to 1998
Overall	60 696	1651	278	1990 to 2009
Overall after edits	60 058	1630	276	1990 to 2009

For the purposes of the present study, data pertained to 449 cows that first calved between the years 1998 and 2008. Individual records available for each cow were: daily milk yield (sum of two milkings) and weekly milk fat and protein yield, milk somatic cell count, liveweight and dry matter intake.

The Netherlands – TGEN and NBZ herds

Data originated from two experimental dairy herds, one located near Lelystad (TGEN) and another one near Leeuwarden (NBZ) in the Netherlands.

Data from the TGEN herd were collected between 1990 and 1998 and, for the purposes of the present study, pertained to 549 cows (Veerkamp *et al.*, 2000). Two-thirds of these cows belonged to a high genetic merit group that participated in the Delta sib-testing program of Holland Genetics and the others were part of a control group. Cows in the latter group were, on average, about half a standard deviation below the former on the Dutch production index reflecting the impact of milk, fat and protein yield on future net profit. All cows were fed *ad libitum* with a complete ration of artificially dried grass, corn silage and concentrates with proportions in the dry matter of 6:5:10. On average, the total mixed ration (64% dry matter) contained 6.87 MJ net energy and 98 g digestible CP/kg of dry matter. Individual records available for each cow were: daily milk yield (sum of two milkings), weekly milk fat and protein yield, milk somatic cell count and liveweight, and five times/week dry matter intake.

Data collected from the NBZ herd were from the period between 2003 and 2004 and pertained to 90 first lactation cows that were participating in a genetic and feeding experiment. Specifically, these cows were divided into two genetic groups with high and low genetic merit for fat and protein production, respectively. Furthermore, cows were split into two diet groups fed a high and a low caloric density ration, respectively. The former comprised 49% corn silage, 30% grass silage and 21% soybeans meal, whereas the low caloric ration included 86% grass silage and 14% concentrates (Beerda *et al.*, 2007; Windig *et al.*, 2008). Individual records available for each cow were: daily milk yield (sum of three milkings) and dry matter intake, and weekly milk fat and protein yield, milk somatic cell count and liveweight.

Table 1 summarises the different datasets from the four experimental herds described above.

Fixed effects

As previously explained, cows belonged to various past and present selection and feeding experiments run in the four herds. Therefore, cows belonged to different genetic and/or diet groups. Specifically, two genetic groups were identified in each one of the four herds. In all cases, cows remained in their respective genetic groups throughout their life. Furthermore, two diet groups were formed in each of Crichton (Scotland) and NBZ (the Netherlands) herds. In both herds, a cow would remain in the same diet groups throughout the entire lactation. In addition, 18 different feeding treatments were identified in the Moorepark (Ireland) dataset. Contrary to Crichton and NBZ, in the Irish herd, a cow might change feeding treatment group during her lactation.

Traits recorded in the four herds

Weekly individual cow records for milk, fat and protein yield, milk somatic cell count, liveweight, dry matter intake and energy intake were extracted from the database of each herd. Records pertained to the daily observation on the day of recording. When multiple records were available within the same week of lactation, weekly values were corresponding arithmetic means. Only first lactation cows were considered in the present study.

Energy intake had already been calculated in each herd, separately, as net energy for the two Dutch herds (TGEN and NBZ) (Beerda *et al.*, 2007) and metabolisable energy for Crichton (Scotland) and Moorepark (Ireland; Emmans, 1994; Friggens *et al.*, 2003). The latter (ME) was converted to net energy (NE) using the following formula (Van Es, 1978):

$$NE = 0.6[1 + (0.004(q-57))]0.9752ME$$

where a 70% value was assumed for q (ME/gross energy); q values between 50% and 75% were also tried but made practically no difference in the results. Energy intake was then expressed as net energy for all cows for the remainder of this study.

Time edits retained records in the first 44 weeks, corresponding to the typical 305-day lactation. In the case of NBZ, only data for the first 15 weeks of lactation were available. Further edits removed records outside certain biological value ranges set by trait. These limits are shown in Table 2.

In addition, a minimum of 5 weekly records were required per cow in order to keep records of her lactation. This edit was modified to 2 weekly records minimum for dry matter

Table 2 Biological limits set for the recorded traits

Trait	Unit of measurement	Acceptable values
Milk yield	kg/d	3 to 90
Fat yield	kg/d	0.1 to 11
Protein yield	kg/d	0.1 to 9
Milk somatic cell count	1000/ml	10 to 10 000
Liveweight	kg/d	300 to 900
Dry matter intake	kg/d	1 to 60
Energy intake	MJ/d	0.5 to 340

and energy intake at Moorepark (Ireland) due to limited recording of these traits in this herd.

After edits, a total of 60 058 weekly records of 1630 cows remained across the four herds. Missing weekly records per cow were predicted with random regression models as described next.

Prediction of missing weekly records

Theoretically, if each one of the 1630 cows in the final dataset had 44 weekly records, a total of 71 720 records would be available. To predict missing weekly records, the following random regression model was used:

$$Y = \text{HTYM} + \text{CYM} + \text{CA} + \text{GG} + \text{DG} + \text{IRL} + \text{MF} + \text{WK} + \text{COW.WK} \quad (1)$$

where Y is the weekly cow record for a trait; HTYM the fixed effect of herd by year–month of record interaction (four herds, 222 year–month classes); CYM the fixed effect of calving year–month interaction (188 classes); CA the fixed effect of calving age (three classes: <704, 704 to 827, >827 days); GG the fixed effect of genetic group (eight classes); DG the fixed effect of diet group (four classes); IRL the fixed effect of Irish diet treatment (18 classes); MF the fixed effect of milking frequency (two or three times); WK the fixed lactation curve modelled with a 4th order polynomial (5th for somatic cell count); and COW.WK the random cow deviation from fixed curve modelled with a 4th order polynomial (5th for somatic cell count)

Each recorded trait was analysed separately. In the case of milk somatic cell count, a log transformation took place before the analysis to ensure normality.

The order of the polynomial was determined by examining the residual variance. The latter did not change significantly ($P > 0.5$) when the order increased from 4 to 5 (5 to 6 for milk somatic cell count).

Effect solutions obtained from model 1 were combined to re-create the phenotypic record for all cow-weeks, including those with missing observations.

In the first instance, a single fixed curve was calculated across the four herds. Subsequently, a separate series of analyses took place where four different curves were fitted (one for each herd). This was achieved by fitting a herd-by-week interaction in both the fixed curve and random animal deviation in model 1.

In all cases, heterogeneous residual variances in different herds were accounted for in model 1 by fitting a different

residual effect per herd. However, phenotypes were all expressed on the same scale, that is, they were not scaled back according to the original variance of each herd. All analyses were conducted using the ASREML 2.0 software (Gilmour *et al.*, 2006).

Derivation of phenotypic traits

At first, energy balance was calculated for each cow and week of lactation. For this purpose, predicted weekly phenotypic records were used to calculate the energy expended by the cow for yield and maintenance, based on the method described by Beerda *et al.* (2007). This method is consistent with the net energy principle used to assess energy intake in the four herds in the present study, as described previously. Energy-corrected milk (ECM) was first estimated on the basis of milk yield (M) and fat (F) and protein (P) percent, derived from milk, fat and protein yields, as follows:

$$\text{ECM} = M[0.337 + (0.116F) + (0.06P)]$$

Energy expended (EE) for yield and maintenance on a certain week was then calculated as follows:

$$\text{EE} = \{[(42.4\text{LWT}^{0.75}) + (44\text{ECM})] [1 + ((\text{ECM} - 15) 0.00165)] + 660\} 0.0069$$

where LWT was the animal's liveweight. Finally, weekly energy balance was calculated as the difference between energy intake and energy expended.

Subsequently, 23 phenotypic lactation traits were derived for all cows, using the predicted weekly records for milk, fat and protein yield, milk somatic cell count, liveweight, dry matter intake, energy intake and energy balance. Total lactation yield (milk, fat and protein) was calculated by multiplying weekly predictions by 7 and then summing them up. Ratio and percent traits were calculated by simple division. Average traits (e.g. milk somatic cell count) were derived as the arithmetic mean of weekly records for the defined period. All lactation traits are summarised in Table 3.

For average milk somatic cell count, two distinct time periods were considered, weeks of lactation 1 to 15 and weeks of lactation 16 to 44, in order to define the two potentially different traits in early and later lactation (Mrode and Swanson, 2003). For all other traits, two lactation periods were considered: a 44-week lactation (consistent with the typical 305-day lactation) and a 15-week lactation, marking the crucial first 100 days of lactation. The latter also corresponds to the period of the animal's recovery from a negative energy state (Coffey *et al.*, 2001; Banos *et al.*, 2005). Furthermore, a 15-week lactation is more appropriate for NBZ cows where no later weekly records were available.

Estimation of genetic parameters

Lactation traits derived in the previous section were analysed with mixed linear models, including the effects of herd, calving year–month and age, genetic and diet group, milking frequency and cow. A pedigree file comprising 8850 animals

Table 3 Traits derived from predicted weekly records

Trait definition	Mean	s.d.
Total milk yield in 44 weeks (kg)	6996.00	1941.29
Total fat yield in 44 weeks (kg)	278.58	76.74
Total protein yield in 44 weeks (kg)	236.74	70.26
Average fat percentage in 44 weeks (%)	4.04	0.57
Average protein percentage in 44 weeks (%)	3.39	0.34
Average fat to protein ratio in 44 weeks	1.19	0.13
Total milk yield in 15 weeks (kg)	2743.07	624.1
Total fat yield in 15 weeks (kg)	108.14	25.7
Total protein yield in 15 weeks (kg)	88.86	21.9
Average fat percentage in 15 weeks (%)	3.97	0.54
Average protein percentage in 15 weeks (%)	3.25	0.35
Average fat to protein ratio in 15 weeks	1.23	0.15
Average somatic cell count in weeks 1 to 15 (1000/ml)	118.94	192.88
Average somatic cell count in weeks 16 to 44 (1000/ml)	99.65	149.45
Average liveweight in 44 weeks (kg)	530.38	60.23
Total dry matter intake in 44 weeks (kg)	4650.91	1448.06
Total dry matter intake in 15 weeks (kg)	1541.96	483.2
Average dry matter intake to milk yield ratio in 44 weeks	0.66	0.20
Average dry matter intake to milk yield ratio in 15 weeks	0.55	0.17
Total energy intake in 44 weeks (MJ)	31 858.46	10 528.16
Total energy intake in 15 weeks (MJ)	10 538.73	3515.27
Average energy balance in 44 weeks (MJ)	-9.29	22.04
Average energy balance in 15 weeks (MJ)	-20.64	25.09

across the four herds was used to derive genetic variance and heritability estimates.

In a separate set of analyses, sire-by-herd interaction was added in the model as a random effect to assess the magnitude of genotype-by-environment (herd) interaction. Models with and without a sire-by-herd interaction effect were compared with the Akaike Information Criterion (AIC), a measure of the relative goodness of fit of a statistical model proposed by Akaike (1974).

The issue of genotype-by-environment interaction was also addressed with a series of multi-trait analyses, where individual traits in the four herds were treated as different but genetically correlated traits.

All calculations were based on the REML method and used the ASREML 2.0 software package (Gilmour *et al.*, 2006).

Results and discussion

Prediction of missing weekly records

Figure 1 illustrates the predicted lactation curves of the seven recorded traits in the four herds (milk, fat and protein yield, milk somatic cell count, liveweight, dry matter intake and energy intake by week of lactation). These results correspond to the overall fixed curve in model 1, representing the average lactation trajectory of each trait across all animals and herds, adjusted for all other effects in the model.

The trait profile depicted in Figure 1 was as expected in all cases. Milk, fat and protein yields showed the well-known initial increase during the first 10 weeks of lactation followed by a gradual decline towards the end of lactation.

The initial increase in production was matched by an increase in dry matter and energy intake, which remained constant afterwards, similar to trends reported in other studies based on experimental (Ordway *et al.*, 2009) and commercial (Vallimont *et al.*, 2010) herd data. Milk somatic cell count showed an inverse milk yield curve, suggestive of the relationship between the two traits.

Very similar results were derived when separate curves were fitted for each herd, suggesting that combining records across herds did not change the lactation profile of the trait. Furthermore, the correlation between predicted phenotypes with this model and the model with a single overall curve was near unity for milk traits and liveweight, and 0.981 to 0.987 for dry matter and energy intake. Figure 2 illustrates two examples: one for a conventional trait (milk yield) and one for a rare trait (dry matter intake). It should be noted that results in Figure 2 were expressed on the same scale for all herds, that is, were not re-scaled according to the original variance of each herd. Therefore, differences between herds shown in Figure 2 do not represent true differences between animals raised in these herds. The purpose of Figure 2 was to compare the shape of the individual within-herd curves with that of the across-herd curves of Figure 1 and not to compare herd production and performance levels.

Genetic parameters of derived traits

Estimates of genetic variance, heritability and proportion of total variance attributed to sire-by-herd interaction for all derived traits are shown in Table 4. Genetic variance estimates were greater ($P < 0.05$) than zero in all cases except




Figure 1 Predicted lactation curves for milk yield (kg), fat yield (kg), protein yield (kg), milk somatic cell count (SCC – log-transformed 1000/ml), liveweight (LWT – kg), dry matter intake (DMI – kg) and energy intake (MJ) by week of lactation; a single curve was fitted across herds.




Figure 2 Predicted individual lactation curves for milk yield (kg) and dry matter intake (DMI – kg) for each of the four herds (Crichton – Scotland, NBZ – the Netherlands, Moorepark – Ireland, TGEN – the Netherlands) by week of lactation; results are expressed on a common scale for all herds.

for average milk somatic cell count in weeks 16 to 44 of lactation, suggesting that diverse experimental data from distinct herds may be merged into a single database amenable to a joint genetic analysis.

In general, heritability estimates shown in Table 4 were consistent with estimates provided in the literature from various studies worldwide (Toshniwal *et al.*, 2008; Urioste *et al.*, 2010; Vallimont *et al.*, 2010; Buttchereit *et al.*, 2011). With the exception of average milk somatic cell count in weeks 16 to 44 of lactation, all heritability estimates were statistically greater than zero ($P < 0.05$). Heritability estimates were also derived separately within herd and, in general, were in the same range

as the across-herd estimates presented in Table 4. For example, within-herd heritability estimates of total milk yield in the first 15 weeks of lactation varied from 0.20 to 0.31, whereas estimates for total dry matter intake in the same period ranged from 0.15 to 0.27 in the four different herds. However, there were few cases, such as liveweight, where a wider range of within-herd heritability estimates was obtained (0.38, 0.21, 0.47 and 0.48 for Crichton, NBZ, Moorepark and TGEN, respectively). This may be attributed to different variance magnitude in the four herds associated with different dataset sizes. For example, NBZ had the lowest estimate and smaller dataset.

Including a sire-by-herd interaction effect in the model of analysis led to a small reduction in heritability estimates (Table 4), suggesting that a minor part of the previously estimated additive genetic variance might be attributed to interaction effects. However, the proportion of total phenotypic variance accounted for by the sire-by-herd interaction effect was always nonsignificantly ($P > 0.05$) different from zero. Furthermore, the difference of AIC between the two models was statistically not greater than zero ($P > 0.05$), suggesting that including a sire-by-herd interaction effect did not improve the fit of the model for any of the traits. This result implies that a joint analysis of data from the different herds that is based on the assumption of no genotype-by-environment interaction is possible, although supporting the notion of merged phenotypes from these four herds being viewed as a single dataset in a genetic analysis.

In most cases, genetic correlations between the same trait in different herds were inestimable or associated with very large standard errors that rendered them not significantly different from zero ($P > 0.05$). The reason is probably the relatively weak genetic links between the four herds. For example, in the Crichton (Scotland) dataset there were 7, 6 and 16 sires in common with NBZ (the Netherlands), Moorepark (Ireland) and TGEN (the Netherlands), respectively. NBZ had 5 and 2 common sires with Moorepark and TGEN, respectively, whereas there were 11 bulls in common in the last two herds. Of course, pedigree relationships and the use of the numerator

Table 4 Estimates of genetic variance and h^2 obtained with an additive model, and proportion of total variance due to h^2_a and to $S \times H$ when the latter was included in the model; standard errors are in parentheses

Trait	Genetic variance	h^2	h^2_a	$S \times H$
Total milk yield in 44 weeks	267 800 (85 940)	0.22 (0.07)	0.17 (0.08)	0.05 (0.03)
Total fat yield in 44 weeks	317.8 (110.5)	0.20 (0.07)	0.16 (0.08)	0.03 (0.02)
Total protein yield in 44 weeks	170.1 (73.76)	0.16 (0.07)	0.12 (0.07)	0.04 (0.03)
Average fat percentage in 44 weeks	0.154 (0.021)	0.68 (0.07)	0.66 (0.08)	0.02 (0.03)
Average protein percentage in 44 weeks	0.030 (0.005)	0.55 (0.07)	0.49 (0.08)	0.05 (0.03)
Average fat to protein ratio in 44 weeks	0.008 (0.001)	0.66 (0.07)	0.66 (0.08)	0.00 (0.00)
Total milk yield in 15 weeks	37 210 (12 840)	0.21 (0.07)	0.17 (0.08)	0.03 (0.03)
Total fat yield in 15 weeks	74.30 (21.55)	0.27 (0.07)	0.22 (0.08)	0.03 (0.03)
Total protein yield in 15 weeks	24.13 (9.95)	0.17 (0.07)	0.13 (0.07)	0.03 (0.02)
Average fat percentage in 15 weeks	0.110 (0.017)	0.58 (0.07)	0.58 (0.07)	0.00 (0.00)
Average protein percentage in 15 weeks	0.029 (0.004)	0.57 (0.07)	0.50 (0.08)	0.05 (0.03)
Average fat to protein ratio in 15 weeks	0.005 (0.001)	0.37 (0.08)	0.37 (0.08)	0.00 (0.00)
Average somatic cell count in weeks 1 to 15	0.118 (0.056)	0.14 (0.06)	0.10 (0.07)	0.03 (0.03)
Average somatic cell count in weeks 16 to 44	0.059 (0.045)	0.09 (0.06)	0.06 (0.06)	0.02 (0.03)
Average liveweight weight in 44 weeks	532.2 (121.1)	0.35 (0.07)	0.30 (0.08)	0.04 (0.03)
Total dry matter intake in 44 weeks	26 020 (12 700)	0.15 (0.07)	0.15 (0.09)	0.00 (0.03)
Total dry matter intake in 15 weeks	6329 (2381)	0.22 (0.08)	0.17 (0.09)	0.03 (0.03)
Average dry matter intake to milk yield ratio in 44 weeks	0.004 (0.001)	0.28 (0.08)	0.23 (0.10)	0.04 (0.03)
Average dry matter intake to milk yield ratio in 15 weeks	0.002 (0.001)	0.21 (0.07)	0.21 (0.07)	0.00 (0.00)
Total energy intake in 44 weeks	1 170 000 (630 600)	0.14 (0.07)	0.12 (0.08)	0.01 (0.03)
Total energy intake in 15 weeks	298 500 (114 200)	0.22 (0.08)	0.15 (0.09)	0.04 (0.03)
Average energy balance in 44 weeks	10.96 (5.14)	0.17 (0.08)	0.13 (0.09)	0.03 (0.03)
Average energy balance in 15 weeks	38.15 (12.46)	0.27 (0.08)	0.27 (0.08)	0.00 (0.00)

h^2 = heritability; h^2_a = additive effects; $S \times H$ = sire-by-herd interaction.

relationship (*A*) matrix in the analysis provided some additional links among these herds. Thus, the average of the off-diagonals of the *A* matrix was 0.010 between Crichton and NBZ, 0.008 between Crichton and Moorepark, 0.013 between Crichton and TGEN, 0.008 between NBZ and Moorepark, 0.011 between NZB and TGEN and 0.012 between Moorepark and TGEN. For comparison, the average of the off-diagonals of the *A* matrix within herd was 0.016, 0.013, 0.015 and 0.022 for Crichton, NBZ, Moorepark and TGEN, respectively.

The only significant ($P < 0.05$) genetic correlation pertained to high heritability traits (average fat and protein percent, and fat-to-protein ratio). Thus, significant genetic correlations were estimated between Crichton and TGEN for fat percentage in 44 weeks (0.98), protein percentage in 44 weeks (0.94), fat percentage in 15 weeks (0.99) and fat-to-protein ratio in 44 weeks (0.98). Additional significant correlations were found between Crichton and NBZ for fat percentage in 44 weeks (0.94) and protein percentage in 15 weeks (0.97), and between NBZ and Moorepark for fat percentage in 44 weeks (0.85) and protein percentage in 15 weeks (0.94). In all these cases, estimated correlations were practically not different from unity, suggesting that trait definition was consistent in the four herds.

Genetic correlations between rare traits in the different herds were mostly inestimable. Among valid estimates, the correlation between Crichton and TGEN (the two herds with the larger number of common bulls) was 0.74 ($P = 0.33$) for average energy balance in 15 weeks, 0.90 ($P = 0.21$) for total energy intake in 15 weeks, 0.81 ($P = 0.28$) for total dry matter

intake in 15 weeks and 0.75 ($P = 0.38$) for average liveweight. Although highly positive, these estimates can only be viewed as potentially indicative values that did not attain statistical significance, thereby meriting no further discussion.

It is possible that conventional traits such as milk yield are more consistently defined and recorded in different herds than traits like energy intake, energy balance and liveweight. This will be a difficult issue to resolve because of potential differences in recording equipment and methods in the various herds. Nevertheless, the utility of merging database is expected to be more pronounced for such difficult-to-record traits. Despite the inestimability of genetic correlations among herds, lack of evident sire-by-herd interaction reported in the present study lends support to this claim.

Impact of increased dataset size on the power of effect detection

Increasing the size of a dataset implies greater statistical power, meaning that effects of lesser magnitude could become detectable in a genetic analysis. To assess the magnitude of this benefit, *post-hoc* power analyses were conducted based on a simple simulation design. Power values of detection for varying effect sizes and an alpha level equal to 0.05 were calculated (Erdfelder *et al.*, 1996), considering first the data size of each individual herd separately and then that of the combined dataset. Single fixed effects whose size was expressed in phenotypic standard deviation units were considered.

Results from this exercise are shown in Figure 3. As expected, power of effect detection was highest for the combined dataset,




Figure 3 Power values of a single fixed effect detection ($P < 0.05$) for different effect sizes by herd (Crichton – Scotland, NBZ – the Netherlands, Moorepark – Ireland, TGEN – the Netherlands) and for the combined database (All), expressed in standard deviation units.

reaching nearly unity for effect size equal to 0.2 s.d. This value corresponds to the minimum difference between two levels of the effect, expressed in standard deviation units that could be detected; examples of the latter may be two alleles of a certain gene or genetic marker, or two levels of a treatment. When analysed separately, the three largest herds (Crichton – Scotland, Moorepark – Ireland and TGEN – the Netherlands) would reach the same power level for an effect size equal to 0.3 to 0.4 s.d. The smallest population (NBZ herd in the Netherlands) would stand to benefit the most from combining data, as by itself would not be able to achieve the same level of power before the effect size reached 0.8 s.d.

Conclusions

Significant genetic variance and heritability was calculated for nearly all traits after merging data from the four different experimental herds and predicting missing records, suggesting that genetic analyses based on a combined database is both feasible and sensible. The assumption of no genotype-by-environment (herd) interaction, which enhances the usefulness of merging datasets, seems to hold but should nonetheless be quantified in other merged datasets.

A combined dataset including all traits described in the present study is currently being considered in genome-wide association analyses. Across herd variance estimates presented here are being used as starting values for the calculation of genetic marker effects on the various traits.

This study was based on first lactation data only. Future studies may also consider multiple lactations and a different trait definition for mature compared with growing animals.

Acknowledgements

This work was jointly financed by a RERAD grant to Scottish Agricultural College and the RobustMilk project, which is financially supported by the European Commission under the Seventh Research Framework Program, Grant Agreement KBBE-211708. The content of this paper is the sole responsibility of the authors, and it does not necessarily represent the views of the European Commission or its services. Ainsley Bagnall and other farm staff are thanked for taking milk

samples and other recordings on the farm and Ross McGinn is thanked for judiciously curating the Langhill–Crichton data over 25 years. Farm staff, technicians and database managers at Teagasc, Moorepark are also gratefully acknowledged.

References

- Akaike H 1974. A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19, 716–723.
- Banos G, Coffey MP and Brotherstone S 2005. Modeling daily energy balance of dairy cows in the first three lactations. *Journal of Dairy Science* 88, 2226–2237.
- Beerda B, Ouweltjes W, Sebek LBJ, Windig JJ and Veerkamp RF 2007. Effects of genotype by environment interactions on milk yield, energy balance, and protein balance. *Journal of Dairy Science* 90, 219–228.
- Buckley F, Dillon P, Rath M and Veerkamp RF 2000. The relationship between genetic merit for yield and live weight, condition score and energy balance of spring calving Holstein–Friesian dairy cows on grass based systems of milk production. *Journal of Dairy Science* 83, 1878–1886.
- Buttcherit N, Stamer E, Junge W and Thaller G 2011. Genetic relationships among daily energy balance, feed intake, body condition score, and fat to protein ratio of milk in dairy cows. *Journal of Dairy Science* 94, 1586–1591.
- Coffey MP, Emmans GC and Brotherstone S 2001. Genetic evaluation of dairy bulls for energy balance traits using random regression. *Animal Science* 73, 29–40.
- Emmans GC 1994. Effective energy: a concept of energy utilization applied across species. *British Journal of Nutrition* 71, 801–821.
- Erdfelder E, Faul F and Buchner A 1996. GPPOWER: a general power analysis program. *Behavior Research Methods, Instruments and Computers* 28, 1–11.
- Friggens NC, Ingvarsten KL and Emmans GC 2003. Prediction of body lipid change in pregnancy and lactation. *Journal of Dairy Science* 87, 988–1000.
- Gilmour AR, Gogel BJ, Cullis BR, Welham SJ and Thompson R 2006. *ASREML user guide, release 2.0*. VSN International Ltd, Hemel Hempstead, UK.
- Horan B, Faverdin P, Delaby L, Rath M and Dillon P 2006. The effect of strain of Holstein–Friesian dairy cows and pasture-based system on grass intake and milk production. *Animal Science* 82, 435–444.
- International Committee for Animal Recording 2011. www.icar.org, Rome, Italy.
- Kennedy E, O'Donovan M, Murphy JP, O'Mara FP and Delaby L 2006. The effect of initial grazing date and subsequent stocking rate on the grazing management, grass dry matter intake and milk production of dairy cows in summer. *Grass Forage Science* 61, 375–384.
- Kennedy J, Dillon P, Faverdin P, Delaby L, Stakelum G and Rath M 2003. Effect of genetic merit and concentrate supplementation on grass intake and milk production with Holstein–Friesian dairy cows. *Journal of Dairy Science* 86, 610–621.
- McCarthy S, Berry DP, Dillon P, Rath M and Horan B 2007. Effect of strain of Holstein–Friesian and feed system on udder health and milking characteristics. *Livestock Science* 107, 1–28.
- McEvoy M, O'Donovan M, Murphy JP, O'Mara F, Rath M and Delaby L 2007. Effect of concentrate supplementation and herbage allowance on milk production performance of spring calving dairy cows in early lactation. *Proceedings of the Irish Agricultural Research Forum, Tullamore, Ireland, 15–16 March 2007*, p. 46.
- Mrode RA and Swanson GJT 2003. Estimation of genetic parameters for somatic cell count in the first three lactations using random regression. *Livestock Production Science* 86, 253–260.
- O'Donovan M and Delaby L 2005. A comparison of perennial ryegrass cultivars differing in heading date and grass ploidy with spring calving dairy cows grazed at two different stocking rates. *Animal Research* 54, 337–350.
- Ordway RS, Boucher SE, Whitehouse NL, Schwab CG and Sloan BK 2009. Effects of providing two forms of supplemental methionine to periparturient Holstein dairy cows on feed intake and lactational performance. *Journal of Dairy Science* 92, 5154–5166.
- Pryce JE, Nielson BL, Veerkamp RF and Simm G 1999. Genotype and feeding system effects and interactions for health and fertility traits in dairy cattle. *Livestock Production Science* 57, 193–201.
- Toshniwal JK, Dechow CD, Cassell BG, Appuhamy J A D R N and Varga GA 2008. Heritability of electronically recorded daily body weight and correlations with yield, dry matter intake, and body condition score. *Journal of Dairy Science* 91, 3201–3210.
- Urioste JI, Franzén J and Strandberg E 2010. Phenotypic and genetic characterization of novel somatic cell count traits from weekly or monthly observations. *Journal of Dairy Science* 93, 5930–5941.

Banos, Coffey, Veerkamp, Berry and Wall

Vallimont JE, Dechow CD, Daubert JM, Dekleva MW, Blum JW, Barlieb CM, Liu W, Varga GA, Heinrichs AJ and Baumrucker CR 2010. Genetic parameters of feed intake, production, body weight, body condition score, and selected type traits of Holstein cows in commercial tie-stall barns. *Journal of Dairy Science* 93, 4892–4901.

Van Es AJH 1978. Feed evaluation for ruminants. I. The systems in use from May 1977–onwards in the Netherlands. *Livestock Production Science* 5, 331–345.

Veerkamp RF, Simm G and Oldham JD 1995. Genotype by environment interaction – experience from Langhill. In *Breeding and feeding the high genetic*

merit dairy cow (ed. TLJ Lawrence, FJ Gordon and A Carson), vol. 19. pp. 59–66. British Society of Animal Science (Occasional Publication), Midlothian, Scotland.

Veerkamp RF, Oldenbroek JK, van der Gaast HJ and van der Werf JHJ 2000. Genetic correlation between days until start of luteal activity and milk yield, energy balance and live weights. *Journal of Dairy Science* 83, 577–583.

Windig JJ, Beerda B and Veerkamp RF 2008. Relationship between milk progesterone profiles and genetic merit for milk production, milking frequency, and feeding regimen in dairy cattle. *Journal of Dairy Science* 91, 2874–2884.