

1 **Food-chain transfer of zinc to aphids**

2 Danielle Sinnett, Environmental and Human Sciences Division, Forest Research, Alice Holt

3 Lodge, Farnham, Surrey, GU10 4LH, UK, Tel: +44 1420 22255, Fax: +44 1420 520180,

4 danielle.sinnett@forestry.gsi.gov.uk

5 Number of words:

6 Number of references: 32

7 Number of tables: 0

8 Figure legends:

9 **Figure 1: Zn concentration in a) the leaf tissue of *A. pseudoplatanus* (n=25) and above-**
10 **ground tissue of *U. dioica* (n=15) exposed for 98 and 54 days respectively and b) *D.***
11 ***platinoïdis* (n=23) and *M. carnosum* (n=15) exposed for 14 and 28 days respectively**
12 **compared to the Zn concentration in Hoagland's solution and c) is the Zn**
13 **concentration in *D. platinoïdis* (n=23) and *M. carnosum* (n=15) compared with the Zn**
14 **concentration in leaf tissue of *A. pseudoplatanus* and above-ground tissue of *U. dioica*.**
15 **(Where [ZnNettle], [ZnLeaf], [ZnAphid] is the concentration of Zn in the tissue of *U.***
16 ***dioica* (mg/kg), the leaf tissue of *A. pseudoplatanus* (mg/kg), *M. carnosum* (mg/kg) and**
17 **the watering solution (mg Zn/l) respectively).**

18 **Figure 2: Zn concentration in the phloem tissue of *U. dioica* (n=15) and *A.***
19 ***pseudoplatanus* (n=15) exposed for 54 and 98 days respectively to Zn in Hoagland's**
20 **solution.**

21 **Food-chain transfer of zinc from contaminated *Urtica dioica* and *Acer***
22 ***pseudoplatanus* L. to *Microlophium carnosum* and *Drepanosiphum***
23 ***platanoidis* Schrank**

24

25 Danielle Sinnett†‡*, Tony R. Hutchings† and Mark E. Hodson‡

26 †Environmental and Human Sciences Division, Forest Research, Alice Holt Lodge, Farnham,
27 Surrey, GU10 4LH, UK.

28 ‡Department of Soil Science, School of Human and Environmental Sciences, University of
29 Reading, Whiteknights, Reading, RG6 6DW, UK.

¹ danielle.sinnett@forestry.gsi.gov.uk, Alice Holt Research Station

31 **Abstract** – This study examines the food-chain transfer of Zn from two plant species, *Urtica*
32 *dioica* (stinging nettle) and *Acer pseudoplatanus* (sycamore maple), into their corresponding
33 aphid species, *Microlophium carnosum* and *Drepanosiphum platanoidis*. The plants were
34 grown in a hydroponic system using solutions with increasing concentrations of Zn from 0.017
35 to 42 mg Zn/l, although *U. dioica* only survived in solution containing up to 18 mg Zn/l.
36 Above-ground tissue total and phloem concentrations in *U. dioica* and *M. carnosum*
37 concentrations increased with increasing Zn exposure ($p < 0.001$). When *U. dioica* were
38 exposed to the 18 mg Zn/l solution the corresponding above-ground plant tissue, phloem and
39 *M. carnosum* concentrations were around 2100, 50 and 131 mg/kg respectively. Although Zn
40 concentrations in *M. carnosum* were lower than total plant concentrations bioaccumulation
41 was taking place as concentrations were greater than those in the phloem which represents
42 the Zn reservoir to which the aphids were exposed. Zn concentrations in *A. pseudoplatanus*
43 also increased with solution concentration from the control to the 9 mg Zn/l solution, after
44 which concentrations remained constant at around 160 mg/kg. Zn concentrations in both *D.*
45 *platanoidis* and the phloem tissue of *A. pseudoplatanus* were not affected by the Zn
46 concentration in the watering solution with concentrations of 6.2 and 375 mg/kg respectively
47 for exposure to solutions of 18 mg Zn/l and above. It appears that *A. pseudoplatanus* is able
48 to regulate Zn, whereas *U. dioica* is not resulting in increasing Zn exposure to the aphids on
49 the latter species. Despite this Zn concentrations in *M. carnosum* were around a third of
50 those in *D. platanoidis*, suggesting that the latter species may have naturally elevated Zn
51 concentrations.

52

53 **Keywords** – stinging nettle, sycamore maple, common nettle aphid, sycamore aphid,
54 contaminated land

55

INTRODUCTION

56 The importance of the impact of contaminated land on terrestrial ecological receptors is
57 increasingly being recognised in the site investigation, risk assessment and remediation
58 process. Many practitioners commonly use an Ecological Risk Assessment (ERA) to
59 determine the potential for harm that a site may pose to ecological receptors and many
60 countries have produced frameworks and guidance for conducting such investigations [1].
61 The ERA process often makes use of a combination of field and laboratory analysis and
62 models to determine the risk to either ecological function or the food-chain transfer of
63 pollutants. The majority of the ecotoxicological tests used in ERA are based on ecological
64 function and use endpoints such as mortality, reproduction and growth. In order to estimate
65 the risk to higher organisms from a contaminated site it is often necessary to use models to
66 predict the pollutant concentrations through the food-chain and relate these to published
67 toxicological endpoints for the species of interest. There are a variety of models available to
68 estimate the food-chain transfer of pollutants (e.g. [2,3]). However, the models are often not
69 species specific, may have been based on aquatic organisms (for example in the case of
70 flying insects), or may only be applicable to a certain group of contaminants [2,3]. This has
71 serious implications for those using such models to estimate risk from contaminated land to
72 ecological receptors. At best it may result in significant gaps in the range of species for which
73 such a risk assessment can be conducted, at worst it may result in an over or under-
74 estimation of the risk leading to either unnecessarily costly remediation or no remediation
75 taking place where it is needed.

76 *Urtica dioica* L. is prevalent in almost all urban ecosystems and is an early coloniser of
77 contaminated land [4,5]. It is extremely important in urban ecosystems as it provides a habitat
78 for a wide range of invertebrates [5,6]. In addition, it is also relatively simple to cultivate,
79 widely available and fast growing [6], and as such, may be a useful species for
80 ecotoxicological testing. *Acer pseudoplatanus* L. is a tree species that has been introduced to
81 the UK, but is commonly found in urban areas [7]. It is an early coloniser [8] and tolerant of a
82 wide range of site conditions [9]. *U. dioica* and *A. pseudoplatanus* both have extremely

83 prevalent species-specific aphids associated with them; *Microlophium carnosum* Buckton and
84 *Drepanosiphum platanoidis* Schrank respectively.

85 The food-chain transfer of metals to a variety of aphids have been assessed in a number of
86 studies (e.g. [10-12]), although these studies have all concentrated on aphids whose hosts
87 are agricultural plant species. Aphids are an important source of food for a large number of
88 other insects, either indirectly for their honeydew (e.g. ants) or directly (e.g. parasitoids and
89 ladybirds) [7,13]. *M. carnosum* is a large aphid (3.3 to 3.8 mm) commonly found on *U. dioica*,
90 primarily on the underside of the leaves and the stem [14], during May to October [5,6]. *D.*
91 *platanoidis* is abundant on the underside of leaves of *A. pseudoplatanus*, during April to
92 October, with population peaks in June and October [8]. Used in conjunction with *U. dioica*
93 and *A. pseudoplatanus*, *M. carnosum* and *D. platanoidis* have the potential to assess the risk
94 of food-chain transfer of metals in urban ecosystems.

95 This study aims to assess the transfer of Zn to *M. carnosum* and *D. platanoidis* from *U. dioica*
96 and *A. pseudoplatanus* grown under hydroponic conditions in order to determine the potential
97 for Zn transfer to aphid predators in urban ecosystems. The study was originally carried out
98 with Cd in addition to Zn, however the small masses of aphids combined with the smaller
99 concentrations of Cd in their tissue meant that Cd concentrations in aphids were often below
100 detection limits and therefore Cd data are not reported here due to the patchy nature of the
101 dataset.

102

MATERIALS AND METHODS

103

Transfer of Zn into Microlophium carnosum and Drepanosiphum platanoidis

104

U. dioica cuttings, taken from Alice Holt Forest, Farnham, UK, and *A. pseudoplatanus* (bare

105

rooted 1+1 stock; Prees Heath Forest Nurseries, Shropshire, UK) were planted individually in

106

1 litre containers filled with perlite. Perlite was used as it has no inherent sorption capacity

107

that could influence Zn availability. Additionally, pores between individual perlite beads

108

ensure an aerobic environment. Pea shingle was placed on the perlite to a depth of 2 cm to

109

minimise evaporation.

110 A fully replicated randomised block experiment with five replicates was set up in a
111 glasshouse. Plants were grown under 16 h of artificial light and 8 h darkness per day (PAR =
112 0.37 mmol/m/s). The temperature of the glasshouse was regulated to 20 °C (± 5 °C).

113 Each container was watered with one of five solutions: control ($\frac{1}{4}$ strength Hoagland's
114 solution for *A. pseudoplatanus* and full strength for *U. dioica* [15]) or one of four Zn treatments
115 in Hoagland's solution. The Hoagland's formulation provided background micronutrient
116 concentrations of 0.02 or 0.08 mg Zn/l and 0.0008 or 0.0032 mg Cu/l for the $\frac{1}{4}$ strength or full
117 strength solutions respectively. Zn amendments were added as $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ to provide
118 concentrations of 0.02 (control), 5, 10, 20 and 50 mg Zn/l. The solution in each container was
119 replaced by mass when necessary.

120 *M. carnosum* were added to the *U. dioica* pots 26 days after planting whilst *D. platanoidis*
121 were added to the *A. pseudoplatanus* pots 84 days after planting. Differences in timing were
122 due to the availability of sufficient aphid populations in the field. Leaves with aphids on them
123 were removed from Alice Holt Forest and placed at the base of each plant. Enough leaves
124 were used so that at least 5 aphids were transplanted to each pot. Each pot was then
125 covered individually with a fine mesh net suspended from the ceiling, this was tied securely
126 around the lip at the top of the pot to prevent the aphids from moving to different plants. At
127 each watering, the netting was loosened round the pot and lifted enough to add the
128 appropriate solution and re-secured.

129 *U. dioica* and *A. pseudoplatanus* were harvested 28 and 14 days respectively, after the
130 aphids had been added. *D. platanoidis* populations appeared to be declining on the *A.*
131 *pseudoplatanus* so these were harvested earlier than *U. dioica* in order to ensure that enough
132 aphid mass was available for analysis. Reproduction rates of *D. platanoidis* vary during the
133 season, being closely linked to the amino-nitrogen content of the leaves and this decline is
134 likely to have been a result of the leaves reaching maturity [8]. The netting was loosened
135 from around the pot and the stem cut, the netting was then closed at the bottom and detached
136 from the ceiling, the netting along with its contents were then placed in the freezer at -20 °C
137 for 2 hours. The plants were then removed from the freezer and the dead aphids collected
138 with a fine brush. *A. pseudoplatanus* were split into their stem, shoot and leaf components.
139 The above-ground tissue of *U. dioica* and leaf and shoot tissues of *A. pseudoplatanus* were

140 washed in deionised water to remove the honeydew, weighed and dried at 70 °C for 24 hours
141 and reweighed. The stem tissues of *A. pseudoplatanus* were discarded as the aphids do not
142 feed on this woody material. The aphids were weighed, dried at 50 °C for 24 hours and
143 reweighed. The aphid and plant material samples were then milled and analysed to
144 determine their Zn concentrations (see below).

145 *Determination of phloem Zn concentrations*

146 In order to understand the different Zn exposures to the aphids a further experiment was set
147 up to determine the phloem Zn concentrations within *U. dioica* and *A. pseudoplatanus*. *U.*
148 *dioica* cuttings and *A. pseudoplatanus* were planted individually in 1 litre containers filled with
149 perlite in the same way as for the aphid exposure experiment. A fully replicated randomised
150 block experiment with five replicates for *U. dioica* and five replicates for *A. pseudoplatanus*
151 was set up in a glasshouse under the same conditions as the aphid exposure experiment.
152 Each container was watered with one of three solutions: control (¼ strength Hoaglands
153 solution for *A. pseudoplatanus* and full strength for *U. dioica* [15]) or one of two Zn treatments
154 in Hoaglands solution. Zn amendments were added as ZnSO₄·7H₂O to provide
155 concentrations of 0.02 (control), 5 and 20 mg Zn/l. The solution in each container was
156 replaced by mass when necessary.

157 *U. dioica* and *A. pseudoplatanus* were harvested after the same duration as the aphid
158 experiment in order to ensure that the plants had been exposed to the Zn solutions for the
159 same time. The method used to determine the concentration of Zn in the phloem tissue was
160 based on that of Thornber and Northcote [16] which extracts the water-soluble material within
161 the phloem. The leaf and shoot tissues of *A. pseudoplatanus* were removed from the stem
162 tissue and discarded. The bark was carefully removed from the stem tissue using a grafting
163 knife and the phloem tissue was then removed, again with a grafting knife. The phloem tissue
164 was weighed and then boiled at 100 °C in 200 ml of deionised water for 3 hours. Following
165 boiling, the samples were centrifuged and the solution removed and filtered through a 0.45
166 µm Whatman filter. It was not possible to separate the phloem tissue of *U. dioica* from the
167 rest of the stem so the entire above-ground biomass was subjected to boiling under the
168 assumption that the water-soluble fraction of the plant material will give an indication of the

169 phloem concentration. The above-ground tissue of *U. dioica* was removed, weighed and
170 boiled at 100 °C in 300 ml of deionised water for 3 hours. The solutions were then analysed
171 to determine their Zn concentrations (see below).

172 *Determination of Zn concentration*

173 The Zn solutions used for watering and the phloem extracts were analysed using a Spectro
174 Flame Inductively Coupled Plasma – Optical Emission Spectrometer (ICP-OES; Spectro
175 Analytical Instruments, West Midlands, UK). The target Zn concentrations in the solutions
176 used for watering of 0.02, 5 and 20 mg Zn/l were found to be 0.017, 4.71 and 17.97 mg Zn/l
177 respectively.

178 Plant samples were prepared for analysis by dry-ashing at 450 °C for 18 hours and wet
179 digestion [17]. Wet digestion was achieved by incubating each sample for 1 hour at 60 °C in
180 0.75 ml concentrated HNO₃, followed by a further 14 hour incubation with 2.25 ml
181 concentrated HCl and heating for 2 hours at 110 °C. After cooling, 0.15 ml of 30 % H₂O₂ was
182 added to each sample followed by heating for 30 minutes at 110 °C. To ensure complete
183 oxidation of all organic matter the H₂O₂ treatment was performed twice. The digested
184 samples were analysed for Zn using the ICP-OES [18].

185 Aphid samples were digested in 1 ml concentrated HNO₃ at 180 °C for 1 hour, after which 1
186 ml of deionised water was added and the sample further digested at 180 °C to dryness. A
187 further 0.01 ml of concentrated nitric acid was added and the sample digested at 60 °C for 1
188 hour. The digested samples were analysed for Zn using the ICP-OES [18].

189 The limit of detection was 0.67 µg/kg for Zn. Bush branches and leaves (NCS DC73349,
190 China National Analysis Centre for Iron and Steel), oriental tobacco leaves (CTA-OTL-1,
191 Commission for Trace Analysis of the Committee for Analytical Chemistry of the Polish
192 Academy of Sciences and Institute of Nuclear Chemistry and Technology, Warsaw, Poland),
193 mussel (CE278, European Commission, Geel, Belgium) and bovine liver (1577b, US
194 Department of Commerce, National Institute of Standards and Technology, Gaithersburg, MD
195 20899, USA) tissues were used as Certified Reference Materials (CRM) with batches of plant
196 and aphid samples as appropriate. Mean recovery from oriental tobacco leaves was 104.9
197 and 98.6 % from the bush branches and leaves and oriental tobacco respectively. Mean

198 recovery from mussel and bovine liver was 92.9 and 93.6 % respectively for the *M. carnosum*
199 samples and 111.8 and 102.6 % respectively for the *D. platanoidis* samples.

200 *Statistical analysis*

201 The plant and aphid Zn uptake data were subjected to general linear regression analysis to
202 assess the significance of changes in plant and aphid concentrations with increasing Zn
203 concentration in hydroponic solutions and plant material respectively, using Genstat version
204 8.1 [19]. Mean values are reported with \pm standard errors throughout.

205 Linear and exponential models of Zn uptake into each of the plant and aphid tissue types
206 compared to that of the solution concentration and, in the case of aphids, the leaf
207 concentrations were fitted using Genstat version 8.1 [19]. A comparison of the residual sum
208 of squares of alternative models relative to the smallest residual mean square was used to
209 determine the most appropriate model. This comparison used for nested models and is
210 referred to an F-distribution with 1, n degrees of freedom where n is the residual degrees of
211 freedom from the exponential model.

212 **RESULTS**

213 Zn concentration in solution had a significant affect on the Zn uptake into the above ground
214 tissue ($F_{1,13}=533.63$; $p<0.001$) of *U. dioica*; no plants survived in the 42 mg Zn/l solution
215 treatment (Figure 1). Zn concentration in both solution and nettle tissue had a significant
216 affect on the Zn concentration in *M. carnosum* ($F_{1,13}=107.95$; $p<0.001$ and $F_{1,13}=77.38$;
217 $p<0.001$ respectively; Figure 1). The concentration of Zn in the phloem extracts from *U.*
218 *dioica* increased significantly with increasing Zn concentration in solution ($F_{1,13}=138.89$;
219 $p<0.001$; $r^2=0.908$).

220 Zn concentration in solution did not have a significant effect on the Zn uptake into either the
221 leaf or shoot tissue of *A. pseudoplatanus*. This is because the Zn concentrations in the
222 tissues reached a plateau between the 9 and 18 mg Zn/l solutions; the exponential model
223 was, however, significant for both leaf ($F_{2,22}=3.57$; $p=0.046$) and shoot ($F_{2,22}=5.43$; $p=0.012$)
224 tissue (Figure 1).

225 There was no significant effect of the concentration of Zn in solution or in the leaf or shoot
226 tissue of *A. pseudoplatanus* on the concentration in *D. platanoidis* using either the linear or
227 exponential models (Figure 1). Similarly, the concentration of Zn in the phloem extract was
228 not significantly related to the concentration of Zn in solution.

229 The concentration of Zn in the above-ground tissue of *U. dioica* were approximately 13 times
230 that in the *A. pseudoplatanus* as a result of exposure to the 18 mg Zn/l solution;
231 2153 ± 68.7 mg/kg compared with 163 ± 20.6 mg/kg. The phloem extract concentrations at this
232 18 mg Zn/l exposure were 48.2 ± 2.4 mg/kg in *U. dioica* and 6.1 ± 1.2 mg/kg in *A.*
233 *pseudoplatanus* (Figure 2). Despite this, the Zn concentration in *M. carnosum* was less than
234 a third of that in *D. platanoidis*; 131.5 ± 11.0 mg/kg compared with 406 ± 21.2 mg/kg. Phloem
235 concentrations of both species were lower than those in above-ground tissue in *U. dioica* or in
236 the leaf and shoot tissues in *A. pseudoplatanus*, this difference increased with increasing Zn
237 concentrations; from 3 up to 17 times lower and 11 up to 25 times lower in *U. dioica* and *A.*
238 *pseudoplatanus* respectively.

239

DISCUSSION

240 Zn concentrations in the above-ground tissue of *U. dioica* increased with Zn exposure,
241 reaching a mean of approximately 2100 mg/kg for the 18 mg Zn/L solution. In *A.*
242 *pseudoplatanus* tissue concentration increased up to the 9 mg Zn/l solution and then
243 remained constant at around 160 mg/kg despite the increasing Zn concentration in solution.
244 Zn concentrations in the above-ground tissue of *U. dioica* have been reported to range
245 between 42 and 52 mg/kg in uncontaminated soils [20]. Leaf concentrations of between 23
246 and 532 (mean 113 mg/kg) have been reported in *U. dioica* growing on dredged sediments
247 with a Zn concentration of between 149 and 1817 (mean 54 mg/kg) [21]. Zn concentrations in
248 *U. dioica* around the Avonmouth smelter have been found to be as high as 3000 mg/kg,
249 although this is likely to have occurred from atmospheric deposition as well as soil uptake
250 [22]. The substantial quantities of Zn that nettles appear to be capable of accumulating make
251 this species an important pathway for Zn in the food-chain. Mertens et al. [23] found Zn
252 concentrations with a mean of 74 mg/kg in *A. pseudoplatanus* grown on dredged sediments
253 with a Zn concentration of 359 mg/kg. The normal range of Zn in plant tissue has been

254 reported to be 27-150 mg/kg with an upper toxic limit of 100-500 mg/kg [24], which suggests
255 that the concentrations reported here for *A. pseudoplatanus* are unlikely to cause a toxic
256 effect.

257 The Zn concentrations in the tissue of *U. dioica* and *A. pseudoplatanus* showed large
258 differences; at the lowest Zn solution concentration the tissue concentration of *A.*
259 *pseudoplatanus* is greater than that of *U. dioica*, but at higher concentrations the reverse is
260 true, increasing from a 3 fold to a 13 fold difference at the highest solution concentration. The
261 relationships between solution and tissue concentration between the species were also
262 different; *U. dioica* having a steep linear relationship whilst for *A. pseudoplatanus* the
263 relationship was exponential with the Zn tissue concentrations reaching a plateau at around
264 160 mg/kg. This suggests different responses to Zn between the two species. *U. dioica* is
265 unable to regulate Zn and continues to accumulate this metal until a toxic concentration is
266 reached and the plant can no longer survive, in the present experiment this must have
267 occurred to plants grown in the 42 mg Zn / L solution. In contrast *A. pseudoplatanus* is able to
268 regulate the Zn concentration in its above-ground tissue and therefore survive in media
269 containing higher concentrations of Zn.

270 Previous studies investigating the transfer of metals into aphids have used wheat grown in
271 sewage sludge amended soils. In these studies the Zn concentrations in the plant tissue
272 were substantially lower (<150 mg/kg) [11,12,25-27] than those found in *U. dioica* in the
273 current study and more comparable to those in *A. pseudoplatanus*. Despite this, the
274 concentrations of Zn in *M. carnosum* reported in the current study are similar to those found in
275 these previous studies, which used different aphid species [11,12,25-27], whereas the
276 concentrations in *D. platanoidis* were generally two to three times greater, even at the lowest
277 Zn solution concentration. All of these studies found that Zn was bioaccumulated in the
278 aphids *Rhopalosiphum padi* and *Sitobian avenae* feeding on wheat. In our study, from the
279 total plant concentrations it appeared that *M. carnosum* was not accumulating Zn as the
280 *U. dioica* bulk tissue concentration from the 18 mg Zn/l solution was around 2100 mg/kg and
281 the aphid concentration was 131 mg/kg. However, the analysis of the phloem tissue of the
282 nettle tissue revealed that *M. carnosum* were accumulating Zn as this concentration was

283 around 50 mg/kg. Zn concentrations in *D. platanoidis* were greater, at around 375 mg/kg,
284 than both the total plant and the phloem concentrations of 160 and 6.2 mg/kg respectively.

285 It has been reported that Zn is concentrated in the stem tissue as well as the roots [28] and is
286 readily transported in the phloem of *A. pseudoplatanus* [29] and wheat [28,30]. Aphids feed
287 directly on the phloem sap [8] and are therefore exposed to the Zn within this solution. The
288 chemical form that Zn takes within the phloem is not well understood [31], although is likely to
289 be in a soluble form, bound to chelators, amino acids and/or organic acids, it is also unclear
290 whether the Zn is transported apoplastically or symplastically [31]. Studies on barley have
291 shown that, whilst most of the Zn in the roots is soluble, that in the leaves is primarily located
292 in the mesophyll cells and, to a lesser extent the epidermal cells; where it is present primarily
293 in the cytoplasm, followed by the chloroplasts. The Zn in the apoplatic solution is mainly (97
294 %) bound to cell walls [31]. The ability of *A. pseudoplatanus* to regulate Zn in its above-
295 ground biomass, and because the transfer of Zn to the phloem is regulated by the
296 requirements of the plant, may explain why, in this species, the concentrations in the phloem
297 are similar regardless of the exposure to the plant or plant tissue concentrations. Water-
298 soluble concentrations of Zn in *U. dioica* are much lower than the total plant concentrations,
299 suggesting that the Zn within this species is also bound within the plant tissue and not readily
300 transported in the phloem. However, the water-soluble concentrations increase with
301 increasing Zn concentration in the watering solution and the plant tissue, further suggesting
302 that *U. dioica* is not able to regulate Zn transport within the plant. This results in increased
303 exposure to aphids as the concentration of Zn in the plant tissue increases.

304 Although it appears that both aphid species bioaccumulated Zn, the concentrations in *M.*
305 *carosum* were smaller than those for *D. platanoidis* despite an increased level of exposure.
306 This may be because the duration of exposure of *D. platanoidis* was double that for *M.*
307 *carosum*. Alternatively, *M. carosum* may be able to regulate Zn; Crawford et al. [10] found
308 that *Aphis fabae* on broad beans (*Vicia faba*) were able to regulate Cu by excretion in
309 honeydew. Unfortunately, it proved impossible to obtain sufficient quantities of honey dew for
310 analysis in this study. The Zn concentrations in *M. carosum* were comparable with those
311 found in other studies, whereas those in *D. platanoidis* were elevated. This, coupled with the
312 fact that the concentrations in *D. platanoidis* were elevated even when *A. pseudoplatanus*

313 was watered with the control solution suggest that this species may simply have naturally
314 greater Zn concentrations compared with other aphid species regardless of the concentration
315 within the plant.

316 The greater Zn concentrations in *D. platanoidis* has important implications, both for the
317 estimation of risk to higher organisms and the modelling of food-chain transfer, particularly
318 given that the Zn tissue concentrations in *A. pseudoplatanus* were substantially lower than
319 those in *U. dioica*. When the ladybird *Coccinella septempunctata*, lacewing *Chysoperla*
320 *carnae* and carabid beetle *Bembidion lampros* were fed aphids with Zn concentrations ranging
321 between 163-249, 104-188 and 60-116 mg/kg respectively their corresponding tissue
322 concentrations were between 184-217, 105-249 and 99-112 respectively [12,26,27]. This
323 suggests that, although only in the lacewing was Zn accumulated, the tissue concentrations of
324 the predators of aphids are likely to reflect the tissue concentrations of their prey. Therefore
325 species feeding on *D. platanoidis* may be exposed to higher concentrations of Zn in their diet
326 than those feeding on other species of aphid. This demonstrates the importance of species
327 specificity in modelling food-chain transfer in terrestrial ecosystems.

328

329 Acknowledgements – The authors wish to thank Sylvia Cowdry for assisting with experimental
330 work and Alberto Morales from the Environmental and Human Science Division Laboratory,
331 Forest Research for sample analysis. This work was funded by EPSRC as part of the
332 Pollutants in the Urban Environment (PurE) research consortium (Grant Number
333 EP/C532651/2).

334

REFERENCES

335 Literature Cited

336

- 337 1. Bryns G, Crane M. Assessing risks to ecosystems from land contamination.
338 Environment Agency, Bristol.
339 Ref Type: Report
- 340 2. USEPA. Multimedia, multipathway and multireceptor risk assessment (3MRA) modeling
341 system. Volume II: Site-based, regional and national data. USEPA, Washington DC.
342 Ref Type: Report
- 343 3. USEPA. TRIM:FaTE Technical Support Document. Volume II: Description of chemical
344 transport and transformation algorithms. USEPA, North Carolina.
345 Ref Type: Report

- 346 4. Edwards SC, MacLeod CL, Lester JN. 1998. The bioavailability of copper and mercury
347 to the common nettle (*Urtica dioica*) and the earthworm *Eisenia fetida* from
348 contaminated dredge spoil. *Water, Air, and Soil Pollution* 102:75-90.
349 Ref Type: Journal
- 350 5. Barta M, Cagán L. 2003. Entomophthoralean fungi associated with the common nettle
351 aphid (*Microlophium carnosum* Buckton) and the potential role of nettle patches as
352 reservoirs for the pathogens in landscape. *Journal of Pest Science* 76:6-13.
353 Ref Type: Journal
- 354 6. Davis BNK. 1991. *Insects on nettles*. The Richmond Publishing Co. Ltd., Slough.
355 Ref Type: Book, Whole
- 356 7. Gilbert OL. 1991. *The Ecology of Urban Habitats*. Chapman and Hall, London.
357 Ref Type: Book, Whole
- 358 8. Dixon T. 2005. *Insect Herbivore-Host Dynamics: Tree dwelling aphids*. Cambridge
359 University Press, Cambridge.
360 Ref Type: Book, Whole
- 361 9. Moffat, A. J. and McNeill, J. D. Reclaiming disturbed land for forestry. [110]. 1994.
362 Edinburgh, HMSO. Forestry Commission Bulletin.
363 Ref Type: Serial (Book, Monograph)
- 364 10. Crawford LA, Hodkinson ID, Lepp NW. 1995. The effects of elevated host-plant
365 cadmium and copper on the performance of the aphid *Aphis fabae* (Homoptera:
366 Aphididae). *Journal of Applied Ecology* 32:528-535.
367 Ref Type: Journal
- 368 11. Merrington G, Winder L, Green I. 1997. The uptake of cadmium and zinc by the bird-
369 cherry oat aphid *Rhopalosiphum padi* (Homoptera: Aphididae) feeding on wheat grown
370 on sewage sludge amended agricultural soil. *Environmental Pollution* 96:111-114.
371 Ref Type: Journal
- 372 12. Green I, Merrington G, Tibbet M. 2003. Transfer of cadmium and zinc from sewage
373 sludge amended soil through a plant-aphid system to newly emerged adult ladybirds
374 (*Coccinella septempunctata*). *Agriculture, Ecosystems and Environment* 99:171-178.
375 Ref Type: Journal
- 376 13. Rotheray GE. 1989. *Aphid predators*. The Richmond Publishing Co. Ltd., Slough.
377 Ref Type: Book, Whole
- 378 14. Kean JM, Müller CB. 2004. Can competition explain local rarity of a nettle aphid?
379 *Ecological Entomology* 29:706-710.
380 Ref Type: Journal
- 381 15. Hoagland DR, Arnon DI. The water-culture method for growing plants without soil.
382 California Agricultural Experiment Station, California.
383 Ref Type: Report
- 384 16. Thornber JP, Northcote DH. 1961. Changes in the chemical composition of a cambial
385 cell during its differentiation into xylem and phloem tissue in trees. *Biochemical Journal*
386 81:449-455.
387 Ref Type: Journal
- 388 17. Chapman HD. 1967. Plant analysis values suggestive nutrient status of selected crops.
389 *Soil Testing and Plant Analysis: Plant Analysis Part II*, Soil Science Society of America,
390 Wisconsin.
391 Ref Type: Book Chapter


- 392 18. Kilbride C, Poole J, Hutchings TR. 2006. A comparison of Cu, Pb, As, Cd, Zn, Fe, Ni
393 and Mn determined by acid extraction/ICP-OES and ex situ field portable X-ray
394 fluorescence analysis. *Environmental Pollution* 143:16-23.
395 Ref Type: Journal
- 396 19. Genstat. 2005. *The Guide to GenStat Release 8.1 Part 2: Statistics*. VSN International,
397 Oxford.
398 Ref Type: Book, Whole
- 399 20. Hobbelen PHF, Koolhaas JE, van Gestel CAM. 2004. Risk assessment of heavy metal
400 pollution for detritivores in floodplain soils in the Biesbosch, the Netherlands, taking
401 bioavailability into account. *Environmental Pollution* 129:409-419.
402 Ref Type: Journal
- 403 21. Tack FM, Verloo MG. 1996. Metal contents in stinging nettle (*Urtica dioica* L.) as
404 affected by soil characteristics. *The Science of the Total Environment* 192:31-39.
405 Ref Type: Journal
- 406 22. Jones DT. 1991. Biological monitoring of metal pollution in terrestrial ecosystems.
407 University of Reading.
408 Ref Type: Thesis/Dissertation
- 409 23. Mertens J, Vervaeke P, De Schrijver A, Luysaert S. 2004. Metal uptake by young trees
410 from dredged brackish sediment: limitations and possibilities for phytoextraction and
411 phytostabilisation. *Science of the Total Environment* 326:209-215.
412 Ref Type: Journal
- 413 24. Kabata-Pendias A, Pendias H. 2001. Elements of Group II. In Kabata-Pendias A,
414 Pendias H, eds, *Trace Elements in Soils and Plants*, 3rd Edition ed, CRC Press Ltd.,
415 Boca Raton, Florida, pp 123-164.
416 Ref Type: Book Chapter
- 417 25. Merrington G, Winder L, Green I. 1997. The bioavailability of Cd and Zn from soils
418 amended with sewage sludge to winter wheat and subsequently to the grain aphid
419 *Sitobion avenae*. *The Science of the Total Environment* 205:245-254.
420 Ref Type: Journal
- 421 26. Winder L, Merrington G, Green I. 1999. The tri-trophic transfer of Zn from the
422 agricultural use of sewage sludge. *The Science of the Total Environment* 229:73-81.
423 Ref Type: Journal
- 424 27. Green ID, Jeffries C, Diaz A, Tibbett M. 2006. Contrasting behaviour of cadmium and
425 zinc in a soil-plant-arthropod system. *Chemosphere* 64:1115-1121.
426 Ref Type: Journal
- 427 28. Haslett BS, Reid RJ, Rengel Z. 2001. Zinc mobility in wheat: Uptake and distribution of
428 zinc applied to leaves or roots. *Annals of Botany* 87:379-386.
429 Ref Type: Journal
- 430 29. Dollard GJ, Lepp NW. 1980. Differential mobility of lead and zinc in phloem tissue of
431 sycamore (*Acer pseudoplatanus* L.). *Zeitschrift fur Pflanzenphysiologie* 97:409-415.
432 Ref Type: Journal
- 433 30. Riesen O, Feller U. 2005. Redistribution of nickel, cobalt, manganese, zinc and
434 cadmium via the phloem in young and maturing wheat. *Journal of Plant Nutrition*
435 28:421-430.
436 Ref Type: Journal

437 31. Grusak MA, Pearson JN, Marentes E. 2007. The physiology of micronutrient
438 homeostasis in field crops. *Field Crops Research* 60:41-56.
439 Ref Type: Journal


440

441

a)



b)



c)




Figure 1: Zn concentration in a) the above-ground tissue of *U. dioica* (n=15) and the leaf tissue *A. pseudoplatanus* (n=25) exposed for 54 and 98 days respectively and b) *M. carnosum* (n=15) and *D. platinoidis* (n=23) exposed for 28 and 14 days respectively compared to the Zn concentration in Hoagland's solution in which the *U. dioica* and *A. pseudoplatanus* were grown and c) *M. carnosum* (n=15) and *D. platinoidis* (n=23) Zn concentrations compared with the Zn concentration in the above-ground tissue of *U. dioica* and the leaf tissue of *A. pseudoplatanus*. (Where [ZnNettle], [ZnLeaf], [ZnAphid], [ZnSoln] is the concentration of Zn in the tissue of *U. dioica* (mg/kg dry weight), the leaf tissue of *A. pseudoplatanus* (mg/kg dry weight), *M. carnosum* (mg/kg dry weight) and the watering solution (mg Zn/l) respectively).




Figure 2: Zn concentration in the phloem tissue of *U. dioica* (n=15; mg/kg wet weight) and *A. pseudoplatanus* (n=15; mg/kg wet weight) exposed for 54 and 98 days respectively to Zn in Hoagland's solution.