

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Free atmospheric CO₂ enrichment did not affect symbiotic N₂-fixation and soil carbon dynamics in a mixed deciduous stand in Wales

M. R. Hoosbeek¹, M. Lukac², E. J. Velthorst¹, and D. L. Godbold³

¹Department of Environmental Sciences, Earth System Science – Climate Change, Wageningen University, P.O. Box 47, 6700AA Wageningen, The Netherlands

²Imperial College London, South Kensington Campus, London SW7 2AZ, UK

³School of the Environment & Natural Resources, Bangor University, Bangor, Gwynedd LL57 2UW, UK

Received: 28 April 2010 – Accepted: 24 May 2010 – Published: 4 June 2010

Correspondence to: M. R. Hoosbeek (marcel.hoosbeek@wur.nl)

Published by Copernicus Publications on behalf of the European Geosciences Union.

4153

Abstract

Through increases in net primary production (NPP), elevated CO₂ is hypothesized to increase the amount of plant litter entering the soil. The fate of this extra carbon on the forest floor or in mineral soil is currently not clear. Moreover, increased rates of NPP can be maintained only if forests can escape nitrogen limitation. In a Free atmospheric CO₂ Enrichment (FACE) experiment near Bangor, Wales, 4 ambient CO₂ and 4 FACE plots were planted with patches of *Betula pendula*, *Alnus glutinosa* and *Fagus sylvatica* on a former arable field. Four years after establishment, only a shallow L forest floor litter layer had formed due to intensive bioturbation. Total soil C and N contents increased irrespective of treatment and species as a result of afforestation. We could not detect an additional C sink in the soil, nor were soil C stabilization processes affected by FACE. We observed a decrease of leaf N content in *Betula* and *Alnus* under FACE, while the soil C/N ratio decreased regardless of CO₂ treatment. The ratio of N taken up from the soil and by N₂-fixation in *Alnus* was not affected by FACE. We infer that increased nitrogen use efficiency is the mechanism by which increased NPP is sustained under elevated CO₂ at this site.

1 Introduction

Using an indirect method, Canadell et al. (2007) estimated the terrestrial carbon (C) sink to account for about a third of total anthropogenic carbon dioxide (CO₂) emissions at present. Forest ecosystems are hypothesized to constitute a large part of this sink and to sequester C due to their regrowth and atmospheric CO₂ fertilization (Houghton, 2003; Janssens et al., 2003; McMahon et al., 2010). In order to test this hypothesis and to assess the strength of this feedback, Free Air CO₂ Enrichment (FACE) experiments in aggrading temperate forests and plantations were initiated. To date, existing experiments have demonstrated that rising atmospheric CO₂ concentrations result in increases in net primary production (NPP) and C storage in forest vegetation, e.g.

4154

(Calfapietra et al., 2003; DeLucia et al., 1999; Gielen et al., 2005; Hamilton et al., 2002; Handa et al., 2006; Karnosky et al., 2003; Liberloo et al., 2009; Norby et al., 2002). Norby et al. (2005) analyzed the response of NPP to elevated CO₂ in four forest FACE experiments and found that its response is well conserved across a broad range of productivity, with a stimulation at the median of 23 ± 2%.

In general, the aboveground biomass contributes its C to the forest litter layer, where it is partially respired and partially incorporated into the mineral soil. Root litter contributes C directly to the mineral soil or, if present, also to the forest litter layer. The extra C taken up due to increased atmospheric CO₂ concentrations may also be stored in forest floor litter, organic O and mineral A horizon. Long term C storage is however thought to primarily take place in mineral soil horizons due to the occurrence of C stabilization mechanisms (Sollins et al., 2006; Six et al., 2002; Von Lützow et al., 2006).

As N availability commonly limits forest productivity, some combination of increased N uptake from the soil and more efficient use of the N assimilated by trees will be necessary to sustain the higher rates of forest NPP at future levels of CO₂. Based on data from four forest FACE sites, Finzi et al. (2007) demonstrated that increases in N uptake rather than N-use efficiency support high rates of temperate forest productivity under elevated CO₂. Nitrogen is also needed for the long term storage of C in stable organic matter fractions in the forest floor and mineral soil. In a meta-analysis based on 65 studies, Van Groenigen et al. (2006) found that soil C content only increases under elevated CO₂ when N is added at rates well above typical atmospheric deposition.

Biological N₂-fixation may be a possible source of N to sustain increased N uptake due to high rates of temperate forest productivity under elevated CO₂ (Vitousek et al., 2002). Although assimilation of N by N₂-fixation is considered to be more costly than uptake of ammonium or nitrate at the plant level, the extra cost might be offset by greater availability of assimilates in high CO₂. In a growth chamber experiment, elevated CO₂ increased dry weight and total nitrogenase activity of *Robinia pseudoacacia* and *Alnus glutinosa* seedlings, supporting the premise that CO₂ enrichment can stimulate symbiotic activity (Norby, 1987). In a chamber experiment with seedlings of *Alnus*

4155

rubra, Arnone III and Gordon (1990) observed a positive feedback loop between N-fixation and photosynthesis in nodulated plants growing under elevated CO₂. Similarly, in a number of open-top chamber experiments, symbiotic N₂-fixing *Alnus glutinosa* trees showed a positive response to elevated CO₂ (Vogel et al., 1997; Temperton et al., 2003).

In 2004 a mixed deciduous forest FACE experiment was initiated near Bangor, Wales, UK. This is the first FACE experiment which includes a symbiotic N₂-fixing tree, offering an opportunity to study the effects of elevated CO₂ on N fixation in forests. Taking into account preliminary biomass growth observations and published results from other forest FACE experiments, we formulated the following hypotheses:

1. In order to sustain higher rates of forest NPP under FACE, additional N is taken up from the soil.
2. FACE stimulates symbiotic N₂-fixation by increasing C availability in *Alnus glutinosa* root nodules, increasing the ratio of N taken up by N₂-fixation to N taken up from the soil.
3. Total soil C content and, to a lesser extent, N content increase due to afforestation.
4. Increased NPP under FACE creates additional C storage in the soil.
5. The additional soil C input due to FACE results in an increase of coarse, fine and micro-aggregate protected particulate organic matter (POM).

20 **2 Methods**

The BangorFACE experiment was established at the Henfaes experimental research area in March 2004. The area is located on the coastal plain about 12 km east of Bangor, near the village of Abergwyngregyn, Wales, UK. The climate is Hyperoceanic, with annual rainfall of about 1000 mm. The soil at Henfaes is a fine loamy brown earth

over gravel (Rheidol series) classified as a Dystric Cambisol in the FAO system (Teklehaimanot et al., 2002). The parent material consists of postglacial alluvial deposits from the Aber river, comprising Snowdonian rhyolitic tuffs and lavas, microdiorites and dolerite in the stone fractions and Lower Paleozoic shale in the finer fractions. The topography consists of a shallow slope of approximately 1–2° on a deltaic fan. The aspect is northwesterly, at an altitude of 13 to 18 m a.s.l. The depth of the water table ranges between 1 and 6 m.

Trees were planted on two adjacent fields, one of which was previously used both as pasture and arable land, whereas the other was used for small scale agroforestry experiments. The experimental plots were 8 m in diameter, the seedlings of *Betula pendula*, *Alnus glutinosa* and *Fagus sylvatica* were planted inside the plots at 80 cm spacing in a hexagonal design. The species were planted in a pattern that created mixtures containing one, two and three species. For the purposes of this study, 4 mixtures have been monitored within each experimental plot; three single species sub-plots and a sub-plot containing the mixture of all tree species. The experimental plots were surrounded by a 10 m buffer strip containing the same species and planted at the same density and pattern. The rest of the plantation was planted with a mixture of tree species at slightly smaller density. In total, 4 ambient and 4 FACE plots were randomly located within the plantation in order to form a complete replicated block design. Carbon enrichment started in April 2005 and was achieved by injecting pure CO₂ through laser-driller holes in tubing mounted on eight masts (Miglietta et al., 2001). The elevated CO₂ concentrations, measured at 1 min intervals, were within 30% deviation from the pre-set target concentration of 580 ppm CO₂ for 75–79% of the time during the photosynthetically active part of 2005–2008. The CO₂ used for enrichment originated from natural gas and had a δ¹³C of –39‰.

Soil samples were taken from each sub-plot in October of years 2004 through 2008. Bulk density samples were taken at 0–10 cm from the A horizon using a bulk density sampler holding 100 cm³ metal rings. Adjacent to these samples, three bulk samples representative for the 0–10 cm depth were taken with a small spade and mixed for C

4157

and N analyses and fractionation. After transportation in a mobile refrigerator, the ring samples were dried at 105 °C for 3 d, while the bulk samples were split in a part that was dried at room temperature and a part that was stored at 4 °C. Bulk densities were calculated based on oven dry weight of the ring samples and ring volume.

Soil texture and pH were only determined for the 2004 samples. After pre-treatment of the samples, the particle size distribution was measured by laser diffraction (Coulter LS230 Grain Sizer; Buurman et al., 1996). Soil pH was measured with a pH meter (Orion 701A) in a 1 M KCl solution suspension.

For C and N analyses, sub-samples of the air-dried bulk samples were crushed by hand and ball milled after roots were removed. No carbonates were present in the soil. Ammonium and nitrate were measured colorimetrically in a 1 M KCl extraction by using an auto analyzer (Buurman et al., 1996). Total C and N were determined with an elemental analyzer (Interscience EA 1108) and expressed as gram C or N per m² per depth increment.

2.1 Isotope analyses

In 2007, leaves and young branches of *Betula* and *Alnus* and soil samples from all sub-plots were collected, dried, milled and prepared and sent for analysis at the Stable Isotope Laboratory at UC Davis (<http://stableisotopefacility.ucdavis.edu>). Results were expressed as δ¹³C (‰) versus the PDB standard and as δ¹⁵N (‰) versus standard air. The fraction of soil C derived from litter input (C_{new}) between October of 2004 and 2007 (f_{new C}) was calculated as (Balesdent et al., 1988; Van Kessel et al., 2000):

$$f_{\text{new C}} = \frac{(\delta^{13}\text{C}_{\text{soil FACE}} - \delta^{13}\text{C}_{\text{soil ambient}})}{(\delta^{13}\text{C}_{\text{new}} - \delta^{13}\text{C}_{\text{soil ambient}})} \quad (1)$$

The new soil C input (g C m⁻²) into the 0–10 cm increment of the FACE plots was calculated as:

$$C_{\text{soil new}} = f_{\text{new C}} \cdot C_{\text{soil FACE}} \quad (2)$$

4158

respectively. These $\delta^{15}\text{N}$ values represent the uptake of N solely from the soil. Domenach et al. (1989) measured the $\delta^{15}\text{N}$ of leaves as -2.6 ± 0.6 in *Alnus* grown with atmospheric N_2 as the sole source of N, which represents the B value and is the measure of isotopic fractionation during N_2 -fixation. In *Alnus*, we observed $\delta^{15}\text{N}$ values of -0.74 (se 0.16) and -0.53‰ (se 0.13) respectively under ambient CO_2 and FACE. Based on the sole soil N versus sole N_2 source, and the observed $\delta^{15}\text{N}$ values in *Alnus*, we estimated the fraction of N uptake in *Alnus* through N_2 -fixation (f_n) to be 0.61 (se 0.02) under ambient and 0.60 (se 0.02) under elevated CO_2 .

3.5 Soil organic matter fractionation

Averaged over the species, the coarse POM C fraction was larger under FACE than under ambient CO_2 , 877 and 732 g C m^{-2} , respectively, however this effect was not significant ($P=0.356$, Table 3). The coarse POM N fractions were about equal under ambient CO_2 and FACE ($P=0.928$). Neither coarse POM C or N fractions were affected by species ($P=0.230$ and $P=0.067$).

Similarly to coarse POM, the fine POM C fraction was larger under FACE than under ambient CO_2 , 193 and 144 g C m^{-2} respectively and this effect was also not significant ($P=0.138$). However, the fine POM N fraction was significantly larger under FACE than under ambient CO_2 , i.e. 16 and 11 g N m^{-2} ($P=0.041$). The fine POM C and N fractions were not affected by species ($P=0.650$ and $P=0.950$).

The micro-aggregate protected POM C fraction was larger under ambient CO_2 than under FACE, i.e. 547 and 449 g C m^{-2} ($P=0.200$). The micro-aggregate protected POM N fraction was also larger under ambient CO_2 (40 and 33 g C m^{-2} ; $P=0.314$), however, again not significantly. Just like the POM fractions, the micro-aggregate protected POM C and N fractions were not affected by species either ($P=0.564$ and $P=0.244$).

4163

4 Discussion

4.1 Change of soil C

The increase of total C in the top 10 cm of the mineral soil during the four year experiment was about equal under ambient CO_2 and FACE, reaching 530 and 555 g C m^{-2} , respectively, which makes the expected additional C sink under elevated CO_2 negligible and, in this experiment, insignificant. Similarly, we did not observe any species effect on soil C, which may in part have been obscured by wind redistribution of above ground litter between the patches of different tree species. In 2007, cross contamination with leaves from other species, based on litterfall data published elsewhere, was about 24–27% of the total litterfall within alder and birch single species patches. The beech leaves remained on branches until the following spring and then slowly shed. The observed increase of soil C, irrespective of treatment, is therefore due to afforestation of the former agricultural fields. Based on the $\delta^{13}\text{C}$ data we estimated the average input of new soil C into the FACE plots to be 494 (se 64) g C m^{-2} between October of 2004 and 2007. This input seems to relate well to the average increase of total soil C in the FACE plots over the same period, i.e. 486 g C m^{-2} (Fig. 1a).

Carbon storage in litter and soil has been assessed at several other forest FACE experiments. For instance, at the Duke Forest and POP-EuroFACE experiments a significant additional C sink was created in the litter layer after six years of elevated CO_2 treatment (Table 4) (Lichter et al., 2005; Hoosbeek and Scarascia-Mugnozza, 2009). However, in Duke Forest the stimulation of organic matter accumulation by FACE ceased after the sixth year, resulting in an average additional C sink of $\sim 30 \text{ g C m}^{-2} \text{ yr}^{-1}$ over the nine year experiment (Lichter et al., 2008). At both sites, characterized by negligible bioturbation, the increase of C in the mineral soil depended solely on C input from roots and on downward leaching of DOC from the litter layers. In these forests it was not enhanced by FACE, i.e. no significant additional C sink was created in the mineral soil. At the Oak Ridge FACE experiment, most of the above ground litter was

4164

incorporated into the mineral soil by bioturbation. This resulted, in combination with C input from root turnover, in a significant additional C sink in the top 5 cm of the mineral soil (Jastrow et al., 2005). At the 0–15 cm increment, however, this FACE effect on soil C was no longer significant. Hoosbeek and Scarascia-Mugnozza (2009) hypothesized that the mixing of above ground litter into the mineral soil by bioturbation may have facilitated the FACE effect at the 0–5 cm increment. At the BangorFACE experiment, we also observed bioturbation which resulted in the “concentration” of above and below ground litter inputs at the top of mineral soil. However, at the BangorFACE site this “concentration effect” did not amplify a possible FACE effect on soil C storage.

4.2 Soil organic matter stabilization

In order to evaluate the effect of elevated CO₂ on soil C dynamics it is not sufficient to only look at changes in total C content, but it is also necessary to assess possible soil C stabilization mechanisms. The stability of SOM is controlled by the chemical structure of the organic matter and the existence of protection offered by the soil matrix and minerals (Baldock and Skjemstad, 2000; Krull et al., 2003; Davidson and Janssens, 2006). Oades (1993) suggested a model of aggregate formation in which micro-aggregates (~100 μm in diameter) are formed within macro-aggregates (> 250 μm in diameter). Fresh litter entering the soil forms sites for microbial activity and nucleation centers for aggregation (Six et al., 2002). This fraction is, in the conceptual model of aggregate formation, represented by coarse POM (> 250 μm). We observed that C and N contents of this coarse POM fraction were not significantly larger under FACE, meaning that the first phase towards SOM stabilization was not significantly enhanced under FACE.

As the organic matter enclosed in the macro-aggregates is decomposed, fine POM and micro-aggregates (53–250 μm) are formed. The fine POM C fraction was not affected by CO₂ treatment or species, implying that the next step towards stabilization was also unchanged. However, the fine POM N fraction was significantly larger under

4165

FACE, suggesting increased microbial activity and N-immobilization. At the smallest scale, the micro-aggregate protected C and N fractions were also not affected by CO₂ treatment and species.

The degree of soil C stabilization was found to vary among FACE experiments with trees. At Duke Forest, the increase of soil C due to forest regrowth occurred entirely within the free light fraction, while the iPOM and mineral associated fractions were not affected by FACE (Lichter et al., 2005). No additional soil C protection and stabilization took place. At Oak Ridge, the protection and stabilization processes in the soil kept up with the extra C input under FACE, i.e. the additional C input due to FACE was protected at the same rate as under ambient CO₂ (Jastrow et al., 2005). At POP-EuroFACE, iPOM and mineral associated C and N fractions increased in macro-aggregates and in newly formed micro-aggregates which indicates that protection and stabilization processes increased due to FACE (Hoosbeek and Scarascia-Mugnozza, 2009). However, at BangorFACE, we observed no FACE or species effect on soil C stabilization mechanisms, leading us to conclude that soil C stabilization processes were not affected by CO₂ treatment or by species.

4.3 Soil N and N uptake

Total soil N was not affected by CO₂ treatment or species and nor was the C/N ratio. However, the interannual variation of soil C and N did not follow the same pattern. In 2004 and 2005 the NPP was still relatively low, but increased in 2006. As a result, in 2006 soil C increased both under ambient CO₂ and FACE, however, the increase under FACE was smaller. At the same time, soil N did not change under ambient CO₂ while there was a decrease under FACE. Based on a higher NPP under FACE, we expected a reverse scenario, i.e. a larger increase of soil C and N under FACE. However, the extra NPP under FACE could have resulted in a larger availability of labile substrate in the soil, which may have caused a priming effect (Hoosbeek et al., 2004). This priming effect may have stimulated the decomposition of older SOM with a relatively low C/N ratio, which may explain the observed limited increase of soil C and a decrease of

4166

N in 2006 under FACE. Usually, a priming effect is a temporal phenomenon, as was observed by Hoosbeek and Scarascia-Mugnozza (2009).

Afforestation of our experimental fields has increased the litter input in comparison with previous crops, resulting in an increase of $\text{NH}_4\text{-N}$ in the soil. This increase was significantly smaller under FACE between 2006–2008, probably due to the combination of increased nitrification, microbial activity or plant uptake under FACE. Soil $\text{NO}_3\text{-N}$ increased after 2005 which may also have been due to the increased turnover of SOM, while larger uptake under FACE may explain lower $\text{NO}_3\text{-N}$. The N concentration in *Alnus* and *Betula* leaves was lower under FACE than under ambient CO_2 , which means that the demand for more N in order to sustain higher NPP under FACE was at least in part met by an increase of the N-use efficiency (NUE). A similar effect was observed by Calfapietra et al. (2007) in three *Populus* species in POP-EuroFACE in central Italy.

Since *Alnus* supports symbiotic N_2 -fixation, we hypothesized that it would be able to gain extra N by increasing the C supply to N_2 -fixing bacteria. We did not observe this effect, the N uptake ratio (N_2 -fixation/soil N) did not change in high CO_2 treatment. *Alnus* growing in elevated CO_2 did not use the extra available NPP (labile C) to increase symbiotic N_2 -fixation in order to meet the higher N demand under higher NPP. Instead, *Alnus* increased its NUE.

Finzi et al. (2007) pointed out that some combination of increased N uptake from the soil and more efficient use of the N already assimilated by trees is necessary to sustain the high rates of forest NPP under FACE. Based on a larger FACE data set including a wider variety of plants, Leakey et al. (2009) concluded that elevated CO_2 increases NUE. At Oak Ridge, the FACE induced soil C accrual was accompanied by a significant increase in soil N, i.e. FACE did not affect the C/N ratio of the mineral soil. Jastrow et al. (2005) postulated that FACE also affected N cycling by some combination of reducing N losses, stimulation of N fixation and increasing N uptake through greater root exploration. During the first three years of the POP-EuroFACE experiment, the increase of NUE was the major mechanism sustaining increased NPP under FACE (Calfapietra et al., 2007). As mentioned before, leaf N content of *Alnus* and *Betula*

4167

decreased under FACE while the soil C/N ratio was not affected by FACE. From this we infer that at Bangor the major mechanism to sustain increased NPP under FACE is also based on increased NUE. This fits well with the conclusions of (Finzi et al., 2007) (Table 4) stating that on the one hand, at sites with N-limited growth, i.e. Duke Forest and Oak Ridge, trees increase N uptake from the soil supporting greater NPP, while on the other hand, at sites without N-limitation, i.e. POP-EuroFACE and BangorFACE established on former agricultural soils, increased N-use efficiency seems to be the major mechanism sustaining increased NPP under FACE.

Acknowledgement. The development of the Bangor FACE site was funded by SRIF. We thank the Forestry Commission Wales and CIRRE for support of the running costs.

References

- Amarger, N., Mariotti, A., Mariotti, F., Durr, J. C., Bourguignon, C., and Lagacherie, B.: Estimate of symbiotically fixed nitrogen in field grown soybeans using variations in ^{15}N natural abundance, *Plant Soil*, 52, 269–280, 1979.
- Arnone III, J. A. and Gordon, J. C.: Effect of nodulation, nitrogen fixation and CO_2 enrichment on the physiology, growth and dry mass allocation of seedlings of *Alnus rubra* Bong., *New Phytol.*, 116, 55–66, 1990.
- Baldock, J. A. and Skjemstad, J. O.: Role of the soil matrix and minerals in protecting natural organic materials against biological attack, *Org. Geochem.*, 31, 697–710, 2000.
- Balesdent, J., Wagner, G. H., and Mariotti, A.: Soil organic matter turnover in long-term field experiments as revealed by carbon-13 natural abundance, *Soil Sci. Soc. A. J.*, 52, 118–124, 1988.
- Buurman, P., Van Lagen, B., and Velthorst, E. J.: *Manual for Soil and Water Analysis*, Backhuys Publishers, Leiden, The Netherlands, 1996.
- Cadish, G., Hairiah, K., and Giller, K. E.: Applicability of the natural ^{15}N abundance technique to measure N_2 fixation in *Arachis hypogaea* grown on an Ultisol, *Neth. J. Agr. Sci.*, 48, 31–45, 2000.
- Calfapietra, C., Gielen, B., Galema, A. N. J., Lukac, M., De Angelis, P., Moscatelli, M. C.,

4168

- Vitousek, P. M., Cassman, K. G., Cleveland, C., Crews, T., Field, C. B., Grimm, N. B., Howarth, R. W., Marino, R., Martinelli, L., Rastetter, E. B., and Sprent, J. I.: Towards an ecological understanding of biological nitrogen fixation, *Biogeochemistry*, 57/58, 1–45, 2002.
- Vogel, C. S., Curtis, P. S., and Thomas, R. B.: Growth and nitrogen accretion of dinitrogen-fixing *Alnus glutinosa* (L.) Gaertn. under elevated carbon dioxide, *Plant Ecol.*, 130, 63–70, 1997.
- Von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., and Flessa, H.: Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions – a review, *Eur. J. Soil Sci.*, 57, 426–445, 2006.

4173

Table 1. Initial soil conditions at the BangorFACE experimental site.

Plot	CO ₂ trmt	Clay %	Silt %	Sand %	pH (KCl)		NH ₄ (g N m ⁻²)		NO ₃ (g N m ⁻²)		C _{total} (g C m ⁻²)		N _{total} (g N m ⁻²)	
					mean	s.e.	mean	s.e.	mean	s.e.	mean	s.e.	mean	s.e.
1	ambient	10	33	57	5.0	0.3	0.09	0.00	0.74	0.07	2753	40	242	6
2	ambient	8	26	67	5.1	0.4	0.10	0.01	0.92	0.09	2569	49	214	8
3	FACE	9	27	64	4.6	0.2	0.10	0.01	0.92	0.12	3068	135	249	10
4	FACE	9	29	62	4.9	0.2	0.07	0.01	0.79	0.19	2898	236	237	27
5	FACE	9	26	65	4.4	0.1	0.08	0.01	0.77	0.44	2171	269	226	32
6	FACE	10	29	61	4.5	0.1	0.10	0.01	0.88	0.23	2786	120	276	12
7	ambient	9	25	66	4.4	0.0	0.08	0.01	0.64	0.06	2808	136	271	9
8	ambient	10	30	60	4.1	0.1	0.08	0.00	0.57	0.05	3191	74	304	8

4174

