

# $\Omega$ -results for Beurling's zeta function and lower bounds for the generalised Dirichlet divisor problem<sup>1</sup>

Titus W. Hilberdink

## Abstract

In this paper we study generalised prime systems for which the integer counting function  $N_{\mathcal{P}}(x)$  is asymptotically well-behaved, in the sense that  $N_{\mathcal{P}}(x) = \rho x + O(x^\beta)$ , where  $\rho$  is a positive constant and  $\beta < \frac{1}{2}$ . For such systems, the associated zeta function  $\zeta_{\mathcal{P}}(s)$  is holomorphic for  $\sigma = \Re s > \beta$ . We prove that for  $\beta < \sigma < \frac{1}{2}$ ,  $\int_0^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt = \Omega(T^{2-2\sigma-\varepsilon})$  for any  $\varepsilon > 0$ , and also for  $\varepsilon = 0$  for all such  $\sigma$  except possibly one value.

The Dirichlet divisor problem for generalised integers concerns the size of the error term in  $N_{k\mathcal{P}}(x) - \text{Res}_{s=1}(\zeta_{\mathcal{P}}(s)^k x^s/s)$ , which is  $O(x^\theta)$  for some  $\theta < 1$ . Letting  $\alpha_k$  denote the infimum of such  $\theta$ , we show that  $\alpha_k \geq \frac{1}{2} - \frac{1}{2k}$ .

*Keywords:* Beurling's generalised primes, Dirichlet divisor problem.

*AMS Mathematics subject classification 2010:* 11N80.

## 1. Introduction

A *generalised prime system* (or *g-prime system*)  $\mathcal{P}$  is a sequence of positive reals  $p_1, p_2, p_3, \dots$  satisfying

$$1 < p_1 \leq p_2 \leq \dots \leq p_n \leq \dots$$

and for which  $p_n \rightarrow \infty$  as  $n \rightarrow \infty$ . From these can be formed the system  $\mathcal{N}$  of *generalised integers* or *Beurling integers*; that is, the numbers of the form

$$p_1^{a_1} p_2^{a_2} \dots p_k^{a_k}$$

where  $k \in \mathbb{N}$  and  $a_1, \dots, a_k \in \mathbb{N}_0$ .<sup>2</sup> Such systems were first introduced by Beurling [2] and have been studied by many authors since then (see in particular [1]). Define the g-integer counting function  $N_{\mathcal{P}}(x)$  and the associated Beurling zeta function, respectively, by

$$N_{\mathcal{P}}(x) = \sum_{n \in \mathcal{N}, n \leq x} 1, \quad \zeta_{\mathcal{P}}(s) = \sum_{n \in \mathcal{N}} \frac{1}{n^s}.$$

(Here,  $\sum_{n \in \mathcal{N}}$  means a sum over all the g-integers, counting multiplicities.) In this paper, we shall be concerned with g-prime systems for which

$$N_{\mathcal{P}}(x) = \rho x + O(x^\beta), \tag{1.1}$$

for some  $\beta < \frac{1}{2}$  and  $\rho > 0$ . Then  $\zeta_{\mathcal{P}}(s)$  is defined and holomorphic for  $\Re s > 1$ , and has an analytic continuation to the half-plane  $\Re s > \beta$  except for a simple pole at  $s = 1$  with residue  $\rho$ . Furthermore,  $\zeta_{\mathcal{P}}(s)$  has *finite order* for  $\Re s > \beta$ ; i.e.  $\zeta_{\mathcal{P}}(\sigma + it) = O(|t|^\lambda)$  for some  $\lambda$  for  $\sigma > \beta$ . Let  $\mu_{\mathcal{P}}(\sigma)$  denote the infimum of all such  $\lambda$ . It is well-known that  $\mu_{\mathcal{P}}(\sigma)$  is non-negative,

<sup>1</sup>Journal of Number Theory **130** (2010) 707-715.

<sup>2</sup>Here,  $\mathbb{N} = \{1, 2, 3, \dots\}$ ,  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ , and  $\mathbb{P} = \{2, 3, 5, \dots\}$  — the set of primes.

decreasing, and convex (and hence continuous) (see, for example, [5]). For  $\mathcal{P} = \mathbb{P}$  (so that  $\mathcal{N} = \mathbb{N}$ ), the Lindelöf Hypothesis is the conjecture that  $\mu_{\mathbb{P}}(\sigma) = \mu_0(\sigma)$  for all  $\sigma$ , where

$$\mu_0(\sigma) = \begin{cases} \frac{1}{2} - \sigma & \text{if } \sigma < \frac{1}{2} \\ 0 & \text{if } \sigma \geq \frac{1}{2} \end{cases}.$$

In [4], it was proven that for all g-prime systems satisfying (1.1),  $\mu_{\mathcal{P}}(\sigma)$  must be *at least* as large as  $\mu_0(\sigma)$ : i.e.  $\mu_{\mathcal{P}}(\sigma) \geq \frac{1}{2} - \sigma$  for  $\sigma \in (\beta, \frac{1}{2})$ . In this paper we prove a stronger result by considering the mean square behaviour of  $\zeta_{\mathcal{P}}(\sigma + it)$ . For  $\sigma > \beta$ , define  $\nu_{\mathcal{P}}(\sigma)$  to be the infimum of numbers  $\lambda$  such that

$$\int_1^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt = O(T^{1+2\lambda}).$$

As in the case of  $\mu_{\mathcal{P}}(\sigma)$ ,  $\nu_{\mathcal{P}}(\sigma)$  is non-negative and convex decreasing (cf. [6], §7.8). Trivially,  $\nu_{\mathcal{P}}(\sigma) \leq \mu_{\mathcal{P}}(\sigma)$ . We show here that  $\nu_{\mathcal{P}}(\sigma) \geq \mu_0(\sigma)$ . In fact we prove slightly more.

### Theorem 1

Let  $\mathcal{P}$  be a g-prime system for which (1.1) holds for some  $\beta < \frac{1}{2}$  and  $\rho > 0$ . Then  $\nu_{\mathcal{P}}(\sigma) \geq \mu_0(\sigma)$  for  $\sigma \in (\beta, \frac{1}{2})$ . Furthermore,

$$\int_0^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt = o(T^{2-2\sigma}) \tag{1.2}$$

can hold for at most one value of  $\sigma$  in this range. In this case  $T^{2\sigma-2} \int_0^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt$  is unbounded for all other values of  $\sigma$ .

*Remark.* For  $\mathcal{P} = \mathbb{P}$ , we have  $\nu_{\mathcal{P}}(\sigma) = \mu_0(\sigma)$ , which shows the first part of Theorem 1 is best possible. However, in this case we have the asymptotic formula

$$\int_1^T |\zeta(\sigma + it)|^2 dt \sim \frac{\zeta(2-2\sigma)}{(2\pi)^{1-2\sigma}(2-2\sigma)} T^{2-2\sigma}$$

for  $0 < \sigma < \frac{1}{2}$ , showing that the exceptional value need not exist. In fact it seems unlikely an exceptional value exists and hence that  $\int_0^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt = \Omega(T^{2-2\sigma})$  for all  $\sigma \in (\beta, \frac{1}{2})$ , but we cannot quite show this. Furthermore it seems plausible that we should have  $\int_0^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt \geq C_{\sigma} T^{2-2\sigma}$  for some  $C_{\sigma} > 0$ .

## 2. Dirichlet divisor problems for g-primes

For a g-prime system satisfying (1.1) (with  $\beta < 1$ ), we can study the equivalent of the Dirichlet divisor problem concerning the error term in the asymptotic formula for the average of the ‘generalised divisor’ function. For  $k \in \mathbb{N}$ , let  $k\mathcal{P}$  denote the g-prime system obtained from  $\mathcal{P}$  by letting every g-prime from  $\mathcal{P}$  be counted  $k$  times. (If an original g-prime has multiplicity  $m$ , then in the new system it will have multiplicity  $km$ .) The Beurling zeta function of  $k\mathcal{P}$  is

$$\zeta_{k\mathcal{P}}(s) = \zeta_{\mathcal{P}}(s)^k.$$

By standard methods using Perron’s formula,

$$N_{k\mathcal{P}}(x) = \text{Res}_{s=1} \left\{ \frac{\zeta_{k\mathcal{P}}(s)^k}{s} x^s \right\} + \Delta_{\mathcal{P},k}(x) = xP_{k-1}(\log x) + \Delta_{\mathcal{P},k}(x),$$

where  $P_{k-1}(\cdot)$  is a polynomial of degree  $k-1$  and  $\Delta_{\mathcal{P},k}(x) = O(x^\theta)$  for some  $\theta < 1$ , depending on  $k$ . Let  $\alpha_k$  denote the infimum of such  $\theta$ . The *generalised Dirichlet divisor problem* is the problem of determining  $\alpha_k$ . Also let  $\beta_k$  denote the infimum of  $\phi$  for which

$$\int_0^x \Delta_{\mathcal{P},k}(y)^2 dy = O(x^{1+2\phi}).$$

Trivially,  $\beta_k \leq \alpha_k$ .

For  $\mathbb{P}$ , it is known that

$$\alpha_k \geq \beta_k \geq \frac{1}{2} - \frac{1}{2k} \quad (2.1)$$

and it is conjectured that there is equality throughout (actually  $\beta_k = \frac{1}{2} - \frac{1}{2k}$  for all  $k$  is equivalent to the Lindelöf Hypothesis — see [6], Theorem 13.4). We use Theorem 1 to show that (2.1) remains true for  $\mathcal{P}$  satisfying (1.1). In fact we have the following two corollaries:

### Corollary 2

Let  $\mathcal{P}$  satisfy (1.1) for some  $\beta < \frac{1}{2}$ . Then for  $\sigma \in (\beta, \frac{1}{2} - \frac{1}{2k})$ ,

$$\int_{-\infty}^{\infty} \frac{|\zeta_{\mathcal{P}}(\sigma + it)|^{2k}}{|\sigma + it|^2} dt \quad (2.2)$$

diverges. Further, if  $\frac{1}{2} - \frac{1}{2k}$  is not the exceptional value in (1.2), then the integral also diverges for  $\sigma = \frac{1}{2} - \frac{1}{2k}$ .

### Corollary 3

Let  $\mathcal{P}$  satisfy (1.1) for some  $\beta < \frac{1}{2}$ . With  $\alpha_k$  and  $\beta_k$  as above,  $\alpha_k \geq \beta_k \geq \max\{\beta, \frac{1}{2} - \frac{1}{2k}\}$ .

### 3. Proofs

*Proof of Theorem 1.* If  $\nu_{\mathcal{P}}(\sigma') < \frac{1}{2} - \sigma'$  for some  $\sigma' \in (\beta, \frac{1}{2})$  then, by continuity of  $\nu_{\mathcal{P}}(\cdot)$ ,  $\nu_{\mathcal{P}}(\sigma) < \frac{1}{2} - \sigma$  throughout some interval around  $\sigma'$  and (1.2) holds for all such  $\sigma$ ; in particular for two such values. We shall show that this is impossible.

Suppose, for a contradiction, that (1.2) holds for  $\sigma = \sigma_0, \sigma_1$  where  $\beta < \sigma_0 < \sigma_1 < \frac{1}{2}$ .

For  $N \geq 1$  let  $\zeta_{N,\mathcal{P}}(s) = \sum_{n \leq N} n^{-s}$ , where the sum ranges over  $n \in \mathcal{N}$ . As was stated in [4] (and shown in [3]), for  $\sigma < \frac{1}{2}$  there exist constants  $c_1, c_2 > 0$  such that for  $R \geq c_1 N$ ,

$$\sum_{r=1}^R \int_0^{2r-1} |\zeta_{N,\mathcal{P}}(\sigma + it)|^2 dt \geq c_2 R^2 N^{1-2\sigma}. \quad (3.1)$$

Also, writing  $s = \sigma + it$ , and following the arguments in [3], we have

$$\zeta_{N,\mathcal{P}}(s) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{\zeta_{\mathcal{P}}(s+w)N^w}{w} dw + O\left(\frac{N^c}{T(c+\sigma-1)}\right) + O\left(\frac{N^{1-\sigma}}{T} \sum_{\substack{\frac{N}{2} < n < 2N \\ n \in \mathcal{N}}} \frac{1}{|n-N|}\right), \quad (3.2)$$

for  $|t| < T$ ,  $c > 1 - \sigma$  and  $N \notin \mathcal{N}$ . We shall put  $c = 1 - \sigma + \frac{1}{\log N}$  and choose  $N$  in such a way that  $(N - \alpha, N + \alpha) \cap \mathcal{N} = \emptyset$ . (As was shown in [4], this is possible for arbitrarily large  $N$  if

$0 < \alpha < \frac{1}{4\rho}$ .) With this choice of  $N$ , the final sum in (3.2) was shown to be  $O(\sqrt{N})$ . As such (3.2) becomes

$$\zeta_{N,\mathcal{P}}(s) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{\zeta_{\mathcal{P}}(s+w)N^w}{w} dw + O\left(\frac{N^{\frac{3}{2}-\sigma}}{T}\right). \quad (3.3)$$

Now put  $\sigma = \sigma_1$  and push the contour in the integral to the left as far as  $\Re w = \sigma_0 - \sigma_1 < 0$ , picking up the residues at  $w = 0$  and  $w = 1 - s$  (since  $|t| < T$ ).

The contribution along the horizontal line  $[\sigma_0 - \sigma_1 + iT, c + iT]$  is, in modulus, less than

$$\frac{1}{2\pi T} \int_{\sigma_0 - \sigma_1}^c N^y |\zeta_{\mathcal{P}}(\sigma_1 + y + i(t+T))| dy.$$

Using the uniform bound  $|\zeta_{\mathcal{P}}(\sigma + it)| = O(t^{\frac{1-\sigma}{1-\beta}+\varepsilon})$ , this is at most a constant times

$$\frac{1}{T} \int_{\sigma_0 - \sigma_1}^{1-\sigma_1} T^{\frac{1-\sigma_1-y}{1-\beta}+\varepsilon} N^y dy + \frac{1}{T} \int_{1-\sigma_1}^{1-\sigma_1+\frac{1}{\log N}} T^\varepsilon N^y dy = O(T^{\frac{\beta-\sigma_0}{1-\beta}+\varepsilon} N^{\sigma_0-\sigma_1}) + O(T^{\varepsilon-1} N^{1-\sigma_1}). \quad (3.4)$$

Similarly on  $[\sigma_0 - \sigma_1 - iT, c - iT]$ .

The integral along  $\Re w = \sigma_0 - \sigma_1$  is at most

$$\begin{aligned} \frac{N^{\sigma_0-\sigma_1}}{2\pi} \int_{-T}^T \frac{|\zeta_{\mathcal{P}}(\sigma_0 + i(t+y))|}{\sqrt{(\sigma_1 - \sigma_0)^2 + y^2}} dy &= O\left(N^{\sigma_0-\sigma_1} \int_1^{2T} \frac{|\zeta_{\mathcal{P}}(\sigma_0 + iy)|}{y} dy\right) \\ &= o(N^{\sigma_0-\sigma_1} T^{\frac{1}{2}-\sigma_0}), \end{aligned} \quad (3.5)$$

using<sup>3</sup> the hypothetical bound  $\int_0^T |\zeta_{\mathcal{P}}(\sigma_0 + it)|^2 dt = o(T^{2-2\sigma_0})$ .

The residues at  $w = 0$  and  $w = 1 - s$  are, respectively,  $\zeta_{\mathcal{P}}(s)$  and  $\rho N^{1-s}/(1-s) = O(\frac{N^{1-\sigma_1}}{|t|+1})$ . Putting (3.3), (3.4), and (3.5) together gives

$$\zeta_{N,\mathcal{P}}(\sigma_1 + it) = \zeta_{\mathcal{P}}(\sigma_1 + it) + O\left(\frac{N^{1-\sigma_1}}{|t|+1}\right) + O(N^{1-\sigma_1} T^{\varepsilon-1}) + o(N^{\sigma_0-\sigma_1} T^{\frac{1}{2}-\sigma_0}) + O\left(\frac{N^{\frac{3}{2}-\sigma_1}}{T}\right),$$

for  $|t| < T$ . (Note that the first  $O$ -term in (3.4) is superfluous since  $\frac{\beta-\sigma_0}{1-\beta} < \frac{1}{2} - \sigma_0$ .) Hence, using  $(a+b+c+d+e)^2 \leq 5(a^2+b^2+c^2+d^2+e^2)$ , we have

$$|\zeta_{N,\mathcal{P}}(\sigma_1 + it)|^2 \leq 5|\zeta_{\mathcal{P}}(\sigma_1 + it)|^2 + O\left(\frac{N^{2-2\sigma_1}}{t^2+1}\right) + O(N^{2-2\sigma_1} T^{2\varepsilon-2}) + o(N^{2\sigma_0-2\sigma_1} T^{1-2\sigma_0}) + O\left(\frac{N^{3-2\sigma_1}}{T^2}\right).$$

Now apply  $\sum_{r=1}^R \int_0^{2r-1} \dots dt$  to both sides to give (for  $2R-1 < T$ )

$$\begin{aligned} \sum_{r=1}^R \int_0^{2r-1} |\zeta_{N,\mathcal{P}}(\sigma_1 + it)|^2 dt &= O\left(\sum_{r=1}^R \int_0^{2r-1} |\zeta_{\mathcal{P}}(\sigma_1 + it)|^2 dt\right) + O\left(\sum_{r=1}^R \int_0^{2r-1} \frac{N^{2-2\sigma_1}}{(t+1)^2} dt\right) \\ &\quad + O(R^2 N^{2-2\sigma_1} T^{2\varepsilon-2}) + O\left(\frac{R^2 N^{3-2\sigma_1}}{T^2}\right) + o(R^2 N^{2(\sigma_0-\sigma_1)} T^{1-2\sigma_0}) \\ &= o(R^{3-2\sigma_1}) + O(RN^{2-2\sigma_1}) + O(R^2 N^{2-2\sigma_1} T^{2\varepsilon-2}) + O\left(\frac{R^2 N^{3-2\sigma_1}}{T^2}\right) + o(R^2 N^{2(\sigma_0-\sigma_1)} T^{1-2\sigma_0}) \end{aligned}$$

---

<sup>3</sup>If  $f \geq 0$  and  $\int_0^T f^2 = o(T^\lambda)$  (some  $\lambda > 1$ ), then  $\int_{T/2}^T \frac{f(y)}{y} dy \leq \frac{2}{T} \int_0^T f \leq \frac{2}{T} \sqrt{T \int_0^T f^2} = o(T^{\frac{\lambda-1}{2}})$ , and  $\int_1^T \frac{f(y)}{y} dy = o(T^{\frac{\lambda-1}{2}})$  follows.

using (1.2) for  $\sigma_1$ . Let  $T = 2R$ . The left-hand side above is at least  $c_2 R^2 N^{1-2\sigma_1}$  by (3.1) if  $R \geq c_1 N$ . Dividing both sides through by  $R^2 N^{1-2\sigma_1}$  gives

$$c_2 \leq o\left(\left(\frac{R}{N}\right)^{1-2\sigma_1}\right) + O\left(\frac{N}{R}\right) + O(NR^{2\varepsilon-2}) + O\left(\frac{N^2}{R^2}\right) + o\left(\left(\frac{R}{N}\right)^{1-2\sigma_0}\right). \quad (3.6)$$

Put  $R = KN$  where  $K \geq c_1$  is a fixed, but arbitrary, constant. Letting  $N \rightarrow \infty$ , the  $o$ -terms both tend to zero as does the middle  $O$ -term. Hence

$$c_2 \leq \frac{A}{K} + \frac{B}{K^2}$$

for some absolute constants  $A, B$ . But  $K$  can be made arbitrarily large, so this gives a contradiction.

For the final part, suppose (1.2) holds for  $\sigma = \sigma_0$  say. If  $\int_0^T |\zeta_{\mathcal{P}}(\sigma' + it)|^2 dt = O(T^{2-2\sigma'})$  for some  $\sigma' \in (\beta, \frac{1}{2})$  with  $\sigma' \neq \sigma_0$ , then (1.2) actually holds for all  $\sigma$  between  $\sigma_0$  and  $\sigma'$ . (This follows from the Phragmen-Lindelöf Theorem for a strip (see [6], §7.8, with  $\varepsilon$  in the place of  $C$ )). This was shown to be impossible, and hence  $T^{2\sigma-2} \int_0^T |\zeta_{\mathcal{P}}(\sigma' + it)|^2 dt$  must be unbounded for all  $\sigma \neq \sigma_0$ . □

Now we apply Theorem 1 to find lower bounds in the Dirichlet divisor problem. Note that Theorem 1 actually shows that given  $\varepsilon > 0$ ,

$$\int_{T/2}^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt = \Omega(T^{2-2\sigma-\varepsilon}),$$

for if it was  $o(T^{2-2\sigma-\varepsilon})$ , then by telescoping it would follow that  $\int_0^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt = o(T^{2-2\sigma-\varepsilon})$  which is false.

*Proofs of Corollaries 2 and 3.* By Hölder's inequality,

$$\int_{T/2}^T |\zeta_{\mathcal{P}}(\sigma + it)|^{2k} dt \geq \frac{2^{k-1}}{T^{k-1}} \left( \int_{T/2}^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt \right)^k,$$

for every  $k \in \mathbb{N}$ . By Theorem 1, given  $\varepsilon > 0$ ,  $\int_{T/2}^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt \geq aT^{2-2\sigma-\varepsilon}$  for some  $a > 0$  and some arbitrarily large  $T$ . Hence for such  $T$ ,

$$\int_{T/2}^T |\zeta_{\mathcal{P}}(\sigma + it)|^{2k} dt \geq a^k T^{k(1-2\sigma)+1-\varepsilon k}.$$

It follows that

$$\int_{T/2}^T \frac{|\zeta_{\mathcal{P}}(\sigma + it)|^{2k}}{|\sigma + it|^2} dt \geq a' T^{k(1-2\sigma)-1-\varepsilon k}$$

for some  $a' > 0$ . But for  $\sigma < \frac{1}{2} - \frac{1}{2k}$ , we have  $k(1-2\sigma) - 1 > 0$ . Hence for  $\varepsilon$  sufficiently small,  $k(1-2\sigma) - 1 - \varepsilon k > 0$  also, and so  $\int_{T/2}^T \frac{|\zeta_{\mathcal{P}}(\sigma + it)|^{2k}}{|\sigma + it|^2} dt \not\rightarrow 0$  as  $T \rightarrow \infty$ , and Corollary 2 follows. Of course, if  $\frac{1}{2} - \frac{1}{2k}$  is not the exceptional value in Theorem 1, then we can take  $\varepsilon = 0$  in the above and the result also holds for  $\sigma = \frac{1}{2} - \frac{1}{2k}$ .

Let  $\gamma_k$  be the infimum of  $\sigma$  (with  $\sigma > \beta$ ) for which  $\int_{-\infty}^{\infty} \frac{|\zeta_{\mathcal{P}}(\sigma+it)|^{2k}}{|\sigma+it|^2} dt$  converges. By Corollary 2,  $\gamma_k \geq \frac{1}{2} - \frac{1}{2k}$ . An identical argument as in the  $\mathcal{P} = \mathbb{P}$  case (see [6], Theorem 12.5) shows that  $\gamma_k = \beta_k$ . (The argument is simply based upon Parseval's formula for Mellin transforms, which in this case is the identity

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{|\zeta_{\mathcal{P}}(\sigma+it)|^{2k}}{|\sigma+it|^2} dt = \int_0^{\infty} \frac{\Delta_{\mathcal{P},k}(x)^2}{x^{1+2\sigma}} dx$$

for  $\sigma$  in some interval  $(\theta, 1)$  with  $\theta < 1$ .) Hence  $\beta_k \geq \frac{1}{2} - \frac{1}{2k}$ . □

#### 4. On the line $\sigma = \frac{1}{2}$

In this article, we have considered the mean-value along vertical lines  $\Re s = \sigma$  with  $\sigma < \frac{1}{2}$ . This raises the question of what happens on the line  $\sigma = \frac{1}{2}$ . For  $\mathcal{P} = \mathbb{P}$ , we have  $\int_0^T |\zeta(\frac{1}{2} + it)|^2 dt \sim T \log T$ , so do we have  $\int_0^T |\zeta_{\mathcal{P}}(\frac{1}{2} + it)|^2 dt = \Omega(T \log T)$  in general? As in the  $\sigma < \frac{1}{2}$  case, we relate the behaviour of the mean-square value at  $\sigma = \frac{1}{2}$  to the behaviour of the mean-square for some  $\sigma = \sigma_0 < \frac{1}{2}$ .

#### Theorem 4

Let  $\mathcal{P}$  be a  $g$ -prime system for which (1.1) holds. If  $\int_1^T \frac{|\zeta_{\mathcal{P}}(\sigma+it)|}{t} dt = o((T \log T)^{\frac{1}{2}-\sigma})$  for some  $\sigma \in (\beta, \frac{1}{2})$ , then  $\int_0^T |\zeta_{\mathcal{P}}(\frac{1}{2} + it)|^2 dt = \Omega(T \log T)$ .

Note that the assumption is implied by  $\int_1^T |\zeta_{\mathcal{P}}(\sigma+it)|^2 dt = o(T^{2-2\sigma}(\log T)^{1-2\sigma})$ .

*Sketch of Proof.* We follow the proof of Theorem 1 as much as possible, this time taking  $\sigma_1 = \frac{1}{2}$ .

Using the argument in [3] for  $\sigma = \frac{1}{2}$ , (3.1) becomes: *there exist constants  $c_1, c_2 > 0$  such that for  $R \geq c_1 N / \log N$ ,*

$$\sum_{r=1}^R \int_0^{2r-1} \left| \zeta_{N,\mathcal{P}}\left(\frac{1}{2} + it\right) \right|^2 dt \geq c_2 R^2 \log N. \quad (4.1)$$

To see this, note that we have

$$\int_0^T \left| \zeta_{N,\mathcal{P}}\left(\frac{1}{2} + it\right) \right|^2 dt = T \sum_{n \leq N}^* \frac{1}{n} + 2 \sum_{n \leq N} \frac{1}{\sqrt{n}} \sum_{m < n} \frac{S_{m,n}(T)}{\sqrt{m}},$$

where  $S_{m,n}(T) = \frac{\sin(T \log(n/m))}{\log(n/m)}$ . (Here  $m, n \in \mathcal{N}$  and the  $*$  indicates that any multiplicities must be squared.) In any case, we have  $\sum_{n \leq N}^* \frac{1}{n} \geq \sum_{n \leq N} \frac{1}{n} \geq k_1 \log N$  for some  $k_1 > 0$ .<sup>4</sup> For  $m \leq \frac{n}{2}$ ,  $|S_{m,n}(T)| \leq 1/\log 2$ , so this part of the double sum is  $O(\sum_{n \leq N} \frac{1}{\sqrt{n}} \sum_{m \leq n/2} \frac{1}{\sqrt{m}}) = O(N)$ . Thus, for some positive constants  $k_1, k_2$ , independent of  $T$  and  $N$ ,

$$\int_0^T \left| \zeta_{N,\mathcal{P}}\left(\frac{1}{2} + it\right) \right|^2 dt \geq k_1 T \log N + 2 \sum_{n \leq N} \frac{1}{\sqrt{n}} \sum_{\frac{n}{2} < m < n} \frac{S_{m,n}(T)}{\sqrt{m}} - k_2 N.$$

Putting  $T = 2r - 1$  for  $r = 1, 2, \dots, R$ , and summing both sides gives, on noticing that  $\sum_{r=1}^R \sin((2r-1) \log \frac{n}{m}) = \frac{\sin^2(R \log n/m)}{\sin(\log n/m)} \geq 0$  since  $0 < \log n/m < \log 2$ ,

$$\sum_{r=1}^R \int_0^{2r-1} \left| \zeta_{N,\mathcal{P}}\left(\frac{1}{2} + it\right) \right|^2 dt \geq k_1 R^2 \log N - k_2 RN,$$

<sup>4</sup>This follows readily from  $N_{\mathcal{P}}(x) \sim \rho x$ .

and (4.1) follows.

In (3.2), we need a better estimate for the final sum. Let  $M \in \mathbb{N}$ . Then, with  $N$  such that  $(N - \alpha, N + \alpha) \cap \mathcal{N} = \emptyset$ ,

$$\begin{aligned} \sum_{\substack{\frac{N}{2} < n < 2N \\ n \in \mathcal{N}}} \frac{1}{|n - N|} &= \sum_{m=1}^M \sum_{\alpha N^{\frac{m-1}{M}} \leq |n-N| < \alpha N^{\frac{m}{M}}} \frac{1}{|n - N|} + O(1) \\ &\leq \frac{1}{\alpha} \sum_{m=1}^M \frac{1}{N^{\frac{m-1}{M}}} \left( N(N + \alpha N^{m/M}) - N(N - \alpha N^{m/M}) \right) + O(1) \\ &= O(N^{1/M}) + O(N^\beta), \end{aligned}$$

using (1.1). Since  $M$  is arbitrary, this is  $O(N^{\beta+\varepsilon})$  for every  $\varepsilon > 0$  in any case. Thus (3.3) becomes

$$\zeta_{N,\mathcal{P}}(s) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{\zeta_{\mathcal{P}}(s+w)N^w}{w} dw + O\left(\frac{N^{\frac{1}{2}+\beta+\varepsilon}}{T}\right).$$

The analysis up to (3.5) remains the same (with  $\sigma_0 = \sigma$  and  $\sigma_1 = \frac{1}{2}$ ) but in (3.5) we use the bound assumed in the statement to give  $o(N^{\sigma-\frac{1}{2}}(T \log T)^{\frac{1}{2}-\sigma})$ . The arguments following (3.5) remain valid and we put  $T = 2R$  again, but this time we divide through by  $R^2 \log N$ . On assuming  $\int_0^T |\zeta_{\mathcal{P}}(\frac{1}{2} + it)|^2 dt = o(T \log T)$ , (3.6) now becomes

$$c_2 \leq o\left(\frac{\log R}{\log N}\right) + O\left(\frac{N}{R \log N}\right) + O\left(\frac{NR^{2\varepsilon-2}}{\log N}\right) + O\left(\frac{N^{1+2\beta+2\varepsilon}}{R^2}\right) + o\left(\left(\frac{R \log R}{N}\right)^{1-2\sigma} \frac{1}{\log N}\right).$$

Put  $R = KN / \log N$  where  $K \geq c_1$  is a fixed, but arbitrary, constant. Letting  $N \rightarrow \infty$ , all the terms tend to zero except the first  $O$ -term. Hence

$$c_2 \leq \frac{A}{K}$$

for some absolute constant  $A$ . As  $K$  can be made arbitrarily large, this gives a contradiction. Hence  $\int_0^T |\zeta_{\mathcal{P}}(\frac{1}{2} + it)|^2 dt = \Omega(T \log T)$ . □

## References

- [1] P. T. Bateman and H. G. Diamond, Asymptotic distribution of Beurling's generalised prime numbers in: *Studies in Number Theory* 6, Prentice-Hall, 1969, pp. 152-212.
- [2] A. Beurling, Analyse de la loi asymptotique de la distribution des nombres premiers généralisés, I, *Acta Math.* **68** (1937), 255-291.
- [3] T. W. Hilberdink, Well-behaved Beurling primes and integers, *J. Number Theory* **112** (2005) 332-344.
- [4] T. W. Hilberdink, A lower bound for the Lindelöf function associated to generalised integers, *J. Number Theory* **122** (2007) 336-341.
- [5] E. C. Titchmarsh, *The Theory of Functions*, Second edition, Oxford University Press, 1986.
- [6] E. C. Titchmarsh, *The Theory of the Riemann Zeta-function*, Second edition, Oxford University Press, 1986.

Titus W. Hilberdink, Department of Mathematics, University of Reading, Whiteknights, PO  
Box 220, Reading RG6 6AX, UK.  
E-mail address: [t.w.hilberdink@reading.ac.uk](mailto:t.w.hilberdink@reading.ac.uk)