

A NOTE ON BESSEL'S INEQUALITY

S.S. DRAGOMIR

ABSTRACT. A monotonicity property of Bessel's inequality in inner product spaces is given.

1. INTRODUCTION

Let X be a linear space over the real or complex number field \mathbb{K} . A mapping $(\cdot, \cdot) : X \times X \rightarrow \mathbb{K}$ is said to be a *positive hermitian form* if the following conditions are satisfied:

- (i) $(\alpha x + \beta y, z) = \alpha(x, z) + \beta(y, z)$ for all $x, y, z \in X$ and $\alpha, \beta \in \mathbb{K}$;
- (ii) $(y, x) = \overline{(x, y)}$ for all $x, y \in X$;
- (iii) $(x, x) \geq 0$ for all $x \in X$.

If $\|x\| := (x, x)^{\frac{1}{2}}$, $x \in X$ denotes the semi-norm associated to this form and $(e_i)_{i \in I}$ is an orthonormal family of vectors in X , i.e., $(e_i, e_j) = \delta_{ij}$ ($i, j \in I$), then one has the following inequality [15]:

$$(1.1) \quad \|x\|^2 \geq \sum_{i \in I} |(x, e_i)|^2 \quad \text{for all } x \in X,$$

which is well known in the literature as Bessel's inequality.

Indeed, for every finite part H of I , one has:

$$\begin{aligned} 0 &\leq \left\| x - \sum_{i \in H} (x, e_i) e_i \right\|^2 = \left(x - \sum_{i \in H} (x, e_i) e_i, x - \sum_{j \in H} (x, e_j) e_j \right) \\ &= \|x\|^2 - \sum_{i \in H} |(x, e_i)|^2 - \sum_{j \in H} |(x, e_j)|^2 + \sum_{i, j \in H} (x, e_i) (e_j, x) \delta_{ij} \\ &= \|x\|^2 - \sum_{i \in H} |(x, e_i)|^2, \end{aligned}$$

for all $x \in X$, which proves the assertion.

The main aim of this paper is to improve this result as follows.

2. RESULTS

The following theorem holds.

Theorem 1. *Let X be a linear space and $(\cdot, \cdot)_2, (\cdot, \cdot)_1$ two hermitian forms on X such that $\|\cdot\|_2$ is greater than or equal to $\|\cdot\|_1$, i.e., $\|x\|_2 \geq \|x\|_1$ for all $x \in X$. Assume that $(e_i)_{i \in I}$ is an orthonormal family in $(X; (\cdot, \cdot)_2)$ and $(f_i)_{i \in J}$ is an*

Date: January 05, 2000.

1991 Mathematics Subject Classification. Primary 26D15; Secondary 46C99.

Key words and phrases. Bessel's inequality, Inner product spaces.

orthonormal family in $(X; (\cdot, \cdot)_1)$ such that for any $i \in I$ there exists a finite $K \subset J$ so that

$$(F) \quad e_i = \sum_{j \in K} \alpha_j f_j, \quad \alpha_j \in \mathbb{K} \quad (j \in K),$$

then one has the inequality:

$$(2.1) \quad \|x\|_2^2 - \sum_{i \in I} |(x, e_i)_2|^2 \geq \|x\|_1^2 - \sum_{j \in J} |(x, f_j)_1|^2 \geq 0,$$

for all $x \in X$.

In order to prove this fact, we require the following lemma.

Lemma 1. *Let X be a linear space endowed with a positive hermitian form (\cdot, \cdot) and $(g_k)_{k=1, \dots, n}$ be an orthonormal family in $(X; (\cdot, \cdot))$. Then*

$$(2.2) \quad \left\| x - \sum_{k=1}^n \lambda_k g_k \right\|^2 \geq \|x\|^2 - \sum_{k=1}^n |(x, g_k)|^2 \geq 0,$$

for all $\lambda_k \in \mathbb{K}$ and $x \in X$ ($k = 1, \dots, n$).

Proof. We will prove this fact by induction over “ n ”.

Suppose $n = 1$. Then we must prove that

$$\|x - \lambda_1 g_1\|^2 \geq \|x\|^2 - |(x, g_1)|^2, \quad x \in X, \quad \lambda_1 \in \mathbb{K}.$$

A simple computation shows that the above inequality is equivalent with

$$|\lambda_1|^2 - 2 \operatorname{Re}(x, \lambda_1 g_1) + |(x, g_1)|^2 \geq 0, \quad x \in X, \quad \lambda_1 \in \mathbb{K}.$$

Since $\operatorname{Re}(x, \lambda_1 g_1) \leq |(x, \lambda_1 g_1)|$, one has

$$\begin{aligned} |\lambda_1|^2 - 2 \operatorname{Re}(x, \lambda_1 g_1) + |(x, g_1)|^2 &\geq |\lambda_1|^2 - 2|\lambda_1| |(x, g_1)| + |(x, g_1)|^2 \\ &\geq (|\lambda_1| - |(x, g_1)|)^2 \geq 0 \end{aligned}$$

for all $\lambda_1 \in \mathbb{K}$ and $x \in X$, which proves the statement.

Now, assume that (2.2) is valid for “ $(n - 1)$ ”. Then we have:

$$\begin{aligned} &\left\| x - \sum_{k=1}^n \lambda_k g_k \right\|^2 \\ &= \left\| (x - \lambda_n g_n) - \sum_{k=1}^{n-1} \lambda_k g_k \right\|^2 \geq \|x - \lambda_n g_n\|^2 - \sum_{k=1}^{n-1} |(x - \lambda_n g_n, g_k)|^2 \\ &= \|x - \lambda_n g_n\|^2 - \sum_{k=1}^{n-1} |(x, g_k)|^2 \geq \|x\|^2 - |(x, g_n)|^2 - \sum_{k=1}^{n-1} |(x, g_k)|^2 \\ &= \|x\|^2 - \sum_{k=1}^n |(x, g_k)|^2, \end{aligned}$$

for all $\lambda_k \in \mathbb{K}$, $x \in X$ ($k = 1, \dots, n$), and the proof of the lemma is complete. \blacksquare

Proof. (Theorem) Let H be a finite part of I . Since $\|\cdot\|_2$ is greater than $\|\cdot\|_1$, we have:

$$\begin{aligned} \|x\|_2^2 - \sum_{i \in H} |(x, e_i)_2|^2 &= \left\| x - \sum_{i \in H} (x, e_i)_2 e_i \right\|_2^2 \\ &\geq \left\| x - \sum_{i \in H} (x, e_i)_2 e_i \right\|_1^2, \quad x \in X. \end{aligned}$$

Since, by (F), we may state that for any $i \in H$ there exists a finite $K \subset J$ with

$$e_i = \sum_{j \in K} (e_i, f_j)_1 f_j,$$

we have

$$\begin{aligned} \left\| x - \sum_{i \in H} (x, e_i)_2 e_i \right\|_1^2 &= \left\| x - \sum_{i \in H} (x, e_i)_2 \sum_{j \in K} (e_i, f_j)_1 f_j \right\|_1^2 \\ &= \left\| x - \sum_{j \in K} \left(\sum_{i \in H} (x, e_i)_2 (e_i, f_j)_1 \right) f_j \right\|_1^2 \end{aligned}$$

for all $x \in X$.

Applying the above lemma for $(\cdot, \cdot) = (\cdot, \cdot)_1$, $(g_k)_{k=\overline{1, n}} = (f_j)_{j \in K}$, we can conclude that

$$\left\| x - \sum_{j \in K} \lambda_j f_j \right\|_1^2 \geq \|x\|_1^2 - \sum_{j \in K} |(x, f_j)_1|^2, \quad x \in X,$$

where

$$\lambda_j = \left(\sum_{i \in H} (x, e_i)_2 (e_i, f_j)_1 \right) \in \mathbb{K} \quad (j \in K).$$

Consequently, we have:

$$\|x\|_2^2 - \sum_{i \in H} |(x, e_i)_2|^2 \geq \|x\|_1^2 - \sum_{j \in K} |(x, f_j)_1|^2 \geq \|x\|_1^2 - \sum_{j \in J} |(x, f_j)_1|^2$$

for all $x \in X$ and H a finite part of I , from where results (2.1).

The proof is thus completed. ■

Corollary 1. Let $\|\cdot\|_1, \|\cdot\|_2 : X \rightarrow \mathbb{R}_+$ be as above. Then for all $x, y \in X$, we have the inequality:

$$(2.3) \quad \|x\|_2^2 \|y\|_2^2 - |(x, y)_2|^2 \geq \|x\|_1^2 \|y\|_1^2 - |(x, y)_1|^2 \geq 0,$$

which is an improvement of the well known Cauchy-Schwartz inequality.

Proof. If $\|y\|_2 = 0$, then (2.3) holds with equality.

If $\|y\|_i \neq 0$, ($i = 1, 2$), then for $\{e_1\} = \left\{ \frac{y}{\|y\|_2} \right\}$, $\{f_1\} = \left\{ \frac{y}{\|y\|_1} \right\}$, the above theorem yields that

$$\frac{\|x\|_2^2 \|y\|_2^2 - |(x, y)_2|^2}{\|y\|_2^2} \geq \frac{\|x\|_1^2 \|y\|_1^2 - |(x, y)_1|^2}{\|y\|_1^2}$$

and since $\|y\|_2 \geq \|y\|_1$, the inequality (2.3) is obtained. ■

Remark 1. For a different proof of (2.3), see also [5].

Now, we will give some natural applications of the above theorem.

3. APPLICATIONS

- (1) Let $(X; (\cdot, \cdot))$ be an inner product space and $(e_i)_{i \in I}$ an orthonormal family in X . Assume that $A : X \rightarrow X$ is a linear operator such that $\|Ax\| \leq \|x\|$ for all $x \in X$ and $(Ae_i, Ae_j) = \delta_{ij}$ for all $i, j \in I$. Then one has the inequality

$$\|x\|^2 - \sum_{i \in I} |(x, e_i)|^2 \geq \|Ax\|^2 - \sum_{i \in I} |(Ax, Ae_i)|^2 \geq 0$$

for all $x \in X$.

The proof follows by the hermitian forms $(x, y)_2 = (x, y)$ and $(x, y)_1 = (Ax, Ay)$ for $x, y \in X$ and for the family $(f_i)_{i \in I} = (e_i)_{i \in I}$.

- (2) If $A : X \rightarrow X$ is such that $\|Ax\| \geq \|x\|$ for all $x \in X$, then, with the previous assumptions, we also have

$$0 \leq \|x\|^2 - \sum_{i \in I} |(x, e_i)|^2 \leq \|Ax\|^2 - \sum_{i \in I} |(Ax, Ae_i)|^2,$$

for all $x \in X$.

- (3) Suppose that $A : X \rightarrow X$ is a symmetric positive definite operator with $(Ax, x) \geq \|x\|^2$ for all $x \in X$. If $(e_i)_{i \in I}$ is an orthonormal family in X such that $(Ae_i, Ae_j) = \delta_{ij}$ for all $i, j \in I$, then one has the inequality

$$0 \leq \|x\|^2 - \sum_{i \in I} |(x, e_i)|^2 \leq (Ax, x) - \sum_{i \in I} |(Ax, e_i)|^2,$$

for all $x \in X$.

The proof follows from the above theorem for the choices $(x, y)_1 = (Ax, y)$ and $(x, y)_2 = (x, y)$, $x, y \in X$. We omit the details.

For other inequalities in inner product spaces, see the papers [1]-[14] and [7]-[6] where further references are given.

REFERENCES

- [1] S.S. DRAGOMIR, A refinement of Cauchy-Schwartz inequality, *G.M. Metod.* (Bucharest), **8**(1987), 94-95.
- [2] S.S. DRAGOMIR, Some refinements of Cauchy-Schwartz inequality, *ibid*, **10**(1989), 93-95.
- [3] S.S. DRAGOMIR and B. MOND, On the Boas-Bellman generalisation of Bessel's inequality in inner product spaces, *Italian J. of Pure and Appl. Math.*, **3** (1998), 29-38.
- [4] S.S. DRAGOMIR and B. MOND, On the superadditivity and monotonicity of Gram's inequality and related results, *Acta Math. Hungarica*, **71** (1-2) (1996), 75-90.
- [5] S.S. DRAGOMIR and B. MOND, On the superadditivity and monotonicity of Schwartz's inequality in inner product spaces, *Contributions Macedonian Acad. Sci. and Arts*, **15** (2) (1994), 5-22.
- [6] S.S. DRAGOMIR, B. MOND and Z. PALES, On a supermultiplicity property of Gram's determinant, *Aequationes Mathematicae*, **54** (1997), 199-204.
- [7] S.S. DRAGOMIR, B. MOND and J.E. PEČARIĆ, Some remarks on Bessel's inequality in inner product spaces, *Studia Univ. "Babes-Bolyai", Math.*, **37** (4) (1992), 77-86.
- [8] S.S. DRAGOMIR and J. SANDOR, On Bessel's and Gram's inequalities in prehilbertian spaces, *Periodica Math. Hungarica*, **29** (3) (1994), 197-205.

- [9] S.S. DRAGOMIR and J. SÁNDOR, Some inequalities in prehilbertian spaces, *Studia Univ. "Babeş-Bolyai"*, Mathematica, 1, **32**, (1987), 71-78.
- [10] W.N. EVERITT, Inequalities for Gram determinants, *Quart. J. Math.*, Oxford, Ser. (2), **8**(1957), 191-196.
- [11] T. FURUTA, An elementary proof of Hadamard theorem, *Math. Vesnik*, 8(**23**)(1971), 267-269.
- [12] C.F. METCALF, A Bessel-Schwartz inequality for Gramians and related bounds for determinants, *Ann. Math. Pura Appl.*, (4) **68**(1965), 201-232.
- [13] D.S. MITRINOVIĆ, *Analytic Inequalities*, Springer-Verlag, Berlin-Heidelberg and New York, 1970.
- [14] C.F. MOPPERT, On the Gram determinant, *Quart. J. Math.*, Oxford, Ser (2), **10** (1959), 161-164.
- [15] K. YOSHIDA, *Functional Analysis*, Springer-Verlag, Berlin, 1966.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, P.O. BOX 14428, MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA

E-mail address: `sever.dragomir@vu.edu.au`

URL: `http://rgmia.vu.edu.au/SSDragomirWeb.html`