

# A COMPLETE MONOTONICITY OF THE GAMMA FUNCTION

FENG QI AND CHAO-PING CHEN

ABSTRACT. The function  $\frac{1}{x} \ln \Gamma(x+1) - \ln x + 1$  is strictly completely monotonic on  $(0, \infty)$ .

The classical gamma function is usually defined for  $\operatorname{Re} z > 0$  by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt. \quad (1)$$

The psi or digamma function, the logarithmic derivative of the gamma function, and the polygamma functions can be expressed [6, p. 16] as

$$\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)} = -\gamma + \int_0^\infty \frac{e^{-t} - e^{-xt}}{1 - e^{-t}} dt, \quad (2)$$

$$\psi^{(m)}(x) = (-1)^{m+1} \int_0^\infty \frac{t^m}{1 - e^{-t}} e^{-xt} dt \quad (3)$$

for  $x > 0$  and  $m \in \mathbb{N}$ , where  $\gamma = 0.57721566490153286 \dots$  is the Euler-Mascheroni constant.

In 1985, D. Kershaw and A. Laforgia [5] showed that the function  $x[\Gamma(1 + \frac{1}{x})]^x$  is strictly increasing on  $(0, \infty)$ , which is equivalent to the function  $\frac{[\Gamma(x+1)]^{1/x}}{x}$  being strictly decreasing on  $(0, \infty)$ . In addition, it was proved that the function  $x^{1-\gamma}[\Gamma(1 + \frac{1}{x})]^x$  decreases for  $0 < x < 1$ , which is equivalent to  $\frac{[\Gamma(1+x)]^{\frac{1}{x}}}{x^{1-\gamma}}$  being increasing on  $(1, \infty)$ .

In [2, 10], it is proved that the function  $f(x) = \frac{[\Gamma(x+1)]^{1/x}}{x+1}$  is strictly decreasing and strictly logarithmically convex in  $(0, \infty)$  and the function  $g(x) = \frac{[\Gamma(x+1)]^{1/x}}{\sqrt{x+1}}$  is strictly increasing and strictly logarithmically concave in  $(0, \infty)$ . Some new proofs for the monotonicity of the function  $x^r[\Gamma(x+1)]^{1/x}$  on  $(0, \infty)$  are given for  $r \notin (0, 1)$ . In addition, if  $s$  is a positive real number, then for all real numbers  $x > 0$ ,

$$\frac{e^{-\gamma}}{[\Gamma(s+1)]^{1/s}} < \frac{[\Gamma(x+1)]^{1/x}}{[\Gamma(x+s+1)]^{1/(x+s)}} < 1, \quad (4)$$

$\lim_{x \rightarrow 0} f(x) = e^{-\gamma}$  and  $\lim_{x \rightarrow \infty} f(x) = e^{-1}$ .

Using monotonicity and inequalities of the generalized weighted mean values [1, 7, 8, 12], the first author proved [9] that the functions  $\left[\frac{\Gamma(s)}{\Gamma(r)}\right]^{1/(s-r)}$ ,  $\left[\frac{\Gamma(s,x)}{\Gamma(r,x)}\right]^{1/(s-r)}$

2000 *Mathematics Subject Classification.* 33B15.

*Key words and phrases.* Gamma function, psi function, completely monotonic function.

The authors were supported in part by NNSF (#10001016) of CHINA, SF for the Prominent Youth of Henan Province (#0112000200), SF of Henan Innovation Talents at Universities, Doctor Fund of Jiaozuo Institute of Technology, CHINA.

and  $\left[\frac{\gamma(s,x)}{\gamma(r,x)}\right]^{1/(s-r)}$  are increasing in  $r > 0$ ,  $s > 0$  and  $x > 0$ . For any given  $x > 0$ , the function  $\frac{s\gamma(s,x)}{x^s}$  is decreasing in  $s > 0$ .

In [3], N. Elezović, C. Giordana and J. Pečarić, among others, verified the convexity with respect to variable  $x$  of the function  $\left[\frac{\Gamma(x+t)}{\Gamma(x+s)}\right]^{1/(t-s)}$  for  $|t-s| < 1$ .

Recall that a function  $f$  is said to be completely monotonic on an interval  $I$  if  $f$  has derivatives of all orders on  $I$  which alternate successively in sign, that is

$$(-1)^n f^n(x) \geq 0 \quad (5)$$

for  $x \in I$  and  $n \geq 0$ . If inequality (5) is strict for all  $x \in I$  and for all  $n \geq 0$ , then  $f$  is said to be strictly completely monotonic. See [11] and references therein.

In this short note, we are about to prove a complete monotonicity result of a function involving the gamma function.

**Theorem 1.** *The function  $\frac{1}{x} \ln \Gamma(x+1) - \ln x + 1$  is strictly completely monotonic on  $(0, \infty)$  and tends to  $\infty$  as  $x \rightarrow 0$  and to 0 as  $x \rightarrow \infty$ .*

*Proof.* It has been shown in [5] that the function  $\frac{[\Gamma(x+1)]^{1/x}}{x}$  is strictly decreasing on  $(0, \infty)$ , then  $f(x) = \frac{1}{x} \ln \Gamma(x+1) - \ln x + 1$  is strictly decreasing on  $(0, \infty)$ . From the asymptotic expansion in [4]:

$$\ln \Gamma(x) = \left(x - \frac{1}{2}\right) \ln x - x + \ln \sqrt{2\pi} + \frac{1}{12x} + O(x^{-3}) \quad \text{as } x \rightarrow \infty,$$

we conclude that  $\lim_{x \rightarrow \infty} f(x) = 0$  and  $\lim_{x \rightarrow 0} f(x) = \infty$ . This implies  $f(x) > 0$  for  $x > 0$ .

Using Leibniz' rule

$$[u(x)v(x)]^{(n)} = \sum_{k=0}^n \binom{n}{k} u^{(k)}(x)v^{(n-k)}(x) \quad (6)$$

we obtain

$$\begin{aligned} f^{(n)}(x) &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{x}\right)^{(n-k)} [\ln \Gamma(x+1)]^{(k)} - \frac{(-1)^{n-1}(n-1)!}{x^n} \\ &= \left(\frac{1}{x}\right)^{(n)} \ln \Gamma(x+1) + \sum_{k=1}^n \binom{n}{k} \left(\frac{1}{x}\right)^{(n-k)} \psi^{(k-1)}(x+1) + \frac{(-1)^n n!}{nx^n} \\ &= \frac{(-1)^n n!}{x^{n+1}} \ln \Gamma(x+1) + \sum_{k=1}^n \frac{n!}{k!} \frac{(-1)^{n-k}}{x^{n-k+1}} \psi^{(k-1)}(x+1) + \frac{(-1)^n n!}{nx^n} \\ &\triangleq (-1)^n \frac{n!}{x^{n+1}} g(x), \end{aligned} \quad (7)$$

and

$$g'(x) = \frac{(-1)^n}{n!} x^n \psi^{(n)}(x+1) + \frac{1}{n}. \quad (8)$$

Using (3) and  $\frac{(n-1)!}{x^n} = \int_0^\infty t^{n-1} e^{-xt} dt$  for  $x > 0$  and  $n \in \mathbb{N}$ , we conclude

$$\frac{1}{x^n} g'(x) = \frac{1}{n!} \int_0^\infty \left(1 - \frac{t}{e^t - 1}\right) t^{n-1} e^{-xt} dt > 0, \quad (9)$$

since  $0 < \frac{t}{e^t-1} < 1$  for  $x > 0$ . Thus, the function  $g$  is strictly increasing and  $g(x) > g(0) = 0$  on  $(0, \infty)$ , which implies  $(-1)^n f^{(n)}(x) > 0$  for  $x > 0$  and  $n = 0, 1, 2, \dots$ . The proof is complete.  $\square$

*Remark 1.* After this paper was finalized, a similar result in [13] was found: The function  $1 + \frac{1}{x} \ln \Gamma(x+1) - \ln(x+1)$  is strictly completely monotone in  $(-1, \infty)$  and tends to 1 as  $x \rightarrow -1$  and to 0 as  $x \rightarrow \infty$ . This property is derived from the following integral representation:

$$\ln \Gamma(x+1) = x \ln(x+1) - x + \int_0^\infty \left( \frac{1}{t} - \frac{1}{e^t-1} \right) e^{-t} \frac{1-e^{-xt}}{t} dt. \quad (10)$$

## REFERENCES

- [1] B.-N. Guo and F. Qi, *Inequalities for generalized weighted mean values of convex function*, Math. Inequal. Appl. **4** (2001), no. 2, 195–202. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 11, 1059–1065. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [2] Ch.-P. Chen and F. Qi, *Monotonicity results for the gamma function*, J. Inequal. Pure Appl. Math. **3** (2003), no. 2, Art. 44. Available online at [http://jipam.vu.edu.au/v4n2/065\\_02.html](http://jipam.vu.edu.au/v4n2/065_02.html). RGMIA Res. Rep. Coll. **5** (2002), suppl., Art. 16. Available online at [http://rgmia.vu.edu.au/v5\(E\).html](http://rgmia.vu.edu.au/v5(E).html).
- [3] N. Elezović, C. Giordano and J. Pečarić, *The best bounds in Gautschi's inequality*, Math. Inequal. Appl. **3** (2000), 239–252.
- [4] C. L. Frenzen, *Error bounds for asymptotic expansions of the ratio of two gamma functions*, SIAM J. Math. **18** (1987), 890–896.
- [5] D. Kershaw and A. Laforgia, *Monotonicity results for the gamma function*, Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **119** (1985), 127–133.
- [6] W. Magnus, F. Oberhettinger, and R. P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer, Berlin, 1966.
- [7] F. Qi, *Generalized abstracted mean values*, J. Inequal. Pure Appl. Math. **1** (2000), no. 1, Art. 4. Available online at [http://jipam.vu.edu.au/v1n1/013\\_99.html](http://jipam.vu.edu.au/v1n1/013_99.html). RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 4, 633–642. Available online at <http://rgmia.vu.edu.au/v2n5.html>.
- [8] F. Qi, *Generalized weighted mean values with two parameters*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **454** (1998), no. 1978, 2723–2732.
- [9] F. Qi, *Monotonicity results and inequalities for the gamma and incomplete gamma functions*, Math. Inequal. Appl. **5** (2002), no. 1, 61–67. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 7, 1027–1034. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [10] F. Qi and Ch.-P. Chen, *Monotonicity and convexity results for functions involving the gamma function*, RGMIA Res. Rep. Coll. **6** (2003), no. 4, Art. 10. Available online at <http://rgmia.vu.edu.au/v6n4.html>.
- [11] F. Qi and S.-L. Xu, *The function  $(b^x - a^x)/x$ : Inequalities and properties*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3355–3359.
- [12] F. Qi and Sh.-Q. Zhang, *Note on monotonicity of generalized weighted mean values*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **455** (1999), no. 1989, 3259–3260.
- [13] H. Vogt and J. Voigt, *A monotonicity property of the  $\Gamma$ -function*, J. Inequal. Pure Appl. Math. **3** (2002), no. 5, Art. 73. Available online at [http://jipam.vu.edu.au/v3n5/007\\_01.html](http://jipam.vu.edu.au/v3n5/007_01.html).

(F. Qi) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

*E-mail address:* [qifeng@jzit.edu.cn](mailto:qifeng@jzit.edu.cn), [fengqi618@member.ams.org](mailto:fengqi618@member.ams.org)

*URL:* <http://rgmia.vu.edu.au/qi.html>

(Ch.-P. Chen) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

*E-mail address:* [chenciaoping@sohu.com](mailto:chenciaoping@sohu.com)