

A Note on Short Intervals Containing Primes

Mehdi Hassani
Narges Sariolgalam

Department of Mathematics
Institute for Advanced Studies in Basic Sciences
Zanjan, Iran

mhassani@iasbs.ac.ir

s-ghalam@iasbs.ac.ir

Abstract

In 1999, P. Dusart showed that for $x \geq 3275$, there exists at least a prime number in the interval $(x, x(1 + \frac{1}{2\ln^2 x})]$ and in 2003, O. Ramaré and Y. Saouter showed that for $x \geq 10726905041$ there exists at least a prime number in the interval $(x(1 - \Delta^{-1}), x]$, in which $\Delta = 28314000$. In this note, we show that for $x \geq 1.17 \times 10^{1634}$, we can yield Ramaré-Saouter's result from Dusart's result.

2000 *Mathematics Subject Classification*: 11A41.

Keywords: Primes.

As usual, suppose \mathbb{P} be the set of all prime numbers. In 1999, P. Dusart [1] showed that for every $x \geq 3275$, we have

$$\mathbb{P} \cap \left(x, x \left(1 + \frac{1}{2\ln^2 x} \right) \right] \neq \phi.$$

In 2003, O. Ramaré and Y. Saouter [2] showed that for $x \geq 10726905041$, we have

$$\mathbb{P} \cap (x(1 - \Delta^{-1}), x] \neq \phi,$$

in which $\Delta = 28314000$. If $L_D(x)$ denotes the length of Dusart's interval, $(x, x(1 + \frac{1}{2\ln^2 x})]$, and $L_{RS}(x)$ denotes the length of Ramaré-Saouter's interval, $(x(1 - \Delta^{-1}), x]$, then clearly we have:

$$L_D(x) = \frac{x}{2\ln^2 x} = O\left(\frac{x}{\ln^2 x}\right),$$

and

$$L_{RS}(x) = \frac{x}{\Delta} = O(x).$$

Also, we observe that

$$\lim_{x \rightarrow \infty} \frac{L_D(x)}{L_{RS}(x)} = 0,$$

and this suggest that for sufficiently large values of x 's, Dusart's interval become shorter than Ramaré-Saouter's interval. In this research report, we show that for $x \geq 1.17 \times 10^{1634}$, we can yield Ramaré-Saouter's interval from Dusart's interval.

Let $d(x) = x(1 + \frac{1}{2\ln^2 x})$. For $x > 0$, $d'(x) > 0$; so, $d^{-1}(x)$, the inverse of the function $d(x)$, is well-define. According to Dusart, for $x \geq 3275$, $\mathbb{P} \cap (x, d(x)] \neq \phi$, and so for such x 's that $d^{-1}(x) \geq 3275$, we have

$$\mathbb{P} \cap (d^{-1}(x), x] \neq \phi.$$

Therefore, $\mathbb{P} \cap (d^{-1}(x), x] \neq \phi$ holds for $x \geq d(3275)$ or for $x \geq 3300$. Now, we search such x 's that $x(1 - \Delta^{-1}) \leq d^{-1}(x)$ or $d(x(1 - \Delta^{-1})) \leq x$ and this is equivalent to

$$x(1 - \Delta^{-1}) \left(1 + \frac{1}{2\ln^2(x(1 - \Delta^{-1}))} \right) \leq x,$$

and since $x > 0$, we yield that for $x \geq \frac{e\sqrt{\frac{\Delta-1}{2}}}{1-\frac{1}{\Delta}} \approx 1.167417545 \times 10^{1634}$, Dusart's interval yields Ramaré-Saouter's interval. This prove our claim at above.

We end this short note with a question about Dusart's interval:

Question. For every $x \in \mathbb{R}$, let

$$n(x) := \#\mathbb{P} \cap \left(x, x(1 + \frac{1}{2\ln^2 x}) \right].$$

Is there some elementary function $f(x)$ such that $n(x) \sim f(x)$, when $x \rightarrow \infty$?

More generally study of $n(x)$ is a nice subject.

Note. All computations in this note done by Maple software.

Acknowledgment. We would like to express our gratitude to G.A. Pirayesh for his nice computational comments.

References

- [1] P. Dusart, Inégalités explicites pour $\psi(X)$, $\theta(X)$, $\pi(X)$ et les nombres premiers, *C. R. Math. Acad. Sci. Soc. R. Can.* **21** (1999), no. 2, 53–59.
- [2] O. Ramaré and Y. Saouter, Short effective intervals containing primes, *Journal of Number Theory*, **98** (2003), 10-33.