



# Durham E-Theses

---

## *The structure and evolution of the Wessex Basin.*

Lake, S.D.

### How to cite:

---

Lake, S.D. (1985) *The structure and evolution of the Wessex Basin.*, Durham theses, Durham University.  
Available at Durham E-Theses Online: <http://etheses.dur.ac.uk/1215/>

### Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in Durham E-Theses
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full Durham E-Theses policy](#) for further details.

THE STRUCTURE AND EVOLUTION OF THE  
WESSEX BASIN

by

STUART DAVID LAKE B.Sc.

VOL II

A thesis submitted to the University of Durham  
for the degree of Doctor of Philosophy.

The copyright of this thesis rests with the author.  
No quotation from it should be published without  
his prior written consent and information derived  
from it should be acknowledged.

Department of Geological Sciences, December 1985.



15. APR. 1986



NORTH  
COOLES FARM


DEVIZES

BRUTON (10Km to west)


WINTERBORNE KINGSTON

WYTCH FARM

ARRETON (50Km to east)




KEY

-  Basin infill
  - TERTIARY SEDIMENTS
  - CRETACEOUS SEDIMENTS
  - JURASSIC SEDIMENTS
  - PERMO-TRIAS SEDIMENTS
-  UPPER CARBONIFEROUS TO PROTEROZOIC VOLCANICS AND SEDS
-  OLDER PRECAMBRIAN BASEMENT
-  MOHO (Max and Min depth)
-  FAULT
-  BOREHOLE CONTROL


Durham University NERC

SCALE

1:100,000



15  
20  
25  
30  
35  
40




WYTCH FARM      ARRETON (50Km to east)


SOUTH

<p>STUART LAKE</p> <p>Sources:-          CHADWICK et al 1983          COLTER &amp; HARVARD 1981          DINGWALL 1971          BAMFORD et al 1976          etc</p>	<p>STRUCTURAL PROFILE ACROSS THE          WESSEX BASIN</p>	 <p>40</p>
---	--	---










001 12W 001 00W 001 00W 000 36W 000 24W 000 12W 000 00E 000 12E 000 24E 000 36E 000 48E 001 00E 001 12E 001 24E 001 36E 001 48E

1123

50 00N

0000 00E

Scale  
1:250,000




# Top Penarth group structural map English Channel

INTERPRETATION BY S.D.LAKE  
based on G.S.I non-exclusive seismic survey

**KEY**

- |     |                                    |     |            |                                      |
|-----|------------------------------------|-----|------------|--------------------------------------|
| ▲   | Reverse faults                     | —◆— | Faults     | } Surface exposure (Geological maps) |
| —◆— | >50 Millisecs throw (TWT)          | —x— | Anticlines |                                      |
| —x— | <50 Millisecs throw (TWT)          | —◆— | Synclines  | } Penarth.G.                         |
| —   | Isochrons (100 Millisec intervals) | —x— | Anticlines |                                      |
| ⋯   | Outcrop of Penarth Group           | —x— | Synclines  |                                      |

© STUART LAKE  
1985 DURHAM UNIVERSITY



COMPOSITE BOUGUER  
GRAVITY ANOMALY  
MAP, WESSEX BASIN


1:500,000


REDRAWN FROM B.G.S SOURCES

- KEY**
- ISOGAL VALUES IN MILLIGAL LINES AT 5mGal INTERVALS
  - ANOMALY HIGH
  - ANOMALY LOW

- SOURCES**
- U.K. DATA
  - IGS 1:250,000 BOUGUER GRAVITY
  - ADDITIONAL UNPUBLISHED DATA
  - FRENCH DATA
  - B.R.G.M. 1:80,000 CARTES
  - GRAVIMÉTRIQUES DE LA FRANCE

International Gravity Formula 1967  
-5mGal applied to B.R.G.M. data to  
convert to I.G.F. 1967 and N.G.R.N. 1973






REFLECTORS (IN SECONDS)

- R1
- R2
- · - · - R3
- ..... R4
- R5
- · - · - R6

- ..... G.S.I NON EXCLUSIVE
- SEISMIC LINES
- · - · - R1 RELATED STRUCTURES
- R2 RELATED STRUCTURES
- · - · - R6 RELATED STRUCTURES
- ▨ 'FLAT' PORTION OF REFLECTOR





# PRE APTIAN SUBCROP MAP

- LITHOLOGICAL BOUNDARIES
- WESTWARD EROSIONAL LIMIT OF LOWER GREENSAND
- WESTWARD LIMIT OF GAULT
- FAULTS

H1	WEALDEN
G14	PURBECK
G13	PORTLAND
G12	KIMMERIDGE CLAY
G9-11	CORNBRASH TO CORALLIAN
G5-8	INFERIOR OOLITE TO FOREST MARBLE
G1-4	LOWER JURASSIC
F	TRIASSIC
E	PERMIAN
B	BASEMENT








LAKE 1985






North

South





**KEY**

- |   |  |                                      |
|---|--|--------------------------------------|
| TERTIARY FORMATIONS                     | PORTLAND, KIMMERIDGE, CORALLIAN          | LIAS                                 |
| CHALK                                   | CORALLIAN (part), OXFORD CLAY, KELLAWAYS | PENARTH GROUP, MERCIA MUDSTONE GROUP |
| UPPER GREENSAND, GAULT, LOWER GREENSAND | GREAT OOLITE GROUP                       | SHERWOOD SANDSTONE GROUP             |
| WEALDEN, PURBECK                        | INFERIOR OOLITE                          | PERMIAN                              |



- KEY**
- major growth faults
  - Aptian/Albian unconformity
  - unconformity
- LITHOGRAPHY**
- Absent
  - mudstone/shale/marl
  - sandstone/clay alternating
  - limestone
  - sandstone
  - limestone/clay
  - shale predominant - limestone local
  - siltstone
  - silty mudstone
  - non sequence
  - silty limestone
  - conglomerates & marginal deposits
  - halite
  - anhydrite
  - dolomite
  - possibly diachronous boundary
  - chert
  - iron ore





# LOCATION MAP

# Figure

## A LYME REGIS

## B BURTON BRADSTOCK

## C KIMMERIDGE




**KEY**


- 123 Locality number
- } Locality area
- A Areas enlarged to 1:50,000 (see insets)

1:250,000

1:633,600



GRID NORTH




SY SZ




### ADDITIONAL JURASSIC LOCALITIES


A LYME REGIS



B BURTON BRADSTOCK




C KIMMERIDGE



#### KEY

- x NO EXPOSURE
- o SMALL EXPOSURE
- UNVISITED
- NO SIGNIFICANT KINEMATIC INDICATORS


1:250,000




Scale

# SHEAR SENSE INDICATORS



A LYME REGIS



B BURTON BRADSTOCK



C KIMMERIDGE




## KEY

- LOCALITY
- DEXTRAL
- - - SINISTRAL
- ⋯ UNKNOWN OFFSET

1:250,000




GRID NORTH  
↑





### ADDITIONAL KINEMATIC INDICATORS




### KEY

- LOCALITY
- CALCITE VIENS
- - - PYRITE VIENS
- \* STROLITE
- (F) FAULT RELATED
- (S) SELENITE COVERED

1:250,000

GRID NORTH  
↑




WNW

solifluxion deposits

Loc A


Channels

-  INFERIOR OOLITE
-  FRIABLE SANDSTONE
-  CEMENTED SANDSTONE
-  QUATERNARY INFILL
-  FRACTURE SOLUTION PIPES
-  VEGETATION
-  JOINTS




11.55

ESE






STRUCTURES IN THE BRIDPORT SANDS. EAST CLIFF, BRIDPORT




S.D.LAKE


1985




# SECONDARY AND TERTIARY JOINTS




A LYME REGIS




B BURTON BRADSTOCK



C KIMMERIDGE



100 M




## KEY

- LOCALITY
- SECONDARY JOINT TRENDS
- - - TERTIARY JOINT TRENDS
- (C) CLAY INFILL
- (Ca) CALCITE INFILL


1:250,000

GRID NORTH




Scale


# JOINTS ALL ORIENTATIONS UNDEFINED





## A LYME REGIS



## B BURTON BRADSTOCK




## C KIMMERIDGE



**KEY**

- LOCALITY (Ca) CALCITE ON SURFACE
- JOINT TREND AND NUMBER (F) FISSURES

1:250,000



SS ST

ST 3 4 5


11A1

# BOREHOLES AND WELLS IN THE WESTERN WESSEX BASIN


1:250,000

### KEY

- DRYHOLE
- ☼ GEOTHERMAL
- OIL WELL
- ABANDONED OIL WELL
- OTHER WELLS, B.G.S
- OFFSHORE B.G.S WELLS
- ⊗ WELLS AND SPRINGS
- CONFIDENTIAL WELL



COMPILED BY S.D.LAKE 1985



Grid N



SX SY

SY SZ

SU SZ




# LANDSAT MSS lineament map path 216:row 24 path 217:row 24 1:250,000

BAND 7 (0.8-1.10 μm NEAR INFRA RED)  
WINTER IMAGERY  
MOSAIC COMPILED by G.S.L  
Interpretation by Stuart Lake

DURHAM UNIVERSITY NERC FEB 1984

**KEY**

- CLEAR LINEAMENTS
- - - - - SUBTLE LINEAMENTS
- ◊ LIMIT OF CLOUD COVER
- ANTICLINAL AXIS
- ◊ SU BURBAN LONDON



# LANDSAT TM lineament map path 202:row 24 1:250,000

BAND 5 (1.55-1.75  $\mu\text{m}$  NEAR INFRARED)


IMAGE TAKEN 4-2-83




Processed on Diad system by N.P.A


Interpretation by Stuart Lake

DURHAM UNIVERSITY N.E.R.C JAN 1984

### KEY

-  CLEAR LINEAMENTS
-  SUBTLE LINEAMENTS
-  ANTICLINAL AXIS
-  SU BURBAN LONDON







**TM LINEAMENT MAP**

--- Major lineament related to growth f.  
- - - Other lineaments


0 5 10 15 20 25 Km  
0 5 10 M



LOWER LIAS Source Mature Early Maturity




KIMMERIDGE Source Mature Early Maturity



SHERWOOD SANDSTONE Reservoir



BRIDPORT SANDSTONE Reservoir



GREAT OOLITE LIMESTONE FORMATION Reservoir

Reservoirs distribution Optimum reservoir location

# HYDROCARBON OCCURENCES IN THE WESSEX BASIN

- KEY**
- OIL WELL
  - GAS WELL
  - OIL AND GAS WELL
  - SIGNIFICANT OIL AND GAS SHOWS
  - ABANDONED DRY WELL
  - ◇ LOCATION OF WELLBORE IN WYTCHE FARM OILFIELD
  - OIL SEEPAGE
  - GAS SEEPAGE
  - ▨ SHERWOOD SANDSTONE RESERVOIR
  - ▨ BRIDPORT S/ST OIL WATER CONTACT
  - ▨ GREAT OOLITE LIMESTONE RESERVOIR
- |      |                                 |
|------|---------------------------------|
| AG   | ARNE G1                         |
| B    | BUSHEY FARM A12                 |
| E    | ENCOMBE 1                       |
| G1   | CHALDON DOWN G1                 |
| G2   | CHALDON DOWN 2                  |
| HR   | CHALDON HERRING 1,G2,G3         |
| K    | BROAD BENCH 1, KIMMERIDGE 1,2,3 |
| K5   | KIMMERIDGE NORTH K5             |
| LHN  | LANGTON HERRING NORTH 1         |
| LHS  | LANGTON HERRING 1               |
| O1-2 | OSMINGTON 1,2                   |
| P    | POXWELL 1,2,3                   |
| R    | RINGSTEAD 1                     |
| RA   | RADIPOLE 1                      |
| S    | STOBOROUGH 1,2                  |
| W    | WAREHAM 1,2,3                   |

1 : 625,000 Scale

VOLUME 2

- Fig.II.1.1 Western Wessex Basin, Solid Geology map, (1:250,000 scale).
- Fig.II.2.1 Previous north-south structural profile across the Wessex Basin, 1:100,000 scale based on data acquired prior to this study.
- Fig.II.2.2 Track lines of seismic reflection profiles based on G.S.I. non-exclusive seismic reflection survey and I.G.S. project 76/8 and 77/1 (part) used to compile a structural map of the Top Penarth Group in the English Channel.
- Fig.II.2.3 Top Penarth Group structural map of the English Channel. Interpretation by S.D. Lake. Based on G.S.I. non-exclusive seismic survey.
- Fig.II.2.4 Composite Bouguer Gravity anomaly map, Wessex Basin, 1:500,000 scale (Redrawn from B.G.S. Sources).
- Fig.II.2.5 Two-way time isochron map of dipping reflectors in Lyme Bay and associated active late Palaeozoic/Mesozoic faulting, (1:250,000 scale).
- Fig.II.3.1 Pre-Aptian subcrop map, (1:250,000 scale).
- Fig.II.3.2 Six serial north-south structural profiles across the Wessex Basin. Scale shown. Compiled from geological maps of the onshore British Isles by Whittaker, 1985.
- Fig.II.3.3 Chrono-lithostratigraphic column. Western Wessex Basin, S. England (Permian - Quaternary).
- Fig.II.5.1 Location map at 1:250,000 scale. Three insets of Lyme Bay, Bradstock and Kimmeridge at 1:50,000 scale.
- Fig.II.5.2 Additional Jurassic localities also visited during research.
- Fig.II.5.3 Shear sense indicators (1:250,000 scale). Three insets of Lyme Bay, Burton Bradstock and Kimmeridge at 1:50,000 scale. Dextral and sinistral movements distinguished (frequency also shown).
- Fig.II.5.4 Additional Kinematic indicators (1:250,000 scale). Three insets of the Lyme Regis, Burton Bradstock and Kimmeridge areas enlarged to 1:50,000 scale. The detailed orientation of calcite and pyrite veins and stylolites are also shown.

Fig.II.5.5 Structures in the Bridport Sands, East Cliff, Bridport.

Fig.II.5.6 Primary Joints 1:250,000 scale. Three insets of Lyme Bay, Bradstock and Kimmeridge at 1:50,000 scale. (Frequency also shown).

Fig.II.5.7 Secondary and Tertiary Joints (1:250,000 scale). Three insets of the Lyme Regis, Burton Bradstock and Kimmeridge areas enlarged to 1:50,000. Clay or Calcite infill along joint planes are also distinguished.

Fig.II.5.8 Joints all orientations undefined (1:250,000 scale). Three insets of the Lyme Regis, Burton Bradstock and Kimmeridge areas enlarged to 1:50,000 scale.

Fig.II.6.1 Hydrocarbon occurrences in the Wessex Basin (1:625,000 scale).

Inset 1. Detailed plan of the Wytch Farm Oilfield.

Inset 2. Detailed plan of the Humbly Grove Oilfield.

5 insets 1) Lower Lias Clay source rocks - showing areas of postulated maturity and early maturity.

2) Kimmeridge Clay Source rock - showing areas of postulated maturity and early maturity.

3) Sherwood Sandstone Reservoir - areas of reservoir distribution and postulated optimum reservoir location are also shown.

4) Bridport Sands reservoir - areas of reservoir distribution and postulated optimum reservoir location are also shown.

5) Great Oolite Limestone Formation reservoir - areas of reservoir distribution and postulated optimum reservoir location are also shown.

Fig.II.A.1 Boreholes and wells in the western Wessex Basin, at 1:250,000 Scale (including exploration license blocks).

Fig.II.A.2 Landsat MSS lineament map, path 216: row 24, path 217: row 24 (1:250,000 scale). Band 7 (0.8 - 1.10 m. Near Infra Red) Winter Imagery. Mosaic compiled by G.S.L. (Interpretation by S.D. Lake).

Fig.II.A.3 Landsat TM lineament map, path 202: row 24, 1:250,000 scale. Band 5 (1.55 - 1.75 m Near Infra Red). Image taken 4.2.83. Processed on Dial system by Nigel Press Associates. Interpretation by S.D. Lake.

Fig.II.A.4 TM lineament map, 1:250,000 scale of Dorset. Detail of the structures along the Dorset coast particularly well exemplified.