



## SEAFDEC/AQD Institutional Repository (SAIR)

Title	Impact assessment of cage culture in Lake Taal, Philippines.
Author(s)	Alcañices, Marilyn M.; Pagulayan, Roberto C.; Mamaril, Augustus C.
Citation	Alcañices, M.M., Pagulayan, R.C. and Mamaril, A.C. (2001). Impact assessment of cage culture in Lake Taal, Philippines ((Abstract only). In CB Santiago, ML Cuvin-Aralar and ZU Basiao (eds.). Conservation and Ecological Management of Philippine Lakes in Relation to Fisheries and Aquaculture (p.153). Southeast Asian Fisheries Development Center, Aquaculture Department, Iloilo, Philippines; Philippine Council for Aquatic and Marine Research and Development, Los Baños, Laguna, Philippines; and Bureau of Fisheries and Aquatic Resources, Quezon City, Philippines.
Issue Date	2001
URL	<a href="http://hdl.handle.net/10862/836">http://hdl.handle.net/10862/836</a>

This document is downloaded at: 2013-07-02 07:21:13 CST



# Impact Assessment of Cage Culture in Lake Taal, Philippines

Marilyn M. Alcañices, Roberto C. Pagulayan and Augustus C. Mamaril

Environmental Science Program, College of Science  
University of the Philippines,  
Diliman, Quezon City

Alcañices MM, Pagulayan RC and Mamaril AC. 2001. Impact assessment of cage culture in Lake Taal, Philippines, p.153. In CB Santiago, ML Cuvin-Aralar and ZU Basiao (eds.). Conservation and Ecological Management of Philippine Lakes in Relation to Fisheries and Aquaculture. Southeast Asian Fisheries Development Center, Aquaculture Department, Iloilo, Philippines; Philippine Council for Aquatic and Marine Research and Development, Los Baños, Laguna, Philippines; and Bureau of Fisheries and Aquatic Resources, Quezon City, Philippines. 187 pp. (Abstract only)

## Abstract

The environmental impact of cage culture on water quality of Lake Taal was assessed from March 1996 through February 1997. Three stations were considered namely: Balas, which serves as station 1 (non-cage area) and Sampaloc and Laurel, stations 2 and 3 (cage areas), respectively. Monthly water samples with two replicates were collected using a van Dorn sampler at 0, 5, 10 and 15-m depths in all stations. Below surface water from the inside of the cages was also collected. Water temperature, water transparency, pH, and conductivity were determined *in situ*. Dissolved oxygen, chloride,  $\text{NO}_3$ ,  $\text{NH}_3$ ,  $\text{PO}_4$ , and total P were analyzed in the laboratory. Phytoplankton density and algal biomass (through chlorophyll a) and primary productivity indices were determined with the light-and-dark bottle method.

Of the water quality parameters, conductivity and DO had significant differences between non-cage and cage areas. Conductivity gave significant difference ( $P < 0.01$ ) between control and cage area during the wet season. Highest conductivity value ( $2100 \mu\text{S/cm}$ ) was observed in station 3. Mean values of DO gave significant differences ( $P < 0.05$ ) in the different stations throughout the study period. A decrease of DO to  $2.5 \text{ mg/l}$  was observed below 10-m depth around the cage areas. Analysis indicates that cage culture leads to oxygen depletion in the water column. The presence of cage structures decreased the flow rate resulting to weak circulation. The reduced water circulation in effect decreased the supply of oxygen and removal of toxic waste metabolites from the vicinity of the fish farm, and reduced the supply of plankton. These results suggest that the impact of cage culture in Lake Taal is minor but can alter the lake ecosystem if not properly managed. Zoning and continuous water quality monitoring are needed.