

Among the postlarvae of *Penaeus*, *P. (Fenneropenaeus) indicus* was dominant followed by *P. (Penaeus) monodon*, *P. (P.) semisulcatus*, *P. (F.) merguensis* and *P. (Melicertus) latisulcatus*. In *Metapenaeus*, postlarvae of *M. monoceros* were abundant followed by *M. dobsoni*, *M. affinis*, *M. brevicornis* and *M. lysianassa*.

Two peaks were observed in the postlarval penaeid prawn population. In *P. (F.) indicus* and *P. (P.) monodon*, the primary peak occurred from January to April and the secondary peak from July to September. In *M. monoceros* and *M. dobsoni*, the primary peak was from March to May and the secondary peak from August to September. The postlarvae of *P. (F.) indicus*, *P. (P.) monodon*, *M. monoceros* and *M. dobsoni* were available throughout the year while the others were seasonal. The distribution of postlarvae in the estuary is related to the type of substratum, salinity and temperature. The postlarval population declined during the northeast monsoon (November-December) and in peak summer (May-June). Their abundance decreased in the lower salinity areas of the upper reaches of the estuary.

### Environmental Physiology of the Prawn *Penaeus (Melicertus) latisulcatus*

A. Ramasamy and Al. Paul Pandian

Centre of Advanced Study in Marine Biology  
Annamalai University  
Parangipettai 608 502, Tamil Nadu, India

There are a number of intrinsic and extrinsic factors which affect the normal routine activity of the prawn. The present study attempts to elucidate the optimum levels of various environmental factors for the culture of prawns.

The salinity tolerance capacity of *Penaeus (Melicertus) latisulcatus* was estimated in 13 different test salinities from 0 to 60 ppt (at 5 ppt increments). The prawns can tolerate a wide salinity range of 20-50 ppt. Maximum survival, however, was between 25 to 45 ppt. The extreme low (0-10 ppt) and high (60 ppt) salinities were highly lethal to the prawns. The change in acclimation temperature from 30 to 35°C increased the upper incipient lethal level from 38.5 to 39.5°C. The prawns acclimated to 30°C tolerated 42°C for 275 sec and 45.5°C for 13 sec, while prawns acclimated to 35°C tolerated 42°C for 505 sec and 46.5°C for 11 sec.

Prawns were acclimated to a salinity of 26 ppt and oxygen consumption was measured at 5, 15, 26, and 38 ppt in a continuous water-flow method. The total oxygen consumption showed an inverse relationship with weight. Oxygen consumption declined with increase in salinity. The resistance of prawns to hydrogen sulphide was tested in 18 different concentrations of sodium sulphide mixed with seawater. The prawns tolerated sodium sulphide concentrations up to 20 mg/l. The dissolved oxygen in the water was found to be reduced to very low levels with the increase in the concentration of sodium sulphide (from 5.9 ml O<sub>2</sub>/l to 0.54 ml O<sub>2</sub>/l). This may cause heavy mortality of the prawns.

### Molt Staging in Adult *Penaeus monodon*

Rosario Pudadera, Jose Llobrera,  
Rose Marie Caballero and Nieves Aquino

Aquaculture Department  
Southeast Asian Fisheries Development Center  
P.O. Box 256, Iloilo City, Philippines

Changes and formation of cuticular layers and setae bordering the uropods and endopodites of the pleopods of adult *Penaeus monodon* were examined under a light microscope. Observations and photographs were made at 0, 12 and 24 hours after molting and every 24 hours thereafter until second molting occurred. Results show that the internal structures of the setae and cuticle undergo marked changes throughout the molt cycle. It was possible to identify the molt stages A, B, C and D. Rapid examination of the molt stages allows the proper timing of eyestalk ablation to induce ovarian maturation.

### Effect of Temperature and Salinity on the Hatching of Eggs and Larval Development of Supgo, *Penaeus monodon*

Edgardo P. Reyes

Aquaculture Department  
Southeast Asian Fisheries Development Center  
P.O. Box 256, Iloilo City, Philippines

Incubation of *Penaeus monodon* eggs and rearing of different larval stages were undertaken at nine temperature-salinity combinations. The eggs, nauplii, zoea and mysis from one spawner kept as stock culture at ambient temperatures of 26-30°C and salinity of 32-33 ppt were exposed to temperature levels of 23, 28 and 33°C and salinity levels of 23, 28 and 33 ppt.

Eggs and nauplii survived the sudden change of temperature and salinity (from ambient to experimental) but the zoea and mysis did not. However, salinities of 23 and 28 ppt in combination with any of the temperature levels produced weak larvae. Highest mean hatching rate was obtained at the temperature-salinity combination of 23°C-33 ppt, followed by 28°C-33 ppt and 33°C-33 ppt. Incubation periods for these treatments were 22, 16 and 14 hr, respectively. Survival rate of nauplius (taken from stock cultures) to first zoeal stage was highest at 28°C-33 ppt, followed by 33°C-33 ppt and 23°C-33 ppt with molting time of 50, 45 and 75 hr, respectively.

The nauplii exposed to 33°C-33 ppt molted to zoea stage within 38 to 40 hr but later died. Those exposed to 23°C-33 ppt and 28°C-33 ppt reached zoea stage within 57 to 60 hr and 48 to 50 hr, respectively. Similarly, the nauplii taken from the stock cultures and reared until postlarval stage (P<sub>1</sub>) under experimental conditions completed the zoea and mysis