

QUT Digital Repository:
<http://eprints.qut.edu.au/>



Turrell, Gavin and Bentley, Rebecca and Thomas, Lyndal and Jolley, Damien and Subramanian, S.V. and Kavanagh, Anne (2009) *A multilevel study of area socio-economic status and food purchasing behaviour*. Public Health Nutrition, 12(11). pp. 2074-2083.

© Copyright 2009 Gavin Turrella, Rebecca Bentley, Lyndal R. Thomas, Damien Jolley, S.V. Subramanian and Anne M. Kavanagh

1 A multilevel study of area socioeconomic status and food purchasing behaviour

2

3

4 Running head: Area socioeconomic status and diet

5

6 **Abstract**

7 **Objective**

8 This study examined the association between area socioeconomic status (SES) and food
9 purchasing behaviour.

10

11 **Setting**

12 Melbourne city, Australia, 2003.

13

14 **Participants**

15 Residents of 2,564 households located in 50 small areas.

16

17 **Design**

18 Data were collected by mail survey (64.2% response rate). Area SES was indicated by the
19 proportion of households in each area earning less than Aus\$400 per week, and
20 individual-level socioeconomic position was measured using education, occupation, and
21 household income. Food purchasing was measured on the basis of compliance with
22 dietary guideline recommendations (for grocery foods) and **variety of fruit and**
23 **vegetable purchase.** Multilevel regression examined the association between area SES
24 and food purchase after adjustment for individual-level demographic (age, sex, household
25 composition) and socioeconomic factors.

26

27 **Results**

28 Residents of low SES areas were significantly less likely than their counterparts in
29 advantaged areas to purchase grocery foods that were high in fibre and low in fat, salt,
30 and sugar; **and they purchased a smaller variety of fruits.** There was no evidence of an
31 association between area SES and vegetable variety.

32

33 **Conclusions**

34 In Melbourne, area SES was associated with some food purchasing behaviours
35 independent of individual-level factors, suggesting that areas in this city may be
36 differentiated on the basis of food availability, accessibility, and affordability, making the
37 purchase of some types of foods more difficult in disadvantaged areas.

38

39

40

41 **Introduction**

42 A large literature has examined the association between individual-level socioeconomic
43 position (SEP) and diet. The findings of this work typically show that socioeconomically
44 disadvantaged groups have diets that are least consistent with recommended intakes of
45 foods and nutrients,¹⁻³ and least in accord with dietary guideline messages that promote
46 foods that are high in fibre and low in fat, salt, and sugar.⁴⁻⁵ Significantly, the poorer
47 dietary intake of disadvantaged groups contributes in part to their higher rates of
48 mortality and morbidity for chronic disease.⁶⁻⁷

49 During the last decade, researchers have increasingly turned their attention to the
50 question of whether place of residence influences diet independently of individual-level
51 factors; and more particularly, whether living in a socioeconomically disadvantaged area
52 is associated with a less healthy diet. Our review of this (small) literature suggested that
53 area studies of diet can be broadly divided into two types that reflect the analytic method
54 used; namely, studies that undertake multivariable analyses using both area- and
55 individual-level variables but without the capacity to statistically integrate the two levels
56 (i.e. contextual studies), and multilevel studies. Six of the former types of study⁸⁻¹³ and
57 five of the latter were identified,¹⁴⁻¹⁸ and key aspects of each are summarised in Table 1.

58 TABLE 1 ABOUT HERE

59 The findings of the contextual studies were reasonably consistent in that they each found
60 some evidence that living in a disadvantaged area was associated with a poorer diet after
61 adjusting for individual-level socioeconomic and demographic factors. These studies
62 however were often based on a small number of areas, and in most cases, the analytic
63 approach did not allow for the partitioning of area- and individual-level sources of
64 variation, hence it was unclear whether differences in diet between advantaged and
65 disadvantaged areas were due to a composition effect (i.e. the clustering of rich and poor
66 people in rich and poor areas) or the environmental characteristics of the areas per se (i.e.
67 a context effect, possibly reflecting area differences in physical infrastructure, services,
68 and facilities). The findings of the multilevel studies, which allow for area- and
69 individual-level variation to be partitioned and quantified, present a somewhat different
70 picture. Of the five identified, only two reported a significant difference in diet between
71 areas after adjustment for individual-level factors.^{15,17} Diez-Roux et al¹⁴ and Ecob and
72 Macintyre¹⁵ found that residents of socioeconomically disadvantaged areas had poorer
73 diets than those in more advantaged areas, although the findings of the former were weak
74 and often not statistically significant. Area SES was not associated with food purchasing

75 behaviour in an Australian study¹⁶ or with dietary intake and food choice in a Dutch
76 study.¹⁸

77 This paper contributes to the literature on areas and diet by reporting on a
78 multilevel study that examined the association between area SES and food purchasing
79 behaviour in the Melbourne metropolitan region (Australia) in 2003. The relationship
80 between SES and diet in Australia (and elsewhere) has to date been investigated
81 primarily using ‘quantitative’ dietary indicators such as mean daily intakes of nutrients,
82 nutrient density levels, or percentage contribution of food to nutrition and energy.⁵ By
83 contrast, few studies have examined the relationship using ‘qualitative’ indicators such as
84 food purchasing behaviour. Clearly, people need to procure food (which usually means
85 purchase it) before it can be consumed and converted into energy and nutrients, and there
86 are a number of compelling reasons why it is important to better understand the factors
87 that influence the food purchasing choices of different socioeconomic groups. First, most
88 people make dietary decisions in relation to food and not nutrients,¹⁹ thus when shopping
89 our food choices are more likely to be influenced by factors such as price, availability,
90 taste preference, and convenience etc than by the vitamin and mineral content of the food.
91 Second, research has shown that the type of food people buy influences dietary quality.²⁰
92 Third, food choice differences between socioeconomic groups translate into concomitant
93 differences in nutrient intake.²¹⁻²² Fourth, knowing about the factors that influence
94 socioeconomic differences in food purchasing is important in assessing the reach and
95 impact of health promotion messages, many of which focus on encouraging people to
96 make healthy food choices when shopping.²³⁻²⁵

97 This study investigates whether residents of socioeconomically advantaged and
98 disadvantaged areas differ in their purchase of grocery foods, fruits, and vegetables.
99 Specifically, three questions are examined:

- 100 1. Do areas vary in their food purchasing profiles?
- 101 2. To what extent does within-area clustering by individual-level SEP account for any
102 observed differences between areas in their food purchasing profiles?
- 103 3. What is the relationship between area SES and food purchasing after adjustment for
104 within-area differences in food purchasing by individual-level SEP?

105
106
107
108
109

110 **Methods**

111 *Geographic scope*

112 This paper is based on data collected as part of the Victorian Lifestyle and
113 Neighbourhood Environment Study (VicLANES), a cross-sectional multilevel
114 investigation of area- and individual-level factors and health-related behaviour. The
115 target population for VicLANES comprised people living in an area extending 20km
116 from the central business district of Melbourne city, the capital of the state of Victoria.

117

118 *Sample design*

119 The sample comprised non-institutionalised residents of private dwellings (households)
120 and Census Collector Districts (CCD). A CCD is the smallest administrative unit used by
121 the Australian Bureau of Statistics (ABS) to collect census data. In urban areas such as
122 Melbourne, a CCD contains an average of 200 private dwellings which are deemed to be
123 relatively homogeneous in terms of their socioeconomic characteristics. Households and
124 CCDs were selected using a stratified two-stage cluster design. At the first stage, all
125 CCDs in the Melbourne metropolitan area were ranked according to the proportion of the
126 households in each CCD with an income of less than Aus\$400 per week. The resultant
127 distribution was stratified into septiles, and a total of 50 CCDs were randomly selected
128 from the low (n=17), middle (n=16) and high income (n=17) strata. At the second stage,
129 we used names and addresses on the Australian Electoral Roll to identify all residents
130 aged 18-74 years in each of the 50 CCDs. Voting is compulsory in Australia for persons
131 aged 18 years and over, so the electoral roll provides near-complete coverage of the
132 resident adult population. A total of 3995 households were then randomly sampled, and
133 the person within each household who was primarily responsible for most of the food
134 shopping was targeted for data collection.

135

136 *Data Collection*

137 The household-level data collection within each CCD occurred between September and
138 December 2003, and was conducted using a mail-survey method described by Dillman.²⁶
139 A total of 2564 usable surveys were returned to yield a final response rate of 64.2%.

140

141 *Measures*

142 Area SES: The septiles forming the sampling strata were used as the basis for measuring
143 area SES. In each of the three strata the average proportion of households earning less
144 than Aus\$400 per week was 7.0% (range 3.5%-8.5%), 15.3% (14.4%-16.7%), and 31.0%

145 (24.1%-59.6%) respectively: these strata were subsequently labelled as high, medium,
146 and low SES. The area-level socioeconomic characteristics of the three strata were
147 further examined using 2001 census data,²⁷ and they differed markedly in terms of their
148 unemployment rate (4.0%, 6.6%, 11.0% respectively), the percentage of employees in
149 unskilled and semi-skilled jobs (7.1%, 13.8%, 20.7%), the percentage of dwellings that
150 were rented from the public housing authority (0.17%, 1.6%, 14.5%), and the percentage
151 of dwellings with no motor vehicle (3.9%, 9.6%, 21.2%).

152

153 Education: Respondents were asked to provide information about whether they had
154 attained further education since leaving school, and if so, the highest qualification
155 completed. Respondent's education was subsequently coded as (1) bachelor degree or
156 higher (the latter included post graduate diploma, masters degree, or doctorate), (2)
157 diploma (associate or undergraduate), (3) vocational (trade or business certificate, or
158 apprenticeship), and (4) no post-school qualifications.

159

160 Occupation: Respondents who were employed at the time of completing the survey were
161 asked to indicate their job title and then to describe the main tasks or duties they
162 performed. This information was subsequently coded to the Australian Standard
163 Classification of Occupations (ASCO).²⁸ For the purposes of this study, the original nine-
164 level ASCO classification was re-coded into three categories: (1) managers/professionals
165 (managers and administrators, professionals, and para-professionals); (2) white collar
166 employees (clerks, salespersons and personal service workers), and (3) blue collar
167 employees (trades-persons, plant and machine operators and drivers, and labourers and
168 related workers). A fourth category, "not in the labour force", comprising the retired,
169 unemployed, students, and those engaged in home duties on a full time basis, was also
170 created.

171

172 Income: Respondents were asked to indicate their total annual household income
173 (including pensions, allowances, and investments) using a 14 category measure that was
174 subsequently re-coded into five groups for analysis: (1) Aus\$78,000 or more, (2)
175 \$52,000-77,999, (3) \$36,400-51,999, (4) \$20,800-36,399, and (5) less than \$20,799.
176 Households in categories 4 and 5 received annual incomes at or below the Australian
177 average in 2000.²⁹

178

179 Confounding: Age in years (centred), sex, and household composition were used as
180 potential confounding variables.

181

182 Food purchasing: As part of the questionnaire, information was sought about the
183 purchase of grocery items, fruits, and vegetables.

184

185 Grocery food purchase: this was examined on the basis of 15 questions, each of which
186 had two or more response options. For example, respondents were asked “When
187 shopping for your household, what type of milk do you usually buy”? The response
188 options included: ‘I do not buy milk’, ‘extra creamy’, ‘full cream’, ‘low fat/trim’,
189 ‘skim/fat free’, plus others. Multiple responses were permitted for each question. The
190 other 14 questions were structured in an identical manner and pertained to bread, rice,
191 pasta, noodles, baked beans, tinned fruit, cheese, yoghurt, beef mince, chicken, tinned
192 fish, cooking oils, butter, and solid cooking fat. In Australia, dietary authorities
193 recommend that people purchase and consume a variety of foods that are relatively high
194 in fibre and low in fat, salt, and sugar²⁵; and consistent with these guidelines, we
195 classified the foods into a recommended and regular category (Table 2). Using this
196 classification, we developed a measure that captured the extent to which peoples’ grocery
197 purchasing patterns were consistent (or not) with dietary guideline recommendations.
198 First, for each food-type (e.g. milk), respondents were assigned the value 1 if they
199 reported usually purchasing only the regular option exclusively (and not any
200 recommended options); they were assigned the value 3 if they reported usually
201 purchasing only the recommended option exclusively (and not any regular options); and
202 they were assigned a value of 2 if they reported usually purchasing a mix of regular and
203 recommended options (e.g. full cream and skim milk). There were a small number of
204 respondents who reported that they never purchased a particular type of food and these
205 were assigned the value 0. In sum, for each of the 15 food-types, respondents were
206 assigned a value of 0, 1, 2 or 3. Second, an initial food purchasing index was created that
207 involved summing the scores for the 15 food-types, with those scoring 0 being excluded
208 at this point. This initial index had a potential range of 15 – 45, with 15 denoting people
209 who purchased the regular option for each food-type, and 45 denoting those who
210 purchased the recommended option for all foods. It is important to note that the
211 respondents included in this initial index reported purchasing all of the 15 food-types.
212 Those scoring 0 for one or more food-types were excluded because their final index score
213 would not accurately reflect their purchasing pattern. For example, someone who

214 purchased all 15 food-types and chose the recommended option for each item would
215 score 45, whereas someone who purchased 13 food-types and chose the recommended
216 option for each item would score 39. Clearly, both people have identical purchasing
217 patterns with respect to the dietary guidelines (i.e. they are making the healthier choice
218 for every food item) but this isn't reflected in their index score. To deal with this issue,
219 and as a way of including the full sample in the analyses, respondents who reported not
220 buying one or more of the food items were included in the index using the following
221 formula: $\text{Index score} = a / 15 - b$. The quantity 'a' represented each respondent's initial
222 score which was derived by summing the values (1, 2 or 3) for each of the food-types.
223 The denominator comprised the constant '15' which represented the number of food-
224 types in the index, and the variable 'b', which represented the number of food-types not
225 purchased by the respondent. In effect, the formula calculated a mean food purchasing
226 score for each respondent. Finally, the index was re-scored to range from 0 – 100, with
227 higher scores indicating a purchasing pattern that was more consistent with dietary
228 guideline recommendations (sample mean 47.6, SD 13.4).

229 TABLE 2 ABOUT HERE

230 Fruit purchasing: this was examined using a question that asked 'When shopping for
231 fresh fruit, how often do you buy these types'? The respondent was instructed to include
232 seasonal fruits, but exclude fruit juice, canned, and dried fruit. The question item-set
233 consisted of 22 fresh fruits selected (mostly) from the food frequency questionnaire used
234 in the 1995 Australian National Nutrition Survey.³¹ For each fruit, respondents were
235 asked to indicate their usual purchasing pattern on the basis of five-point scales:
236 1='Never buy', 2='Rarely Buy', 3='Sometimes buy', 4='Nearly Always Buy' and
237 5='Always buy'.

238 Using these items we created an index that measured variety of fruit purchased.
239 For each fruit item, respondents reporting 'never' or 'rarely' buy were scored 0, and those
240 reporting any of the other three options were scored 1. The items were then summed,
241 with the resultant index score for each respondent indicating the variety of fruits
242 purchased (sample mean 14.2, SD 4.1). Importantly, the variety score does not reflect the
243 range of fruits purchased on any particular shopping trip, but rather, the types that are
244 purchased at least sometimes over the course of many shopping episodes depending on
245 factors such as seasonality, price, and quality. As the variety index was essentially a
246 count-measure and non-normally distributed it was categorised into quartiles, with Q1
247 denoting high variety and Q4 low variety.

248

249 **Vegetable purchasing: Respondents were asked to indicate how often they**
250 **purchased 25 vegetables, including fresh and frozen, but excluding canned or dried**
251 **vegetables. A purchasing index measuring vegetable variety was constructed using**
252 **an identical format and method to that used for fresh fruit. The mean variety score**
253 **for vegetables for the sample was 18.5 (SD 4.1).**

254

255 *Analysis*

256 Table 3 presents descriptive statistics for each of the measures used in this analysis.
257 From the 2564 questionnaires that were returned, missing data were identified for
258 education (n=106, 4.1%), occupation (n=83, 3.2%), income (n=903, 35.2%), sex (n=4,
259 0.16%), age (n=5, 0.19%), and household composition (n=55, 2.2%). In total, the
260 proportion of the sample with completely observed data for all the variables examined
261 (complete cases) was 57%. We have not reported results obtained by analysing only the
262 complete cases because of the potential bias and loss of precision associated with the
263 large proportion of missing income data: instead, we used multiple imputation. We
264 imputed all missing data under a missing at random (MAR) assumption and adopted an
265 inclusive strategy for the imputation model³²⁻³⁴ Ten datasets with imputed values for
266 missing items on each variable were estimated using the command ‘Imputation by
267 Chained Equations (ICE)’ in Stata 9.2.³⁵

268

TABLE 3 ABOUT HERE

269 The grocery data were analysed as a two-level random intercept model in Stata. We
270 specified three models that directly addressed the three research questions identified
271 earlier. Model 1 (baseline) quantified the extent of area-level variation in food-purchasing
272 behaviour conditional on the confounders. Here, the substantive interest was on the
273 random term, which if significant, indicated that food purchasing patterns differed
274 between the 50 CCDs. For this and subsequent models we also calculated an intraclass
275 correlation (ICC) by dividing the between-CCD variance by the total variance, and this is
276 interpreted as the proportion of the total variation in food purchasing behaviour that is
277 between the CCDs. Model 2 extends Model 1 by adding education, occupation, and
278 income as fixed effects, and examined the extent to which they account for variation in
279 food purchasing between the CCDs. Model 3 then extended Model 2 by including the
280 measure of area SES as a fixed effect: here the focus is on whether area SES is associated
281 with food purchasing independently of within-area variation in age, sex, household
282 composition and individual-level SEP.

283 **Variety of fruit and vegetable purchase was examined using a two-level**
284 **ordered multinomial logit-link model. ‘High’ variety (quartile 1) was denoted the**
285 **reference category, hence positive regression coefficients for any of the predictor**
286 **variables indicate a greater odds of purchasing a lower variety of fruits and**
287 **vegetables. Three models were specified. Model 1 (baseline) quantified the extent of**
288 **area-level variation in fruit and vegetable variety conditional on the confounders.**
289 **Model 2 added education, occupation, and income, and Model 3 included area SES.**
290 **The results are presented as odds ratios and their 95% confidence intervals.**

291

292 **Results**

293 Table 4 presents the findings of the multilevel analyses which examined the independent
294 contribution of area- and individual-level socioeconomic factors to grocery food
295 purchase. In Model 1, the area-level random terms was statistically significant ($p=0.033$),
296 indicating that the average grocery purchasing score was different (beyond chance)
297 across the 50 CCD. Of the total variability in grocery purchase, 1.5% occurred between
298 CCD and 98.5% between individuals. Model 2 adds the fixed (average) effects for
299 education, occupation and income: this attenuated the between-area variation by 59.8%,
300 and the random term was no longer significant ($p=0.241$). Education and income were
301 associated with grocery purchase: respondents with no post-school qualifications and
302 those living in low income households scored significantly lower on the index. No
303 significant occupational effects were observed. Model 3 adds the fixed effect for area
304 SES and the coefficients indicate that residents of medium and low-SES areas scored
305 significantly lower on the grocery purchasing index than their counterparts from high-
306 SES areas.

307

TABLE 4 ABOUT HERE

308 Table 5 presents the findings of the ordered multilevel logistic regression analysis which
309 examined the contribution of area- and individual-level socioeconomic factors to variety
310 of fruit and vegetable purchasing. Fruit variety scores were significantly different
311 ($p=0.01$) across the 50 CCDs (Model 1). After adjustment for education, occupation and
312 income (Model 2) the between-area variation in fruit variety was attenuated by 50.0%
313 and remained marginally statistically significant ($p=0.06$). Respondents with no post-
314 school qualifications had 1.72 (95% CI 1.25-2.38) times higher odds of purchasing a
315 lower variety of fruits. The corresponding odds for respondents from low income families
316 was 1.69 (95% CI 1.11-2.57). Model 3 adds the measure of area SES which made no
317 appreciable difference to the between-CCD variation (relative to Model 2) although the

318 random term was no longer statistically significant ($p=0.11$). The coefficients for area
319 SES show that residents of low SES areas had significantly higher odds of purchasing a
320 lower variety of fruits than residents in the high SES areas (OR 1.30, 95% CI 1.00-1.67).
321 Independent of area SES, respondents with lower levels of education, and residents of
322 lower income households, had significantly higher odds of purchasing a more limited
323 variety of fruits than their higher status counterparts.

324 Vegetable variety scores did not differ significantly across the 50 CCD (Model 1)
325 and the inclusion of education, occupation, and income further attenuated the CCD
326 variation (Model 2). Respondents with no post-school qualifications had a significantly
327 higher odds of purchasing a lower variety of vegetables relative to those with a bachelor
328 degree (OR 1.36 95% CI 1.08-1.72). There was no association between vegetable variety
329 and occupation, income or area SES (Model 3).

330 TABLE 5 ABOUT HERE

331

332 **Discussion**

333 In metropolitan Melbourne in 2003 area SES was associated with the purchase of grocery
334 foods and fruit variety. Compared with their counterparts in high SES areas, residents of
335 low SES areas were less likely to buy groceries that were high in fibre and low in fat, salt,
336 and sugar; and they purchased a smaller variety of fruits. These findings are broadly
337 consistent with the results of multilevel studies conducted in the US¹⁴ and Scotland¹⁵;
338 however, they are at odds with multilevel research conducted in the Netherlands¹⁸ and in
339 Brisbane, Australia.¹⁶ Reconciling these differences, and hence being able to generalise
340 about the relationship between area SES and diet, is difficult. In part, these difficulties
341 stem from the limited evidence-base (i.e. the small number of multilevel studies) and
342 methodological issues such as differences in the conceptualisation and measurement of
343 diet, the individual-level variables used as confounders, and the number and size of the
344 area-units used.¹⁵ The inconsistencies between study findings however, are likely to be
345 more than a methodological artefact, and may reflect “real” historical, cultural, political,
346 socioeconomic, and geospatial differences between countries (e.g. US and Australia) and
347 between regions within the same country (e.g. Brisbane and Melbourne). At present, the
348 mixed findings of the small number of multilevel studies do not provide a sufficiently
349 reliable basis on which to make a general call for area-level public health interventions to
350 improve conditions in deprived areas to facilitate the procurement of foods that are
351 conducive to a healthy diet: rather, any “call” may have to be specific and tailored to each
352 particular geographic and spatial context.

353 A large literature documents an association between individual-level SEP and
354 diet, and most of this work has focused on socioeconomic differences in food and
355 nutrient intakes.⁵ These studies usually find that socioeconomically disadvantaged groups
356 have intakes that are consistent with their higher rates of diet-related chronic disease.¹⁻³
357 To some extent at least, the results of this food purchasing study extend and complement
358 the findings of the intake studies by showing that those of low SEP are less likely to buy
359 grocery foods that accord with diet-related health promotion messages and dietary
360 guidelines. In addition, low socioeconomic groups had a significantly higher odds of
361 purchasing a lower variety of fruits and vegetables.

362

363 **Study Limitations**

364 First, survey non-response tends to be higher in disadvantaged areas³⁶ and among
365 individuals of low SEP.³⁷ Non-response in the VicLANES study was 35.8%, hence the
366 sample probably under-represents the disadvantaged areas and individuals, and over-
367 represents the advantaged, and the observed socioeconomic differences in food
368 purchasing are likely to be an under-estimate of the actual differences in the Melbourne
369 population.

370 Second, as with most multilevel studies³⁸ our use of a CCD to represent a
371 neighborhood was made for reasons of sampling and analytic convenience rather than
372 being underpinned by an explicit theory linking area SES and food purchasing; hence
373 associations among these variables are likely to be underestimated.

374 Third, our finding of an association between area SES and food purchase might be
375 confounded by individual-level socioeconomic factors not included in the models. This
376 said however, we included the three most widely used indicators of a person's
377 socioeconomic characteristics,³⁹ and given the correlation among these indicators⁴⁰ it is
378 likely that education, occupation, and income were capturing most of the unmeasured
379 influences of other socioeconomic factors excluded from the models. Alternatively, it
380 may be that the inclusion of these individual-level measures resulted in 'over-adjustment'
381 which argues for the possibility of an even stronger contextual effect on food purchase
382 than was observed in this study. If education, occupation and household income represent
383 part of the pathway via which area SES influences food procurement, then modelling
384 individual-level socioeconomic variables may inappropriately attenuate the variation that
385 is more correctly attributable to area disadvantage.⁴¹

386

387

388 **Conclusion**

389 In the Melbourne metropolitan region in 2003, differences between advantaged and
390 disadvantaged areas in their purchasing profiles for grocery foods and fruits, and
391 ‘healthier’ purchasing in higher SES areas, suggest that the areas may be differentiated on
392 the basis of food availability, accessibility, and affordability, making the purchase of
393 some types of foods more difficult for people living in disadvantaged areas. To date, the
394 between- and within-country (multilevel) evidence linking area-disadvantage and diet is
395 both sparse and inconsistent. Methodological issues notwithstanding, this might suggest
396 that area deprivation is not universally associated with poorer access to healthy food.
397 Cummins and Macintyre⁴² reached a somewhat similar conclusion based on their review
398 of the literature on food environments and obesity. A challenge for future area-based
399 dietary research is to identify those ecologic characteristics (e.g. urban design, shopping
400 infrastructure, and transport services) that promote equality of access to healthy food, and
401 those characteristics that make its attainment difficult.

402 **References**

- 403 1. Metcalf P, Scragg R, Davis P. Dietary intakes by different markers of socioeconomic
404 status: results of a New Zealand workforce survey. *New Zealand Medical Journal*.
405 2006; **26**: U2127.
- 406 2. Shahar D, Shai L, Vardi H, Shahar A, Fraser D. Diet and eating habits in high and
407 low socioeconomic groups. *Nutrition* 2005;**21**: 559-566.
- 408 3. De Irala-Estevez J, Groth M, Johansson L, Oltersdorf U, Prattala R, Martinez-
409 Gonzalez MA. A systematic review of socioeconomic differences in food habits in
410 Europe: consumption of fruit and vegetables. *European Journal of Clinical Nutrition*.
411 2000;**54**: 706-714.
- 412 4. Turrell G. Compliance with the Australian dietary guidelines in the early 1990's:
413 have population-based health promotion programs been effective? *Nutrition and*
414 *Health* 1997;**11**: 271-288.
- 415 5. Turrell G, Hewitt B, Patterson C, Oldenburg B, Gould T. Socioeconomic differences
416 in food purchasing behaviour and suggested implications for diet-related health
417 promotion. *Journal of Human Nutrition Dietetics*. 2002; **15**:355-364.
- 418 6. James WPT, Nelson M, Leather S. The contribution of nutrition to inequalities in
419 health. *British Medical Journal* 1997;**314**: 1545-1549.
- 420 7. Davey Smith G, Brunner E. Socioeconomic differentials in health: the role of
421 nutrition. *Proceedings of the Nutrition Society* 1997;**56**: 75-90.
- 422 8. Diehr P, Koepsell T, Cheadle A, Psaty BM, Wagner E, Curry S. Do communities
423 differ in health behaviors? *Journal of Clinical Epidemiology* 1993;**46**: 1141-1149.
- 424 9. Forsyth A, Macintyre S, Anderson A. Diets for disease? Intraurban variation in
425 reported food consumption in Glasgow. *Appetite* 1994; **22**:259-274.
- 426 10. Karvonen S, Rimpela A. Socio-regional context as a determinant of adolescents'
427 health behaviour in Finland. *Social Science and Medicine* 1996; **43**:1467-1474.
- 428 11. Ellaway A, Macintyre S. Does where you live predict health related behaviours? A
429 case study in Glasgow. *Health Bulletin*. 1996; **54**: 443-446.
- 430 12. Karvonen S, Rimpela AH. Urban small area variation in adolescents' health
431 behaviour. *Social Science and Medicine*. 1997; **45**: 1089-1098.
- 432 13. Shohaimi S, Welch A, Bingham S, Luben R, Day N, Wareham N, Khaw K-T.
433 Residential area deprivation predicts fruit and vegetable consumption independently
434 of individual educational level and occupational social class: a cross-sectional
435 population study in the Norfolk cohort of the European Prospective Investigation into

- 436 Cancer (EPIC-Norfolk). *Journal of Epidemiology and Community Health*. 2004; **58**:
437 686-691.
- 438 14. Diez-Roux AV, Nieto FJ, Caulfield L, Tyroler HA, Watson RL, Szklo M.
439 Neighbourhood differences in diet: the Atherosclerosis Risk in Communities (ARIC)
440 study. *Journal of Epidemiology and Community Health*. 1999; **53**: 55-63.
- 441 15. Ecob R, Macintyre S. Small area variations in health related behaviours; do these
442 depend on the behaviour itself, its measurement, or on personal characteristics?
443 *Health and Place*. 2000; **6**: 261-274.
- 444 16. Turrell G, Blakely T, Patterson C, Oldenburg B. A multilevel analysis of
445 socioeconomic (small area) differences in household food purchasing behaviour.
446 *Journal of Epidemiology and Community Health*. 2004; **58**: 208-215.
- 447 17. Ball K, Crawford D, Mishra. G. Socioeconomic inequalities in women's fruit and
448 vegetable intakes: a multilevel study of individual, social, and environmental
449 mediators. *Public Health Nutrition*. 2006; **9**: 623-630.
- 450 18. Giskes K, Turrell G, van Lenthe FJ, Brug J, Mackenbach JP. A multilevel study of
451 socioeconomic inequalities in food choice behaviour and dietary intake among the
452 Dutch population: the GLOBE study. *Public Health Nutrition* 2006; **9**: 75-83.
- 453 19. Crotty P, Rutishauser I, Cahill M. From food to nutrients and back. *Proceeding of the*
454 *Nutrition Society of Australia*. 1991; **16**: 38.
- 455 20. Shimakawa T, Sorlie P, Carpenter MA, Dennis B, Tell GS, Watson R, Williams D.
456 Dietary intake patterns and sociodemographic factors in the Atherosclerosis Risk in
457 Communities Study. *Preventive Medicine* 1994; **23**: 769-780.
- 458 21. Smith A, Baghurst K. Public Health implications of dietary differences between
459 social status and occupational category groups. *Journal of Epidemiology and*
460 *Community Health*. 1992; **46**: 409-416.
- 461 22. Syrette JA, Baghurst KI, Record SJ. Sociodemographic determinants of the
462 contribution of cereal foods to nutrient intake in the Australian population. *Food*
463 *Australia*. 1990; **42**: 330-337.
- 464 23. Dixon H, Borland R, Segan C. Public reaction to Victoria's '2 Fruit 'n' 5 Veg every
465 day' campaign and reported consumption of fruit and vegetables. *Preventive*
466 *Medicine*. 1998; **27**: 572-582.
- 467 24. Krebs-Smith SM, Scott Kantor L. Choose a variety of fruit and vegetables daily:
468 understanding the complexities. *Journal of Nutrition*. 2001; **131**: 487S-501S.
- 469 25. Strategic Inter-Governmental Nutrition Alliance (SIGNAL). *Eat Well Australia: An*
470 *agenda for action for public health nutrition 2001-2010*. Canberra, National Public

- 471 Health Partnership, 2001.
472 http://www.health.vic.gov.au/nhpa/resources/nu_eatwell.htm
- 473 26. Dillman DA. *Mail and internet surveys: the tailored design method*. Chichester,
474 Wiley, 2000.
- 475 27. Australian Bureau of Statistics. *CDATA 2001. Your Census at Work CD-ROM (2nd*
476 *Release)*. Cat. No. 2019.0.30.001, Canberra, Australian Bureau of Statistics, 2003.
- 477 28. Australian Bureau of Statistics. *Australian Standard Classification of Occupations,*
478 *Second Edition*. Catalogue No. 1220.0. Canberra, Australian Government Publishing
479 Service, 1997.
- 480 29. Australian Bureau of Statistics. *Income Distribution 1999-2000*. Catalogue No.
481 6523.0. Canberra, Australian Government Publishing Service, 2001.
- 482 30. Turrell G. Educational differences in dietary guideline food practices: are they
483 associated with educational differences in food and nutrition knowledge? *Australian*
484 *Journal of Nutrition and Dietetics*. 1997; **54**: 25-33.
- 485 31. McLennan W. *National Nutrition Survey User's Guide*. Canberra, Australian
486 Government Publishing Service, 1998.
- 487 32. Collins LM, Schafer JS, Kam C. A comparison of inclusive and restrictive strategies
488 in modern missing data procedures. *Psychological Methods*. 2001; **6**: 330-351.
- 489 33. Rubin DB. Multiple imputation after 18+ years. *Journal of the American Statistical*
490 *Association*. 1996; **91**: 473-489.
- 491 34. Meng XL. Multiple-imputation inferences with uncongenial sources of input.
492 *Statistical Science*. 1994; **9**: 538-573.
- 493 35. Stata Statistical Software, Release 9. College Station, Stata Corporation, Texas, 2003.
- 494 36. Turrell G, Sanders AE, Slade GD, Marcenes W, Spencer AJ. The contribution of
495 area-level disadvantage and individual-level socioeconomic position to self-reported
496 oral health. *Community Dentistry and Oral Epidemiology*. 2007; **35**: 195-206.
- 497 37. Turrell G, Najman JM. Collecting food-related data from low socioeconomic groups:
498 how adequate are our current research designs? *Australian Journal of Public Health*.
499 1995; **19**: 410-416.
- 500 38. Boyle MH, Wilms JD. Place effects for areas defined by administrative boundaries.
501 *American Journal of Epidemiology*. 1999; **149**: 577-585.
- 502 39. Dutton T, Turrell G, Oldenburg B. *Measuring Socioeconomic Position in Population*
503 *Health Monitoring and Health Research*. Health Inequalities Monitoring Series No.
504 3. Brisbane, Australia, Queensland University of Technology, Brisbane, 2005.

- 505 40. Turrell G, Hewitt B, Patterson C, Oldenburg B. Measuring socioeconomic position in
506 dietary research: is choice of socioeconomic indicator important? *Public Health*
507 *Nutrition*. 2003; **6**: 191-200.
- 508 41. Diez Roux AV. Estimating neighbourhood health effects: the challenges of causal
509 inference in a complex world. *Social Science and Medicine*. 2004; **58**:1953-1960.
- 510 42. Cummins S, Macintyre S. Food environments and obesity – neighbourhood or
511 nation? *International Journal of Epidemiology*. 2006; **35**: 100-104.
- 512
- 513
- 514
- 515
- 516
- 517
- 518
- 519

Table 1: Studies examining the association between areal-level socioeconomic disadvantage and diet

Study and Country	Sample	Dietary Outcome	Significant area differences in diet (after adjustment for individual-level factors)	Significant association between area disadvantage and diet (after adjustment for individual-level factors)
<i>Contextual (non-multilevel) studies</i>				
Diehr et al, ⁸ US	Areas N=15 Individuals N=7863 adults 18+	Percentage of calories from fat	Yes (but very small)	Areas with higher unemployment rates had higher percentage of calories from fat
Forsyth et al, ⁹ Scotland	Areas N=4 Individuals N=691 adults 40 & 60 years	Reported consumption of vegetables, potatoes, fruit, fish, bread, cereals, cakes, pastries & biscuits, confectionary, savoury snacks, meat, spreads, sugar, milk, & soft drinks	Area differences in consumption of fruits, vegetables, fried or roasted potatoes, white bread, brown/wholemeal bread, meat (all), processed meat, poultry, shallow fat frying	Residents of disadvantaged areas were less likely to use polyunsaturated fatty acids, brown/wholemeal bread, & natural fruit juice; and more likely to add sugar to drinks & consume soft drink
Karvonen et al, ¹⁰ Finland	Areas N=460 Individuals N=9121 adolescents 16 & 18 years	Percentage using high-fat milk and butter daily	Yes	Consumption of high-fat products was higher in areas with fewer services (boys); and lower in less educated areas (girls)
Ellaway et al, ¹¹ Scotland	Areas N=4, Individuals N=691 adults 40 & 60 years	'Healthy' and 'Less healthy' food consumption indexes	Yes	Not explicitly stated; however, bivariate analyses suggested that residents of disadvantaged areas consumed a less healthy diet
Karvonen et al, ¹² Finland	Areas N=33 Individuals N=1048 adolescents aged 16 & 18 years	Abstaining from use of milk and fat-containing spreads	Could not be determined	Among girls, rates of abstinence from dietary fat were higher in areas with lower rates of prolonged unemployment; no area effects for boys
Shohaimi et al, ¹³ England	Areas (not reported) Individuals N=22 562 adults 39-79 years	Mean intakes (g/day) of fruits (n=11) and vegetables (n=26) combined	Could not be determined	Residents of disadvantaged areas (Townsend Index) were more likely to report lower intakes of fruits and vegetables

Table 1 Continued: Studies examining the association between area-level socioeconomic disadvantage and diet

Study and Country	Sample	Dietary Outcome	Significant area differences in diet (after adjustment for individual-level factors)	Significant association between area disadvantage and diet (after adjustment for individual-level factors)
<i>Multilevel studies</i> Diez-Roux et al, ¹⁴ US	Areas (not reported) Individuals N=13 095 adults 45-64 years	Energy adjusted daily intakes of fruits, vegetables, meats, and fish. Daily intake of saturated and polyunsaturated fat, and cholesterol. Keys score: extent to which diet increases serum cholesterol	No	Lower income neighborhoods typically had lower energy adjusted intakes of fruits, vegetables, fish, and increased intake of meats, but the associations were weak and often not significant. Inconsistent associations between neighborhood income and intakes of fats and cholesterol.
Ecob et al, ¹⁵ Scotland	Areas N=52 Individuals N=3036 persons 15, 35 & 55 years	Consumption of five foods (fresh fruits and vegetables, wholemeal bread versus white, soft margarine vs butter, & low fat milk vs full cream) scored as a 'good' diet (i.e. consumes 4+ healthy options) and 'bad' diet (no healthy options)	Yes ('bad' diet only)	Residents of disadvantaged areas (Carstairs-Morris deprivation index) were significantly more likely to have a 'bad' diet and less likely to have a 'good' diet.
Turrell et al, ¹⁶ Australia	Areas N=50 Individuals N=970 adults 18-94 years	Three indexes measuring purchase of fruits, vegetables, and grocery foods	No	Advantaged and disadvantaged areas did not differ significantly in their food purchasing patterns.
Ball et al, ¹⁷ Australia	Areas N=45 Individuals N=1347 women 18-65 years	Servings of fruits and vegetables each day (1, 2, 3-4, 5 or more)	Fruit: no. Vegetables: yes	Not reported
Giskes et al, ¹⁸ The Netherlands	Areas N=85 Individuals N=1339 adults 25-79 years	Index measuring food choices consistent with Dutch dietary guidelines; fruit consumption (servings per day); breakfast consumption (days per week); total and saturated fat intake (% of energy)	No	Advantaged and disadvantaged areas did not differ significantly in their grocery food purchasing patterns, their consumption of fruit and propensity to skip breakfast, nor in terms of their total and saturated fat intake

Table 2: Classification of grocery food-types into ‘recommended’ and ‘regular’ categories

Food-type	Recommended	Regular
Bread	Wholemeal, multigrain, white high in fibre, rye, soy and linseed	White
Rice	Wholemeal or brown	Plain white and other white rice (basmati, jasmine, Arborio)
Pasta	Wholemeal or brown	Other pasta (white, spinach, herb)
Noodles	Wholemeal or brown	Other noodles (white, egg, spinach)
Baked Beans	Salt-reduced or unsalted	Regular salt
Tinned Fruit	In natural juice	In syrup
Cheese	Reduced Fat (25% less fat), low fat (<10% fat)	Full fat
Milk	Reduced fat, low fat, high calcium, high calcium skim, high iron, high protein, reduced lactose, no cholesterol, soy or soy & linseed (Skim)	Extra Creamy, full cream, soy or soy & linseed (full cream)
Yoghurt	Low-Fat (plain and fruit)	Full fat (plain and fruit)
Beef Mince	Lean (Trim/Premium)	Regular (Choice/Fine Grade)
Chicken (uncooked)	Without skin, with skin (and remove before eating)	With skin (and eat skin)
Tinned Fish	In water/spring water	In oil or brine
Vegetable Oil	Canola, sunflower, safflower, olive, corn, soybean, peanut or sesame, grape seed or macadamia	Blended oils, coconut oil, palm oil
Butter	Salt-reduced, unsalted	Regular salt
Solid Cooking Fat	Cooking margarine, solidified oil	Solid animal fat (lard, beef dripping), vegetable shortening, Ghee or butter (and use for cooking)

Table 3: Descriptive statistics for the socioeconomic and demographic variables and the measures of food-purchasing behaviour (Melbourne city, Australia, 2003)

N=2564	N	%
Area-Disadvantage		
Low	914	35.7
Medium	895	34.9
High	755	29.5
Education		
Bachelor degree or higher	815	31.8
Diploma	290	11.3
Vocational	393	15.3
No post-school qualifications	1006	41.6
Occupation		
Professionals	861	33.6
White collar	485	18.9
Blue collar	140	5.5
Not in the labour force	1078	42.0
Income		
Aus\$78,000 or more	702	27.4
\$52,000 - \$77,999	605	23.6
\$36,400 - \$51,999	398	15.5
\$20,800 - \$36,399	391	15.3
\$20,799 or less	468	18.3
Sex		
Female	2181	85.1
Male	383	14.9
Household composition		
1 adult, no children	427	16.7
1 adult, 1 or more children	207	8.1
2 or more adults, no children	911	35.5
2 or more adults, 1 or more children	1019	39.7
Age (mean, SD)	49.0	13.5

Table 4: Area- and individual-level socioeconomic effects on the purchase of grocery foods consistent with dietary guideline recommendations (Melbourne city, Australia, 2003)

	Groceries ^{a, b, c}					
	Model 1		Model 2		Model 3	
	β	se	β	se	β	se
Intercept	43.0	1.5	41.5	1.1	42.8	1.1
<i>Area SES</i>						
High					--	
Medium					-2.09	0.70**
Low					-2.43	0.76***
<i>Education</i>						
Bachelor Degree or higher			--		--	
Diploma			0.35	0.99	0.45	0.98
Vocational			0.22	0.93	0.20	0.93
No post-school qualifications			-1.73	0.79**	-1.54	0.78*
<i>Occupation</i>						
Professionals			--		--	
White collar			-0.09	0.89	-0.07	0.88
Blue collar			0.41	1.38	0.64	1.37
Not in the labour force			-0.85	0.80	-0.83	0.80
<i>Income</i>						
A\$78,000 or more			--		--	
\$52,000 - \$77,999			-0.86	0.81	-0.70	0.80
\$36,400 - \$51,999			-1.08	0.94	-0.78	0.93
\$20,800 - \$36,399			-2.47	0.99*	-2.06	0.99*
\$20,799 or less			-2.98	1.05**	-2.31	1.06*
<i>Random effects</i>						
Area variance	2.54	1.2	1.02	0.9	0.182	0.7
p-value for area variance	0.033		0.241		0.784	
Intra-class correlation (%)	1.5		0.60		0.10	

a. Model 1: Baseline model adjusted for age, sex, and household composition,

b. Model 2: Model 1 plus education, occupation and income

c. Model 3: Model 2 plus area SES

p-value significant at * ≤ 0.05 , ** ≤ 0.01 , *** ≤ 0.001

Table 5: Area- and individual-level socioeconomic effects on variety of fruit and vegetable purchasing (Melbourne city, Australia, 2003) ^a

	Fruit Variety ^{b, c, d}				Vegetable Variety							
	Model 1		Model 2		Model 3		Model 1		Model 2		Model 3	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
<i>Area SES</i>												
High			1.00							1.00		
Medium			1.07	0.84-1.37						0.88	0.70-1.11	
Low			1.30	1.00-1.67						1.06	0.83-1.35	
<i>Education</i>												
Bachelor Degree or higher	1.00		1.00				1.00		1.00			
Diploma	1.27	0.84-1.91	1.26	0.94-1.68			1.08	0.78-1.48	1.07		0.79-1.45	
Vocational	1.44	0.99-2.11	1.45	1.11-1.89			1.01	0.75-1.36	1.01		0.76-1.34	
No post-school qualifications	1.72	1.25-2.38	1.70	1.35-2.14			1.36	1.06-1.75	1.36		1.08-1.72	
<i>Occupation</i>												
Professionals	1.00		1.00				1.00		1.00			
White collar	0.85	0.59-1.22	0.85	0.66-1.10			1.02	0.77-1.35	1.02		0.78-1.33	
Blue collar	0.94	0.54-1.64	0.93	0.62-1.37			1.14	0.74-1.76	1.15		0.76-1.73	
Not in the labour force	0.91	0.66-1.27	0.91	0.72-1.15			1.13	0.88-1.46	1.12		0.88-1.43	
<i>Income</i>												
A\$78,000 or more	1.00		1.00				1.00		1.00			
\$52,000 - \$77,999	1.26	0.90-1.75	1.25	0.99-1.58			0.19	0.93-1.56	1.21		0.95-1.55	
\$36,400 - \$51,999	1.09	0.74-1.60	1.07	0.81-1.40			0.88	0.65-1.18	0.88		0.66-1.17	
\$20,800 - \$36,399	1.43	0.95-2.13	1.39	1.04-1.85			1.12	0.81-1.53	1.12		0.83-1.52	
\$20,799 or less	1.69	1.11-2.57	1.59	1.18-2.16			1.17	0.84-1.63	1.16		0.84-1.60	
Random effects												
Area variance & se	0.08	0.03	0.04	0.02	0.04	0.02	0.03	0.02	0.02	0.02	0.01	0.02
p-value for area variance	0.01		0.06		0.11		0.11		0.30		0.41	

a. **High variety (quartile 1) was denoted the reference category, hence odds ratios greater than one indicates an increased likelihood of purchasing a lower variety of fruits and vegetables**

b. Model 1: Baseline model adjusted for age, sex, and household composition

c. Model 2: Model 1 plus education, occupation and income

d. Model 3: Model 2 plus area SES