



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Jiang, H., Liu, F., Turner, I., & Burrage, K. (2012) Analytical solutions for the multi-term time-space Caputo-Riesz fractional advection-diffusion equations on a finite domain. *Journal of Mathematical Analysis and Applications*, 389(2), pp. 1117-1127.

This file was downloaded from: <http://eprints.qut.edu.au/51507/>

© Copyright 2012 Elsevier

This is the author's version of a work that was accepted for publication in the *Journal of Mathematical Analysis and Applications*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Journal of Mathematical Analysis and Applications*, Volume 389, Issue 2, (2012), Pages 1117–1127. DOI: 10.1016/j.jmaa.2011.12.055

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://dx.doi.org/10.1016/j.jmaa.2011.12.055>

Analytical solutions for the multi-term time-space Caputo-Riesz fractional advection-diffusion equations on a finite domain

H. Jiang^{a,b}, F. Liu^{b,*}, I. Turner^b, K. Burrage^{b,c}

^a*Department of Mathematical and Information Sciences, Qinghai Normal University, Xining 810008, China*

^b*Mathematical Sciences, Queensland University of Technology, GPO Box 2434, Brisbane, Qld. 4001, Australia*

^c*Oxford University Department of Computing Science and OCISB, Oxford University, Oxford, OX1 3QD, U.K.*

Abstract

Generalized fractional partial differential equations have now found wide application for describing important physical phenomena, such as subdiffusive and superdiffusive processes. However, studies of generalized multi-term time and space fractional partial differential equations are still under development.

In this paper, the multi-term time-space Caputo-Riesz fractional advection diffusion equations (MT-TSCR-FADE) with Dirichlet nonhomogeneous boundary conditions are considered. The multi-term time fractional derivatives are defined in the Caputo sense, whose orders belong to the intervals $[0, 1]$, $[1, 2]$ and $[0, 2]$, respectively. These are called respectively the multi-term time-fractional diffusion terms, the multi-term time-fractional wave terms and the multi-term time-fractional mixed diffusion-wave terms. The space fractional derivatives are defined as Riesz fractional derivatives. Analytical solutions of three types of the MT-TSCR-FADE are derived with Dirichlet boundary conditions. By using Luchko's Theorem (Acta Math. Vietnam, 1999), we proposed some new techniques, such as a spectral representation of the fractional Laplacian operator and the equivalent relationship between fractional Laplacian operator and Riesz fractional derivative, that enabled the derivation of the analytical solutions for the multi-term time-space Caputo-Riesz fractional advection-diffusion equations.

Key words: Multi-term time-space Caputo-Riesz fractional advection-diffusion equations; Multivariate Mittag-Leffler function; nonhomogeneous initial-boundary-value problem; fractional Laplacian operator; analytical solution.

1 Introduction

In recent years, fractional partial differential equations have attracted great attention [16,26]. The fractional advection-diffusion equation is presented as a useful approach for the description of transport dynamics in complex systems that are governed by anomalous diffusion and non-exponential relaxation patterns [23]. The time, space and time-space fractional advection-dispersion equations have been used to describe important physical phenomena that arise in amorphous, colloid, glassy and porous media, in fractals and percolation clusters, comb structures, dielectrics and semiconductors, biological systems, polymers, random and disordered media, geophysical and geological processes (see [3,8,19,23,24,41]) and in groundwater hydrology to model the transport of passive tracers carried by fluid flow in a porous medium [44].

It is noteworthy to mention recent important progress in numerically simulating fractional advection-diffusion equations. Meerschaert and Tadjeran [21] presented numerical methods to solve the one-dimensional fractional advection-diffusion equation with a Riemann-Liouville fractional derivative on a finite domain. Roop [34] investigated the numerical approximation of the variational solution to the fractional advection-diffusion equation on boundary domains in \mathbb{R}^2 . Liu et al. [11] transformed the space fractional advection-diffusion equation into a system of ordinary differential equations (method of lines) that was then solved using backward differentiation formulas. Liu et al. [12] also considered a space-time fractional advection-diffusion equation with a Caputo time fractional derivative and Riemann-Liouville space fractional derivatives. Shen et al. [36] discussed the fundamental solution and discrete random walk model for the time-space Riesz fractional advection-dispersion equation. Shen et al. [37] also investigated the fundamental solution and numerical solution of the Riesz fractional advection-dispersion equation. Shen et al. [38] proposed numerical approximations and solution techniques for the space-time Riesz-Caputo fractional advection-diffusion equation.

Analytical solutions of fractional advection-diffusion / fractional advection-diffusion-wave equations are of fundamental importance in describing and understanding dispersion phenomena, since all the parameters are expressed in a mathematically closed form and therefore the influence of individual parameters on pollutant concentration can be easily examined [25]. Also, the analytical solutions make it easy to obtain asymptotic behaviours of the solutions, that are usually difficult to obtain through numerical calculations.

The fundamental solution for the fractional diffusion-wave equation in one space dimension was obtained by Mainardi [18]. Agrawal [1] analyzed the

* Corresponding author.

Email address: f.liu@qut.edu.au (F. Liu).

fractional diffusion-wave equation in a half-line and a bounded domain $[0, L]$. Gorenflo and Mainardi [7] studied the time-fractional, spatially one-dimensional diffusion-wave equation on the spatial half-line with zero initial conditions. They considered the Dirichlet and Neumann boundary conditions and proved that the Dirichlet-Neumann map is given by a time-fractional differential operator whose order is one-half the order of the time-fractional derivative. Mainardi and Paradisi [20] employed the time-fractional diffusion-wave equation to study the propagation of stress waves in viscoelastic media relevant to acoustics and seismology. Several initial and boundary-value problems for the time-fractional diffusion-wave equation were investigated by Povstenko [27–31]. He also considered the generalized Cattaneo-type equations with time- and space-fractional derivatives [32]. Qi and Jian also presented the exact solution of the fractional Cattaneo equation [33]. Golbabai and Sayevand [5] investigated fractional advection dispersion equation by means of the homotopy perturbation method with consideration of a promising scheme to calculate nonlinear terms. Zhou and Jiao [45] discussed the nonlocal Cauchy problem for the fractional evolution equations in an arbitrary Banach space and obtained various criteria on the existence and uniqueness of mild solutions. By applying fractional calculus to model the behavior of cells and tissues, Magin et al. [17] unraveled the inherent complexity of individual molecules and membranes in a way that leads to an improved understanding of the overall biological function and behavior of living systems. Shakhmurov and Shahmurova [35] studied the boundary value problems for linear and nonlinear degenerate elliptic differential operator equations of a second order. By using these results the existence, uniqueness and the maximal regularity of boundary value problems for nonlinear degenerate parabolic differential operator equations are established. Wang et al. [42] developed a fast characteristic finite difference method for the efficient solution of space-fractional transient advection-diffusion equations in one space dimension. Meerschaert et al. [22] derived explicit strong solutions and stochastic analogues for distributed-order time-fractional diffusion equations on bounded domains, with Dirichlet boundary conditions.

Gejji and Bhalekar [4] considered a multi-term fractional diffusion-wave equation along with the homogeneous/non-homogeneous boundary conditions and this equation has been solved using the method of separation of variables. Based on orthogonal polynomials of the Laguerre type, Stojanovic [39] found solutions for the diffusion-wave problem in one dimension with n -term time fractional derivatives whose orders belong to the intervals $(0, 1)$, $(1, 2)$ and $(0, 2)$, respectively. Equations in this paper can be solved in one-dimension by the method of the approximation of the tempered convolution. This method transfers the diffusion-wave problem into the corresponding infinite system of linear algebraic equations through the coefficients, which are uniquely solvable under some relations between the coefficients with index zero. Luchko [15] considered the initial-boundary-value problems for the generalized time-fractional diffusion equation over an open bounded domain. Based on an appropriate

maximum principle, Luchko [15] derived some *a priori* estimates for the solution and then its uniqueness is established using the Fourier method of the separation of variables. The time-dependent components of the solution are given in terms of the multinomial Mittag-Leffler function.

Time domain wave-equations for lossy media obeying a frequency power-law were discussed by Kelly et al. [10]. Mathematically, the power-law frequency dependence of the attenuation coefficient cannot be modeled with standard dissipative partial differential equations with integer-order derivatives. Kelly et al. [10] considered the Szabo wave equation with the n -term time fractional derivatives whose orders belong to the intervals $(1, 2]$ or $[2, 3)$, respectively. Treeby et al. [40] proposed modeling power-law absorption and dispersion for acoustic propagation using the fractional Laplacian.

However, many practical problems also involve space-fractional derivatives, in particular, Riesz fractional derivatives [37,38]. The main purpose of this paper is to derive the analytical solutions of the multi-term time-space Caputo-Riesz fractional advection-diffusion equations (MT-TSCR-FADE) with nonhomogeneous Dirichlet boundary conditions. The multi-term time fractional derivatives are defined in the Caputo sense, whose orders belong to the intervals $[0, 1]$, $[1, 2]$ and $[0, 2]$, respectively. These are known as multi-term time fractional diffusion terms, multi-term time fractional wave terms and multi-term time fractional mixed diffusion-wave terms.

In this paper, the derived analytical solutions are based on Luchko's Theorem [13–15]. We proposed some new techniques, such as a spectral representation of the fractional Laplacian operator and the equivalent relationship between the fractional Laplacian operator and the Riesz fractional derivative, that enabled derivation of the analytical solutions for the multi-term time-space Caputo-Riesz fractional advection-diffusion equations. As far as we are aware there are no research papers in the published literature written on this topic.

The remainder of this paper is organized as follows. In section 2, we give some relevant definitions and lemmas. The analytical solutions of the MT-TSCR-FADE with multi-term time-fractional diffusion terms and multi-term time-fractional wave terms are derived in sections 3 and 4, respectively. The analytical solution of the MT-TSCR-FADE with multi-term time-fractional mixed wave-diffusion terms is given in section 5, and the conclusions of the work are presented in section 6.

2 Background Theory

For convenience, we introduce the following definitions and theorems that are used throughout this paper.

The MT-TSCR-FADE can be written in the following form:

$$P(D_t)u(x, t) = k_\beta \frac{\partial^\beta u(x, t)}{\partial |x|^\beta} + k_\gamma \frac{\partial^\gamma u(x, t)}{\partial |x|^\gamma} + f(x, t), \quad (1)$$

in a finite domain $0 < x < L, t > 0$. Here x and t are the space and time variables, and k_β, k_γ are positive constants, $0 < \beta < 1, 1 < \gamma \leq 2$.

The operator $P(D_t)u(x, t)$ is defined as

$$P(D_t)u(x, t) = \left(D_t^\alpha + \sum_{i=1}^s a_i D_t^{\alpha_i} \right) u(x, t), \quad (2)$$

$0 \leq \alpha_s < \dots < \alpha_1 < \alpha \leq 1$ or $1 \leq \alpha_s < \dots < \alpha_1 < \alpha \leq 2$ or $0 \leq \alpha_s < \dots < \alpha_{h_0-1} \leq 1 < \alpha_{h_0} < \dots < \alpha_1 < \alpha \leq 2$, and $a_i \in \mathbb{R}, i = 1, \dots, n, n \in \mathbb{N}, D_t^{\alpha_i}$ is a Caputo fractional derivative of order α_i with respect to t , which is defined as follows:

$$D_t^{\alpha_i} f(t) = \begin{cases} \frac{1}{\Gamma(m-\alpha_i)} \int_0^t \frac{f^{(m)}(\tau)}{(t-\tau)^{1+\alpha_i-m}} d\tau, & m-1 < \alpha_i < m, \\ \frac{d^m}{dt^m} f(t), & \alpha_i = m \in \mathbb{N}. \end{cases} \quad (3)$$

The space fractional derivatives $\frac{\partial^\beta u(x, t)}{\partial |x|^\beta}$ and $\frac{\partial^\gamma u(x, t)}{\partial |x|^\gamma}$ are Riesz space-fractional derivatives of order β, γ , respectively, that are defined by Gorenflo and Mainardi [6] in Definition 1.

Definition 1 (See [6]) The Riesz fractional operator for $n \in \mathbb{N}, n-1 < \beta \leq n$, on a finite interval $0 \leq x \leq L$ is defined as

$$\frac{\partial^\beta}{\partial |x|^\beta} u(x, t) \equiv D^\beta u(x, t) = -c_\beta ({}_0D_x^\beta + {}_x D_L^\beta) u(x, t), \quad (4)$$

where the coefficient $c_\beta = \frac{1}{2 \cos(\frac{\beta\pi}{2})}, \beta \neq 1$, and

$${}_0D_x^\beta u(x, t) = \frac{1}{\Gamma(n-\beta)} \frac{\partial^n}{\partial x^n} \int_0^x \frac{u(\xi, t) d\xi}{(x-\xi)^{\beta+1-n}}, \quad (5)$$

$${}_x D_L^\beta u(x, t) = \frac{(-1)^n}{\Gamma(n - \beta)} \frac{\partial^n}{\partial x^n} \int_x^L \frac{u(\xi, t) d\xi}{(x - \xi)^{\beta+1-n}}, \quad (6)$$

are the left-side and right-side Riemann-Liouville fractional derivatives, respectively. In particular, when $n = 2$, ${}_0 D_x^\beta u(x, t) = {}_x D_L^\beta u(x, t) = \frac{\partial^2}{\partial x^2} u(x, t)$.

If $0 \leq \alpha_s < \dots < \alpha_1 < \alpha \leq 1$, Equation (1) is a generalized time and space fractional advection-diffusion equation with multi-term time fractional diffusion terms. When we discuss its solutions, the initial condition is given as follows:

$$u(x, 0) = \phi(x), \quad 0 < x < L. \quad (7)$$

In this case, if $a_i = 0, i = 1, 2, \dots, s, \alpha = 1$, then equation (1) becomes a space fractional advection-diffusion equation with the Riesz space fractional derivatives, which was discussed by Yang et al. [43].

If $1 \leq \alpha_s < \dots < \alpha_1 < \alpha \leq 2$, then, equation (1) becomes a generalized time and space fractional advection-diffusion equation with multi-term time fractional wave terms. In this case, the initial conditions are given as follows:

$$u(x, 0) = \phi(x), \quad u_t(x, 0) = \psi(x), \quad 0 < x < L. \quad (8)$$

If $0 \leq \alpha_s < \dots < \alpha_{h_0} \leq 1 < \alpha_{h_0-1} < \dots < \alpha \leq 2$, then equation (1) becomes a generalized time and space fractional advection-diffusion equation, which we refer to as a multi-term time fractional mixed wave-diffusion equation. In this case, the initial conditions are also given by (8).

Definition 2. (See [13]) A real or complex-valued function $f(x)$, $x > 0$, is said to be in the space C_α , $\alpha \in \mathbb{R}$, if there exists a real number p , $p > \alpha$, such that

$$f(x) = x^p f_1(x),$$

where $f_1(x)$ is in $C[0, \infty)$.

Definition 3. (See [13]) A function $f(x)$, $x > 0$, is said to be in the space C_α^m , $m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$, if and only if $f^m \in C_\alpha$.

Lemma 1. For a function $u(x, t)$ defined on a finite domain $[0, L; 0, T]$, and $u(0, t) = u(L, t) = 0$, the following equality holds:

$$-(-\Delta)^{\frac{\beta}{2}} u(x, t) = -c_\beta [{}_0 D_x^\beta u(x, t) + {}_x D_L^\beta u(x, t)] = \frac{\partial^\beta}{\partial |x|^\beta} u(x, t), \quad (9)$$

where the coefficient $c_\beta = \frac{1}{2 \cos(\frac{\beta\pi}{2})}$, $0 < \beta < 1$, or $1 < \beta < 2$, and the space fractional derivative $\frac{\partial^\beta u(x,t)}{\partial |x|^\beta}$ is a Riesz space-fractional derivative of order β respectively.

Proof. See [43].

Definition 4. (See [13]) A multivariate Mittag-Leffler (n dimensional) function is defined as

$$E_{(a_1, \dots, a_n), b}(z_1, \dots, z_n) \equiv \sum_{k=0}^{\infty} \sum_{\substack{l_1 + \dots + l_n = k \\ l_1 \geq 0, \dots, l_n \geq 0}} \frac{k!}{l_1! \dots l_n!} \frac{\prod_{i=1}^n z_i^{l_i}}{\Gamma(b + \sum_{i=1}^n a_i l_i)}, \quad (10)$$

in which $b > 0$, $a_i > 0$, $|z_i| < \infty$, $i = 1, \dots, n$.

Lemma 2. Let $\mu > \mu_1 > \dots > \mu_n \geq 0$, $m_i - 1 < \mu_i \leq m_i$, $m_i \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $d_i \in \mathbb{R}$, $i = 1, \dots, n$. The initial value problem

$$\begin{cases} (D^\mu y)(x) - \sum_{i=1}^n d_i (D^{\mu_i} y)(x) = g(x), \\ y^{(k)}(0) = c_k \in \mathbb{R}, \quad k = 0, \dots, m-1, \quad m-1 < \mu \leq m, \end{cases} \quad (11)$$

where the function $g(x)$ is assumed to lie in C_{-1} , if $\mu \in \mathbb{N}$, and in C_{-1}^1 , if $\mu \notin \mathbb{N}$, and the unknown function $y(x)$ is to be determined in the space C_{-1}^m , has the solution

$$y(x) = y_g(x) + \sum_{k=0}^{m-1} c_k u_k(x), \quad x \geq 0, \quad (12)$$

where

$$y_g(x) = \int_0^x t^{\mu-1} E_{(\cdot), \mu}(t) g(x-t) dt, \quad (13)$$

and

$$u_k(x) = \frac{x^k}{k!} + \sum_{i=l_k+1}^n d_i x^{k+\mu-\mu_i} E_{(\cdot), k+1+\mu-\mu_i}(x), \quad k = 0, \dots, m-1, \quad (14)$$

fulfills the initial conditions $u_k^{(l)}(0) = \delta_{kl}$, $k, l = 0, \dots, m-1$. The function

$$E_{(\cdot), \beta}(x) = E_{\mu-\mu_1, \dots, \mu-\mu_n, \beta}(d_1 x^{\mu-\mu_1}, \dots, d_n x^{\mu-\mu_n}) \quad (15)$$

is a particular case of the multivariate Mittag-Leffler function (see [13]) and the natural numbers $l_k, k = 0, \dots, m - 1$, are determined from the condition

$$\begin{cases} m_{l_k} \geq k + 1, \\ m_{l_k+1} \leq k. \end{cases} \quad (16)$$

In the case $m_i \leq k, i = 1, \dots, n$, we define $l_k := 0$, and if $m_i \geq k + 1, i = 1, \dots, n$, then $l_k := n$.

Proof. See [13].

Definition 5. [9] Suppose the Laplacian $(-\Delta)$ has a complete set of orthonormal eigenfunction φ_n corresponding to eigenvalues λ_n^2 on a bounded region \mathcal{D} , i.e., $(-\Delta)\varphi_n = \lambda_n^2\varphi_n$ on a bounded region \mathcal{D} ; $\mathcal{B}(\varphi) = 0$ on $\partial\mathcal{D}$, where $\mathcal{B}(\varphi)$ is one of the standard three homogeneous boundary conditions. Let

$$\mathcal{F} = \{f = \sum_{n=1}^{\infty} c_n \varphi_n, c_n = \langle f, \varphi_n \rangle, \sum_{n=1}^{\infty} |c_n|^2 |\lambda_n|^\beta < \infty, \}, \quad (17)$$

then for any $f \in \mathcal{F}$, $(-\Delta)^{\frac{\beta}{2}}$ is defined by

$$(-\Delta)^{\frac{\beta}{2}} f = \sum_{n=1}^{\infty} c_n (\lambda_n)^\beta \varphi_n. \quad (18)$$

Lemma 3. (See [9]) Suppose the one-dimensional Laplacian $(-\Delta)$ defined with Dirichlet boundary condition at $x = 0$ and $x = L$ has a complete set of orthonormal eigenfunctions φ_n corresponding to eigenvalues λ_n^2 on a boundary region $\Omega = [0, L]$, if $(-\Delta)\varphi_n = \lambda_n^2\varphi_n$ then, the eigenvalues are given by $\lambda_n^2 = \frac{n^2\pi^2}{L^2}$ for $n = 1, 2, \dots$ and the corresponding eigenfunctions are nonzero scalar multiples of $\varphi_n = \sqrt{\frac{2}{L}} \sin(\frac{n\pi x}{L})$.

3 Analytical solution of the MT-TSCR-FADE with multi-term time fractional diffusion terms

In this section, we consider the analytical solution of the MT-TSCR-FADE (1) with multi-term time fractional diffusion terms. In this case, $0 \leq \alpha_s < \dots < \alpha_1 < \alpha \leq 1$ in Eq. (1). The equation satisfies initial condition (7) and

the following Dirichlet boundary conditions

$$\begin{cases} u(0, t) = \mu_1(t), & t \geq 0, \\ u(L, t) = \mu_2(t), & t \geq 0, \end{cases} \quad (19)$$

where $\mu_1(t), \mu_2(t)$ are nonzero smooth functions with order-one continuous derivatives. In order to solve the problem with nonhomogeneous boundary conditions, we firstly transform the nonhomogeneous condition into a homogeneous boundary condition. Let

$$u(x, t) = W(x, t) + V(x, t), \quad (20)$$

where

$$V(x, t) = \frac{\mu_2(t) - \mu_1(t)}{L}x + \mu_1(t), \quad (21)$$

satisfies the boundary conditions

$$\begin{cases} V(0, t) = \mu_1(t), \\ V(L, t) = \mu_2(t). \end{cases} \quad (22)$$

Using Lemma 1 and substituting (20) into (1), the function $W(x, t)$ is the solution of the following homogeneous problem:

$$\begin{cases} P(D_t)W(x, t) + k_\beta(-\Delta)^{\frac{\beta}{2}}W(x, t) \\ + k_\gamma(-\Delta)^{\frac{\gamma}{2}}W(x, t) = f_1(x, t), & 0 < x < L, t > 0, \\ W(x, 0) = \phi_1(x), & 0 \leq x \leq L, \\ W(0, t) = W(L, t) = 0, & t \geq 0, \end{cases} \quad (23)$$

with

$$\begin{cases} f_1(x, t) = -P(D_t)V(x, t) + k_\beta \frac{\partial^\beta}{\partial |x|^\beta} V(x, t) + k_\gamma \frac{\partial^\gamma}{\partial |x|^\gamma} V(x, t) \\ + f(x, t), \\ \phi_1(x) = \phi(x) - \frac{\mu_2(0) - \mu_1(0)}{L}x - \mu_1(0), \end{cases} \quad (24)$$

using Definition 1 and applying the fractional derivative formulas.

We assume that the solution of the homogeneous equation in (23) has the form

$$W(x, t) = X(x)T(t). \quad (25)$$

Substituting this expression for $W(x, t)$ in (23), we obtain the Sturm-Liouville problem

$$\begin{aligned} -k_\beta(-\Delta)^{\frac{\beta}{2}}X(x) - k_\gamma(-\Delta)^{\frac{\gamma}{2}}X(x) + \lambda X(x) &= 0, \\ X(0) = 0, \quad X(L) &= 0, \end{aligned} \quad (26)$$

and a fractional ordinary linear differential equation with Caputo derivatives for $T(t)$, namely

$$P(D_t)T(t) + \lambda T(t) = 0, \quad (27)$$

$$T(0) = \bar{\phi}_0(x), \quad \frac{dT(0)}{dt} = \bar{\phi}_1(x), \quad (28)$$

where the parameter λ is a positive constant.

The Sturm-Liouville problem given by (26) and (27) has eigenvalues

$$\lambda_n^2 = \frac{n^2\pi^2}{L^2}, \quad n = 1, 2, \dots \quad (29)$$

and corresponding eigenfunctions

$$X_n(x) = \sin\left(\frac{n\pi x}{L}\right), \quad n = 1, 2, \dots \quad (30)$$

Based on Definition 4 and Lemma 3, we set

$$W(x, t) = \sum_{n=1}^{\infty} c_{n1}(t) \sin\left(\frac{n\pi x}{L}\right). \quad (31)$$

In order to determine $c_{n1}(t)$, we expand $f_1(x, t)$ as a Fourier series using the eigenfunction $\{\sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right)\}_{n=1}^{\infty}$ as

$$f_1(x, t) = \sum_{n=1}^{\infty} f_{n1}(t) \sin\left(\frac{n\pi x}{L}\right), \quad (32)$$

where

$$f_{n1}(t) = \frac{2}{L} \int_0^L f_1(x, t) \sin\left(\frac{n\pi x}{L}\right) dx. \quad (33)$$

Using Definition 3 and substituting (31) and (32) into (23), we obtain

$$\begin{aligned} P(D_t) \sum_{n=1}^{\infty} c_{n1}(t) \sin\left(\frac{n\pi x}{L}\right) + k_{\beta} \sum_{n=1}^{\infty} c_{n1}(t) \lambda_n^{\beta} \sin\left(\frac{n\pi x}{L}\right) \\ + k_{\gamma} \sum_{n=1}^{\infty} c_{n1}(t) \lambda_n^{\gamma} \sin\left(\frac{n\pi x}{L}\right) = \sum_{n=1}^{\infty} f_{n1}(t) \sin\left(\frac{n\pi x}{L}\right), \end{aligned} \quad (34)$$

where the λ_n are given in Lemma 3.

Then, we obtain

$$P(D_t)c_{n1}(t) + k_{\beta}c_{n1}(t)\lambda_{n1}^{\beta} + k_{\gamma}c_{n1}(t)\lambda_{n1}^{\gamma} = f_{n1}(t). \quad (35)$$

Using the initial condition

$$W(x, 0) = \phi_1(x), \quad 0 \leq x \leq L, \quad (36)$$

we have

$$\sum_{n=1}^{\infty} c_{n1}(0) \sin\left(\frac{n\pi x}{L}\right) = \phi_1(x) \quad 0 < x < L, \quad (37)$$

and therefore

$$c_{n1}(0) = \frac{2}{L} \int_0^L \phi_1(x) \sin\left(\frac{n\pi x}{L}\right) dx \quad n = 1, 2, \dots. \quad (38)$$

According to Lemma 2, we obtain

$$c_{n1}(t) = \int_0^t G_{\alpha}^m(\tau) \tau^{\alpha-1} f_{n1}(t - \tau) d\tau + c_{n1}(0) u_0(t), \quad (39)$$

where

$$G_\eta^n(t) = E_{(\nu_1, \dots, \nu_s, \alpha), \eta}(-a_1 t^{\nu_1}, \dots, -a_s t^{\nu_s}, -\kappa_n t^\alpha), \quad (40)$$

$$\kappa_n = k_\beta \lambda_n^\beta + k_\gamma \lambda_n^\gamma, \quad \nu_i = \alpha - \alpha_i, \quad i = 1, \dots, s, \quad (41)$$

$$u_0(t) = 1 - \kappa_n t^\alpha G_{1+\alpha}^n(t). \quad (42)$$

Thus, the solution of problem (23) is given by

$$W(x, t) = \sum_{n=1}^{\infty} c_{n1}(t) \sin\left(\frac{n\pi x}{L}\right), \quad (43)$$

where $c_{n1}(t)$ is given in (39).

Finally, the analytical solution of the MT-TSCR-FADE (1) with multi-term time fractional diffusion terms is

$$u(x, t) = \frac{(\mu_2(t) - \mu_1(t))x}{L} + \mu_1(t) + \sum_{n=1}^{\infty} \left[\int_0^t G_\alpha^n(\tau) \tau^{\alpha-1} f_{n1}(t - \tau) d\tau + c_{n1}(0) u_0(t) \right] \sin\left(\frac{n\pi x}{L}\right). \quad (44)$$

4 Analytical solution of the MT-TSCR-FADE with multi-term time-fractional wave terms

In this section, we consider the analytical solution of the MT-TSCR-FADE with multi-term time-fractional wave terms. In this case, $1 \leq \alpha_s < \dots < \alpha_1 < \alpha \leq 2$ in Eq. (1). The boundary and initial conditions are (19) and (8), respectively.

In a similar manner as presented in Section 3, in order to solve the problem with nonhomogeneous boundary conditions, we firstly transform the nonhomogeneous problem into a homogeneous problem.

Using Lemma 1 and substituting (20) into (1), the function $W(x, t)$ is the solution of the problem:

$$\begin{cases} P(D_t)W(x, t) + k_\beta (-\Delta)^{\frac{\beta}{2}} W(x, t) \\ + k_\gamma (-\Delta)^{\frac{\gamma}{2}} W(x, t) = f_1(x, t), & 0 < x < L, t > 0, \\ W(x, 0) = \phi_1(x), \frac{\partial W(x, 0)}{\partial t} = \psi_1(x), & 0 \leq x \leq L, \\ W(0, t) = W(L, t) = 0, & t \geq 0, \end{cases} \quad (45)$$

with

$$\begin{cases} f_1(x, t) = -P(D_t)V(x, t) + k_\beta \frac{\partial^\beta}{\partial |x|^\beta} V(x, t) + k_\gamma \frac{\partial^\gamma}{\partial |x|^\gamma} V(x, t) \\ \quad + f(x, t), \\ \phi_1(x) = \phi(x) - \frac{\mu_2(0) - \mu_1(0)}{L} x - \mu_1(0), \\ \psi_1(x) = \psi(x) - \frac{\mu'_2(0) - \mu'_1(0)}{L} x - \mu'_1(0). \end{cases} \quad (46)$$

Similarly, based on Definition 4, we set

$$W(x, t) = \sum_{n=1}^{\infty} c_{n2}(t) \varphi_n(x) \quad (47)$$

to obtain

$$P(D_t)c_{n2}(t) + k_\beta c_{n2}(t) \lambda_{n1}^\beta + k_\gamma c_{n2}(t) \lambda_{n1}^\gamma = f_{n1}(t). \quad (48)$$

Using the initial conditions

$$W(x, 0) = \phi_1(x), \quad \frac{\partial W(x, 0)}{\partial t} = \psi_1(x), \quad 0 \leq x \leq L, \quad (49)$$

we have

$$\begin{cases} \sum_{n=1}^{\infty} c_{n2}(0) \sin\left(\frac{n\pi x}{L}\right) = \phi_1(x) & 0 < x < L, \\ \sum_{n=1}^{\infty} c'_{n2}(0) \sin\left(\frac{n\pi x}{L}\right) = \psi_1(x) & 0 < x < L, \end{cases} \quad (50)$$

and obtain

$$\begin{cases} c_{n2}(0) = \frac{2}{L} \int_0^L \phi_1(x) \sin\left(\frac{n\pi x}{L}\right) dx & n = 1, 2, \dots, \\ c'_{n2}(0) = \frac{2}{L} \int_0^L \psi_1(x) \sin\left(\frac{n\pi x}{L}\right) dx & n = 1, 2, \dots. \end{cases} \quad (51)$$

For each value of n , (48) and (51) constitutes a fractional initial value problem.

According to Lemma 2, we obtain

$$c_{n2}(t) = \int_0^t G_\alpha^m(\tau) \tau^{\alpha-1} f_{n1}(t - \tau) d\tau + c_{n2}(0) u_0(t) + c'_{n2}(0) u_1(t), \quad (52)$$

where

$$u_0(t) = 1 - \kappa_n t^\alpha G_{1+\alpha}^n(t), \quad (53)$$

$$u_1(t) = t - \kappa_n t^{1+\alpha} G_{2+\alpha}^n(t). \quad (54)$$

Thus, the solution of problem (45) is given by

$$W(x, t) = \sum_{n=1}^{\infty} c_{n2}(t) \sin\left(\frac{n\pi x}{L}\right), \quad (55)$$

where $c_{n2}(t)$ is given in (52).

Finally, the analytical solution of the MT-TSCR-FADE (1) with multi-term time fractional wave terms is

$$u(x, t) = \sum_{n=1}^{\infty} \left[\int_0^t G_\alpha^n(\tau) \tau^{\alpha-1} f_{n1}(t-\tau) d\tau + c_{n2}(0)u_0(t) + c'_{n2}(0)u_1(t) \right] \\ \times \sin\left(\frac{n\pi x}{L}\right) + \frac{(\mu_2(t) - \mu_1(t))x}{L} + \mu_1(t). \quad (56)$$

5 Analytical solutions of the MT-TSCR-FADE with multi-term time-fractional mixed diffusion-wave terms

In this section, we consider the analytical solution of the MT-TSCR-FADE with multi-term time-fractional mixed diffusion-wave terms. In this case, $0 \leq \alpha_s < \dots < \alpha_{h_0} \leq 1 < \alpha_{h_0-1} < \dots < \alpha_1 < \alpha \leq 2$ in Eq. (1). The boundary and initial conditions are (19) and (8), respectively.

Similarly to the analysis in Section 4, based on Definition 4, we set

$$W(x, t) = \sum_{n=1}^{\infty} c_{n3}(t) \varphi_n(x) \quad (57)$$

to obtain

$$P(D_t)c_{n3}(t) + k_\beta c_{n3}(t)\lambda_n^\beta + k_\gamma c_{n3}(t)\lambda_n^\gamma = f_{n1}(t). \quad (58)$$

According to Lemma 2, we have

$$c_{n3}(t) = \int_0^t G_\alpha^n(\tau) \tau^{\alpha-1} f_{n1}(t-\tau) d\tau + c_{n3}(0)u_0(t) + c'_{n3}(0)u_1(t), \quad (59)$$

where

$$c_{n3}(0) = \frac{2}{L} \int_0^L \phi_1(x) \sin\left(\frac{n\pi x}{L}\right) dx \quad n = 1, 2, \dots, \quad (60)$$

$$c'_{n3}(0) = \frac{2}{L} \int_0^L \psi_1(x) \sin\left(\frac{n\pi x}{L}\right) dx \quad n = 1, 2, \dots, \quad (61)$$

$$u_0(t) = 1 - \kappa_n t^\alpha G_{1+\alpha}^n(t), \quad (62)$$

$$u_1(t) = t - \sum_{i=h_0}^s a_i t^{1+\nu_i} G_{2+\nu_i}^n(t) + \kappa_n t^{1+\alpha} G_{2+\alpha}^n(t). \quad (63)$$

Thus, the solution of problem (45) is

$$W(x, t) = \sum_{n=1}^{\infty} c_{n3}(t) \sin\left(\frac{n\pi x}{L}\right), \quad (64)$$

where $c_{n3}(t)$ is given in (59).

Finally, the analytical solution of the MT-TSCR-FADE (1) with multi-term time-fractional diffusion-wave terms is

$$u(x, t) = \sum_{n=1}^{\infty} \left\{ \int_0^t G_\alpha^n(\tau) \tau^{\alpha-1} f_{n1}(t-\tau) d\tau + c_{n3}(0) u_0(t) + c'_{n3}(0) u_1(t) \right\} \sin\left(\frac{n\pi x}{L}\right) + \frac{(\mu_2(t) - \mu_1(t))x}{L} + \mu_1(t). \quad (65)$$

Special case 1: The time-space fractional telegraph equation:

$$D_t^\alpha u(x, t) + a D_t^{\frac{\alpha}{2}} u(x, t) = k_\beta (-\Delta)^{\frac{\beta}{2}} + k_\gamma (-\Delta)^{\frac{\gamma}{2}} + f(x, t), \quad (66)$$

with initial conditions

$$u(x, 0) = \phi(x), \quad u_t(x, 0) = \psi(x), \quad 0 < x < L, \quad (67)$$

and boundary conditions:

$$u(0, t) = \mu_1(t), \quad t \geq 0, \quad u(L, t) = \mu_2(t), \quad t \geq 0, \quad (68)$$

where $1 < \alpha \leq 2$, $0 < \beta < 1$ and $1 < \gamma \leq 2$.

From (65), we obtain the solution of equations (66)-(67):

$$u(x, t) = \sum_{n=1}^{\infty} \left\{ \int_0^t G_{\alpha}^n(\tau) \tau^{\alpha-1} f_{n1}(t - \tau) d\tau + c_{n3}(0) u_0(t) + c'_{n3}(0) u_1(t) \right\} \sin\left(\frac{n\pi x}{L}\right) + \frac{(\mu_2(t) - \mu_1(t))x}{L} + \mu_1(t) \quad (69)$$

where

$$u_0(t) = 1 + \kappa_n t^{\alpha} G_{1+\alpha}^n(t), \quad (70)$$

$$u_1(t) = t - at^{1+\frac{\alpha}{2}} G_{2+\frac{\alpha}{2}}^n(t) + \kappa_n t^{1+\alpha} G_{2+\alpha}^n(t), \quad (71)$$

$$G_{\alpha}^n(\tau) = E_{(\frac{\alpha}{2}, \alpha), \alpha}(-a\tau^{\frac{\alpha}{2}}, -\kappa_n \tau^{\alpha}), \quad (72)$$

$$G_{1+\alpha}^n(t) = E_{(\frac{\alpha}{2}, \alpha), 1+\alpha}(-at^{\frac{\alpha}{2}}, -\kappa_n t^{\alpha}), \quad (73)$$

$$G_{2+\frac{\alpha}{2}}^n(t) = E_{(\frac{\alpha}{2}, \alpha), 2+\frac{\alpha}{2}}(-at^{\frac{\alpha}{2}}, -\kappa_n t^{\alpha}), \quad (74)$$

$$G_{2+\alpha}^n(t) = E_{(\frac{\alpha}{2}, \alpha), 2+\alpha}(-at^{\frac{\alpha}{2}}, -\kappa_n t^{\alpha}), \quad (75)$$

here $f_{n1}(t - \tau)$, G_{η}^n , κ_n , $c_{n3}(0)$, $c'_{n3}(0)$ and $E_{(\frac{\alpha}{2}, \alpha), \eta}$ are as given in (33), (40), (41), (60), (61) and (10), respectively.

Special case 2: The time-fractional telegraph equation:

When $\kappa_{\beta} = 0$, $\kappa_{\gamma} = k$, $\gamma = 2$ in equation (66), we obtain the solution of the time-fractional telegraph equation:

$$u(x, t) = \sum_{n=1}^{\infty} \left\{ \int_0^t E_{(\frac{\alpha}{2}, \alpha), 1+\alpha}(-a\tau^{\frac{\alpha}{2}}, -\frac{k\pi^2 n^2}{L^2} \tau^{\alpha}) \tau^{\alpha-1} f_{n1}(t - \tau) d\tau + c_{n3}(0) u_0(t) + c'_{n3}(0) u_1(t) \right\} \sin\left(\frac{n\pi x}{L}\right) + \frac{(\mu_2(t) - \mu_1(t))x}{L} + \mu_1(t) \quad (76)$$

where

$$u_0(t) = 1 - \frac{k\pi^2 n^2}{L^2} t^{\alpha} E_{(\frac{\alpha}{2}, \alpha), 1+\alpha}(-a\tau^{\frac{\alpha}{2}}, -\frac{k\pi^2 n^2}{L^2} \tau^{\alpha}),$$

$$u_1(t) = t - at^{1+\frac{\alpha}{2}} E_{(\frac{\alpha}{2}, \alpha), 2+\frac{\alpha}{2}}(-a\tau^{\frac{\alpha}{2}}, -\frac{k\pi^2 n^2}{L^2} \tau^{\alpha}) - \frac{k\pi^2 n^2}{L^2} t^{1+\alpha} E_{(\frac{\alpha}{2}, \alpha), 2+\alpha}(-a\tau^{\frac{\alpha}{2}}, -\frac{k\pi^2 n^2}{L^2} \tau^{\alpha}). \quad (77)$$

This result is in accord with the result obtained in [2]

6 Conclusions

In this paper, we have proposed some new analytic techniques to solve three types of MT-TSCR-FADE with nonhomogeneous Dirichlet boundary conditions. By using Luchko's Theorem and the equivalent relationship between the Laplacian operator and the Riesz fractional derivative, we have used the spectral representation of the fractional Laplacian operator to derive analytical solutions. As far as we are aware there are no research papers in the published literature written on this topic. The methods and techniques discussed in this paper can also be applied to solve other types of multi-term fractional partial differential equations with fractional Laplacian, such as, the Szabo wave equation with the n -term time fractional derivatives and the power-law absorption and dispersion equation for acoustic propagation with the fractional Laplacian.

ACKNOWLEDGEMENTS

The first author wishes to acknowledge the funding received by the China Scholarship Council to support this work. They also thank QUT for the resources made available to them doing the work. The authors wish to thank the referees for their constructive comments and suggestions.

References

- [1] O.P. Agrawal, Solution for a fractional diffusion-wave equation defined in a bounded domain, *Nonlinear Dynamics*, 29, (2002), 145-155.
- [2] J. Chen, F. Liu, V. Anh, Analytical solution for the time-fractional telegraph equation by the method of separating variables, *J. Math. Anal. Appl.*, 338, (2008), 1364-1377.
- [3] V.V. Gafiychuk and B.Yo. Datsko, Pattern formation in a fractional reaction diffusion system, *Physica*, A365, (2006), 300-306.
- [4] V. Daftardar-Gejji and S. Bhalekar, Boundary value problems for multi-term fractional differential equations, *J. Math. Anal. Appl.* 345 (2008) 754-765.
- [5] A. Golbabai and K. Sayevand, Analytical modelling of fractional advection-dispersion equation defined in a bounded space domain, *Math. Comput. Modelling*, 53, (2011), 1708-1718.
- [6] R. Gorenflo, F. Mainardi, Random walk models for space-fractional diffusion processes, *Fract. Calc. Appl. Anal.*, 1, (1998), 167-191.
- [7] R. Gorenflo and F. Mainardi, Signalling problem and Dirichlet-Neumann map for time-fractional diffusion-wave equation, *Matimyas Matematika*, 21, (1998), 109-118.

- [8] R. Hilfer,(Ed.) Applications of Fractional Calculus in Physics, World Scientific, Singapore,(2000).
- [9] M. Ilić, F. Liu, I. Turner and V. Anh, Numerical approximation of a fractional-in-space diffusion equation, *Fractional Calculus and Applied Analysis*, 8(3), (2005), 323–341.
- [10] J.F. Kelly and R.J. McGough, Analytical time-domain Greens functions for poerw-law media, *J. Acoust. Soc. Am.*, 124, (2008), 2861-2872.
- [11] F. Liu, V. Anh and I.Turner, Numerical solution of the space Fokker-Planck equation, *J. Comput. Appl. Math.* 166, (2004), 290-219.
- [12] F. Liu, P. Zhuang, V. Anh, I. Turner and K. Burrage, Stability and convergence of difference methods for the space-time fractional advection-diffusion equation, *Appl. Math. Comp.*, 191, (2007), 12-20.
- [13] Y. Luchko and R. Gorenflo, An operational method for solving fractional differential equations with the Caputo derivatives, *Acta Math. Vietnam*, 24, (1999), 207-233.
- [14] Y. Luchko, Maximum Principle for the fractional differential equations and its application, *Proceedings of FDA'10, The 4th IFAC Workshop Fractional Differentiation and Applications*. Badajoz, Spain, October 18-22, 2010.
- [15] Y. Luchko, Initial-boundary-value problems for the generalized multi-term time-fractional diffusion equation, *J. Math. Anal. Appl.*, 374, (2011), 538-548.
- [16] A.A. Kilbas, N.H.M. Srivastava and J.J. Trujillo, *Theory and applications of fractional differential equation*, Elsevier, Amsterdam, 2006.
- [17] R. Magin, M.D. Ortigueira, I. Podlubny and J. Trujillo, On the fractional signals and systems, *Signal Processing*, 91, (2011), 350-371.
- [18] F. Mainardi, The fundamental solutions for the fractional diffusion-wave equation, *Appl. Math. Lett.*, 9, (1996), 23-28.
- [19] F. Mainardi, Fractional calculus: some basic problems in continuum and statistical mechanics. In: A. Carpinteri and F.Mainardi (eds.): *Fractals and Fractional Calculus in Continuum Mechanics*, (1997), 291-348, Springer, New York.
- [20] F. Mainardi and P. Paradisi, Fractional diffusive waves, *J. Comput. Acoustics*, 9, (2001), 1417-1436.
- [21] M. Meerschaert and C. Tadjeran, Finite difference approximations for fractional advection-dispersion flow equation, *J. Comput. Appl. Math.*, 172, (2004), 65-77.
- [22] M. Meerschaert, E. Nane and P. Vellaisamy, Distributed-order fractional diffusions on bounded domains, *J. Math. Anal. Appl.*, 379, (2011), 216-228.
- [23] R. Metzler and J. Klafter, The random walk's guide to anomalous diffusion:a fractional dynamics approach, *Phys. Rep.*, 339, (2000), 1-77.

- [24] R. Metzler and J. Klafter, The restaurant at the end of the random walk: recent developments in the description of anomalous transport by fractional dynamics, *J. Phys. A: Math. Gen.*, 37, (2004), 161-208.
- [25] Y.S. Park, J. Jin. Baik, Analytical solution of the advection-diffusion equation for a ground-level finite area source, *Atmospheric Environment*, 42, (2008), 9063C9069.
- [26] I. Podlubny, *Fractional differential equations*, Academic Press, San Diego, 1999.
- [27] Y.Z. Povstenko, Two-dimensional axisymmetric stresses exerted by instantaneous pulses and sources of diffusion in an infinite space in a case of time-fractional diffusion equation, *Int. J. Solids Struct.*, 44, (2009), 2324-2348.
- [28] Y.Z. Povstenko, Signaling problem for time-fractional diffusion-wave equation in a half-plane, *Fractional Calculus Appl. Anal.*, 11, (2008), 329-352.
- [29] Y.Z. Povstenko, Fundamental solution to three dimensional diffusion-wave equation and associated diffusive stresses, *Chaos, Solitons Fractals*, 36, (2008), 961-972.
- [30] Y.Z. Povstenko, Analysis of fundamental solutions to fractional diffusion-wave equation in polar coordinates, *Sci. Issues Jan Dlugosz Univ. Czestochowa, Mathematics*, XIV, (2009), 97-104.
- [31] Y.Z. Povstenko, Signaling problem for time-fractional diffusion-wave equation in a half-space in the case of angular symmetry, *Nonlinear Dynamics*, 55, (2010), 593-605.
- [32] Y.Z. Povstenko, Theories of thermoelasticity based on space-time-fractional Cattaneo-type equations, *Proceedings of FDA'10. The 4th IFAC Workshop Fractional Differentiation and Its Applications*, Badajoz, Spain, October 18-20, 2010.
- [33] H. Qi and, X. Jiang, Solutions of the space-time fractional Cattaneo diffusion equation, *Physica A*, 390 (2011) 1876-1883.
- [34] J.P. Roop, Computational aspects of FEM approximation of fractional advection dispersion equation on boundary domains in \mathbb{R}^2 , *Journal of computational and Applied Mathematics*, 193(1), (2006), 243-268.
- [35] V.B. Shakhmurov and A. Shahmurova, Nonlinear abstract boundary value problems modelling atmospheric dispersion of pollutants, *Nonlinear Analysis: Real World Applications*, 11, (2010), 932-951.
- [36] S. Shen, F. Liu and V. Anh, Fundamental solution and discrete random walk model for a time-space fractional diffusion equation of distributed order, *J. Appl. Math. and Computing*, 28, (2008), 147-164.
- [37] S. Shen, F. Liu, V. Anh, and I. Turner, The fundamental solution and numerical solution of the Riesz fractional advection-dispersion equation, *IMA J. Appl. Math.*, 73, (2008), 850 - 872.

- [38] S. Shen, F. Liu and V. Anh, Numerical approximations and solution techniques for the space-time Riesz-Caputo fractional advection-diffusion equation, *Numerical Algorithm*, 56(3), (2011), 383-403.
- [39] M. Stojanovic, Numerical method for solving diffusion-wave phenomena, *Journal of Computational and Applied Mathematics*, Volume 235, Issue 10, 15 March (2011), 3121-3137.
- [40] B.E. Treeby and B.T. Cox, Modeling power law absorption and dispersion for acoustic propagation using the fractional Laplacian, *J. Acoust Soc. Am.*, 127, (2010), 2741-2748.
- [41] V.V. Uchaikin, *Method of fractional derivatives*, Artishok, UI'janovask, 2008 (in Russian).
- [42] K. Wang and H. Wang, A fast characteristic finite difference method for fractional advection-diffusion equations, *Advances in Water Resources*, 34 (2011), 810-816.
- [43] Q. Yang, F. Liu, I. Turner, Numerical methods for fractional partial differential equations with Riesz space fractional derivatives, *Applied Mathematical modeling*, 34, (2010), 200-218.
- [44] Y. Zhang, D.A. Benson and D.M. Reeves, Time and space nonlocalities underlying fractional-derivative models: Distinction and literature review of field applications, *Adv. Water Resour.*, 32 (2009) 561-581.
- [45] Y. Zhou and F. Jiao, Nonlocal Cauchy problem for fractional evolution equations, *Nonlinear Analysis: Real World Applications*, 11, (2010), 4465-4475.