



ELSEVIER

Veterinary Microbiology 86 (2002) 313–324

**veterinary
microbiology**

www.elsevier.com/locate/vetmic

The effect of concurrent infections with *Pasteurella multocida* and *Ascaridia galli* on free range chickens

C. Dahl^a, A. Permin^{a,*}, J.P. Christensen^a, M. Bisgaard^a,
A.P. Muhairwa^a, K.M.D. Petersen^a, J.S.D. Poulsen^b,
A.L. Jensen^b

^aDepartment of Veterinary Microbiology, The Royal Veterinary and Agricultural University,
4 Stigbøjlen, 1870 Frederiksberg C, Denmark

^bCentral Laboratory, The Royal Veterinary and Agricultural University, 3 Grønnegårdsvej,
1870 Frederiksberg C, Denmark

Received 20 August 2001; received in revised form 3 January 2002; accepted 26 January 2002

Abstract

Pasteurella multocida and *Ascaridia galli* are observed with high prevalences in free range chickens in Denmark, but the impact is unknown. A study was carried out to examine the interaction between *A. galli* and *P. multocida* in chickens and the impact on production.

Five groups, each with 20 18-week-old Lohmann Brown chickens were infected. Group 1 was orally infected with 1000 ± 50 embryonated *A. galli* eggs. Group 2 received 10^4 cfu *P. multocida* intratracheally. Group 3 was infected with *A. galli* and subsequently with *P. multocida*. Group 4 was infected with *P. multocida* followed by *A. galli*. Group 5 was the control. The study ran for 11 weeks where clinical manifestations, weight gain and egg production were recorded. Excretion of *P. multocida* was determined on individual basis and blood smears were made for differential counts. At the end of the study pathological lesions and the number of adult worms, larvae and eggs in the faeces were recorded.

The birds were more severely affected when infected with both pathogens compared to single infections with *A. galli* or *P. multocida*, respectively. A lower weight gain and egg production was observed with dual infections. *A. galli* infection followed by a secondary *P. multocida* infection resulted in more birds with pathological lesions and continued *P. multocida* excretion.

In conclusion a negative interaction between *A. galli* and *P. multocida* was observed and it is postulated that free range chickens are at higher risk of being subjected to outbreaks of fowl cholera when they are infected with *A. galli*. © 2002 Elsevier Science B.V. All rights reserved.

Keywords: Chicken; Epidemiology; *Pasteurella multocida*; *Ascaridia galli*; Concurrent infections; Interactions

* Corresponding author. Tel.: +45-3528-3763; fax: +45-3528-3762.

E-mail address: ape@kvl.dk (A. Permin).

1. Introduction

During the last five decades commercial poultry production has almost exclusively been based on confined production systems with a high level of biosecurity. However, during the last 10 years consumers have shown an increased interest in products from free range and organic production systems. Table egg production from these systems has increased significantly and now accounts for almost one-third of all table eggs produced in Denmark (Anon., 2000). In contrast to confined production systems in which high levels of biosecurity can be obtained, free range production is characterised by a high risks of acquiring infections. A recent study in Denmark has shown that the flock prevalence of *Ascaridia galli* is 100% in free range/organic systems, whereas it is only about 25% in confined indoor deep litter production systems (Permin et al., 1999). *Pasteurella multocida* infections are also common in non-confined poultry production systems (Christensen et al., 1999). Recent investigations showed that about 80% of the diagnosed cases of pasteurellosis in Danish poultry appeared in flocks that had access to outdoor areas (Christensen et al., 1998). Studies on the interactions between *A. galli* and *Escherichia coli* in chickens have indicated that subclinical *A. galli* infections, may have an immunosuppressive effect, allowing *E. coli* to establish itself (Permin et al., 2002). Likewise, a synergistic effect of migrating *Ascaris suum* larvae and *E. coli* was observed in piglets (Adedeji et al., 1989) and of salmonellosis on subsequent infections with *A. suum* in adult swine (Wade and Gaafar, 1981). A similar effect may exist for *P. multocida* infections in *A. galli* infected flocks. The aim of the present study was to examine interactions between *A. galli* and *P. multocida* and their impact on disease manifestations, weight gain, egg production and excretion of *A. galli* and *P. multocida*.

2. Materials and methods

2.1. Experimental animals

One hundred 16-week-old Lohmann Brown female chickens with no history of fowl cholera in previous or present flocks on the farm were used for each experiment. The chickens were randomly distributed into five groups each consisting of 20 individually identified animals. Two weeks before the onset of the study all birds were examined for the presence of *P. multocida* by taking swabs from the cloaca, dissolving the swabs in saline water and subsequently injecting white balb/cJ mice with the solution as described by Muhairwa et al. (2000). Furthermore, a faecal sample from each bird was examined for the presence of parasites using the McMaster concentration techniques (Permin and Hansen, 1998). Regardless of the parasite status all the birds were treated with flubendazole (7 mg/kg live weight per 3 days) to ensure no interference from a low grade parasite infection.

2.2. Housing

Each group was placed in a 6 m² cleaned and disinfected henhouse with nesting boxes and perches. The floor was covered with sawdust. The birds received 11–14 h of light

(10–15 lux) a day depending on the age of the chickens (Anon., 2001). The birds had free access to water and commercial feed containing 18% protein (layers mesh).

2.3. Experimental design

The experiment was a $2 \times 2 + 1$ cohort study (Thrusfield, 1995) in which group 1 was infected with *A. galli*, group 2 with *P. multocida*, while groups 3 and 4 were infected with both pathogens but in different orders. The group 5 represented uninfected controls. The study had a duration of 11 weeks.

2.4. Infections

A. galli eggs were prepared according to Permin et al. (1997b). A clinical isolate of *P. multocida*, serotype A3 originating from an outbreak of fowl cholera among wild birds was used for inoculation (Christensen et al., 1998). Chickens in group 1 were infected with 1000 ± 50 infective *A. galli* eggs orally. In group 2 all birds were infected with 10^4 cfu *P. multocida* intratracheally. Group 3 was initially infected with 1000 ± 50 infective *A. galli* eggs per bird and 4 days later with 10^4 cfu *P. multocida* intratracheally. Group 4 was first infected with 10^4 cfu *P. multocida* intratracheally and 4 days later with 1000 ± 50 infective *A. galli* eggs. Group 5 was sham infected orally with normal tap water (Table 1).

2.5. Parameters

All groups were observed on a daily basis for clinical signs. Eggs were collected, counted and weighed on a daily basis while the birds were weighed only once a week. Blood samples were taken 1 week before the start of the experiment at the age of 17 weeks and subsequently at the age of 18, 21, 24, 28 and 29 weeks. Blood smears for differential counts of the white blood cells were made according to Permin and Hansen (1998). At the age of 20, 22, 25 and 28 weeks, swabs were taken from the cloaca and pharynx of each chicken to examine for the presence of *P. multocida* as described by Muhairwa et al. (2000).

At the end of the experiment all surviving birds were killed and subjected to post-mortem examinations for pathological changes. From each spleen swabs were streaked onto blood agar plates and checked for growth of *P. multocida*. The intestines were removed to check for the presence of adult and larval forms of *A. galli*. The intestines were

Table 1
Experimental design including group size, infection and infection doses

Group	Group size	Infection	Dosage
1	20	<i>A. galli</i>	1000 ± 50 eggs
2	20	<i>P. multocida</i>	10^4 cfu
3	20	<i>A. galli</i> + <i>P. multocida</i>	1000 ± 50 eggs + 10^4 cfu
4	20	<i>P. multocida</i> + <i>A. galli</i>	10^4 cfu + 1000 ± 50 eggs
5	20	Control	Sham infected with water

opened in a longitudinal section and rinsed with water over two sieves, the smallest with a mesh aperture of 38 µm. The sieve retentate was examined for adult worms and larvae using a stereo microscope at 40× magnification. Faecal samples were taken from the cloaca and parasite eggs were counted using a concentrated McMaster technique (Permin and Hansen, 1998).

2.6. Statistical methods

Data was stored and statistical analyses were carried out using the statistical software program GraphPad Prism[®] version 3.0 from GraphPad Software (2000). Analysis of variance, chi-square and unpaired *t*-tests were used for normal distributed data. Otherwise a Kruskal–Wallis test was used. The level of significance was set as $p < 0.05$.

3. Results

3.1. Pretrial examinations

The swabbings showed that the birds were free of infections with *P. multocida*. Furthermore, the birds were free of infections with parasites, i.e. *A. galli*, *Heterakis gallinarum* and *Capillaria* spp. (Permin et al., 1999).

3.2. Clinical signs

Depression was observed in the birds infected with *P. multocida*, i.e. groups 2–4 during the first week after infection with *P. multocida*. After inoculation the chickens became quiet and passive and their feathers ruffled. Feed and water consumption were also depressed. The chickens were most affected the first week after the infection, after which an improvement was seen. Differences in clinical signs between groups only infected with *P. multocida* and groups infected with both *P. multocida* and *A. galli* were not observed. Increasing mortality, starting shortly after inoculations with *P. multocida*, was observed for groups 2 (three birds), 3 (six birds) and 4 (three birds) and continued up to 4 weeks after inoculation. These chickens tested positive for *P. multocida* in spleen and liver when swabs were streaked onto blood agar plates. No indications of clinical disease were observed in groups 1 and 5.

3.3. Weight gain

The average weight gain for all groups are shown in Fig. 1. For groups 2–4 a drop in weight gain was observed immediately after they had been infected with *P. multocida*. Both the control group and the *A. galli* group showed weight gains during the whole observation period and had the highest weight gain while the doubly infected groups had the lowest. However, the groups were not easily distinguished from each other. During the first week a significant difference was seen between groups infected with *P. multocida* and groups which had not been infected with bacteria. Groups 1 and 3 were only infected with *A. galli*




Fig. 1. Weight gain for all five groups ($N = 20$ per group) until week 10.

at this time and could not be separated from the control group ($p > 0.05$). Groups infected with *P. multocida* could not be differentiated from each other at this point of the trial.

During the second week group 3 dropped in the weight gain immediately after infection with *P. multocida*. At this time no significant differences between the three groups infected with *P. multocida* were observed. In the final week a significant difference between the *A. galli* group and the two double infected groups as well as between the control group and the three groups infected with *P. multocida* were observed ($p < 0.05$).

3.4. Eggs

All groups started laying within 3 days after the first infection at the age of 18 weeks (Table 2). Differences observed between groups as to the age at start of lay were not significant ($p > 0.05$). However, at the end of the experiment a significant number (at 90% level) of hens were not in lay in groups 2 (three birds), 3 (six birds) and 4 (three birds) ($p < 0.06$). Significant differences were observed in the numbers of eggs per hen per week for groups 1 and 5 (the *A. galli* group and the control group) compared to groups 2–4

Table 2
Egg production data for all groups

	Group 1: <i>A. galli</i>	Group 2: <i>P. multocida</i>	Group 3: <i>A. galli</i> + <i>P. multocida</i>	Group 4: <i>P. multocida</i> + <i>A. galli</i>	Group 5: control
Age at the first egg in the group (days)	130	130	131	130	132
Age when all the chickens were in lay (days)	145	158	157	159	143
Hens remaining at week 29	20	17	14	17	20
Hens in lay of remaining animals (in %) at end of trial (week 29)	20 (100%)	14 (82%)	8 (57%)*	14 (82%)	20 (100%)

* Significantly lower (at 90% level) compared to other groups ($p < 0.06$).




Fig. 2. Number of eggs per hen per week.

infected with *P. multocida* (Fig. 2). Finally, significant differences in single egg weights between groups were not observed during the whole experiment.

3.5. Differential counts

Double infected groups had the lowest percentage of lymphocytes. Significant differences between the control group and the *A. galli* group were not observed. The *A. galli* group however differed significantly from the double infected groups during the weeks 3, 6 and 10. Significant differences between groups 2 and 3 as well as between the control group and the double infected groups were noticed in week 3. Statistically significant differences were also seen between groups 2 and 3, 3 and 4 as well as 3 and 5 in week 6 and between groups 4 and 5 in week 10.

Double infected groups had the highest percentage of heterophil granulocytes compared to other groups. A statistical significant difference in per cent heterophil granulocytes was observed between the *A. galli* group and the control group versus the double infected groups as well as between groups 2 and 3.

Differences in per cent eosinophilic granulocytes were observed between groups 1 and 4 in first week. At the time of inoculation differences between groups 1–3 and the control group as well as between groups 2 and 4 were observed. In the sixth week group 1 separated itself from the rest of the groups significantly ($p < 0.05$). In the final week there were statistical differences between the *A. galli* group and groups 2, 4 and 5 as well as between groups 3 and 4.

Basophilic granulocytes and monocytes were only represented in small percentages making it difficult to evaluate the results.

3.6. Excretion of *P. multocida*

The excretion of *P. multocida* in the various groups is given in Table 3. Birds from groups 2 and 3 inoculated with *P. multocida* and *A. galli* and *P. multocida*, respectively, did not

Table 3
Chickens testing positive for *P. multocida* during experiment

Week	Group 1: <i>A. galli</i> ^a		Group 2: <i>P. multocida</i>		Group 3: <i>A. galli</i> + <i>P. multocida</i>		Group 4: <i>P. multocida</i> + <i>A. galli</i>		Group 5: control ^a	
	Pos./all	%	Pos./all	%	Pos./all	%	Pos./all	%	Pos./all	%
2	370/10	0	9/17	53	11/17	65*	6/18	33	370/10	0
4	370/10	0	1/17	6	1/14	7	1/17	6	370/10	0
7	370/10	0	0/17	0	1/14	7*	0/17	0	370/10	0
10	370/10	0	1/17	6	2/14	14*	0/17	0	370/10	0

^a Swabs were taken from four animals and pooled into one sample, i.e. five samples from each group before inoculation into mice.

* Significantly more ($p < 0.05$).

eliminate the infection. Significantly more birds were positive in group 3 infected with both the pathogens. Birds in group 4 infected with *P. multocida* and subsequently with *A. galli* eliminated the infection 7 weeks after the infection. There were no positive samples for *P. multocida* in groups 1 and 5 throughout the trial.

3.7. Bacteriological investigation of the spleens

All spleens sampled negative for *P. multocida* at the end of the trial.

3.8. *A. galli* eggs, larvae and worms

In Table 4 EPG (eggs per gram of faeces), larvae and adult worm counts are shown. Fecundity (EPG/adult female worm) and percentage of infected chickens are also given. Parasites were only recorded in the *A. galli* infected groups, i.e. groups 1, 3 and 4. A chi-square test showed significant lower percentage of infected birds in group 3 compared to groups 1 and 4. A significant lower EPG was seen in group 4 compared to group 1 ($p < 0.05$). No differences were seen in EPG neither between groups 1 and 3 nor between

Table 4
Parasitological findings at slaughter (\pm S.D.)

	Group 1: <i>A. galli</i>	Group 2: <i>P. multocida</i>	Group 3: <i>A. galli</i> + <i>P. multocida</i>	Group 4: <i>P. multocida</i> + <i>A. galli</i>	Group 5: control
EPG	652.6 \pm 802.0	–	676.7 \pm 845.1	137.5 \pm 232.8*	–
Larvae	0.6 \pm 1.3	–	0.2 \pm 0.6	0.2 \pm 0.7	–
Female worms	4.4 \pm 5.4	–	4.2 \pm 1.1	2.1 \pm 2.4	–
Male worms	1.6 \pm 2.5	–	1.4 \pm 0.6	1.2 \pm 2.4	–
Larvae + worms	6.6 \pm 7.4	–	5.9 \pm 5.9	3.6 \pm 4.7	–
Fecundity (EPG/females)	178.0 \pm 2480	–	90.0 \pm 123.0	61.0 \pm 97.0	–
Percentage infected chickens	100	–	79*	94	–

* Significantly lower ($p < 0.05$).

Table 5

Total number of chickens (in %) with pathological findings at end of experiment

	Group 1: <i>A. galli</i>	Group 2: <i>P. multocida</i>	Group 3: <i>A. galli</i> + <i>P. multocida</i>	Group 4: <i>P. multocida</i> + <i>A. galli</i>	Group 5: control
Poor body condition	–	–	1 (5)	1 (5)	–
Dehydrated	–	–	–	1 (5)	–
Pneumonia/lung necrosis	–	5 (25)	4 (20)	3 (15)	–
Pleuritis	–	3 (15)	2 (10)	1 (5)	–
Aerosacculitis	–	–	–	1 (5)	–
Liver necrosis	–	4 (20)	–	–	–
Spleen necrosis	–	3 (15)	–	1 (5)	–
Uremia	–	–	1 (5)	–	–
Peritonitis	–	–	–	–	1 (5)
Bursitis presternalis	–	–	–	–	1 (5)
Not in lay	–	1 (5)	4 (20)	1 (5)	–
Mortality due to <i>P. multocida</i>	–	3 (15)	6 (30)	3 (15)	–
Total number of chickens with pathological changes	0/20 (0)	5/17 (29)	8/14 (57)*	4/17 (23)	2/20 (10)

* Significantly more chickens with pathological changes compared to other groups ($p < 0.05$).

groups 3 and 4 ($p > 0.05$). Statistical significantly differences between groups 1, 3 and 4 in larvae or worms were not shown ($p > 0.05$). Neither were significant changes in fecundity seen with *t*-test ($p > 0.05$) although a trend showed lower fecundity in groups 3 and 4.

3.9. Pathological changes

Table 5 shows the pathological lesions observed at the end of the experiment. Lesions were seen in the three groups infected with *P. multocida*. In the double infected group 3 significantly more chickens had pathological lesions compared to the group only infected with *P. multocida*. No pathological lesions due to *P. multocida* were observed in the *A. galli* group or the control group.

4. Discussion and conclusions

A pronounced effect of concurrent infections with *P. multocida* and *A. galli* was observed on weight gain, egg production, differential counts, excretion of *P. multocida*, parasitic findings at post-mortem and pathological changes.

Clinical signs associated with the *P. multocida* infected groups included depression, anorexia, ruffled feathers and mortality as the main features as reported previously by Rhoades (1964), Nagi et al. (1990) and Petersen et al. (2001) and was apparently not enhanced by the parasitic infection. The *A. galli* group and the control group showed, as expected, no clinical signs. Low level infections with *A. galli* may not cause clinical symptoms as reported by Permin et al. (1998b). Mortality occurred in the three groups infected with *P. multocida*, but no statistical differences were seen between the groups.

The observed mortality was considered low, probably due to the low *P. multocida* dose (Matsumoto et al., 1991; Toth and Siegel, 1993; Petersen et al., 2001). A trend however indicated that the group infected with *A. galli* followed by *P. multocida* had a higher mortality. These findings are in accordance with findings by Permin et al. (2002) who demonstrated that chickens infected with *E. coli* and *A. galli* reacted more severely compared to chickens only infected with *E. coli*.

Weight gain was depressed in the group infected with *P. multocida* as a single infection and in the two groups additionally infected with *A. galli* either as a primary or secondary infection. Acute *P. multocida* infections are known to cause anorexia and subsequent weight depression as it has also been described for *A. galli* (Rhoades, 1964; Ikeme, 1971; Nagi et al., 1990; Permin et al., 1998a). The two double infected groups never reached a weight as high as the other groups, and at the end of the experiment especially the weight gain of the group first infected with *A. galli* and subsequently with *P. multocida* was lower compared to all the other groups. This could be due to an immune suppressive effect of *A. galli* supporting the establishment of *P. multocida* in the host (Roepstorff et al., 1999).

All groups started laying within 1–3 days after infection, e.g. at the age of 19 weeks, indicating that neither of the infections had an impact on point of lay (Anon., 2001). Significantly fewer eggs were laid per hen per week in the groups infected with *P. multocida* compared to the *A. galli* group and the control group. This indicates that the acute *P. multocida* infection had a negative effect on the egg production which is in accordance with previous observations (Campi et al., 1990). Infections with *A. galli* may decrease egg production in layers (Permin et al., 1998a); however, the missing effect observed in this study might be related to the relatively high age of chickens when infected (Kerr, 1955).

In the double infected groups the percentage of lymphocytes were significantly lower compared to the *A. galli* and the control groups. These results suggest that the immune system was depressed in double infected groups compared to groups only infected with one pathogen and the control group. Furthermore, the results indicate that a primary *A. galli* infection followed by a secondary *P. multocida* infection affect the birds more seriously than a single bacterial infection. When the birds are infected with *P. multocida* first and *A. galli* last, an inhibiting effect on the immune system may be seen, but it cannot be distinguished from a single bacterial infection, underlining that the order of multiple infections might have an important impact on the outcome. Likewise the double infected groups had higher heterophilic counts compared to the other groups suggesting that the parasitic infection enhances the heterophilic response. The double infected groups did not seem to have an eosinophilic reaction towards the parasites as they did not distinguish themselves from the control group, whereas the *A. galli* infected group had a significantly higher eosinophilic count coincident with the prepatent time of *A. galli* (Kerr, 1955; Ikeme, 1971; Permin et al., 1998b).

A decreasing number of *P. multocida* carriers was seen in all *P. multocida* infected groups during the experiment. However, chickens in groups 2 and 3 did not clear themselves totally for the infection as the birds in group 4. An infection dose of approximately 10^4 cfu per chicken is considered low (Matsumoto et al., 1991; Toth and Siegel, 1993). The low infections dose was chosen to allow us study the epidemiology

of *P. multocida* in relation to *A. galli* infections. *P. multocida* carriers were observed even 10 weeks after inoculation, especially in the group primary infected with *A. galli*. Under free range conditions, where parasitic infections always are present (Permin et al., 1998b), chickens remain carriers of *P. multocida* for a longer period, having epidemiological implications. Cannibalism (i.e. stress) seems to have a similar effect on the carrier status of *P. multocida* infected chickens (Permin et al., unpublished results).

At the end of the experiment none of the spleens tested positive for *P. multocida*. This is in agreement with previous studies by Petersen et al. (2001) who showed that *P. multocida* was eliminated from the spleens of chickens within 48 h. In contrast, all *A. galli* infected groups harboured larval and adult stages of *A. galli* at slaughter. Significantly fewer chickens were infected in the group first infected with *A. galli* and subsequently with *P. multocida*. Generally, the establishment rate in individual chickens depends on the size of the infection dose (Ackert et al., 1931; Permin et al., 1997a). Furthermore, in experimental and natural infections all chickens become infected (Permin et al., 1998b, 1999). The reason for some of the animals being negative in this study is not known, but might possibly be related to a cross-over effect of the concurrent infection. A recent paper by Pritchard and Brown, 2001 has indicated that although cellular response mechanisms of bacteria and parasites are related to each their pathway (Th2 for parasites and Th1 for bacteria) there is a balance between the two pathways. Thus a parasite infection might favour the Th2 cell development and indirectly suppress the establishment of bacteria or vice versa. A significantly lower EPG was observed in the group infected first with *P. multocida* and then with *A. galli*. This could imply that *P. multocida* has a suppressive effect on the livelihood and establishment of the parasite, since a trend showing a lower worm burden in this group was observed. This might also be explained as a result of an enhanced immune defence triggered by the *P. multocida* infection.

Significantly more birds were observed with pathological lesions in the group first infected with *A. galli* and subsequently with *P. multocida*. This might be a result of an inhibited defence system in the birds first infected with the parasite, facilitating the establishment of bacteria in the host (Pritchard and Brown, 2001).

Health aspects are rarely discussed in relation to welfare of free range chickens (Craig and Swanson, 1994). However, free range chickens are often subclinically infected with *A. galli* (Matter and Oester, 1989; Permin et al., 1998b, 1999) and *P. multocida* (Christensen et al., 1998). It is therefore postulated that free range chickens are at higher risk of being subjected to serious outbreaks of fowl cholera when they are permanently infected with *A. galli*. The fact that poultry in organic systems are not treated against gastrointestinal parasites contributes to this postulation.

In conclusion a negative interaction between *A. galli* and *P. multocida* was observed. Combined infections had a more severe effect on the birds compared to single infections for reasons which remains to be investigated. Carrier animals were also common in this group. Even a subclinical *A. galli* infection seems to enhance the establishment of *P. multocida*. If *A. galli* represents a secondary infection, the clinical picture was quite equal to that of the single *P. multocida* infection. However, *P. multocida* affected *A. galli* establishment. Since multiple infections are common under non-confined conditions basic studies on interaction of infections are greatly needed. Understanding the mechanisms behind the interactions will improve the possibilities of prevention.

Acknowledgements

We would like to thank Rikke Frahm Lundvig, Pernille Ginsbo, Pia Mortensen, Stina Holm, Niels Midtgaard, Jørgen Olesen, Johnny Jensen, Thomas Bernau Kristensen and Rene Bülow for technical assistance during the experiment. Financial support by the Council for Development Research (Danida) through the project “Multiple infections in free range poultry” is highly appreciated.

References

- Ackert, J.E., Graham, G.L., Nolf, L.O., Porter, D.A., 1931. Quantitative studies on the administration of variable numbers of nematode eggs (*Ascaridia lineata*) to chickens. *Trans. Am. Microsc. Soc.* 50, 206–214.
- Adedeji, S.O., Ogunba, E.O., Dipeolu, O.O., 1989. Synergistic effect of migrating *Ascaris* larvae and *Escherichia coli* in piglets. *J. Helminthol.* 26, 17–32.
- Anon., 2000. The Danish Poultry Council. Statistics on www.poultry.dk.
- Anon., 2001. Layer Management Program. Lohmann Brown. Lohmann Tierzucht.
- Campi, T.W., Carpenter, T.E., Hird, D.W., Snipes, K.P., Hirsh, D.C., 1990. Fowl cholera in California multiplier breeder turkeys: 1985–1986. *Avian Dis.* 34, 928–933.
- Christensen, J.P., Dietz, H.H., Bisgaard, M., 1998. Phenotypic and genotypic characters of isolates of *Pasteurella multocida* obtained from back-yard poultry and from two outbreaks of avian cholera in avifauna in Denmark. *Avian Pathol.* 27, 373–381.
- Christensen, J.P., Petersen, K.M.D., Hansen, H.C., Bisgaard, M., 1999. Forekomst af fjerkrækolera i dansk avifauna og fjerkræproduktion—mulige smitemæssige sammenhænge. *Dansk Vet. Tidsskr.* 82, 342–346.
- Craig, J.V., Swanson, J.C., 1994. Review: welfare perspectives on hens kept for egg production. *Poult. Sci.* 73, 921–938.
- Ikeme, M.M., 1971. Effects of different levels of nutrition and continuing dosing of poultry with *Ascaridia galli* eggs on the subsequent development of parasite populations. *Parasitology* 63, 233–250.
- Kerr, K.B., 1955. Age of chickens and the rate of maturation of *Ascaridia galli*. *J. Parasitol.* 41, 233–235.
- Matsumoto, M., Strain, J.G., Engel, H.N., 1991. The fate of *Pasteurella multocida* after intratracheal inoculation into turkeys. *Poult. Sci.* 70, 2259–2266.
- Matter, F., Oester, H., 1989. Hygiene and welfare implications of alternative husbandry systems for laying hens. In: Faure, J.M., Mills, A.D. (Eds.), *Proceedings of the Third European Symposium on Poultry Welfare*, Tours, France.
- Muhairwa, A.P., Christensen, J.P., Bisgaard, M., 2000. Investigations on the carrier rate of *Pasteurella multocida* in healthy commercial poultry flocks and flocks affected by fowl cholera. *Avian Pathol.* 29, 133–142.
- Nagi, A.A., Youssef, M.S., Mousa, S., Bayoumi, A.H., 1990. Clinico-pathological and bacteriological studies on avian pasteurellosis. *Assiut Vet. Med. J.* 22, 88–94.
- Permin, A., Hansen, J.W., 1998. Epidemiology, Diagnosis and Control of Poultry Parasites. Food and Agriculture Organization of the United Nations, Rome, Italy, pp. 74–105, 111–115.
- Permin, A., Bojesen, M., Nansen, P., Bisgaard, M., Frandsen, F., Pearman, M., 1997a. *Ascaridia galli* populations in chickens following single infections with different dose levels. *Parasitol. Res.* 83, 614–617.
- Permin, A., Pearman, M., Nansen, P., Bisgaard, M., Frandsen, F., 1997b. An investigation on different media for embryonation of *Ascaridia galli* eggs. *Helminthologia* 34, 75–79.
- Permin, A., Nansen, P., Bisgaard, M., Frandsen, F., 1998a. *Ascaridia galli* infections in free range layers fed different levels of protein. *Brit. Poult. Sci.* 39, 441–445.
- Permin, A., Nansen, P., Bisgaard, M., Frandsen, F., 1998b. Investigations on the infection and transmission of *Ascaridia galli* in free range chickens kept at different stocking rates. *Avian Pathol.* 27, 382–389.
- Permin, A., Bisgaard, M., Frandsen, F., Pearman, M., Nansen, P., Kold, J., 1999. The prevalence of gastrointestinal helminths in different poultry systems. *Brit. Poult. Sci.* 40, 439–443.
- Permin, A., Christensen, J.P., Bisgaard, M., 2002. Consequences of concurrent *Ascaridia galli* and *Escherichia coli* infections in chickens. *Avian Pathol.*, in press.

- Petersen, K.M.D., Christensen, J.P., Permin, A., Bisgaard, M., 2001. Virulence studies in domestic and free-living birds of a clone of *Pasteurella multocida* ssp. *multocida* obtained from an outbreak of fowl cholera in wild birds. *Avian Pathol.* 30, 27–31.
- Pritchard, D.I., Brown, A., 2001. Is *Necator americanus* approaching a mutualistic symbiotic relationship with humans? *Trends Parasitol.* 17, 169–172.
- Rhoades, K.R., 1964. The microscopic lesions of acute fowl cholera in mature chickens. *Avian Dis.* 8, 658–665.
- Roepstorff, A., Nørgaard-Nielsen, G., Permin, A., Simonsen, H.B., 1999. Male behaviour and male hormones in *Ascaridia galli* infected hens. In: Proceedings of the XIX Symposium of the Scandinavian Society for Parasitology, Iceland, May 8–11.
- Thrusfield, M., 1995. *Veterinary Epidemiology*, 2nd Edition. Blackwell Science Publications, Oxford.
- Toth, T.E., Siegel, P.B., 1993. Cellular defence of the avian respiratory system: dose–response relationship and duration of response in intratracheal stimulation of avian respiratory phagocytes by a *Pasteurella multocida* bacterin. *Avian Dis.* 37, 756–762.
- Wade, W.F., Gaafar, S.M., 1981. Effects of salmonellosis on subsequent infections with *Ascaris suum* in swine. *Vet. Parasitol.* 8, 309–317.