

Kaolin protects olive fruits from *Bactrocera oleae* (Gmelin) infestations unaffected olive oil quality

Enzo Perri, Nino Iannotta, Innocenzo Muzzalupo, Anna Russo, Maria Anna Caravita, Massimiliano Pellegrino, Attilio Parise, Paolo Tucci
C.R.A. Institute for Olive Growing - 87036 Rende, Cosenza, ITALY
Department of Chemicals, University of Calabria - 87036 Rende, Cosenza, ITALY

Abstract: The efficacy of the processed kaolin “Surround WP” to control olive fruit fly, *Bactrocera oleae* Gmelin, field infestations was investigated in east Calabria. The preliminary results showed that fruit infestation levels were significantly reduced on kaolin-treated trees compared with untreated trees. The promising results of these experiments points to the feasibility of using particle film technology composed of a non-toxic material, to avoid olive fly damage as an alternative to the applications of rotenone in organic orchards. Finally, kaolin treatment unaffected the nutritional and sensory quality parameters of the corresponding virgin olive oils obtained by a laboratory scale olive mill, thus satisfying the present quality requirements.

Key words: Kaolin, *Bactrocera oleae*, olive, olive oil quality.

Introduction

Alternatives to synthetic chemical pesticide as well as organic pesticide are today required to reduce the heart pollution, improve the safety condition and to ensure safer foods. Kaolin-based particle film is a new tool to protect fruits (Mazor M. And Erez A., 2004) and olives (Sour and Makee, 2004) from fly infestations. Therefore, the efficacy of the kaolin to control olive fruit fly, *Bactrocera oleae* Gmelin, field infestations was investigated in east Calabria.

Materials and Methods

A kaolin-based particle film formulation commercially available under the trade name Surround WP Crop Protectant (Engelhard Corporation, Iselin, NJ, USA) was used. The effectiveness of the kaolin to control *B. Oleae* was investigated in two olive groves located at Strongoli (KR), in the east part of Calabria. The experiment was carried out during two years on bearing 50-year-old oil olive trees (cv. Carolea). In each trial, 20 trees were chosen for their homogeneity in terms of canopy and production. Treatments consisted of single trees arranged in a completely random design with three replications. Rotenone at the first grove and Surround WP at the second grove were sprayed one time in 2004 year and two times in 2005 year, on 22 september and 14 and 28 september, respectively. The olive oils were obtained by a laboratory scale olive mill (Toscana Enologica Mori, Tavernelle Val di Pesa, SI, Italy). Olive oils were analysed by EU Official Methods (Reg. CE 1989/2003).

Paper presented at 2nd European Meeting of the IOBS/WPRS Study group Integrated Protection of olive crops, Florence, 26-28 October 2005; To be published in *IOBC/WPRS Bulletin*.




Figure 1. Evaluation of *Bactrocera oleae* mean percentage total and active infestations on olive trees sprayed with kaolin particle film and unsprayed control.

Results and discussion

There was a significant difference between kaolin based particle film and control treatments in the mean percentage of olive attacked by *Bactrocera oleae* in the second year on 5 october, the day of the olive harvest and olive oil production, seven days after the second spray (Table 1).

Table 1. Mean percentages of olive fruits active infestations

Mean percentage	
Rotenone active infestation	20a
Control active infestation	20a
Kaolin active infestation	14b
Tukey test, P<0.05	

Moreover, the promising results of these experiments points to the feasibility of using particle film technology composed of a non-toxic material, to avoid olive fly damage as an alternative to the applications of rotenone in organic orchards. Finally, kaolin treatment unaffected the nutritional and sensory quality parameters of the corresponding virgin olive oils obtained by a laboratory scale olive mill, thus satisfying the present nutritional (table 2) and sensory (Figures 2 and 3) quality requirements.

Table 2. Mean of the main olive oil quality parameters

	Kaolin	Crontrol
Tocopherol	250.60	260.21
Total phenols	193.43	170.13
C18:1	75.868	75.625
C16:0	13.513	14.565
Delta K	-0.002	-0.003
K 270	0.094	0.113
K 232	1.667	1.603
Peroxide Index	5	5
Free Acidity	0.428	0.428




Figure 2. The sensory profiling of olive oil from Kaolin treated olives.




Figure 3. The sensory profiling of olive oil from control olives.

References

- Caleca, V., Lo Verde, G., Lo Verde, V. & Rizzo, R. 2005: Il caolino e la mosca: come controllare quella dell'olivo e quella della frutta. *Bioagricoltura (maggio-giugno 2005)*: 28-31.
- Caleca, V., Lo Verde, V. & Palumbo Piccionello, M. 2005: Prove di efficacia del caolino nel controllo di *Ceratitis capitata* (Wiedemann) in pescheti biologici. *Atti XX Congresso Nazionale Italiano di Entomologia, Perugia-Assisi 13-18 Giugno 2005*: 228.
- Caleca, V., Rizzo, R., Corcella, M. & Battaglia, I. 2005: Prove sull'efficacia del caolino e dell'idrossido di rame nel controllo di *Bactrocera oleae* (Gmelin). *Atti XX Congresso Nazionale Italiano di Entomologia, Perugia-Assisi 13-18 Giugno 2005*: 229.
- Caleca, V. & Rizzo, R. 2005: Tests on the effectiveness of kaolin and copper hydroxide in the control of *Bactrocera oleae* (Gmelin) - Abstracts 2nd European Meeting of the IOBC/WPRS Study Group "Intergrated Protection of Olive Crops", Florence, Italy, October 26-28 2005: 25.
- Knight, A. L., Unruh, T. R., Christianson, B. A., Puterka, G. J., & Glenn, D. M. 2000: Effects of kaolin-based particle films on oblique banded leafroller, *Choristoneura rosaceana* Harris, (Lepidoptera: Tortricidae). *Journal of Economic Entomology* 93: 744-749.
- Liang, G. & Liu, T. X. 2002: Repellency of a kaolin particle film, Surround, and a mineral oil, sunspray oil, to silverleaf whitefly (Homoptera: Aleyrodidae) on melon in the laboratory. *Journal of Economic Entomology* 95: 317-324.
- Lo Verde, G. & Lo Verde, V. 2005: Controllo di *Ceratitis capitata* (Wiedemann) con l'uso del caolino nell'agrumicoltura biologica - *Atti XX Congresso Nazionale Italiano di Entomologia, Perugia-Assisi 13-18 Giugno 2005*: 234.
- Saour, G. 2005: Morphological assessment of olive seedlings treated with kaolin-based particle film and biostimulant. *Advances in Horticultural Science* 19 (4): 193-197.
- Saour, G., & Makee, H. 2003: Effects of kaolin particle film on olive fruit yield, oil content and quality. *Advances in Horticultural Science* 17 (4): 204-206.

Saour, G., & Makee, H. 2004: A Kaolin-based particle film for suppression of the Olive Fruit Fly *Bactrocera oleae* Gmelin (Dip., Tephritidae) in Olive Groves. *Journal of Applied Entomology* 128: 28-31

Showler, A. T. 2002: Effects of kaolin-based particle film application on boll weevil (Coleoptera: Curculionidae) injury to cotton. *Journal of Economic Entomology* 95: 754-762.