

## Article (refereed) - postprint

---

Woodcock, B.A.; Bullock, J.M.; Shore, R.F.; Heard, M.S.; Pereira, M.G.; Redhead, J.; Ridding, L.; Dean, H.; Sleep, D.; Henrys, P.; Peyton, J.; Hulmes, S.; Hulmes, L.; Sároszpataki, M.; Saure, C.; Edwards, M.; Genersch, E.; Knäbe, S.; Pywell, R.F.. 2017. **Country-specific effects of neonicotinoid pesticides on honey bees and wild bees**. *Science*, 356 (6345). 1393-1395. [10.1126/science.aaa1190](https://doi.org/10.1126/science.aaa1190)

2017 @ The Authors

This version available <http://nora.nerc.ac.uk/518181/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

**This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.**

The definitive version is available at <http://science.sciencemag.org/>

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

# Country-specific effects of neonicotinoid pesticides on honeybees and wild bees

**Authors:** B.A.Woodcock<sup>1\*</sup>, J.M.Bullock<sup>1</sup>, R.F.Shore<sup>2</sup>, M.S.Heard<sup>1</sup>, M.G.Pereira<sup>2</sup>, J.Redhead<sup>1</sup>,  
L.Ridding<sup>1</sup>, H.Dean<sup>1</sup>, D.Sleep<sup>2</sup>, P.Henrys<sup>2</sup>, J.Peyton<sup>1</sup>, S.Hulmes<sup>1</sup>, L.Hulmes<sup>1</sup>, M.Sárosipatáki<sup>3</sup>,  
C.Saure<sup>4</sup>, M.Edwards<sup>5</sup>, E.Genersch<sup>6</sup>, S.Knäbe<sup>7</sup> & R.F.Pywell<sup>1</sup>

## Affiliations:

<sup>1</sup> NERC Centre for Ecology & Hydrology, Oxfordshire OX10 8BB, UK.

<sup>2</sup> NERC CEH, Lancaster Environment Centre, Lancaster LA1 4AP, UK.

<sup>3</sup> Szent-István University, 2103 Gödöllő, Hungary.

<sup>4</sup> Am-Heidehof 44, 14163 Berlin, Germany.

<sup>5</sup> Leaside, Carron Lane, West Sussex GU29 9LB, UK

<sup>6</sup> Institute for Bee Research, 16540 Hohen-Neuendorf, Germany

<sup>7</sup> Eurofins, Ecotox-GmbH, 75223 Niefern-Öoschelbronn, Germany

\*Correspondence: B.A.Woodcock ([bawood@ceh.ac.uk](mailto:bawood@ceh.ac.uk))

## Abstract:

**Neonicotinoid seed dressings have caused concern world-wide. We use large field experiments to assess effects of neonicotinoid-treated crops on three bee species across three countries (Hungary, Germany and the UK). Winter-sown oilseed rape was grown commercially with either seed coatings containing neonicotinoids (clothianidin or thiamethoxam) or no seed treatment (control). For honeybee we found both negative (Hungary and UK) and positive (Germany) effects during crop flowering. In Hungary, negative effects on honeybees (associated with clothianidin) persisted over winter and resulted in smaller colonies in the following spring (24% declines). In wild bees (*Bombus terrestris* and *Osmia bicornis*), reproduction was negatively correlated with neonicotinoid residues. These findings point to neonicotinoids causing a reduced capacity of bee species to establish new populations in the year following exposure.**

30 **One Sentence Summary:**

31 Honeybee and wild bee exposure to neonicotinoid pesticides reduces their ability to establish  
32 populations.

33

34 **Main Text:**

35 Global declines in honeybees and wild bees have been linked to pathogens, climate  
36 change, habitat fragmentation and pesticide use (1-3). The potential threat from neonicotinoid  
37 seed coatings applied to flowering crops has been the subject of considerable debate (4-9).  
38 Neonicotinoids have been shown to increase mortality in honeybees by impairing their homing  
39 ability (4) and to reduce the reproductive success of bumblebees (5, 8, 10) and solitary bees (8,  
40 11), while other studies have identified no effects (8, 12, 13). There is limited information from  
41 replicated studies on longer-term survival of honeybee colonies following exposure (see (12)).  
42 Landscape-scale experiments under real world agricultural conditions are needed to integrate  
43 spatial, temporal and species-specific variation to understand the impacts of neonicotinoids on  
44 bees (8, 12, 14-16). Such studies should explore the impacts of different neonicotinoid  
45 formulations, land use and regional climate. In a large-scale experiment spanning three European  
46 countries, we tested the hypotheses that: (i) exposure to seed treatments containing  
47 neonicotinoids affected the reproductive potential of managed and wild bee species and (ii) if  
48 such effects differ between countries.

49 At each of 33 sites (Germany=9, Hungary=12, UK=12) an average of 63.1 ha (SE±2.8  
50 ha) of winter-sown oilseed rape (OSR) was established in 2014 (Fig.1 & S1, Table S1). We  
51 clustered sites into triplets (>3.2 km between sites) and randomly allocated sites to one of three  
52 treatments: 1) Clothianidin applied at 11.86-18.05 g ha<sup>-1</sup> a.i. with a fungicide (Thiam and  
53 prochloraz) and non-systemic pyrethroid (beta-cyfluthrin) (trade name Modesto); 2)  
54 Thiamethoxam applied at 10.07-11.14 g ha<sup>-1</sup> a.i. and combined with the fungicides fludioxonil  
55 and metalaxyl-M (trade name Cruiser); 3) Control oilseed rape receiving a commercial fungicide  
56 (Thiam and Dimethomorph in Germany & Hungary, Thiam and Prochloraz in the UK), but no  
57 neonicotinoid seed treatment. All treatments received typical commercial inputs of pesticide  
58 (e.g. Lambda-cyhalothrin) and fertilizer, with these standardized across a triplet. Standardized  
59 colonies of honeybees (*Apis mellifera*) and wild bees (bumblebee *Bombus terrestris* and solitary  
60 bee *Osmia bicornis*) were introduced to each site. For honeybees we quantified the impacts of

61 the treatments on colony viability during the crop flowering period and in the year following  
62 exposure (hive survival and overwintering worker, brood and storage cell numbers).  
63 Overwintering fitness defines the multi-year persistence of honeybees. For *B. terrestris* we  
64 measured impacts on within-year reproductive output (colony weight gain, and worker, queen  
65 and drone production) and for *O. bicornis* the number of reproductive cells produced (Table S2).  
66 Neonicotinoids can be persistent and widespread in agro-ecosystems (17, 18), so we quantified  
67 residues both in the nests of bee species and those expressed in the crop.

68 We found that neonicotinoid seed treatment affected the inter-annual viability of  
69 honeybee colonies following the winter period in a country-specific manner. In Hungary worker  
70 numbers were 24% lower where clothianidin was compared to the control (treatment×country:  
71  $\chi^2_6=1.47$ ,  $p=0.01$ , explained variance=59.4%; Fig.2), with no significant effect of thiamethoxam.  
72 Clothianidin was more likely to be expressed in the crop where it was applied as a seed  
73 treatment, which identified a mechanism of exposure to the bees ( $\chi^2_2=6.46$ ,  $p=0.04$ ), but this was  
74 not so for thiamethoxam (Table S3). In the UK high hive mortality precluded a formal statistical  
75 analyses of overwintering worker numbers. However, median worker numbers were zero for all  
76 four clothianidin-treated sites, but above zero for two of the control and one of the thiamethoxam  
77 sites (Table S2; Fig.2). Worker numbers following the winter in Germany showed no treatment  
78 effect (Table S4). Overwintering honeybee brood, stored hive products (pollen and nectar) and  
79 the likelihood of hives surviving the winter were not affected by seed treatments (Table S3).

80 Neither *B. terrestris* queen nor *O. bicornis* egg cell production were directly affected by  
81 the seed treatments or its interaction with country (Table S5). However, they were negatively  
82 correlated with peak ( $\chi^2_1=2.09$ ,  $p=0.03$ , explained variance=13.5%; Fig.3a) and median  
83 ( $\chi^2_1=4.34$ ,  $p=0.04$ , explained variance=0.8%; Fig.3b) neonicotinoid nest residues (combined  
84 clothianidin, thiamethoxam and imidacloprid). Imidacloprid was not applied as part of the study  
85 and its presence is most likely a result of environmental contamination from previous widespread  
86 agronomic use (17, 18). Residues of neonicotinoids detected in stored hive products did not  
87 differ in response to seed treatments for any bee species (Table S6). This may be due to the  
88 amalgamation of stored hive products at the site level for residue analysis, which may have  
89 obscured within site heterogeneity in residues. The negative correlation for *B. terrestris* queen  
90 production remained significant when we excluded sites with imidacloprid residues ( $\chi^2_1=2.14$ ,  
91  $p=0.02$ ), although this was not the case for *O. bicornis* ( $\chi^2_1=0.05$ ,  $p=0.81$ ). Country-specific

92 responses to neonicotinoid seed treatment were found for *B. terrestris* drone production, with  
93 positive and negative effects from exposure to thiamethoxam in Germany and the UK  
94 respectively (treatment×country:  $\chi^2_6=13.1$ ,  $p=0.04$ , explained variance=13.6%; Fig.2).

95 We also found seed treatment effects during the crop flowering period that lasted between  
96 3 to 6 weeks (Table S4 & 5). Significant interactions between seed treatment and country were  
97 identified for peak worker ( $\chi^2_6=16.6$ ,  $p<0.01$ , explained variance=45.3%), egg cell ( $\chi^2_6=4.13$ ,  
98  $p=0.01$ , explained variance=49.9%) and combined pollen and nectar storage cell ( $\chi^2_6=40.5$ ,  
99  $p<0.001$ , explained variance=53.6%) numbers. These responses describe within-year colony  
100 performance. Neonicotinoid exposure resulted in both negative (Hungary and UK) and positive  
101 (Germany) effects on colony size (see Fig.2; pairwise treatment comparison given in Table S4 &  
102 5). *Bombus terrestris* worker and peak colony weight showed no seed treatment response.

103 Our quantification of neonicotinoid effects on the inter-annual viability of honeybees and  
104 wild bee populations represents a fundamental advance in our understanding of the impacts of  
105 these pesticides. For solitary bees and bumblebees (queen production) neonicotinoid impacts  
106 were associated with the residues found in nests rather than the experimental seed treatments.  
107 For *B. terrestris* the few treatment effects and the presence of imidacloprid in stored pollen and  
108 nectar (Table S7-S9) suggests that negative impacts of neonicotinoids may be driven by  
109 persistence of residues in the wider landscape, rather than current management alone (18, 19).  
110 The EU moratorium meant that no neonicotinoids were applied to oilseed in the surrounding  
111 landscapes during the experiment, so such residues may originate from previous agricultural use  
112 leading to expression in non-target plants (17-19), guttation fluids or contaminated water (19,  
113 20). While the reproductive potential of *O. bicornis* was also negatively affected by  
114 neonicotinoid residues in nests, the explained variation of these effects was small. However, a  
115 failure to detect small population changes may be due to limited experimental replication  
116 restricting statistical power. Our results suggest that even if their use were to be restricted, as in  
117 the recent EU moratorium, continued exposure to neonicotinoid residues resulting from their  
118 previous widespread use has the potential to impact negatively wild bee persistence in  
119 agricultural landscapes (14, 18, 19).

120 Taken together, our results suggest that exposure to neonicotinoid seed treatments can  
121 have negative effects on the inter-annual reproductive potential of both wild and managed bees,

122 but that these effects are not consistent across countries. The country-specific responses of  
123 honeybees and bumblebees strongly suggests that the effects of neonicotinoids are a product of  
124 interacting factors (20-23). This study has identified between country differences in the use of  
125 oilseed rape crop as a forage resources by bees (affecting exposure to crop residues) and  
126 incidence of disease within hives. Both factors were higher for Hungarian and UK honeybees  
127 (Table S10 & S11). Overall neonicotinoid residues were detected infrequently and rarely  
128 exceeded 1.5 ng g<sup>-1</sup> w/w. As such, direct mortality effects caused by exposure to high  
129 concentrations of neonicotinoids are likely to be rare (Table S12). However, our results suggest  
130 that exposure to low levels of neonicotinoids may cause reductions in hive fitness that are  
131 influenced by a number of interacting environmental factors. Such interacting environmental  
132 factors can amplify the impact of honeybee worker losses (e.g. through sub-lethal toxicity  
133 effects) and reduce longer-term colony viability (4, 16). Importantly, our common experimental  
134 approach applied across three countries revealed varying impacts and may explain the  
135 inconsistent results of previous studies conducted in single countries or at few sites (4, 5, 8, 12,  
136 13, 15).

137

## 138 **References and Notes:**

- 139 1. A. J. Vanbergen, Insect Pollinators Initiative, Threats to an ecosystem service: pressures on  
140 pollinators. *Front. Ecol. Environ.* **11**, 251 (2013).
- 141 2. S. G. Potts *et al.*, Global pollinator declines: trends, impacts and drivers. *TREE* **25**, 345 (2010).
- 142 3. R. Winfree, R. Aguilar, D. P. Vázquez, G. Lebuhn, M. A. Aizen, A meta-analysis of bees'  
143 responses to anthropogenic disturbance. *Ecology* **90**, 2068 (2009).
- 144 4. M. Henry *et al.*, A common pesticide decreases foraging success and survival in honey bees.  
145 *Science* **336**, 348 (2012).
- 146 5. P. R. Whitehorn, S. O'Connor, F. L. Wäckers, D. Goulson, Neonicotinoid pesticide reduces  
147 Bumble Bee colony growth and queen production. *Science* **336**, 351 (2012).
- 148 6. J. E. Cresswell *et al.*, Differential sensitivity of honey bees and bumble bees to a dietary  
149 insecticide (imidacloprid). *Zoology* **115**, 365 (2012).
- 150 7. B. A. Woodcock *et al.*, Impacts of neonicotinoid use on long-term population changes in wild  
151 bees in England *Nat. Comm.* **7**, 12459 (2016).
- 152 8. M. Rundlöf *et al.*, Seed coating with a neonicotinoid insecticide negatively affects wild bees.  
153 *Nature* **521**, 77 (2015).

- 154 9. G. E. Budge *et al.*, Evidence for pollinator cost and farming benefits of neonicotinoid seed  
155 coatings on oilseed rape. *Sci. Report.* **5**, 12547 (2015).
- 156 10. D. Goulson, Neonicotinoids impact bumblebee colony fitness in the field; a reanalysis of the  
157 UK's Food & Environment Research Agency 2012 experiment. *PeerJ* **3**, e854 (2015).
- 158 11. C. Sandrock *et al.*, Sublethal neonicotinoid insecticide exposure reduces solitary bee reproductive  
159 success. *Agric. Forst. Entomol.* **16**, 119 (2014).
- 160 12. G. C. Cutler, C. D. Scott-Dupree, M. Sultan, A. D. McFarlane, L. Brewer, A large-scale field  
161 study examining effects of exposure to clothianidin seed-treated canola on honey bee colony  
162 health, development, and overwintering success. *PeerJ* **2**, 2167 (2014).
- 163 13. G. C. Cutler, C. D. Scott-Dupree, A field study examining the effects of exposure to  
164 neonicotinoid seed-treated corn on commercial bumble bee colonies. *Ecotox.* **23**, 1755 (2014).
- 165 14. B. A. Woodcock *et al.*, Replication, effect sizes and identifying the biological impacts of  
166 pesticides on bees under field conditions. *J. Appl. Ecol.* **53**, 1358 (2016).
- 167 15. E. Pilling, P. Campbell, M. Coulson, N. Ruddle, I. Tornier, A Four-Year Field Program  
168 Investigating Long-Term Effects of Repeated Exposure of Honey Bee Colonies to Flowering  
169 Crops Treated with Thiamethoxam. *PLoS ONE* **8**, e77193 (2013).
- 170 16. M. Henry *et al.*, Reconciling laboratory and field assessments of neonicotinoid toxicity to  
171 honeybees. *Proc. R. Soc. Lond. Ser. B.* **282**, (2015).
- 172 17. A. Jones, P. Harrington, G. Turnbull, Neonicotinoid concentrations in arable soils after seed  
173 treatment applications in preceding years. *Pest Manag. Sci.* **70**, 1780 (2014).
- 174 18. C. Botías *et al.*, Neonicotinoid residues in wildflowers, a potential route of chronic exposure for  
175 bees. *Environ. Sci. Tech.* **49**, 12731 (2015).
- 176 19. D. Goulson, An overview of the environmental risks posed by neonicotinoid insecticides. *J. Appl.*  
177 *Ecol.* **50**, 977 (2013).
- 178 20. A. Fairbrother, J. Purdy, T. Anderson, R. Fell, Risks of neonicotinoid insecticides to honeybees.  
179 *Environ. Toxicol. Chem.* **33**, 719 (2014).
- 180 21. FERA, Neonicotinoid Pesticides and Bees. Report to Syngenta Ltd. *The Food and Environment*  
181 *Research Agency* (2013).
- 182 22. F. Sanchez-Bayo *et al.*, Are bee diseases linked to pesticides? - A brief review. *Environ. Int.* **89-**  
183 **90**, 7 (2016).
- 184 23. C. R. Archer, C. W. W. Pirk, G. A. Wright, S. W. Nicolson, Nutrition affects survival in African  
185 honeybees exposed to interacting stressors. *Funct. Ecol.* **28**, 913 (2014).
- 186 24. European Commission. (EC, <http://ec.europa.eu/eurostat/web/agriculture/data/database>) (2016).
- 187 25. A. Imdorf, G. Buehlmann, L. Gerig, V. Kilchenmann, H. Wille, A Test of the method of  
188 estimation of brood areas and number of worker bees in free-flying colonies. *Apidologie* **18**, 137  
189 (1987).

- 190 26. D. Kleijn et al., Delivery of crop pollination services is an insufficient argument for wild  
191 pollinator conservation. *Nat. Commun.* **6**, 7414 (2015).
- 192 27. B. Magnusson, T. Näykki, H. Hovind, K. M., Nordtest report 537. Handbook for calculation of  
193 measurement uncertainty, (Nordic Innovation, Norway, 2012).
- 194 28. T. Blacquiere, G. Smagghe, C. A. M. van Gestel, V. Mommaerts, Neonicotinoids in bees: a  
195 review on concentrations, side-effects and risk assessment. *Ecotox.* **21**, 973 (2012).
- 196 29. EFSA, Conclusion on the peer review of the pesticide risk assessment for bees for the active  
197 substance clothianidin/thiamethoxam/imidacloprid, *EFSA* (2013).
- 198 30. R Core Development Team, R: Version 3.2.1 A Language and Environment for Statistical  
199 Computing. R Foundation for Statistical Computing. URL <http://cran.r-project.org>. (2015).

200

201 **Acknowledgments:** Data in supporting online material. Funded by Syngenta Ltd. and Bayer  
202 CropScience (P.Campbell, M.Miles, C.Maus, D.Holah, M.Coulson). Wild pollinator  
203 work supported by NERC CEH National Capability funding (NEC05829). Thanks to  
204 K.Jaekel, P.Fisher, M.Nowakowski, R.Hails, P.Scrimshaw, N.Mitschunas, P.Nuttall,  
205 M.McCracken, S.Ball, J.Webb, B.Sutherland, R.Freckleton, T.Tscharntke, J.Memmott,  
206 K.Norris, Brigitta Raffa and Dóra Vaskor.

207

## 208 **Supplementary content**

209 References (24-30)

210 Materials and methods

211 Figs.S1-2

212 Tables S1-12

213

214 **Fig.1. Location of the 33 experimental sites in the UK, Hungary and Germany.** See Fig S2  
215 for a diagrammatic representation of the experimental setup.

216

217 **Fig.2. Summary effect sizes for the response of honeybees and wild bees to the**  
218 **neonicotinoid seed treatments.** An effect size represents the difference between the mean  
219 population response for a given seed treatment and the control within a country, with this  
220 difference divided by the pooled standard deviation. Where: \* indicates a significant differences  
221 between the control and seed treatment (either TMX=thiamethoxam, CTD=clothianidin)  
222 determined from the predicted marginal means of the model  $y \sim seed\ treatment * country +$




223 *block/country*'. † indicates where UK colony survival was too low for a formal analysis. Note  
224 effect sizes differ between countries.

225

226 **Fig.3. Wild bee reproductive success in response to neonicotinoid nest residues.** Separate  
227 graphs are shown for the response of *B. terrestris* queen production and *O. bicornis* reproductive  
228 cell production to neonicotinoid residues found in nests. The significance of these relationship is  
229 based on a likelihood ratio test comparison of H0: '*y ~ country*' and H1: '*y ~*  
230 *Neonicotinoid+country*'. Neonicotinoid residues are based on summed concentrations of  
231 clothianidin, thiamethoxam and imidacloprid. *Expl.Var*=Explained variance.

232

233



**Honeybee**  
**Oilseed rape flowering**

**Germany**

**Hungary**

**UK**

Worker numbers

Egg cells

Larval cells

Pupal cells

Storage cells

■ CTD

□ TMX

**Post winter**

Hive survival

Worker numbers

Brood cells

Storage cells

**Wild bees**


*O. bicornis*

*B. terrestris* queen

*B. terrestris* drone


*B. terrestris* worker

*B. terrestris* hive weight



**Effect size**

a) *Bombus terrestris* queen production



b) *Osmia bicornis* reproductive cells

