

Original citation:

Brown, Gavin and Wemyss, Michael. (2017) Gopakumar--Vafa invariants do not determine flops. Communications in Mathematical Physics.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/93531>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 4.0 International license (CC BY 4.0) and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/4.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk



Gopakumar–Vafa Invariants Do Not Determine Flops

Gavin Brown¹, Michael Wemyss² 

¹ Mathematics Institute, Zeeman Building, University of Warwick, Coventry CV4 7AL, UK.

E-mail: G.Brown@warwick.ac.uk

² The Mathematics and Statistics Building, University of Glasgow, University Place, Glasgow G12 8SQ, UK.

E-mail: michael.wemyss@glasgow.ac.uk

Received: 14 July 2017 / Accepted: 12 September 2017

© The Author(s) 2017. This article is an open access publication

Abstract: Two 3-fold flops are exhibited, both of which have precisely one flopping curve. One of the two flops is new and is distinct from all known algebraic D_4 -flops. It is shown that the two flops are neither algebraically nor analytically isomorphic, yet their curve-counting Gopakumar–Vafa invariants are the same. We further show that the contraction algebras associated to both are not isomorphic, so the flops are distinguished at this level. This shows that the contraction algebra is a finer invariant than various curve-counting theories, and it also provides more evidence for the proposed analytic classification of 3-fold flops via contraction algebras.

1. Introduction

Flopping neighbourhoods are one of the most elementary building blocks of higher dimensional algebraic geometry, and even in dimension three they exhibit a very rich structure. Over the past thirty years the invariants attached to such curves have become increasingly fine, from the trichotomy in the normal bundle [L81], to the *length* in [K89], to the ADE identification in [KaM, K94], to the association of a finite tuple of integers via the Gopakumar–Vafa (=GV) invariants [K08]. At each stage, the produced invariant is strictly finer than the last, with the GV invariants linking to Donaldson–Thomas theory and all other modern curve counting notions (see e.g. [PT]).

On the other hand, contraction algebras were introduced in [DW1], partially to provide a new curve invariant, but mainly to unify the homological approaches to derived symmetries and twists [B02, C02, T07]. With their roots in homological algebra, and because they are an algebra as opposed to a number, this additional structure allows us to use contraction algebras to establish and control many geometric processes [DW2, W14], whilst at the same time recover the GV and other invariants [DW1, T14, HT] in a variety of natural ways.

M.W. was supported by EPSRC Grant EP/K021400/2.

In this paper, we use the algebra structure to show that the contraction algebra is a strictly finer invariant than that of Gopakumar–Vafa. This is in some ways surprising: the GV invariants are indeed enough to classify Type A flops [R83]. The trick is to use noncommutativity. We produce two flops, and we show that their contraction algebras are not isomorphic, although both have the same dimension. Aside from the issue of actually constructing such an example, which we come back to below, we remark here that the isomorphism problem is delicate and is in general also difficult. Deciding when two finite dimensional algebras are not isomorphic is tricky, especially in the situation here, when by design all the standard numerical information attached to each is the same.

The main result is the following, where g is the standard Laufer flop [L81].

Theorem 1.1 (2.3, 4.4, 4.7). *Consider the flopping contractions $f: X \rightarrow \text{Spec } R$ and $g: Y \rightarrow \text{Spec } L$ constructed in 2.1 and 2.2. Then the following statements hold.*

- (1) R is not analytically (or algebraically) isomorphic to L .
- (2) The Gopakumar–Vafa invariants associated to f and g are the same.
- (3) The contraction algebras associated to f and g are not isomorphic.

It is conjectured in [DW1, 1.4] that contraction algebras are the analytic classification of 3-fold flops. Whilst the new flop f may look like it comes out of the blue, we found it during our systematic attempts to approach the conjecture based on an explicit gluing via a superpotential; on the noncommutative side, the example is much clearer. Indeed, the flop f was constructed by *assuming* the above conjecture is true and working backwards, thus the results in this paper add some weight to the conjecture. We also remark that there are tables of data that numerically suggest, but do not quite yet prove, that different flops having the same GV invariants is actually quite typical behaviour.

It is perhaps worth explaining the heuristic reason as to why the noncommutativity of the contraction algebra helps, rather than hinders, distinguishing the two flops above. Whilst algebraically the two commutative curves $x^3 - y^2$ and $x^3(x + 1) - y^2$ are different, analytically around the origin, their coordinate rings are isomorphic. Set $\widehat{\mathbb{A}}_\lambda := \mathbb{C}\langle\langle x, y \rangle\rangle / x y - \lambda y x$, where implicitly we consider the closure of all ideals. Then the above famous algebro-geometric curve example is precisely the statement that

$$\frac{\widehat{\mathbb{A}}_1}{x^3 - y^2} \cong \frac{\widehat{\mathbb{A}}_1}{x^3(x + 1) - y^2}. \quad (1.A)$$

The third part of 1.1 turns out to be equivalent to establishing the more surprising statement that there is no such isomorphism in the quantum plane, namely

$$\frac{\widehat{\mathbb{A}}_{-1}}{x^3 - y^2} \not\cong \frac{\widehat{\mathbb{A}}_{-1}}{x^3(x + 1) - y^2}. \quad (1.B)$$

Heuristically, noncommutativity gives the flexibility to distinguish: since y no longer commutes with x , it no longer commutes with $\sqrt{x + 1}$, so we should expect the isomorphism in (1.A) to break down. It turns out that the left hand side of (1.B) is the contraction algebra of g [DW1], and we show in Example 4.1 and Remark 4.8 that the right hand side of (1.B) is the contraction algebra of f . The proof of (1.B) is somewhat more involved than this heuristic argument; we give a direct proof in 4.7, but it is also possible to give a computer algebra verification by adapting the Shirayanagi algorithm [S].

2. The Two Flops

Here the two main examples are introduced. All the calculations in 2.1 are easy enough to be done by hand, but to allay any possibilities of error, we have included in Appendix A computer algebra codings that can be used to independently check all claims.

Example 2.1. (The new flop R) Consider the hypersurface $R := \mathbb{C}[u, v, x, y]/(f)$, where

$$f := u^2 + v^2(x + y) + x(x^2 + xy^2 + y^3).$$

By the Jacobi criterion, R has a unique isolated singular point, at the origin. Being a hypersurface in \mathbb{A}^4 , clearly $\text{Spec } R$ is a Gorenstein 3-fold. We next verify that it is the base of a simple flopping contraction, by constructing a small resolution. The same calculation shows that R is cD_4 , although this can also be verified at once from the above equation.

To construct a small resolution, blowup the reflexive ideal $I := (vx - uy, xy^2 + v^2, x^2y + uv)$ to obtain a projective birational morphism

$$X \rightarrow \text{Spec } R.$$

Here we summarise the calculation by hand; the computation using Singular is summarised in the Appendix Sect. A.1. The blowup X is covered by two affine open charts, the first of which is given by the smooth hypersurface

$$U_1 := \text{Spec } \mathbb{C}[x_3, x_4, y_1, y_2]/(x_3(y_1^2 + 1) + x_4y_1^2 + y_2^2)$$

with map to the base

$$\begin{array}{c} (x_3, x_4, y_1, y_2) \in U_1 \\ \downarrow \\ (x_3x_4y_1 + x_4^2y_1 + x_3y_2, x_3y_1 - x_4y_2, x_3, x_4) \in \text{Spec } R. \end{array}$$

Above the origin of $\text{Spec } R$ consists of all points $(0, 0, y_1, y_2)$ of U_1 such that the defining relation of U_1 holds, so necessarily $y_2^2 = 0$. Thus the fibre above the origin is a single curve, with scheme multiplicity two. The second open chart is given by the smooth hypersurface

$$U_2 := \text{Spec } \mathbb{C}[x_2, x_4, y_0, y_2]/(x_2y_0^3 + x_4y_0^2y_2 + x_2y_0 + x_4y_2 + y_2^2 + x_4)$$

with map to the base

$$\begin{array}{c} (x_2, x_4, y_0, y_2) \in U_2 \\ \downarrow \\ (-x_2x_4y_0^2 - x_4^2y_0y_2 + x_2y_2, x_2, x_2y_0 + x_4y_2, x_4) \in \text{Spec } R. \end{array}$$

Here the fibre above the origin consists of $(0, 0, y_0, y_2)$ such that $y_2^2 = 0$, which again is a curve. It is an easy check to see that the reduced fibre above the origin glues via

$$(0, 0, y_1, 0) \leftrightarrow (0, 0, y_1^{-1}, 0)$$

and so is \mathbb{P}^1 . It follows that $X \rightarrow \text{Spec } R$ is a smooth flopping contraction, and thus R is cDV . Since by the above calculation the scheme fibre has multiplicity two, we deduce that this must be a cD_4 flop [KaM, K94].

Example 2.2 (The standard Laufer flop L). Consider $L := \mathbb{C}[u, v, x, y]/(g)$, where

$$g := u^2 + v^2y - x(x^2 + y^3).$$

This has a unique singular point at the origin, and indeed $\text{Spec } L$ is the base of the standard Laufer flop. Blowing up the reflexive ideal $(x^2 + y^3, vx + uy, ux - vy^2)$ gives a projective birational morphism

$$Y \rightarrow \text{Spec } L$$

where Y is smooth. The reduced scheme fibre above the origin is \mathbb{P}^1 , and the full scheme fibre has multiplicity two. This was the first known example of a cD_4 flop [L81, KaM, R83].

Remark 2.3. R is not analytically isomorphic to L , and hence also $R \not\cong L$ algebraically. This can be seen directly by computing the Tjurina numbers of both (see e.g. Sect. A.1), but it also follows from the non-isomorphism of the contraction algebras later in 4.7.

3. GV Invariants and Contraction Algebras

The GV invariants of both the flopping contractions $X \rightarrow \text{Spec } R$ and $Y \rightarrow \text{Spec } L$ in the previous section are determined by their contraction algebras [T14], and this section briefly reviews these notions.

3.1. Contraction algebra background. Throughout, consider a general 3-fold flopping contraction $f: U \rightarrow \text{Spec } \mathfrak{R}$, where U is smooth, $f^{-1}(0) := C$ and $C^{\text{red}} \cong \mathbb{P}^1$, and for simplicity assume that \mathfrak{R} is complete local. To this data, one can associate the *contraction algebra* A_{con} , which can be defined [DW1, §3] as the representing object of the noncommutative deformation functor

$$\text{Def}: \mathbf{Art}_1 \rightarrow \mathbf{Sets},$$

where \mathbf{Art}_1 is the category of augmented finite dimensional \mathbb{C} -algebras. By definition Def sends

$$(\Gamma, \mathfrak{n}) \mapsto \left\{ (\mathcal{F}, \phi, \delta) \left| \begin{array}{l} \mathcal{F} \in \text{Qcoh } U \\ \phi: \Gamma \rightarrow \text{End}_U(\mathcal{F}) \text{ is a } \mathbb{C}\text{-algebra homomorphism} \\ - \otimes_{\Gamma} \mathcal{F}: \text{mod } \Gamma \rightarrow \text{Qcoh } U \text{ is exact} \\ \delta: (\Gamma/\mathfrak{n}) \otimes_{\Gamma} \mathcal{F} \xrightarrow{\sim} \mathcal{O}_{\mathbb{P}^1}(-1) \end{array} \right. \right\} / \sim$$

where the equivalence relation \sim is outlined in detail in [DW1, 2.4]. For the purpose of this paper, the following information suffices.

Theorem 3.1. *Consider a complete local flopping contraction $U \rightarrow \text{Spec } \mathfrak{R}$, as above. Then the following statements hold.*

- (1) A_{con} is a finite dimensional algebra.
- (2) A_{con} is not commutative if and only if C is a $(-3, 1)$ -curve. Moreover, in this case A_{con} can be presented as

$$A_{\text{con}} \cong \frac{\mathbb{C}\langle\langle x, y \rangle\rangle}{(\delta_x W, \delta_y W)}$$

for some superpotential W , where $(\delta_x W, \delta_y W)$ denotes the closure of the ideal generated by the formal derivatives $\delta_x W$ and $\delta_y W$.


(3) Suppose that C is a $(-3, 1)$ -curve. If $X \in \text{CM } \mathfrak{R}$ is a non-free rank two module, and $\text{Ext}_{\mathfrak{R}}^1(X, X) = 0$, then $A_{\text{con}} \cong \underline{\text{End}}_{\mathfrak{R}}(X)$ and $\dim_{\mathbb{C}} A_{\text{con}} = \dim_{\mathbb{C}} \text{Ext}_{\mathfrak{R}}^2(X, X)$.

Proof. Part (1) is [DW1, 2.13(1)], and the first statement of part (2) is [DW1, 2.13]. The fact that A_{con} is a superpotential algebra is a consequence of that it is a factor of an NCCR [DW1, §3], which since \mathfrak{R} is complete, is a superpotential algebra [V10].

For part (3), by [W14, 4.14] there are only two non-free indecomposable CM \mathfrak{R} -modules M_i for which $\text{Ext}_{\mathfrak{R}}^1(M_i, M_i) = 0$, and the rank of each M_i equals the length of the flopping curve. Since the curve is a $(-3, 1)$ -curve, the length is strictly greater than one. Hence X cannot split into two rank one summands, as then there would be a rank one rigid non-free CM \mathfrak{R} -module. We conclude that X is indecomposable, so it must be isomorphic to one of the M_i . By definition, A_{con} is $\underline{\text{End}}_{\mathfrak{R}}(M_i)$, and hence is isomorphic to $\underline{\text{End}}_{\mathfrak{R}}(X)$. The last statement regarding the dimension is then [DW1, 5.2]. \square

Later, 3.1(3) will be used to calculate the contraction algebra, and also to compute its dimension, without requiring knowledge of its algebra structure.

3.2. *GV invariants.* Each flopping contraction f of length ℓ has an associated tuple of integers (n_1, \dots, n_ℓ) called the *Gopakumar–Vafa invariants*. These can be defined as follows. As in [BKL, §2.1], there exists a flat deformation



for some Zariski open neighbourhood T of $0 \in \mathbb{A}^1$, such that

- The central fibre $g_0: X_0 \rightarrow Y_0$ is isomorphic to the formal fibre \widehat{f} of f .
- All other fibres $g_t: X_t \rightarrow Y_t$ for $t \in T \setminus \{0\}$ are flopping contractions whose exceptional locus is a disjoint union of $(-1, -1)$ -curves.

Regarding the flopping curve C of \widehat{f} as a curve in the central fibre of $\mathcal{X} \rightarrow T$, and thus as a curve in \mathcal{X} , then the GV invariant n_j is defined to be the number of g_t -exceptional $(-1, -1)$ -curves C' with curve class $j[C]$, i.e. for every line bundle \mathcal{L} on \mathcal{X} ,

$$\deg(\mathcal{L}|_{C'}) = j \deg(\mathcal{L}|_C).$$

The following is [T14, 1.1], and will be used to deduce the GV invariants later.

Theorem 3.2 (Toda). *Suppose that $f: U \rightarrow \text{Spec } \mathfrak{R}$ is a complete local flopping contraction of a single length ℓ $(-3, 1)$ -curve, where U is smooth. Then $n_1 = \dim_{\mathbb{C}} A_{\text{con}}^{\text{ab}}$ and*

$$\dim_{\mathbb{C}} A_{\text{con}} = \dim_{\mathbb{C}} A_{\text{con}}^{\text{ab}} + \sum_{j=2}^{\ell} j^2 \cdot n_j,$$

where $n_j \in \mathbb{Z}_{\geq 1}$ are the Gopakumar–Vafa invariants associated to the curve.

4. GV Invariants Do Not Determine Flops

This section computes the contraction algebras for the flopping contractions 2.1 and 2.2, and as a corollary shows that the GV invariants attached to both flops are the same. The two contraction algebras are then shown not to be isomorphic, and so the flops are distinguished at this finer level.

4.1. *Calculation of contraction algebras.* Write Λ_{con} for the contraction algebra associated to the standard Laufer flop $Y \rightarrow \text{Spec } L$ in 2.2. The following is known.

Example 4.1. With notation as in the introduction,

$$\Lambda_{\text{con}} \cong \widehat{\Lambda}_{\text{con}} \cong \frac{\mathbb{C}\langle x, y \rangle}{(xy + yx, x^3 - y^2)} = \frac{\mathbb{A}_{-1}}{x^3 - y^2},$$

where the first isomorphism is [DW1, 2.17] and the second is [DW1, 1.3]. Thus Λ_{con} is a 9-dimensional not-commutative ring, given by superpotential $W = x^4 - xy^2$.

The calculation of the contraction algebra associated to the new flop uses a very similar method to the above example. To set notation, write Γ_{con} for the contraction algebra associated to $X \rightarrow \text{Spec } R$ in 2.1.

Proposition 4.2. *There is an isomorphism*

$$\Gamma_{\text{con}} \cong \frac{\mathbb{C}\langle a, b \rangle}{ab + ba, -a^2 + b^3 + aba}.$$

This is a nine-dimensional not-commutative ring.

Proof. The most direct method to prove this is to specialise the universal flop

$$u^2 + v^2Y + x^2\alpha + 2\beta xv + (\alpha Y - \beta^2)\gamma^2$$

given in [AM, (46)] at $\alpha = x$, $Y = x + y$, $\beta = 0$ and $\gamma = y$ to give

$$u^2 + v^2(x + y) + x^3 + x(x + y)y^2,$$

which equals f in 2.1. Consider the cokernel M of the matrix Φ in the following matrix factorisation

$$R^4 \xrightarrow{\Psi := \begin{pmatrix} u & x & v & y \\ -x^2 & u & -xy & v \\ -vx - vy & xy + y^2 & u & -x \\ -x^2y - xy^2 & -vx - vy & x^2 & u \end{pmatrix}} R^4 \xrightarrow{\Phi := \begin{pmatrix} u & -x & -v & -y \\ x^2 & u & xy & -v \\ vx + vy & -(xy + y^2) & u & x \\ x^2y + xy^2 & vx + vy & -x^2 & u \end{pmatrix}} R^4.$$

Then by [AM, §4] $\text{End}_R(R \oplus M)$ is presented as the path algebra of the quiver

$$\begin{array}{c} d = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \quad a = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -x & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -x & 0 \end{pmatrix} \\ R \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} M \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} \\ c = (u \ -x \ -v \ -y) \quad b = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -(x+y) & 0 & 0 & 0 \\ 0 & x+y & 0 & 0 \end{pmatrix} \end{array}$$

where the relations are determined by matrix multiplication, up to the column space of the matrix Ψ . For example, using the above matrices, it can be seen directly that $a \circ b + b \circ a = 0$, and that

$$-a^2 + b^3 + a \circ b \circ a - b \circ d \circ c - d \circ c \circ b = \begin{pmatrix} x & 0 & -y & 0 \\ u & 0 & -v & 0 \\ xy+y^2 & 0 & x & 0 \\ -vx-vy & 0 & -u & 0 \end{pmatrix},$$

which belongs to the column space of Ψ , and thus is zero. Factoring out the vertex corresponding to R in the above presentation, and killing all arrows that factor through it, it follows that $\Gamma_{\text{con}} = \underline{\text{End}}_R(M)$ can be presented as $\mathbb{C}\langle a, b \rangle$ subject to at least the relations $ab+ba = 0$ and $-a^2+b^3+aba = 0$. Since these relations give an algebra which is nine-dimensional (either by the Diamond Lemma, or Sect. A.2), and Γ_{con} is nine-dimensional (see Sect. A.2), it follows that these are all the relations that are needed. \square

Remark 4.3. It is not obvious from the presentation in 4.2, but it is nevertheless true (see 4.8) that Γ_{con} is a superpotential algebra.

The following is the main result of this subsection.

Corollary 4.4. *The GV invariants attached to the flops in 2.1 and 2.2 are the same.*

Proof. Via the explicit presentations in 4.1 and 4.2, we see immediately using Toda’s formula 3.2 that $n_1 = 5$ and $n_2 = 1$ in both cases. \square

Remark 4.5. It is not necessary to compute the algebra structures in order to deduce that the GV invariants of the two flops are the same. Using 3.1(3), the dimension of both Λ_{con} and Γ_{con} can be seen to be nine directly, without knowing the algebra structure. We outline the code in Appendix Sect. A.2. Then, since both are cD_4 flops, neither contraction algebra is commutative by 3.1(2). The abelianization of any not-commutative contraction algebra must be at least four dimensional, being the factor of $\mathbb{C}[[x, y]]$ by two relations in which each word is quadratic or higher. Hence by Toda’s formula 3.2, since the only possibility of writing 9 as a sum of squares is $9 = 5 \cdot 1^2 + 1 \cdot 2^2$, it follows the GV invariants of both flops must be the same, namely $n_1 = 5$ and $n_2 = 1$. These numerics are how this example was discovered.

4.2. *The contraction algebras are not isomorphic.* The proof that Λ_{con} is not isomorphic to Γ_{con} requires the following preparatory lemma.

Lemma 4.6. *With notation as above, the following statements hold.*

- (1) In Γ_{con} , we have $a^3 = 0$ and $b^6 = 0$.
- (2) Λ_{con} has basis $\{1, y, x, y^2, yx, x^2, y^2x, yx^2, y^2x^2\}$.

Proof. (1) Multiplying the defining equation $a^2 = b^3 + aba$ by a on the right, and by a on the left, it follows that

$$ab^3 + a^2ba = a^3 = b^3a + aba^2.$$

Using the defining equation $ab = -ba$ repeatedly,

$$ab^3 - a^3b = a^3 = -ab^3 + a^3b$$

Since the left hand side is the negative of the right hand side, it follows that $a^3 = 0$. Then, simply squaring both sides of the defining equation $a^2 - aba = b^3$ gives

$$b^6 = (a^2 - aba)^2 = a^4 - a^3ba - aba^3 - aba^2ba,$$

which is zero, using the relation $ab = -ba$ together with the fact that $a^3 = 0$.

(2) The first method to establish this is just to use the Diamond Lemma directly, and indeed the stated basis is exactly the one used in [DW1, 3.14]. The second method, using magma, is outlined in the Appendix (Sect. A.2). \square

Theorem 4.7. Λ_{con} is not isomorphic to Γ_{con} .

Proof. Let $\psi: \Gamma_{\text{con}} \rightarrow \Lambda_{\text{con}}$ be an arbitrary isomorphism; we aim for a contradiction. As $\Lambda_{\text{con}}, \Gamma_{\text{con}} \in \mathbf{Art}_1$, in both cases their Jacobson radical is their path ideal. Hence under the isomorphism ψ , necessarily the generators a and b must map to the Jacobson radical, and so by 4.6(2) we may write

$$\psi(a) = \lambda_1 y + \lambda_2 x + \lambda_3 y^2 + \cdots + \lambda_8 y^2 x^2$$

$$\psi(b) = \mu_1 y + \mu_2 x + \mu_3 y^2 + \cdots + \mu_8 y^2 x^2$$

for some scalars λ_1, \dots, μ_8 . Now by 4.6(1), $\psi(a)^3 = \psi(a^3) = \psi(0) = 0$, thus

$$(\lambda_1 y + \lambda_2 x + \lambda_3 y^2 + \cdots + \lambda_8 y^2 x^2)^3 = 0$$

in Λ_{con} . Multiplying out the left hand side, and using the relations of Λ_{con} , we can express the left hand side in terms of the basis 4.6(2) of Λ_{con} (see e.g. Sect. A.2). Doing this,

$$(\lambda_2^3)y^2 + (\lambda_1^2\lambda_2 + 3\lambda_2^2\lambda_5)y^2x + (\lambda_1\lambda_2^2)yx^2 + 3(\lambda_1^2\lambda_5 + \lambda_2^2\lambda_3 + \lambda_2\lambda_5^2)y^2x^2 = 0.$$

Being a basis, all coefficients must be zero. Hence $\lambda_2 = 0$. This in turn implies that:

- $\lambda_1 \neq 0$. This is since a belongs to the Jacobson radical but not the Jacobson radical squared, hence so does $\psi(a)$. As $\lambda_2 = 0$ above, necessarily $\lambda_1 \neq 0$.
- $\lambda_5 = 0$. This is a consequence of the coefficient $\lambda_1^2\lambda_5 + \lambda_2^2\lambda_3 + \lambda_2\lambda_5^2$ being zero, together with the fact that $\lambda_2 = 0$ and $\lambda_1 \neq 0$.

Then, observing that $\psi(ab + ba) = 0$ since $ab + ba = 0$, we see that

$$\psi(a)\psi(b) + \psi(b)\psi(a) = 0.$$

Again, multiplying out the above expressions (using $\lambda_1 = \lambda_5 = 0$, see e.g. Sect. A.2), expressing in terms of the basis of Λ_{con} gives

$$2(\lambda_1\mu_1)y^2 + 2(\lambda_3\mu_2)y^2x + 2(\lambda_1\mu_5)yx^2 \\ + 2(\lambda_1\mu_7 + \lambda_3\mu_5 - \lambda_4\mu_4 + \lambda_6\mu_2 + \lambda_7\mu_1)y^2x^2 = 0.$$

Since $\lambda_1 \neq 0$, necessarily $\mu_1 = \mu_5 = 0$. Again, since b belongs to the radical but not the radical squared, $\mu_2 \neq 0$.

Finally, since $-\psi(a)^2 + \psi(b)^3 + \psi(a)\psi(b)\psi(a) = 0$, multiplying out and expressing in terms of the basis of Λ_{con} (again see e.g. Sect. A.2), using $\lambda_2 = \lambda_5 = \mu_1 = \mu_5 = 0$, we see

$$(-\lambda_1^2 + \mu_2^3)y^2 + (-\lambda_1^2\mu_2)y^2x + (-2\lambda_1\lambda_7 + \lambda_4^2 + 3\mu_2^2\mu_3)y^2x^2 = 0$$

Hence $\lambda_1^2\mu_2 = 0$, which is a contradiction. Thus the isomorphism ψ cannot exist. \square

Remark 4.8. Although we don't strictly need this to show 4.7, in the notation of the introduction it turns out, e.g. using the Shirayanagi algorithm [S], that

$$\Gamma_{\text{con}} \cong \frac{\widehat{\mathbb{A}}_{-1}}{x^4 + x^3 - y^2}.$$

Thus Γ_{con} is a Jacobi algebra, given by the superpotential $W = x^5 + x^4 - xy^2$. Combining this fact with 4.7 justifies the non-isomorphism (1.B) in the introduction.

We next show that commutative deformations cannot determine flopping neighbourhoods. This requires the following.

Proposition 4.9. $\Lambda_{\text{con}}^{\text{ab}} \cong \Gamma_{\text{con}}^{\text{ab}}$.

Proof. By simply commuting variables in the presentations from 4.1 and 4.2,

$$\Lambda_{\text{con}}^{\text{ab}} \cong \frac{\mathbb{C}[x, y]}{xy, x^2 + y^3} \quad \text{and} \quad \Gamma_{\text{con}}^{\text{ab}} \cong \frac{\mathbb{C}[a, b]}{ab, a^2(b-1) + b^3} = \frac{\mathbb{C}[a, b]}{ab, -a^2 + b^3}$$

where the last equality holds simply since $ab = 0$ implies $a^2b = 0$. The above two rings are visibly isomorphic. \square

Corollary 4.10. *Flopping neighbourhoods are not determined by the commutative deformations of the reduced flopping curve.*

Proof. The commutative deformations of $\mathcal{O}_{\mathbb{P}^1}(-1)$ are given by the abelianization of the contraction algebra [DW1, 3.2]. Since $\Lambda_{\text{con}}^{\text{ab}} \cong \Gamma_{\text{con}}^{\text{ab}}$ by 4.9, the commutative deformations of the reduced flopping curves in both flopping contractions are the same. However, the flops are not analytically isomorphic by 2.3 or 4.7. \square

Acknowledgements. The authors would like to thank Agata Smoktunowicz and Natalia Iyudu for many helpful discussions, and for sharing with us a direct proof that the algebra in 4.1 and the algebra in 4.8 are not isomorphic.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Appendix A. Code for Verification

In this appendix we list computer algebra code which can be used to independently verify the claims made in the main text.

A.1. Code for the flops.

- R has a unique singular point at the origin. Singular [DGPS]:

```
LIB"homolog.lib";
ring r = 0, (u, v, x, y), dp;
ideal i = u2 + v2 * (x + y) + x * (x2 + xy2 + y3);
minAssGTZ(radical(slocus(std(i))));
```

- $I := (vx - uy, xy^2 + v^2, x^2y + uv)$ is a rank one reflexive R -module. Using Macaulay2 [GS]:

```
loadPackage"Divisor";
R = QQ[u, v, x, y]/ideal(u^2+v^2*(x+y)+x*(x^2+x*y^2+y^3));
i = ideal(v*x - u*y, x*y^2 + v^2, x^2*y + u*v);
isReflexive(i);
```

Similarly $(x^2 + y^3, vx + uy, ux - vy^2)$ is a rank one reflexive L -module.

- Blowup of Spec R at the ideal I . Singular:

```
LIB"homolog.lib";
LIB"resolve.lib";
ring r = 0, (u, v, x, y), dp;
ideal i = u^2 + v^2 * (x + y) + x * (x^2 + x*y^2 + y^3);
ideal Z = vx - uy, xy^2 + v^2, x^2y + uv;
list blow = blowUp(i, Z);
blow;
def Q = blow[1];
setring Q;
sT;
bM;
dim_slocus(sT);
elim1(sT, x(1));
} Chart 1
def Q = blow[2];
setring Q;
sT;
bM;
dim_slocus(sT);
} Chart 2
```

This returns the defining equations of each chart, the map to the base, and also the fact that each chart is smooth.

- Milnor and Tjurina numbers of R are 12 and 10 respectively. Singular:

```
LIB"sing.lib";
ring r = 0, (u, v, x, y), dp;
poly f = u^2 + v^2 * (x + y) + x * (x^2 + x*y^2 + y^3);
milnor(f); tjurina(f);
```

Complete locally the Milnor number of R drops to 11, whilst the Tjurina number is still 10. Singular:

```
LIB"sing.lib";
ring r = 0, (u, v, x, y), ds;
poly f = u^2 + v^2 * (x + y) + x * (x^2 + x*y^2 + y^3);
milnor(f); tjurina(f);
```

- The Milnor and Tjurina numbers of L are coded similarly, and are both 11 in both dp and ds ordering.

A.2. Code for the contraction algebras.

- For the new flop R , using 3.1(3) the dimension of Γ_{con} is nine. Singular:

```
LIB"homolog.lib"
ring r = 0, (u, v, x, y), dp;
ideal i = u^2 + v^2 * (x + y) + x * (x^2 + x*y^2 + y^3);
qring S = std(i);
module Ma = [vx - uy], [xy^2 + v^2], [x^2y + uv];
module M = syz(Ma);
module X = prune(syz(M));
depth(X); vdim(Ext(1, X, X)); vdim(Ext(2, X, X));
```

This can be easily adapted, using the reflexive module $(x^2 + y^3, vx + uy, ux - vy^2)$ of L , to show that Λ_{con} also has dimension nine.

- The dimension and basis of Γ_{con} . The code for Λ_{con} is similar. Magma [BCP]:

```
K := Rationals();
F < a, b > := FreeAlgebra(K, 2);
I := ideal < F | a*b + b*a, -a^2 + b^3 + a*b*a >;
R := F/I;
W, Wf := VectorSpace(R);
[W.i@@Wf: i in [1..Dimension(W)]];
```

- Expressing products in terms of the basis in the proof of 4.7. Magma:

```

K := RationalS();
k3 < l1, l2, l3, l4, l5, l6, l7, l8, m1, m2, m3, m4, m5, m6, m7, m8 >
:= RationalFunctionField(K, 16);
F < x, y >:= FreeAlgebra(k3, 2);
I := ideal < F | x3 - y2, x * y + y * x >;
T := F/I;
W, Wf := VectorSpace(T);
A := l1 * y + l2 * x + l3 * y2 + l4 * y * x + l5 * x2 + l6 * y2 * x
+ l7 * y * x2 + l8 * y2 * x2;
Wf(A3);

```

Output shows that $l_2 = l_5 = 0$. Setting these to be zero:

```

K := RationalS();
k3 < l1, l3, l4, l6, l7, l8, m1, m2, m3, m4, m5, m6, m7, m8 >
:= RationalFunctionField(K, 14);
F < x, y >:= FreeAlgebra(k3, 2);
I := ideal < F | x3 - y2, x * y + y * x >;
T := F/I;
W, Wf := VectorSpace(T);
A := l1 * y + l3 * y2 + l4 * y * x + l6 * y2 * x + l7 * y * x2 + l8 * y2 * x2;
B := m1 * y + m2 * x + m3 * y2 + m4 * y * x + m5 * x2 + m6 * y2 * x
+ m7 * y * x2 + m8 * y2 * x2;
Wf(A * B + B * A);

```

Output shows that $m_1 = m_5 = 0$. Setting these to be zero:

```

K := RationalS();
k3 < l1, l3, l4, l6, l7, l8, m2, m3, m4, m6, m7, m8 >
:= RationalFunctionField(K, 12);
F < x, y >:= FreeAlgebra(k3, 2);
I := ideal < F | x3 - y2, x * y + y * x >;
T := F/I;
W, Wf := VectorSpace(T);
A := l1 * y + l3 * y2 + l4 * y * x + l6 * y2 * x + l7 * y * x2 + l8 * y2 * x2;
B := m2 * x + m3 * y2 + m4 * y * x + m6 * y2 * x + m7 * y * x2 + m8 * y2 * x2;
Wf(-A2 + B3 + A * B * A);

```

Output shows that $l_1^2 m_2 = 0$.

References

- [AM] Aspinwall, P.S., Morrison, D.R.: Quivers from matrix factorizations. *Commun. Math Phys.* **313**, 607–633 (2012)
- [BCP] Bosma, W., Cannon, J., Playoust, C.: The Magma algebra system. I. The user language. *J. Symb. Comput.* **24**, 235–265 (1997). <http://www.magma.usyd.edu>
- [B02] Bridgeland, T.: Flops and derived categories. *Invent. Math.* **147**(3), 613–632 (2002)
- [BKL] Bryan, J., Katz, S., Leung, N.: Multiple covers and the integrality conjecture for rational curves in Calabi–Yau threefolds. *J. Algebraic Geom.* **10**(3), 549–568 (2001)
- [C02] Chen, J.-C.: Flops and equivalences of derived categories for threefolds with only terminal Gorenstein singularities. *J. Differ. Geom.* **61**(2), 227–261 (2002)
- [DGPS] Decker, W., Greuel, G.-M., Pfister, G., Schönemann, H.: SINGULAR 4-1-0—A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2017)
- [DW1] Donovan, W., Wemyss, M.: Noncommutative deformations and flops. *Duke Math. J.* **165**(8), 1397–1474 (2016)
- [DW2] Donovan W., Wemyss, M.: Contractions and deformations. [arXiv:1511.00406](https://arxiv.org/abs/1511.00406)
- [GS] Grayson, D.R., Stillman, M.E.: Macaulay2, a software system for research in algebraic geometry. <http://www.math.uiuc.edu/Macaulay2/> (2017)
- [HT] Hau, Z., Toda, Y.: Contraction algebra and invariants of singularities. *Int. Math. Res. Notices* **2017**, 1–26. <https://doi.org/10.1093/imrn/rnw333> (2017)

- [K08] Katz, S.: Genus zero Gopakumar–Vafa invariants of contractible curves. *J. Differ. Geom.* **79**, 185–195 (2008)
- [KaM] Katz, S., Morrison, D.R.: Gorenstein threefold singularities with small resolutions via invariant theory for Weyl groups. *J. Algebraic Geom.* **1**, 449–530 (1992)
- [K94] Kawamata, Y.: General hyperplane sections of nonsingular flops in dimension 3. *Math. Res. Lett.* **1**, 49–52 (1994)
- [K89] Kollár, J.: Flops. *Nagoya Math. J.* **113**, 15–36 (1989)
- [L81] Laufer, H.B.: CP^1 as an exceptional set, Recent developments in several complex variables (Proc. Conf., Princeton Univ., Princeton, N.J., 1979), pp. 261–275, *Ann. of Math. Stud.*, **100**, Princeton University Press, Princeton, N.J., 1981
- [PT] Pandharipande, R., Thomas, R.P.: $13/2$ ways of counting curves. In: *Moduli Spaces*, LMS Lecture Note Series, vol. 411. Cambridge University Press, pp. 282–333 (2014)
- [R83] Reid, M.: Minimal models of canonical 3-folds, *Algebraic varieties and analytic varieties* (Tokyo, 1981). *Adv. Stud. Pure Math.*, vol. 1. North-Holland, Amsterdam, pp. 131–180 (1983)
- [S] Shirayanagi, K.: Decision of algebra isomorphisms using Gröbner base, *Computational algebraic geometry* (Nice, 1992). *Progr. Math.*, vol. 109. Birkhäuser Boston, Boston, MA, pp. 255–265 (1993)
- [T07] Toda, Y.: On a certain generalization of spherical twists. *Bull. Soc. Math. France* **135**(1), 119–134 (2007)
- [T14] Toda, Y.: Noncommutative width and Gopakumar–Vafa invariants. *Manuscr. Math.* **148**(3–4), 521–533 (2015)
- [V10] Van den Bergh, M.: Calabi–Yau algebras and superpotentials. *Sel. Math. (N.S.)* **21**(2), 555–603 (2015)
- [W14] Wemyss, M.: Flops and clusters in the homological minimal model program. *Invent. Math.* (2017). <https://doi.org/10.1007/s00222-017-0750-4>

Communicated by N. Nekrasov