

**DESK REFERENCE**

# **A Histological Atlas of Florida Surgeonfish (*Acanthuridae*)**



G. Christopher Tilghman  
Ruth Francis-Floyd  
RuthEllen Klinger

## Introduction

Acanthurids are an important group of reef fishes, because of their impact as herbivores on reef ecology (Carpenter 1986), and their popularity as display animals. The acanthurids are extremely abundant in tropical and subtropical waters. Three species are found in the waters of the Florida Keys: *Acanthurus coeruleus* (blue tang), *A. bahianus* (ocean surgeon), and *A. chirurgus* (doctorfish). Along with parrotfish (*Scaridae*) and damselfish (*Pomacentridae*), these animals are primarily herbivores, and together they form the largest portion of fish biomass of reef systems (Thresher 1980). Such fishes feed on many species of small benthic algae, which are generally poorly digested, although consistent differences in food quality may occur among major taxa or growth forms of algae (Montgomery and Gerking 1980). There has been documentation of selective feeding in schooling surgeonfish responding to these differences, but nonselective feeding behavior is probably more common in wandering species (Randall 1967, Montgomery 1980, Montgomery et al. 1989, Roberts 1987).

Tangs are browsers, with lips and dentition for snipping off the tips and branches of algae. However, anatomical differences between species, particularly with the digestive tract is evident, leading one to suspect varying dietary preferences between these species. The blue tang, *A. coeruleus*, has a long thin-walled digestive tract, while the doctorfish, *A. chirurgus*, and ocean surgeonfish, *A. bahianus*, have sand-filled, muscular gizzard-like stomachs (Tilghman et al. 2001). Herbivorous fish are not known to produce cellulase or other enzymes to digest cell wall components (Lobel 1981). However, they are capable of digesting the materials inside plant cells if they have developed mechanisms to break the cell walls. This can be done in two ways. One method is Trituration, which occurs through chewing, or the material is be masticated in a muscular, gizzard-like stomach. An alternative strategy is the use of acidic stomach secretions, typically secreted by thin-walled stomachs (Lobel 1981). Acanthurids found in the Florida Keys possess both stomach types (Tilghman et al. 2001), so it is likely that they utilize both mechanisms to break algal cell walls.

This histological atlas focuses on *A. coeruleus* and includes major organs and tissues. Particularly note the stomach tissues of both species, which illustrate the difference in digestive strategies of the Caribbean Acanthurids. *Acanthurus chirurgus* was intentionally left out of this atlas, as its tissues are identical to those of *A. bahianus*.

The surgeonfish used for this atlas were obtained from the Florida Keys between 1999 - 2000. They ranged in size from 68 - 318 mm total length. Tissues were collected immediately after euthanasia with methane tricaine sulfonate (MS-222) and fixed with commercially available 10% buffered formalin (Fisher Scientific, Atlanta, GA). Calcified structures found in skin and gill tissues were decalcified with Cal-Ex® (Fisher Scientific, Atlanta, GA) prior to histological processing. The tissues were then embedded in paraffin, cut into five micron sections and stained with 7211 hematoxylin and eosin with phloxine (Richard Allen Scientific Co., Kalamazoo, MI). Photographs were taken with an Olympus DP-11 digital camera (Olympus Optics).

# Table of Contents

Introduction .....	1
Blue Tang Kidney .....	2
Blue Tang Gills .....	3
Blue Tang Intestine .....	4
Blue Tang Hepatopancreas .....	5
Blue Tang Spleen .....	5
Blue Tang Fin .....	6
Blue Tang Skin .....	6, 7
Blue Tang Stomach .....	8
Blue Tang Brain .....	9
Ocean Surgeon Kidney .....	10
Ocean Surgeon Gills .....	11
Ocean Surgeon Intestine .....	12
Ocean Surgeon Hepatopancreas .....	13
Ocean Surgeon Spleen .....	14
Ocean Surgeon Fin .....	15
Ocean Surgeon Skin .....	15
Ocean Surgeon Stomach .....	16
Ocean Surgeon Heart .....	17



© J.E. Randall

## Blue Tang

### **Acanthurus coeruleus**

**Size:** 125-275 mm

**ID:** This acanthurid is distinguished by its bright blue coloration. For this reason, it is the most sought after Atlantic surgeonfish for the aquarium trade.

**Class:** Actinopterygii / Ray-finned fishes

**Order:** Perciformes / Perch-like fishes

**Family:** Acanthuridae / Surgeonfishes

**Genus Species:** Acanthurus coeruleus / Blue Tang

**Habitat:** Reef. The blue tang is found from New York to São Paulo (Roca et al., 2002), and throughout the Caribbean and Gulf of Mexico.

**Food:** Primarily herbivorous; feed on brown and red benthic algae. (Tilghman et al., 2001)



© J.E. Randall

## Ocean Surgeon

### **Acanthurus bahianus**

**Size:** 125-275 mm

**ID:** These fish are distinguished from their coexisting acanthurids by their gray coloration, white caudal peduncle, and a highly lunate caudal fin.

**Class:** Actinopterygii / Ray-finned fishes

**Order:** Perciformes / Perch-like fishes

**Family:** Acanthuridae / Surgeonfishes

**Genus Species:** Acanthurus bahianus / Ocean Surgeonfish

**Habitat:** Reef. The ocean surgeonfish *Acanthurus bahianus* is found throughout the tropical western Atlantic from North America to southern Brazil (Roca et al., 2002). It also occurs in the oceanic islands of Bermuda and in the southern Atlantic islands of Fernando de Noronha, Atol das Rocas, Trindade, Ascension and St. Helena (Randall 1957, Floeter et al., 2001).

**Food:** Primarily herbivorous; feed on green and brown algae as well as small invertebrates. This species also ingests large amounts of sand, presumably to aid in the digestion of algae. (Tilghman et al., 2001)



© Luiz A. Rocha

## Doctorfish

### **Acanthurus chirurgus**

**Size:** 125-275 mm

**ID:** Distinctive vertical bars, white caudal peduncle

**Class:** Actinopterygii / Ray-finned fishes

**Order:** Perciformes / Perch-like fishes

**Family:** Acanthuridae / Surgeonfishes

**Genus Species:** Acanthurus chirurgus / Doctorfish


**Habitat:** Reef

**Food:** Primarily herbivorous; feed on green and brown algae as well as small invertebrates. This species also ingests large amounts of sand, presumably to aid in the digestion of algae. (Tilghman et al., 2001)


# Blue Tang

*Acanthurus coeruleus*

## Kidney



**Kidney** (100x). a) tubule. b) interrenal tissue. c) glomerulus.



**Kidney** (250x). a) proximal tubule segment. b) melanomacrophage center. c) collecting tubule. d) distal tubule segment. e) interrenal tissue. The apparent fewer number of melanomacrophage centers is not specific to this species and can be attributed to a number of factors.


The kidney is not divided into grossly distinct anterior and posterior sections as it is in many fish species. Function is evident at the microscopic level.

Most teleosts have anterior and posterior kidneys, with the anterior involved in excretory function and posterior acting in the hematopoietic role. In hematoxylin and eosin stained specimens, basophilic interrenal tissue is hematopoietic.


# Blue Tang

*Acanthurus coeruleus*

## Gills



**Gill (100x).** a) secondary lamella. b) cartilage of primary lamella




**Gill (250x).** a) epithelium. b) mucus cell. c) central venous sinus. d) cartilage.


# Blue Tang

## *Acanthurus coeruleus*

### Intestine



**Anterior section of intestine** (100x). a) lumen. b) goblet cell. c) circular muscle layer. d) longitudinal muscle layer. e) submucosa. f) mucosa.




**Intestine** (250x). a) lamina epitheliasis. b) submucosa. c) goblet cell. d) mucosa. e) muscularis circularis. f) muscularis longitudinalis.

# Blue Tang


*Acanthurus coeruleus*

## Hepatopancreas



**Hepatopancreas** (250x). (a) hepatic duct. Note the basophilic pancreatic tissue surrounding the ducts. (b) hepatic tissue. (c) pancreatic tissue.

## Spleen




**Spleen** (250x). a) red pulp. b) white pulp. c) melanomacrophage center.



# Blue Tang


## *Acanthurus coeruleus*

### Fin



**Fin** (100x). a) epidermis. b) bone. c) stratum spongiosum. d) melanin layer

### Skin




**Skin** (100x). a) scale. b) epithelium. c) muscularis. d) dermis.


# Blue Tang

## *Acanthurus coeruleus*

### Skin



**Lateral section of skin with lateral line canal (250x).** a) canal lumen. b) goblet cell. c) scale. d) muscle. e) loose connective tissue. f) sensory epithelium. The lateral line system of acanthurids is often the sight of extensive erosion in captive fish. Much speculation as to the etiology of these lesions has been discussed, but some anecdotal information as well as a few studies suggest that nutrition may be involved.



**Lateral section of skin with lateral line canal (250x).** a) sensory epithelium. b) loose connective tissue. c) muscle. d) scale. e) canal pore.


# Blue Tang

## *Acanthurus coeruleus*

### Stomach



**Stomach (100x).** Note the relatively low muscularis-to-mucosa size ratio in this species. a) muscularis. b) gastric gland. c) submucosa. d) lumen. e) mucosa. This species undoubtedly utilizes acidic stomach secretions to break the cell walls of their algal forage, as do other acanthurids that have similar stomach morphology (Lobel). In contrast to *A. bahianus*, wild-caught *A. coeruleus* stomachs are found full of algae, with little or no sand. Grossly, they are very thin-walled with relatively large luminal volumes.



**Stomach (250x).** a) muscularis. b) submucosa. c) gastric gland. d) goblet cell. e) mucosa.


# Blue Tang

## *Acanthurus coeruleus*

### Brain



**Sagittal section of the head (6.3x).** Notice the orientation of the brain anterior to the eye, which appears rotated anteriorly compared to other teleosts' brains. Area of inset is magnified in figure 15. (Negative image used to increase contrast.)




**Sagittal section of the brain (40x).** a) cerebellum. b) olfactory bulb. c) nasal epithelium. d) optic tectum. e) torus semicircularis.


# Ocean Surgeon

## *Acanthurus bahianus*

### Kidney



**Kidney** (100x)



**Kidney** (250x). a) melanomacrophage center. b) proximal tubule. c) distal tubule. d) hematopoietic tissue.


The apparent fewer number of melanomacrophage centers is not specific to this species and can be attributed to a number of factors. The kidney is not divided into grossly distinct anterior and posterior sections as it is in many fish species. Function is evident at the microscopic level.

Most teleosts have anterior and posterior kidneys, with the anterior involved in excretory function and posterior acting in the hematopoietic role. In hematoxylin and eosin stained specimens, basophilic interrenal tissue is hematopoietic.


# Ocean Surgeon

## *Acanthurus bahianus*

### Gills



**Gills** (100x). a) primary lamella. b) secondary lamellae.




**Gills** (250x). a) primary lamella cartilage. b) secondary lamella. The gills' function as an environmental interface make it one of the first sights of pathology development. Hyperplastic or hemorrhagic secondary lamellae are often indicative of environmental or pathogenic problems.


# Ocean Surgeon

## *Acanthurus bahianus*

### Intestine



**Anterior section of intestine (100x).** a) lumen. b) goblet cell. c) circular muscle layer. d) longitudinal muscle layer. e) submucosa. f) mucosa.




**Logitudinal section of the anterior intestine (250x).** a) muscularis longitudinalis. b) muscularis circularis. c) lamina propria. d) lamina epitheliasis. The gut length to total length ratio in these fish is about 2 : 1, and is suggestive of their primarily herbivorous diet. The thin-walled, tightly coiled intestines of these fish are filled with sand (in wild-caught fish) and are often teeming with various flora and fauna that may aid in the digestion of their algal forage.


# Ocean Surgeon

## *Acanthurus bahianus*

### Hepatopancreas



**Hepatopancreas** (100x). a) bile duct. Note basophilic pancreatic tissue (b), surrounding the hepatic ducts. Hepatic tissue (c) is eosinophilic.




**Hepatopancreas** (250x). a) pancreatic tissue. b) hepatic portal vein. c) hepatic tissue. The hepatic tissue is often interspersed with melanomacrophage centers, which are amber-colored areas. Smaller than normal hepatocytes may be indicative of starvation or other problems. Excess fat deposition in the hepatopancreas can reduce liver function, and is often seen in improperly fed aquarium specimens. A liver sample that floats in formalin is abnormally high in fat content.




# Ocean Surgeon

## *Acanthurus bahianus*

### Spleen



**Spleen (250x).** a) melanomacrophage center. b) red pulp. c) white pulp. As in the liver, splenic tissue is often interspersed with melanomacrophage centers. This organ's function as a hematopoietic center and blood filter make it a good place to look for pathogens. Gross granulomatous lesions may sometimes be seen after chronic exposure to bacterial or parasitic agents.




**Spleen with heavy nematode infestation (100x).** a) red pulp. b) white pulp. c) melanomacrophage center. d) nematode (cross section).

# Ocean Surgeon


## *Acanthurus bahianus*

### Fin



**Fin** (100x). a) epithelium. b) cartilage. c) melanophores. Fins are a common site of parasitism. Most of these can be seen grossly, however, and histological exam is often unnecessary. In acanthurids, monogenes and turbellarians are most often seen here. Fin spines in acanthurids with advanced head and lateral line erosion syndrome are often thickened and deformed.

### Skin




**Skin** (100x). a) muscle. b) dermis. c) scale (ctenoid type). d) epidermis. The skin is often the sight of pathology in fishes, since its role is a barrier to the environment and the pathogenic challenges therein. Most often, skin pathology can be determined by microscopic study of scrapings, however internal or systemic conditions that lead to skin pathology may only be observed in histological sections.


# Ocean Surgeon

## *Acanthurus bahianus*

### Stomach



**Stomach (250x).** The actual size of the muscularis (a) is ~ 4x that of the submucosa (c) in this species, and is even more pronounced in larger specimens. b) gastric gland. d) lumen.




**Stomach (250x).** a) submucosa. b) gastric gland. c) lumen.


# Ocean Surgeon

*Acanthurus bahianus*

## Heart



**Ventricle (250x)**



**Bulbus arteriosus (250x)**

## References

- Tilghman, G.C., Klinger-Bowen, R., Francis-Floyd, Ruth. 2001. Feeding electivity indices in surgeonfish (Acanthuridae) of the Florida Keys. *Aquarium Sciences and Conservation*. 3: 215-223.
- Rocha, L.A., Bass, A.L., Robertson, R., Bowen, B.W. 2002. Adult habitat preferences, larval dispersal, and the comparative phylogeography of three Atlantic surgeonfishes (Teleostei: Acanthuridae). *Molecular Ecology*. 11: 243-252.
- Floeter, S.R., Guimarães, R.Z.P., Rocha, L.A. 2001. Geographic variation in reef-fish assemblages along the Brazilian coast. *Global Ecology and Biogeography*. 10: 423-431.
- Randall, J.E. 1956. A revision of the surgeon fish genus *Acanthurus*. *Pacific Science*. 10:159-235.
- Carpenter, R.C. 1986. Partitioning herbivory and its effects on coral reef algal communities. *Ecological Monographs* 56: 343-363.
- Lobel, P.S. 1981. Trophic biology of herbivorous reef fishes: alimentary pH and digestive capabilities. *Journal of Fish Biology* 19: 365-397.



UNIVERSITY OF  
**FLORIDA**

Florida Sea Grant, University of Florida, PO Box 110409,  
Gainesville, FL, 32611-0409, (352) 392-2801, [www.flseagrant.org](http://www.flseagrant.org).