

JH
11
U33
No. 434

NOAA Technical Memorandum NMFS-SEFSC-434




RICE LIBRARY
NATIONAL MARINE FISHERIES SERVICE
BEAUFORT, NORTH CAROLINA 28516-9722

PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE HISTORY STAGES OF GERREID
FISHES OF THE WESTERN CENTRAL ATLANTIC


BY

ALLYN B. POWELL AND MICHAEL D. GREENE



U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southeast Fisheries Science Center
Beaufort Laboratory
101 Pivers Island Road
Beaufort, NC 28516-9722

May 2000



**PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE HISTORY STAGES OF GERREID
FISHES OF THE WESTERN CENTRAL ATLANTIC**

BY

ALLYN B. POWELL AND MICHAEL D. GREENE

**U.S. DEPARTMENT OF COMMERCE
William M. Daley, Secretary**

**National Oceanic and Atmospheric Administration
D. James Baker, Under Secretary for Oceans and Atmosphere**

**National Marine Fisheries Service
Penelope D. Dalton, Assistant Administrator for Fisheries**

May 2000

This Technical Memorandum series is used for documentation and timely communication of preliminary results, interim reports, or similar special-purpose information. Although the memoranda are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.

NOTICE

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or material mentioned in this publication. No reference shall be made to NMFS or to this publication furnished by NMFS, in any advertising or sales promotion which would imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein or which has as its purpose any intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

This report should be cited as follows:

Powell, A. B. and M. D. Greene. 2000. Preliminary guide to the identification of the early life history stages of gerreid fishes of the western central Atlantic. NOAA Technical Memorandum NMFS-SEFSC-434, 7 p.

W. J. Richards, Editor. NOAA Fisheries, 75 Virginia Beach Drive, Miami, FL

This report will be posted on the Bethune Cookman College NOAA Cooperative web site later in 2000 at URL: <http://208.152.233.21/NOAA/> and will also appear on the SEFSC web site at URL: <http://www.sefsc.noaa.gov/> It will be a chapter entitled Gerreidae in the "Guide to the early life history stages of fishes of the western central Atlantic".

Copies may be obtained by writing:

The authors at
NOAA Fisheries
101 Pivers Island Road
Beaufort, NC 28516-9722

National Technical Information Center
5825 Port Royal Road
Springfield, VA 22161
(800) 553-6847 or (703) 605-6000
<<http://www.ntis.gov/numbers.htm>>

Gerreidae: Mojarras

A.B. Powell & M. D. Greene

The family Gerreidae contains four genera and 13 species that occur in the western central North Atlantic. Adult gerreids are small to medium size fishes that are abundant in coastal waters, bays, and estuaries in tropical and warm temperate regions and sometimes occur in freshwaters. They are generally associated with grassy or open bottoms, but not with reefs. Gerreids are silvery fishes, with deeply forked tails, and extremely protrusible mouth that points downward when protracted. They apparently feed on bottom-dwelling organisms and at least one species (*Eucinostomus gula*) shows a distinct transition, during the juvenile period, from a planktivore (exclusively copepods) to a carnivore that includes a diet of almost solely polychaetes (Carr & Adams, 1973; Robins and Ray, 1987; Murdy et al., 1997)

Gerreids eggs are small and pelagic. *Gerres japonicus* (from Japanese waters) eggs are 0.60-0.63 mm in diameter with a single oil globule 0.18-0.20 mm in diameter (Mito, 1963). Larvae from this species hatches at 1.4 mm. Eggs from an Atlantic gerreid, *Diapterus rhombeus* measured 0.70-0.72 mm in diameter with a single oil globule 0.18-0.20 mm in diameter (Rass, 1972). Eggs from the genera *Eucinostomus* and *Eugerres* have not been described. Morphologically, larval gerreids are moderately deep bodied, have a pointed head in the postflexion stage, large eyes, short coiled gut, and a space between the anus and anal fin. Just prior to the flexion stage the prominent premaxillary process begins to form and by the early postflexion stage is well developed and the mouth becomes protractile. Head spination is not well developed. Small preopercular spines develop during the flexion stage, but are not conspicuous until the postflexion stage. Small supracleithral spines might form in some species (Leis & Rennis, 1983). Gerreid larval pigment typically occurs on the ventral midline, tip of notochord and eventually covers the base of the caudal fin, dorsal and anterior portions of the gut, hindgut, ventrum of the gut, anterior to the cleithral symphysis, and dorsal midline at the caudal peduncle that continues along the second dorsal fin ray with development (Leis & Rennis, 1983; Watson, 1996).

Gerreids can generally be distinguished from other percoid taxa by meristic characters , such as myomere and fin ray counts, and morphological characters such as the short preanal length, lack of preopercle spines prior to late flexion or early postflexion and weak spines thereafter, and pigment characters discussed above.

Descriptions of the early stages of gerreids in the western central North Atlantic are rare and counts of meristic characters are not useful to distinguish species of *Eucinostomus*, except *E. lefroyi*, which has two anal spines, nor to distinguish *Gerres* from *Eucinostomus* (Table Gerreidae 1). Descriptions of *Eucinostomus* sp (p). from the California Current region (Watson, 1996) indicate a hatching size of <2.1 mm and a flexion length of approximately 4.0-5.5 mm. Pigmentation of preflexion larvae consist of 18-20 melanophores on the ventral midline that decrease to 12-13 at beginning of flexion; dorsally on gas bladder and hindgut, and anterior gut pigment after 2.4 mm; one melanophore on the anterior and posterior of the hindgut; a pair over midbrain after 2.4 mm. Flexion larvae have 2-4 melanophores on hypural margin; ventral midline decreases to 6-8; posterior portion of dorsal midline at 4.6 mm; melanophore anterior to the cleithral symphysis; paired embedded hindbrain melanophores (4.7 mm) and an additional pair occurs on the midbrain surface (4.8 mm); 0-1 melanophore at dorsal area of preopercle. Melanophores on postflexion larvae increase over the brain, along the fin bases , and on gut and gas bladder (Watson, 1997). A 9.5 mm SL *Eucinostomus* tentatively identified as *E. lefroyi* appears to lack the intensity of pigment on the fin bases compared to the California *Eucinostomus*. *Gerres* larvae from Australian waters resemble *Eucinostomus* except they have less ventral midline melanophores during the preflexion stage, and head pigment does not develop till after postflexion. (Lies & Rennis, 1983). Furthermore , larger postflexion *Gerres* larvae exhibit lateral pigment in the caudal region. Australian *Gerres* hatch at <2.1 mm, and caudal flexiion occurs 3.5-5.0 mm (Leis & Rennis, 1983). In the species account, the life history and meristic data is for *Eucinostomus argenteus* and *E. gula* and 3 illustrations are provided of examples of *Eucinostomus* from our area.

Table Gerreidae 1. Meristic characters for the gerreid species that occur in the western central North Atlantic. All species have 10+14 vertebrae, 9+8 principal caudal fin rays, and I,5 pelvic fin rays. Meristic counts were obtained from Curran (1942), Dahlberg (1975), Hildago and Carrió (1983), Deckert and Greenfield (1987) and Murdy et al. (1987). Values given are the mode and, in parentheses, the range.

Species	Fin rays		
	D	A	GR
<i>Diapterus auratus</i>	IX,10 (VII,10-IX,11)	III,8 (III,8-III,9)	12-13 (12-15)
<i>D. rhombeus</i>	IX,10	II,9 (II,8-II,10)	16-18
<i>Eucinostomus argenteus</i>	IX,10	III,7	8
<i>E. gula</i>	IX,10	III,7	8
<i>E. harengulas</i>	IX,10	III,7	8
<i>E. havana</i>	IX,10	III,7	8
<i>E. jonesi</i>	IX,10	III,7	8
<i>E. lefroyi</i>	IX,10	II,8	8
<i>E. melanopterus</i>	IX,10	III,7	9
<i>Eugerres brasiliianus</i>	IX,10 (IX,10-11)	III,7 (III,7-8)	11-12 (11-13)
<i>E. mexicanus</i>	IX,10 (VIII-X,9-11)	III,8 (III,7-8)	14-15 (13-16)
<i>E. plumieri</i>	IX,10 (VIII-IX,10-11)	III,8 (III,7-9)	14-16 (13-17)
<i>Gerres cinereus</i>	IX,10	III,7	8

Table Gerreidae 2. Distributions of gerreid species that occur in the western central North Atlantic. Distribution data were obtained from Robins and Ray (1986), Deckert and Greenfield (1987), Boschung (1992) and Murdy et al. (1997).

Species	Distribution
<i>Diapterus auratus</i>	Atlantic coast of US from NC south to Florida and through Gulf of Mexico; Central America coast south to Brazil; West Indies.
<i>D. rhombeus</i>	West Indies; Mexico south along Central American coast; northern South America to Brazil.
<i>Eucinostomus argenteus</i>	NJ, Bermuda, and n. Gulf of Mexico to se. Brazil. Always in shallow water; enters fresh water ; absent from reefs.
<i>E. gula</i>	Mass., Bermuda and n. Gulf of Mexico to Argentina. Shallow water, except coral reefs; enters fresh water in limestone regions.
<i>E. harengulas</i>	Northern and eastern Gulf of Mexico; enters fresh water.
<i>E. havana</i>	Most common in Bahamas and West Indies; Bermuda, FL, and Bahamas to Brazil. Inshore habitats but does not appear to enter brackish water.
<i>E. jonesi</i>	Bermuda and Florida to Brazil.
<i>E. lefroyi</i>	NC, Bermuda and n. Gulf of Mexico to Brazil.
<i>E. melanopterus</i>	Georgia to Brazil; Gulf of Mexico; West Indies; absent from Bahamas, favors continental lagoons and bays, and enters fresh water.
<i>Eugerres brasilianus</i>	West Indies and Atlantic coast from Belize to southern Brazil; enters rivers.
<i>E. mexicanus</i>	Atlantic slope of southern Mexico and northern Guatemala; restricted to fresh waters.
<i>E. plumieri</i>	Atlantic coast of US from SC to FL; west coast of FL; West Indies; Mexico, south along Central American coast to Columbia, entering rivers.
<i>Gerres cinereus</i>	Bermuda, FL and Gulf of Mexico to se. Brazil; common in coastal waters, seagrass beds, areas near reefs and mangrove channels.

Literature Cited

- Boschung, H. T. 1992. Catalogue of freshwater and marine fishes of Alabama. *Bull. Alabama Mus. Nat. Hist.* (14): 266 p.
- Carr and Adams, 1973. Food habits of juvenile marine fishes occupying seagrass beds in the estuarine zone near Crystal River, Florida. *Trans. Amer. Fish. Soc.* 102: 511-540.
- Curran, H. W. 1942. A systematic revision of the gerrid fishes referred to the genus *Eucinostomus*, with a discussion of their distribution and speciation. Ph.D. Diss., Univ. Mich., Ann Arbor. Michigan. 183 p.
- Dahlberg, M. D. 1975. Guide to the coastal fishes of Georgia and nearby states. Univ. Georgia Press, Athens, Georgia. 187 p.
- Deckert, G. D. and D.W. Greenfield. 1987. A review of the western Atlantic species of the genera *Diapterus* and *Eugerres* (Pisces: Gerreidae). *Copeia* 1987: 182-194.
- Hidalgo, M. B. and E. G. Carrió. 1983. Estudio morfométrico de *Eugerres brasiliensis* (Cuvier in Cuvier et Valenciennes, 1830). *Rev. Invest. Mar.* 4: 63-90. [In Spanish].
- Leis, J. M. And D.S. Rennis. 1983. The larvae of Indo-Pacific coral reef fishes. New South Wales Univ. Press, Kensington. 269 p.
- Mito, S. 1963. Pelagic fish eggs from Japanese waters--III. Percina. *Jpn. J. Ichthol.* 11: 39-64. [In Japanese].
- Murdy, E. O., R. S. Birdsong and J. Musick. 1997. Fishes of Chesapeake Bay. Smithsonian Press, Washington, D.C. 324 p.
- Rass, T. S. 1972. Ichthyoplankton from Cuban waters. Pelagic fish eggs. *Tr. Inst. Okeanol. Akad. Nauk SSSR* 93: 5-41. [In Russian].
- Robins, C. R. and G. C. Ray, 1986. A field guide to Atlantic coast fishes of North America. Houghton Mifflin Co., Boston. 354 p.
- Watson, W. 1996. Gerreidae: mojarras. Pages 998-1001 in H. G. Moser (ed.). The early stages of fishes in the California Current region. *Calif. Coop. Ocean. Fish. Invest., Atlas No. 33.* 1505 p.

FAMILY GERREIDAE

*Eucinostomus argenteus/gula**

MERISTICS

Vertebrae:	
Precaudal	10
Caudal	14
Total	24
Number of Fin Spines and Rays:	
First Dorsal	IX
Second Dorsal	10
Total	19
Anal	III, 7
Pectoral	
Pelvic	I,5
Caudal	
Dorsal Secondary	10
Principal	9+8
Ventral Secondary	9-10

LIFE HISTORY

Range: *E. argenteus* – NJ, Bermuda & n. Gulf of Mexico to se. Brazil. *E. gula* – Mass, Bermuda & n. Gulf of Mexico to Argentina

Habitat: Shallow water except reefs

ELH Pattern:

Spawning:

 Season: postflexion larvae in June & November in NC

 Mode: oviparous, planktonic eggs & larvae

LITERATURE

Johnson 1984; Murdy et al.

EARLY LIFE HISTORY DESCRIPTION

EGGS: Unknown

LARVAE:

Pigmentation: Large stellate in U-shaped pattern on mid-hind brain; embedded in hind brain; generally 2 melanophores behind eye; over gut & generally 1 anteriorly on gut; embedded pair on hyoidal musculature opposite anterior gut pigment; generally distinct embedded pigment on gut at base of pectoral fin; 1-3 at base of dorsal fin; mainly 2 post dorsal midline; 3-8 at base of anal fin; 3-6 post anal midline; hypural plate; 1-2 ventral gut; 2-3 on isthmus.

Diagnostic Characters: Myomeres 24; short preanal length; weak preopercle spines; pigmentation.

*All larvae used in this description were collected at Beaufort Inlet, NC in June and November. The species are either *E. argenteus* or *E. gula*.

ILLUSTRATIONS

Top: Postflexion larva 11.8 mm SL Beaufort Inlet, NC. Either *E. argenteus* or *E. gula* (original, M. D. Greene)

Middle: Postflexion larva 9.5 mm SL Gulf of Mexico. Tentatively *E. lefroyi* (original, J. Javech)

Bottom: Postflexion larva 10.9 mm SL Bahamas, (original, W. Laroche)

GERREIDAE

Eucinostomus spp.

