

**BALANOKARPOL AND AMPELOPSIN H, TWO OLIGORESVERATROLS FROM
STEM BARK OF *Hopea odorata* (DIPTEROCARPACEAE)**

Sri Atun, Nurfina Aznam, Retno Arianingrum, Masatake Niwa

Two oligoresveratrol, namely balanokarpol (1) and ampelopsin H (2) had been isolated from the steam bark of *Hopea odorata* (Dipterocarpaceae). The structure of this compound were elucidated based on physical and spectroscopic data (MS, ¹H and ¹³C NMR 1D and 2D). The activity of these compounds was evaluated against the 2-deoxyribose degradation induced by the hydroxyl radical generated via a Fenton-type reaction. The result of this study showed that activity each compounds as radical hydroxyl scavenger of balanokarpol, and ampelopsin H with an IC₅₀ 1802,3 and 4840,0 µg/ml. respectively. Each compound showed low activity. Vitamin C (IC₅₀ 83,9 µg/ml) and butylated hydroxyl toluene (1328,0 µg/ml) used as positif control. These results suggest that oligoresveratrols from stem bark of *H. odorata* may be useful as potential sources of natural antioxidants.

Keyword: balanokarpol; ampelopsin h; antioxidant; dipterocarpaceae

FMIPA, 2007 (PEND. KIMIA)