

# Practical Guidelines to Minimize Language and Cognitive Confounds in the Diagnosis of CAPD:

## A Brief Tutorial

Gail D. Chermak<sup>a</sup>

Doris-Eva Bamiou<sup>b</sup>

Vasiliki (Vivian) Iliadou<sup>c</sup>

Frank E. Musiek<sup>d</sup>

<sup>a</sup> Department of Speech and Hearing Sciences, Elson S. Floyd College of Medicine, Washington State University Health Sciences Spokane, WA, USA; <sup>b</sup> UCL Ear Institute, London; <sup>c</sup> Clinical Psychoacoustics Lab, 3rd Psychiatric Department, Neuroscience Sector, Medical School, Aristotle University of Thessaloniki; <sup>d</sup> Neuroaudiology Lab, University of Arizona, Tucson, USA.

**Key Words:** Central auditory processing disorder; confounds, differential diagnosis; central auditory nervous system

**Acronyms:** 6- CIT (Cognitive Impairment Test); ADHD (attention deficit hyperactivity disorder); CAPD (central auditory processing disorder); CANS (central auditory nervous system); DPT (Duration Pattern Test); DR (directed report); FPT (Frequency Pattern Test); FR (free report); LiSN-S (Listening in Spatialized Noise-Sentences Test); MLR (Middle Latency Response); MMSE (Mini-Mental State Exam); MoCA (Montreal Cognitive Assessment)

**Corresponding Author:** Gail D. Chermak, PhD, Department of Speech and Hearing Sciences, Elson S. Floyd College of Medicine, Washington State University Health Sciences Spokane, PO Box 1495, Spokane, WA 99210-1495; [chermak@wsu.edu](mailto:chermak@wsu.edu)

## Abstract

Objective: Provide audiologists with strategies to minimize confounding cognitive and language processing variables and accurately diagnose central auditory processing disorder (CAPD).

Design: Tutorial.

Study Sample: None

Results: Strategies are reviewed to minimize confounding cognitive and language processing variables and accurately diagnose CAPD.

Conclusions: Differential diagnosis is exceedingly important and can be quite challenging. Distinguishing between two or more conditions presenting with similar symptoms or attributes requires multidisciplinary, comprehensive assessment. To ensure appropriate interventions, the audiologist is a member of the multidisciplinary team responsible for determining whether there is an auditory component to other presenting deficits or whether one condition is responsible for the symptoms seen in another. Choice of tests should be guided both by the symptoms of the affected individual, as established in an in depth interview and case history, the individual's age and primary language, and by the specific deficits reported to be associated with specific clinical presentations. Knowing which tests are available, their strengths and limitations, the processes assessed, task and response requirements, and the areas of the central auditory nervous system (CANS) to which each test is most sensitive provides the audiologist with critical information to assist in the differential diagnostic process.

## Chermak      Minimize Confounds CAPD Diagnosis

Central auditory processing disorder (CAPD) is a disorder of the central auditory nervous system (CANS) (i.e. beyond the level of the auditory nerve), which presents as deficits in the perceptual processing of auditory stimuli and in the underlying neurobiological activity that gives rise to the electrophysiological auditory potentials. The predominant deficits characterizing CAPD manifest in the auditory modality (AAA, 2010). Functional deficits associated with CAPD include: difficulty understanding spoken language in competing message or noise backgrounds, in reverberant acoustic environments, or when rapidly presented; difficulty localizing the source of an auditory signal; misunderstanding messages; responding inconsistently or inappropriately; frequently requesting repetitions; difficulty with subtle intonation and prosodic cues; difficulty with similar sounding words; difficulty following complex auditory directions/commands; difficulty “hearing” on the phone; difficulty learning songs, nursery rhymes; poor musical and singing skills and/or appreciation of music; difficulty learning foreign language or novel speech materials, especially technical language; listening difficulties; and academic difficulties (e.g., reading, spelling, and /or learning problems (Musiek & Chermak, 2014). Clearly, many of these behavioral deficits also are seen in individuals diagnosed with other disorders, including peripheral hearing loss, language disorders, and cognitive disorders. However, these functional deficits observed in individuals with CAPD derive from bottom-up (i.e., sensory, data-driven) processing deficits, although they bidirectionally interact with language and cognitive deficits. Auditory discrimination difficulties and difficulties processing rapid spectrotemporal acoustic changes may lead to poor speech sound representation, which has been linked to poor reading and spelling skills, as well as to difficulties in phonological awareness skills (Bellis, 2002; Kraus et al, 1996; Tierney & Kraus, 2013; Power et al, 2013). Poor speech perception gives rise to “fuzzy” or “un

## Chermak      Minimize Confounds CAPD Diagnosis

derspecified” lexical and phonological representations and to weak verbal short term memory (Studdert-Kennedy, 2002).

CAPD frequently is diagnosed in children and adults presenting other comorbid or primary diagnoses, including peripheral hearing loss (Khavarghazalani et al, 2016), confirmed CANS deficits (e.g., multiple sclerosis- Musiek et al, 2005; Musiek & Weihing, 2011), traumatic brain injury (Saunders et al, 2015; Gallun et al, 2012), cerebrovascular accidents (e.g., Stroke-Koohi et al, 2016), psychiatric conditions (e.g., schizophrenia- Iliadou & Iakovides, 2003; Iliadou et al, 2013; Ramage et al, 2015;), cognitive deficits (e.g., Alzheimer’s disease- Quaranta et al, 2015; Griffiths et al, 2012), and developmental disorders (e.g., attention deficit hyperactivity disorder [ADHD], dyslexia, language disorder). Sharma et al (2009) reported comorbidity of CAPD, language impairment, and reading disorders in 47% of a sample of school-age children (7-12 years) diagnosed with CAPD. Iliadou et al (2009) reported CAPD co-morbid with developmental dyslexia in 25% of cases. Atypical N1-P2 (i.e., flatter slopes) to tones, vowels, and CVs was seen in 38% of 6-12 year-old children with specific language impairment or specific reading disability (McArthur et al, 2009). Subcortical differentiation of stop consonants (speech auditory brainstem response) correlated with phonological awareness, reading, and speech-in- noise perception in children 8-13 years (Banai et al, 2009). It should be noted that the fundamental relationships among auditory processing, language, and cognitive measures may differ in typical compared to non-typically developing populations (e.g., Grube, Cooper, Kumar, Kelly, & Griffiths, 2014), and as a function of age as well (e.g., Besser, Festen, Goverts, Kramer, & Pichora-Fuller, 2015).

The co-morbidity of CAPD with neurodevelopmental disorders, as well as with acquired central nervous system conditions (e.g., traumatic brain injury, cerebral vascular accidents, etc.)

## Chermak      Minimize Confounds CAPD Diagnosis

can be most easily explained on the basis of brain organization and function. Our current understanding suggests that perceptual, language, and cognitive functions rely on extensive networks of shared neurophysiologic and vascular substrate. Networks are nonmodular, temporally coupled, interfacing, polymodal or multisensory, overlapping, interconnected, and synchronized (Buschman & Miller, 2007; Musiek et al, 2005; Price et al, 2005; Willott, 1991). The brain is organized in terms of multipurpose representations that different regions support. A given brain region can be useful for many different functions— e.g., regions within the brain’s putative perceptual system also play a role in (attention) and memory (Lopez-Aranda et al, 2009). For example, proper functionality of the posterior temporal gyrus is found to be critical for both short-term memory and language (Leff et al, 2009; Acheson et al, 2011). To distinguish between two or more conditions presenting with similar symptoms or attributes requires multidisciplinary, comprehensive assessment. To ensure that the most appropriate interventions are employed, the audiologist is a member of the multidisciplinary team responsible for determining whether there is an auditory component to other presenting deficits or whether one condition is responsible for the symptoms seen in another.

Differential diagnosis is exceedingly important and can be quite challenging. To further illustrate this challenge consider the potential sources of speech understanding deficits in noise, which can result from peripheral (cochlear) auditory issues, central auditory issues, cognitive issues (including working memory deficits, attention deficits), language issues, and age (i.e., younger children and older adults experience greater difficulty understanding speech in noise). Among individuals with CAPD, auditory processing deficits may vary in different cohorts of affected individuals. Choice of tests should be guided by the symptoms of the affected individual,

## Chermak      Minimize Confounds CAPD Diagnosis

as established by an in-depth interview and case history, the individual's age and primary language, and by the specific deficits reported to be associated with specific clinical presentations (e.g. Cumming, Wilson, & Goswami, 2015; Goll, Crutch, Loo, Rohrer, Frost et al, 2010). In selecting tests, the audiologist must recognize that one symptom (e.g., understanding speech in noise) can result from many underlying central auditory processing deficits (e.g., temporal processing, localization, spatial release from masking, performance with competing/degraded auditory signals), as well as language processing or cognitive issues. The audiologist should select a battery of tests that assess the multiple central auditory processes potentially implicated in a patient's presenting complaints. Speech-in-noise or competition and dichotic listening tests would be obvious first line tests to examine difficulties hearing in noise. Similarly, compressed speech (a type of monaural low-redundancy measure) would be helpful to examine complaints of difficulty following rapid speech. The masking level difference (MLD), the Listening in Spatialized Noise-Sentences Test (LiSN-S) (Cameron & Dillon, 2007; Cameron et al, 2009), and localization tasks would be useful to examine difficulties localizing sound and spatial processing difficulties. Frequency pattern and duration pattern tests (Musiek, 1994), as well as the complex auditory brainstem response (cABR) may provide insights regarding music appreciation difficulties. We discuss below tests and strategies to examine central auditory processing to minimize the influence of cognition and language. When cognitive or language processing deficits are suspected or confirmed, the audiologist should consider the use of auditory evoked potentials rather than behavioral auditory tests, interpreted within the context of multidisciplinary evaluations. Knowing which tests are available, their strengths and limitations, the processes assessed, task and response requirements, and the areas of the CANS to which each test is most sensitive provides the

## Chermak      Minimize Confounds CAPD Diagnosis

audiologist with critical information to assist in the differential diagnostic process. A summary of recommended tests and their strengths and weaknesses is provided in Table 1. The remainder of this brief tutorial provides strategies to minimize confounding cognitive and language processing variables and accurately differentially diagnose CAPD. The audiologist and other members of the multidisciplinary team must determine whether a central auditory processing deficit (e.g., temporal resolution deficit) is adversely impacting attention, whether a more global attention deficit (e.g., faulty temporal synchronization of attention) is impeding central auditory processing, or both issues are impacting the patient's processing of acoustic signals. Similarly, the audiologist in cooperation with the speech-language pathologist and psychologist must determine whether reported listening difficulties result from an auditory problem, a language processing disorder, a cognitive deficit (e.g., attention, executive function, or working memory), or some combination of these deficits. While there is some disagreement between professional associations in the United States and European associations (e.g., the British Society of Audiology [BSA, 2011]) as to the standard test battery to be used to diagnose CAPD, the two major sets of guidelines published by the American Speech-Language-Hearing Association (ASHA) and the American Academy of Audiology (AAA) present consistent positions and recommendations (i.e., AAA, 2010; ASHA, 2005). Notwithstanding the potential interactions among auditory, linguistic, and cognitive factors, it is our position, as well as that of professional associations in the United States that listening difficulties seen in CAPD result primarily from auditory perceptual deficiencies rather than from global attention deficits or behavioral regulation deficits (AAA, 2010). This was underscored recently by Gyldenkaerne et al (2014) who reported a weak correlation between auditory processing and cognitive tests, concluding that poor auditory processing

## Chermak      Minimize Confounds CAPD Diagnosis

certainly is not the mere reflection of attention deficits. Similarly, Weihing et al (2015) concluded that central auditory tests are not likely governed by cognition given the results of their factor analysis in which central auditory tests loaded on separate factors, suggesting a unique variance associated with each test. Indeed, a number of studies have examined the relationship between CAPD and cognitive function; however, the studies use different auditory and cognitive test batteries, and some studies rely on auditory tests with no documented sensitivity to CAPD. Some studies conclude that cognitive disorders and CAPD exist independently (e.g., Tillery, Katz, & Keller, 2000; Sharma, Purdy, & Kelly, 2009; Rosen, Cohen, & Vanniasegaram, 2010), while others suggest that CAPD is itself a cognitive disorder (e.g., Moore, Ferguson, Edmondson-Jones, Ratib, & Riley, 2010; Ferguson, Hall, Riley, & Moore, 2011). The reader is referred to Tomlin, Dillon, Sharma and Rance (2015) and Weihing et al (2015) for reviews of studies examining the interaction between cognition and central auditory processing.

### Controlling Extraneous Variables and Test Selection

In general, confounds can be minimized by controlling extraneous variables. The audiologist must select central auditory tests that are appropriate to the patient's age, education, linguistic background, and cognitive ability. One should consider task requirements, including the modalities used, and the task's language and cognitive demands. The patient's cultural and linguistic backgrounds also are important as these factors can affect response time and accuracy of verbal responses (Pimentel & Inglebret, 2014). For example, individuals from some cultural backgrounds may pause for an extended period before responding to a stimulus item (Wallace, Inglebret, & Friedlander, 1997). (Not surprisingly, the strategies and approaches reviewed here may be useful as well when testing non-native English speakers.) Selecting central auditory tests that



## Chermak      Minimize Confounds CAPD Diagnosis

are less influenced by peripheral hearing loss (e.g., frequency patterns, duration patterns) and considering the influence of medications (e.g., central nervous system stimulants or depressants) is important as well. Peripheral auditory function should be evaluated thoroughly prior to the central auditory evaluation (Iliadou et al, 2015). Perhaps the single most important step in successful audiologic assessment of the child with diagnosed and medically-managed ADHD is to ensure that the child received an effective dose of medication immediately before the test session, since for children with diagnosed ADHD who are treated medically, valid audiologic assessment is difficult without medication (Chermak et al, 1999). It is not uncommon, however, to see children diagnosed with ADHD who are not taking prescribed medication who nonetheless still perform within normal limits on central auditory processing tests, thereby confirming the absence of comorbid CAPD.

Behavioral tests using nonverbal or simple speech stimuli (e.g., frequency or duration patterns, gaps-in-noise, dichotic digits), as well as tests that minimize memory load, and employ a simple response mode (e.g., gap detection, masking level difference) reduce the influence of language and cognitive factors. The linguistic load of instructions also should be minimized to ensure the patient understands the task requirements. One might question why speech stimuli are used at all, given the potential confound with language. With few exceptions (e.g., Pediatric Speech Intelligibility Test [Jerger & Jerger, 1984]), most central auditory tests are not appropriate for use with children under seven years of age due to the extremely challenging nature of the tests and the large performance variability seen among children younger than seven years, despite normal auditory processing function. Young (7-9 years old) patients may attend less to non-speech tasks, which are more abstract than speech tasks, which would inject a confound of

## Chermak      Minimize Confounds CAPD Diagnosis

inattention for non-speech stimuli. Moreover, speech and language signals provide access to different CANS processing mechanisms than do non-speech/language stimuli (Grossman et al, 2010). Speech stimuli have greater ecological validity given the degree of temporal processing required for accurate perception of spoken language, which is significantly greater than required for perception of non-speech sounds (Fitch et al, 1997; Griffiths et al, 1999; Shannon et al, 1995; Zatorre & Belin, 2001). Finally, processing of speech signals may be more vulnerable to CANS dysfunction and central auditory deficits may only be revealed with speech tasks (e.g., Benavidez et al, 1999; Johnson et al, 2005; Russo et al, 2005). The reader is referred to the American Academy of Audiology Clinical Practice Guidelines (2010) for an overview, and to Musiek & Chermak (2014) for a more expanded description of the central auditory tests mentioned in this tutorial.

### Test Design, Intra-Test Comparisons, Psychometric Functions, and Performance Interpretation

Intra-subject comparisons (e.g., interaural differences), which control in part for non-auditory factors, also maximize the potential to assess auditory function by holding contributions from cognitive or language systems constant between ears. Designing tests that rely on tasks that differ in the demands placed on central auditory processing skills, but differ minimally in the demands they place on cognitive abilities are more likely to produce performance more reflective of auditory function. The LiSN-S (Cameron & Dillon, 2007; Cameron et al, 2009) employs this strategy by measuring the ability of the auditory system to take advantage of the spatial separation of the target (simple sentences) and competing sounds (children's stories). Other intra-test comparisons (e.g., performance-intensity functions in which one varies presentation levels or response mode (e.g., humming v. labeling in the Frequency Pattern Test [FPT] and Duration

## Chermak      Minimize Confounds CAPD Diagnosis

Pattern Test [DPT]) also offer insights (Musiek, Baran, & Pinheiro, 1990; Musiek & Pinheiro, 1987). If a patient can perform at some intensity level or can hum, but not label, the audiologist knows that the subject understands the directions and the task. Assessing performance in non-manipulated conditions (e.g., monaural versus dichotic, non-filtered versus filtered, etc.) ensures that performance deficits seen on central auditory tests are due to the acoustic manipulations rather than to lack of familiarity with the language and/or significantly reduced memory skills. The use of electrophysiological measures also assists the audiologist in sorting out potential confounds. For example, the P300 is absent with lesions in auditory (temporal-parietal junction) areas (Knight et al, 1989), while damage to lateral parietal cortex or prefrontal cortex does not interfere with P300 generation. Similarly, Kileny et al (1987) demonstrated the effects of temporal lobe, but not frontal lobe lesions, on the middle latency response (MLR).

The influence of cognition on central auditory test performance is rather minimal. Correlations between central auditory tasks and attention are weak, accounting for minimal shared variance, which indicates that CAPD does not merely reflect attention deficits (Weihing et al, 2015). For some central auditory tests, no significant correlations are seen with attention, indicating that attention may play no significant role in some central auditory test outcomes (Gyldenkaerne et al, 2014; Shinn et al, 2005). Nonetheless, in addition to following the approaches recommended above (i.e., ensuring that the patient understands the directions and the task, minimizing language load, and using a simple response mode) there are several other strategies that will minimize the cognitive demands of the task and thereby lessen the potential for cognitive confounds, allowing a more singular focus on auditory capacities.

## Chermak      Minimize Confounds CAPD Diagnosis

Before administering a central auditory test, the audiologist should ensure that the task does not exceed the patient's working memory capacity. For example, one should confirm that the patient's digit span exceeds the threshold level of four in a noncompeting condition (i.e., non-manipulated condition) prior to presenting the stimuli in the dichotic, competing condition. This provides at least one within-patient control for possible confounding effects of cognition on central auditory test performance and assists in ensuring that any observed dichotic deficit is due more to the auditory processing of the dichotic competition rather than to generalized memory problems. In addition, as noted above, designing tests (e.g., LiSN-S) that rely on tasks that differ in the demands placed on central auditory processing skills, but differ minimally in the demands placed on cognitive abilities should produce performance more reflective of auditory function.

Dichotic testing offers an interesting opportunity to differentiate cognitive from auditory effects. Specifically, the directed report (DR) or divided-attention condition in which the patient repeats stimuli in one or both ears in a specified order correlates most closely with cognitive function, as this condition requires the listener to recruit additional cognitive (e.g., attention, memory) mechanisms to accurately perform the task. The free report (FR) condition in which the patient repeats stimuli directed to both ears, usually in any order, is less susceptible to cognitive confound, even though the paradigm may require the listener to report a greater number of stimuli (Cowell & Hugdahl, 2000; Hallgren et al, 2001; see Hugdahl et al, 2009 for review). While the role of cognition might not be directly extrapolated across dichotic tasks using different stimuli (e.g., Shinn et al., 2005), performance decrements in the FR condition are more likely to be a consequence of true auditory perceptual asymmetries than are performance decrements in the DR condition. Nonetheless, the laterality indexes for the DR of dichotic digits and (non-rhyme)

## Chermak      Minimize Confounds CAPD Diagnosis

words follow a similar developmental trajectory (Moncrieff, 2011), and similar to what is observed for CVs (Hugdahl et al., 2009), directing attention to one ear strongly modulates the laterality index for digits and this modulation is cognitive dependent (Bouma & Gootjes, 2011). These findings support our assertion that for a given dichotic task, the DR mode depends more on cognitive resources than the FR condition. It should also be noted that the DR condition involves an additional degree of auditory complexity in that it requires lateralization of the stimuli. This suggests that while DR is more cognitively dependent than FR, DR requires transfer of sounds across the corpus callosum and therefore still provides an index of sensory demands (Rauschecker & Tian, 2000). Because the two report conditions tap into different underlying auditory processes (i.e., binaural integration v. binaural separation), and cognitive factors confound to different degrees, the use of both FR and DR conditions for the same stimuli, may, when administered and interpreted appropriately, provide useful insight into the specific auditory processes (i.e., binaural integration v. binaural separation) that are impacted by a given central auditory disorder. Jerger and Martin (2006) argued that using divided and directed attention dichotic listening procedures incorporating the same test stimuli may be helpful in sorting out the relative contributions of cognitive and auditory specific factors influencing interaural asymmetry, suggesting that auditory-specific deficits should lead to poor dichotic performance under both modes of administration (Martin et al., 2007).

### Language and Cognitive Screening

In addition to peripheral hearing testing prior to central auditory testing, it is helpful for the audiologist to have some sense of the language and cognitive function of patients seen for central

## Chermak      Minimize Confounds CAPD Diagnosis

auditory evaluation. Patients should be at least screened, if not more fully assessed in the multi-disciplinary evaluation of patients referred for CAPD. For children, such screens might include the Clinical Evaluation of Language Fundamentals, 5th edition (CELF-5) (Wiig, Semel, & Secord, 2013), the auditory backward and forward digit span subtest for working memory (Wechsler, 2014), auditory vigilance (attention) (e.g., Integrated Visual and Auditory Continuous Performance Test (IVA-CPT) (Sandford & Turner 2009), and nonverbal intelligence (e.g., Test of Nonverbal Intelligence (4th edition) TONI-4 (Brown, Sherbenou, & Johnsen, 2010). These screening tests must be administered by the member of the multidisciplinary team in whose scope of practice such testing falls.). For adults, cognitive screening measures include: the Montreal Cognitive Assessment (MoCA) (Nasreddine et al, 2005); Mini-Mental State Exam (MMSE) (Folstein et al, 1975), and the 6- CIT (Cognitive Impairment Test) Kinghill Version (Brooke & Bullock, 1999). Results of any of these cognitive screens should be interpreted with caution when CAPD or hearing loss is suspected as their results may be showing deficits that may be at least partly explained by these auditory disorders (Bush et al, 2015; Lin, 2011). This potential confound underscores the importance of collaborating with the physician or psychologist that is administering the cognitive tests to ensure that the environment in which these cognitive tests are administered is as optimal acoustically as possible.

## Use of Questionnaires

Questionnaires do not predict risk for CAPD and present only weak to moderate correlations with only certain central auditory measures in children (Wilson et al, 2011). Questionnaires may be used to highlight concerns about a child, identify additional areas that require assessment, or

## Chermak      Minimize Confounds CAPD Diagnosis

supplement the diagnostic process, but not to determine whether a diagnostic central auditory processing assessment is warranted, nor do questionnaires differentially diagnose disorders (Tomlin et al, 2015; Iliadou & Bamiou, 2012). Questionnaires lack sensitivity for a number of reasons that include: 1) lack of relevant range of items; 2) qualitatively similar symptoms of difficulty could result from many underlying causes besides CAPD; and 3) respondents' estimates of severity may be imprecise, and/or they might not be able to reliably gauge severity, and/or responses may be influenced by other factors, including respondents' interpretation of the question or their desire for a diagnosis (Tomlin et al, 2015). However, adult studies in both neurological (e.g., Bamiou et al, 2012) and non-neurological CAPD populations (Bamiou et al, 2015) show stronger correlations between self-reported listening difficulties and central auditory processing tests than those reported in pediatric studies. More recently developed questionnaires (e.g., Evaluation of Children's Listening and Processing Skills [ECLiPS]) may ultimately prove to be of greater utility in the diagnostic process (Barry, Tomlin, Moore, & Dillon, 2015), pending additional research with a large number of children diagnosed with CAPD, rather than only referred for central auditory evaluation or suspected of CAPD, but not ultimately diagnosed with CAPD.

### Use of Multimodal Analog Measures

Some have suggested the need for multimodal analog tests to differentially diagnose CAPD. Research has demonstrated, however, that the addition of visual tasks offers little to augment differential diagnosis (Bellis et al, 2011). The reader is referred to several key publications for in-depth discussion of the limited contributions of multimodal tests for the differential diagnosis of CAPD, as well as the difficulties in designing such tests (Bellis et al, 2011; Chermak, 2013; Chermak & Bellis, 2014; Dillon et al, 2014; Moore & Ferguson, 2014; Musiek et al, 2005).

### Summary and Conclusions

The purpose of this brief tutorial was to provide audiologists with strategies to minimize confounding cognitive and language processing variables and accurately differentially diagnose CAPD. We suggest that audiologists select central auditory tests that are appropriate to the patient's age, education, linguistic and cultural background, and cognitive ability and select central auditory tests with low language and cognitive demands. Prior to central auditory testing, we recommend that other professionals be engaged to assess an individual's basic cognitive function and language skills (the latter being particularly important with children). In addition to selecting tests with minimal cognitive and linguistic demands, we suggest intra-subject (e.g., inter-aural) and intra-test (e.g., hum v. label) comparisons to further minimize confounds. Proper test selection and use of the strategies recommended here will help the audiologist and other members of the multidisciplinary team determine the source(s) of listening difficulties and other functional deficits. The audiologist in cooperation with the speech-language pathologist and psychologist must determine whether reported listening difficulties result from an auditory problem, a language processing disorder, a cognitive deficit (e.g., attention, executive function, or working memory), or some combination of these deficits. The strategies and approaches reviewed here are useful to minimize confounds; however, they are not intended as a substitute for appropriately normed tests with documented sensitivity, validity, and reliability with such populations. When behavioral audiometric findings remain incomplete, inconclusive, or invalid despite the implementation of the approaches recommended in this tutorial, the audiologist may rely more on electrophysiological techniques, as well as findings provided by other multidisciplinary team



## Chermak      Minimize Confounds CAPD Diagnosis

members to determine the relative contributions of cognition, language processing, and auditory processing to educational, workplace, and everyday functional listening demands, and develop intervention plans. The reader is referred to Chermak (2013) and Chermak and Bellis (2014) for detailed discussions of differential diagnosis of CAPD and attention deficit hyperactivity disorder.

## References

- Acheson, D.J., Hamidi, M., Binder, J.R., Postle, B.R. 2011. A common neural substrate for language production and verbal working memory. *J Cognitive Neurosci*, 23(6), 1358-1367.
- American Academy of Audiology. Clinical Practice Guidelines for the Diagnosis, Treatment, and Management of Children and Adults with Central Auditory Processing Disorder. Available at: <http://www.audiology.org/publications-resources/document-library/central-auditory-processing-disorder>.
- American Speech-Language-Hearing Association. 2005. (Central) auditory processing disorders. Available at <http://www.asha.org/policy/PS2005-00114.htm> accessed February 2017
- Bamiou, D-E., Iliadou, V., Zanchetta, S., Spyridakou, C. 2015. What can we learn about auditory processing from adult hearing questionnaires? *J Am Acad Audiol*, 26 (10), 824-837.
- Bamiou, D.E., Werring, D., Cox, K., Stevens, J., Musiek, F.E., Brown, M., & Luxon, L. 2012. Patient-reported auditory functions after stroke of the central auditory pathway. *Stroke*, 43, 1285-1289.
- Banai, K., Hornickel, J., Skoe, E., Nicol, T., Zecker, S., Kraus, N. 2009. Reading and subcortical auditory function. *Cereb Cortex*, 19(11), 2699-2707.
- Barry, J.G., Tomlin, D., Moore, D.R., Dillon, H. (2015). Use of questionnaire-based measures in the assessment of listening difficulties in school-aged children. *Ear Hear*, 36(6), e300-e313.
- Bellis, T.J. 2002. Considerations in Diagnosing Auditory Processing Disorders in School-Aged Children. *SIG 16 Perspectives on School-Based Issues*, 3(3), 3-9.
- Bellis, T.J. (ed.) 2011. Assessment and management of central auditory processing disorders in the educational setting: From science to practice. Plural Publishing.

## Chermak      Minimize Confounds CAPD Diagnosis

Bellis, T.J., Billiet, C.R., Ross, J. 2011. The utility of visual analogs of central auditory tests in the differential diagnosis of (central) auditory processing disorder and attention deficit hyperactivity disorder. *J Am Acad Audiol*, 22(8), 501-514.

Benavidez, D.A., Fletcher, J.M., Hannay, H.J., Bland, S.T., Caudle, S.E. et al. 1999. Corpus callosum damage and interhemispheric transfer of information following closed head injury in children. *Cortex*, 35(3), 315-336.

Besser, J., Festen, J.M., Goverts, S.T., Kramer, S. E., Pichora-Fuller, M.K. 2015. Speech-in-speech listening on the LiSN-S test by older adults with good audiograms depends on cognition and hearing acuity at high frequencies. *Ear Hear*, 36(1), 24-41.

Bouma, A., Gootjes, L. 2011. Effects of attention on dichotic listening in elderly and patients with dementia of the Alzheimer type. *Brain Cogn*, 76(2), 286-293.

British Society of Audiology. 2011. Position statement: Auditory processing disorder (APD).

Retrieved from <http://www.thebsa.org.uk/> accessed November 2016.

Brooke, P., Bullock, R. 1999. Validation of the 6 Item Cognitive Impairment Test. *Int J Geriatr Psychiatry*, 14, 936-940.

Brown, L., Sherbenou, R., Johnsen, K. 2010. Test of Non-Verbal Intelligence, 4th ed. Austin, TX: Pro-Ed.

Buschman, T.J., Miller, E.K. 2007. Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. *Science*, 315(5820), 1860-1862.

Bush, A.L.H., Lister, J.J., Lin, F.R., Betz, J., Edwards, J.D. 2015. Peripheral hearing and cognition: evidence from the Staying Keen in Later Life (SKILL) Study. *Ear Hear*, 36(4), 395-407.

Chermak      Minimize Confounds CAPD Diagnosis

Cameron S., Brown D., Keith R., Martin J., Watson C. et al. 2009. Development of the North American Listening in Spatialized Noise-Sentences Test (NA LiSN-S): Sentence Equivalence, Normative Data, and Test-Retest Reliability Studies. *J Am Acad Audiol*, 20, 128-146.

Cameron S. Dillon H. 2007. Development of the Listening in Spatialized Noise-Sentences Test (LISN-S). *Ear Hear*, 28, 196-211.

Cameron, S., Glyde, H., Dillon, H. 2012. Efficacy of the LiSN & Learn auditory training software: randomized blinded controlled study. *Audiol Res*, 2(1).

Chermak, G.D. 2013. Differential diagnosis of central auditory processing disorder and attention deficit hyperactivity disorder. *ENT Audiol News*, 22(5), 100-102.

Chermak, G.D., Bellis, T.J. 2014. Differential diagnosis of central auditory processing disorder and attention deficit hyperactivity disorder. In F.E. Musiek & G.D. Chermak (Eds.), *Handbook of central auditory processing disorder: Vol. 1. Auditory neuroscience and diagnosis* (2<sup>nd</sup> ed.). San Diego: Plural Publishing, pp. 557-590.

Chermak, G.D., Hall, J.W., Musiek, F.E. 1999. Differential diagnosis and management of central auditory processing disorder and attention deficit hyperactivity disorder. *J Am Acad Audiol*, 10(6), 289-303.

Cowell, P., Hugdahl, K. 2000. Individual differences in neurobehavioral measures of laterality and interhemispheric function as measured by dichotic listening. *Dev Neuropsychol*, 18(1), 95-112.

Cumming, R., Wilson, A., Goswami, U. 2015. Basic auditory processing and sensitivity to prosodic structure in children with specific language impairments: a new look at a perceptual hypothesis. *Front Psychol*, 6, 972.

Chermak      Minimize Confounds CAPD Diagnosis

Dillon, H., Cameron, S., Tomlin, D., Glyde, H. 2014. Comments on "factors influencing tests of auditory processing: a perspective on current issues and relevant concerns" by Tony Cacace and Dennis McFarland. *J Am Acad Audiol*, 25 (7), 699-703.

Ferguson, M., Hall, R., Riley, A., Moore, D.R. 2011. Communication, listening, cognitive and speech perception skills in children with auditory processing disorder (APD) or specific language impairment (SLI). *Journal of Speech, Language and Hearing Research*, 54(1), 211-227.

Fitch, R.H., Miller, S., Tallal, P. 1997. Neurobiology of speech perception. *Annu Rev Neurosci*, 20(1), 331-353.

Folstein, M.F., Folstein, S.E., McHugh, P.R. 1975. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*, 12(3), 189-198.

Gallun, F.J., Lewis, M.S., Folmer, R.L., Diedesch, A.C., Kubli, L.R. et al. 2012. Implications of blast exposure for central auditory function: A review. *J Rehabil Res Dev*, 49(7), 1059-1074.

Goll, J.C., Crutch, S.J., Loo, J.H., Rohrer, J.D., Frost, C. et al. 2010. Non-verbal sound processing in the primary progressive aphasia. *Brain*, 133(1), 272-285.

Griffiths, T.D. 1999. Human complex sound analysis. *Clin Sci*, 96(3), 231-234.

Griffiths, T.D., Bamiou, D.E., Warren, J.D. 2012. Disorders of the auditory brain. In Palmer A. & Rees A. (Eds) *The Oxford Handbook of Auditory Science: The Auditory Brain*. Oxford University Press. doi:10.1093/oxfordhb/9780199233281.013.0020.

Grossmann, T., Oberecker, R., Koch, S.P., Friederici, A.D. 2010. The developmental origins of voice processing in the human brain. *Neuron*, 65, 852-858.

Grube, M., Cooper, F.E., Kumar, S., Kelly, T., Griffiths, T.D. 2014. Exploring the role of auditory analysis in atypical compared to typical language development. *Hear Res*, 308, 129-140.

Chermak      Minimize Confounds CAPD Diagnosis

Gyldenkaerne, P., Dillon, H., Sharma, M., Purdy, S.C. 2014. Attend to this: The relationship between auditory processing disorders and attention deficits. *J Am Acad Audiol*, 25, 676-687.

Hällgren, M., Larsby, B., Lyxell, B., Arlinger, S. 2001. Cognitive effects in dichotic speech testing in elderly persons. *Ear Hear*, 22(2), 120-129.

Hugdahl, K., Westerhausen, R., Alho, K., Medvedev, S., Laine, M. et al. 2009. Attention and cognitive control: unfolding the dichotic listening story. *Scand J Psychol*, 50(1), 11-22.

Iliadou, V.V., Apalla, K., Kaprinis, S., Nimatoudis, I., Kaprinis, G. et al. 2013. Is central auditory processing disorder present in psychosis? *Am J Audiol*, 22(2), 201-208.

Iliadou, V., Bamiou, D.E. 2012. Psychometric evaluation of children with auditory processing disorder (APD): Comparison with normal-hearing and clinical non-APD groups. *J Speech Lang Hear Res*, 55(3), 791-799.

Iliadou, V., Bamiou, D.E., Kaprinis, S., Kandylis, D., Kaprinis, G. 2009. Auditory processing disorders in children suspected of learning disabilities—A need for screening? *Int J Pediatr Otorhinolaryngol*, 73(7), 1029-1034.

Iliadou, V., Chermak, G.D., Bamiou, D.E. 2015. Differential diagnosis of speech sound disorder (phonological disorder): Audiological assessment beyond the pure-tone audiogram. *J Am Acad Audiol*, 26(4), 423-435.

Iliadou, V., Iakovides, S. 2003. Contribution of psychoacoustics and neuroaudiology in revealing correlation of mental disorders with central auditory processing disorders. *Ann Gen Hosp Psychiatry*, 2(1), 5.

Jerger, S., & Jerger, J. 1984. Pediatric Speech Intelligibility Test: Manual for administration. St. Louis, MO: Auditec.

Chermak      Minimize Confounds CAPD Diagnosis

Jerger J., Martin, J. 2006. Dichotic listening tests in the assessment of auditory processing disorders. *Audiol Med*, 4, 25–34.

Martin, J., Jerger, J., & Mehta, J. 2007. Divided-attention and directed-attention listening modes in children with dichotic deficits: An event-related potential study. *J Am Acad Audiol*, 18, 34–53.

Johnson, K.L., Nicol, T.G., Kraus, N. 2005. Brain stem response to speech: a biological marker of auditory processing. *Ear Hear*, 26(5), 424-434.

Khavarghazalani, B., Farahani, F., Emadi, M., Hosseni Dastgerdi, Z. 2016. Auditory processing abilities in children with chronic otitis media with effusion. *Acta Otolaryngol*, 136(5), 456-459.

Chermak      Minimize Confounds CAPD Diagnosis

Kileny, P., Paccioretti, D., Wilson, A.F. 1987. Effects of cortical lesions on middle-latency auditory evoked responses (MLR). *Electroencephalogr Clin Neurophysiol*, 66(2), 108-120.

Knight, R.T., Scabini, D., Woods, D.L., Clayworth, C.C. 1989. Contributions of temporal-parietal junction to the human auditory P3. *Brain Res*, 502(1), 109-116.

Koenigs, M., Acheson, D.J., Barbey, A. K., Solomon, J., Postle, B.R., et al. 2011. Areas of left perisylvian cortex mediate auditory-verbal short-term memory. *Neuropsychologia*, 49(13), 3612-3619.

Koohi, N., Vickers, D., Chandrashekar, H., Tsang, B., Werring, D., et al. 2016. Auditory rehabilitation after stroke: Treatment of auditory processing disorders in stroke patients with personal frequency-modulated (FM) systems. *Disabil Rehabil*, 1-8.

Kraus, N., McGee, T.J., Carrell, T.D., Zecker, S.G., Nicol, T.G., et al. 1996. Auditory neurophysiologic responses and discrimination deficits in children with learning problems. *Science*, 273(5277), 971-973.

Chermak      Minimize Confounds CAPD Diagnosis

Leff, A.P., Schofield, T.M., Crinion, J.T., Seghier, M.L., Grogan, A. et al. 2009. The left superior temporal gyrus is a shared substrate for auditory short-term memory and speech comprehension: Evidence from 210 patients with stroke. *Brain*, 132(12), 3401-3410.

Lin, F. R. 2011. Hearing loss and cognition among older adults in the United States. *J Gerontol A Biol Sci Med Sci*, 66(10), 1131-1136.

López-Aranda, M.F., López-Téllez, J.F., Navarro-Lobato, I., Masmudi-Martín, M., Gutiérrez, A. et al. 2009. Role of layer 6 of V2 visual cortex in object-recognition memory. *Science*, 325(5936), 87-89.

Martin, J., Jerger, J., & Mehta, J. 2007. Divided-attention and directed-attention listening modes in children with dichotic deficits: An event-related potential study. *J Am Acad Audiol*, 18, 34–53.

McArthur, G., Atkinson, C., Ellis, D. 2009. Atypical brain responses to sounds in children with specific language and reading impairments. *Dev Sci*, 12(5), 768-783.

Moncrieff, D.W. 2011. Dichotic listening in children: age-related changes in direction and magnitude of ear advantage. *Brain Cogn*, 76(2), 316-322.

Moore, D., Ferguson, M. 2014. It is neither necessary nor desirable to test for abnormalities in other modalities when diagnosing auditory processing disorder (APD). *J Am Acad Audiol*, 25(7), 697-698.

Moore, D., Ferguson, M., Edmondson-Jones, A., Ratib, S., Riley, A. 2010. Nature of auditory processing disorder in children. *Pediatrics*, 126(2), E382-90.

Musiek, F. 1994. Frequency (pitch) and duration pattern tests. *J Am Acad Audiol*, 5, 265-286.

Musiek, F.E., Baran, J.A., Bellis, T.J., Chermak, G.D., Hall, J.W. et al. 2010. Diagnosis, treatment and management of children and adults with central auditory processing disorder. *Ameri*



Chermak      Minimize Confounds CAPD Diagnosis

can Academy of Audiology Clinical Practice Guidelines. Available at : [http://audiology-  
web.s3.amazonaws.com/migrated/CAPD%20Guidelines%208-  
2010.pdf\\_539952af956c79.73897613.pdf](http://audiology-<br/>web.s3.amazonaws.com/migrated/CAPD%20Guidelines%208-<br/>2010.pdf_539952af956c79.73897613.pdf)

Musiek, F.E., Baran, J.A., Pinheiro, M.L. 1990. Duration pattern recognition in normal subjects and patients with cerebral and cochlear lesions. *Audiology*, 29, 304-313.

Musiek, F.E., Bellis, T.J., Chermak, G.D. 2005. Nonmodularity of the central auditory nervous system: Implications for (central) auditory processing disorder. *Am J Audiol*, 14(2), 128-138.

Musiek, F.E., Chermak, G.D. 2014. *Handbook of Central Auditory Processing Disorder Volume 1: Auditory Neuroscience and Diagnosis, 2nd ed.* San Diego: Plural Publishing.

Musiek, F.E., Pinheiro, M. 1987. Frequency patterns in cochlear, brainstem and cerebral lesions. *Audiology*, 26, 78-79.

Musiek, F.E., Weihing, J. 2011. Perspectives on dichotic listening and the corpus callosum. *Brain Cogn*, 76(2), 225-232.

Nasreddine, Z.S., Phillips, N.A., Bédirian, V., Charbonneau, S., Whitehead, V. et al. 2005. The Montreal Cognitive Assessment, MoCA: A brief screening tool for mild cognitive impairment. *J Am Geriatr Soc*, 53(4), 695-699.

Pimentel, J., Inglebret, E. 2014. Evidence-based practice and treatment efficacy. In Chermak, G.D., & Musiek, F.E. (Eds.), *Handbook of central auditory processing disorder, Volume 2: Comprehensive intervention (2nd ed.)*. San Diego, CA: Plural Publishing, pp.39-64.

Power, A.J., Mead, N., Barnes, L., Goswami, U. 2013. Neural entrainment to rhythmic speech in children with developmental dyslexia. *Front Hum Neurosci*, 7(777), 1-19.

Chermak      Minimize Confounds CAPD Diagnosis

Price, C., Thierry, G., Griffiths, T. 2005. Speech-specific auditory processing: where is it? *Trends Cogn Sci*, 9(6), 271-276.

Quaranta, N., Coppola, F., Casulli, M., Barulli, O., Lanza, F. et al. 2015. The prevalence of peripheral and central hearing impairment and its relation to cognition in older adults. *Audiol Neurotol*, 19, 10-14.

Ramage, E., Weintraub, D.M., Vogel, S.J., Sutton, G.P., Ringdahl, E.N. et al. 2015. Preliminary evidence for reduced auditory lateral suppression in schizophrenia. *Schizophr Res*, 162(1-3), 269-275.

Rauschecker, J.P., Tian, B. 2000. Mechanisms and streams for processing of "what" and "where" in auditory cortex. *Proc Natl Acad Sci USA*, 97(22), 11800-118006.

Rosen, S., Cohen, M., Vanniasegaram, I. 2010. Auditory and cognitive abilities of children suspected of auditory processing disorder (APD). *International Journal of Pediatric Otorhinolaryngology*, 74(6), 594-600.

Russo, N.M., Nicol, T.G., Zecker, S.G., Hayes, E.A., Kraus, N. 2005. Auditory training improves neural timing in the human brainstem. *Behav Brain Res*, 156(1), 95-103.

Sandford J., Turner A. 2009. Integrated Visual and Auditory Continuous Performance Test Interpretation Manual. Richmond, VA: BrainTrain.

Saunders, G.H., Frederick, M.T., Arnold, M., Silverman, S., Chisolm, T.H. et al. 2015. Auditory difficulties in blast-exposed veterans with clinically normal hearing. *J Rehabil Res Dev*, 52(3), 343-360.

Semel E., Wiig E., Secord W. 2003. Clinical Evaluation of Language Fundamentals, 4th Edition (CELF-4). San Antonio, TX: The Psychological Corporation.

Chermak      Minimize Confounds CAPD Diagnosis

Shannon, R.V., Zeng, F.G., Kamath, V., Wygonski, J., Ekelid, M. 1995. Speech recognition with primarily temporal cues. *Science*, 270(5234), 303-304.

Sharma, M., Purdy, S.C., Kelly, A.S. 2009. Comorbidity of auditory processing, language, and reading disorders. *J Speech Lang Hear Res*, 52(3), 706-722.

Shinn, J., Baran, J., Moncrieff, D., Musiek, F. 2005. Differential Attention Effects on Dichotic Listening. *J Am Acad Audiol*, 16, 205-218.

Studdert-Kennedy, M. 2002. Deficits in phoneme awareness do not arise from failures in rapid auditory processing. *Read Writ*, 15(1-2), 5-14.

Tierney, A., Kraus, N. 2013. Music training for the development of reading skills. Changing brains: Applying brain plasticity to advance and recover human ability. *Prog Brain Res*, 207, 209-241.

Tillery, K.M., Katz, J., Keller, W.D. .2000. Effects of methylphenidate (Ritalin) on auditory performance in children with attention and auditory processing disorders. *Journal of Speech, Language, and Hearing Research*, 43, 893-901.

Tomlin, D., Dillon, H., Sharma, M., Rance, G. 2015. The impact of auditory processing and cognitive abilities in children. *Ear Hear*, 36(5), 527-542.

Wallace, G., Inglebret, E., Friedlander, R. 1997. American Indians: Culture, communication, and clinical considerations. In G.I. Wallace (Ed.), *Multicultural neurogenics: A resource for speech-language pathologists* (pp. 193-225). Tucson, AZ: Communication Skill Builders.

Wechsler, D. (2014). *The Wechsler Intelligence Scale for Children (WISC) (5<sup>th</sup> ed.)*. Bloomington, MN: Pearson.

Chermak      Minimize Confounds CAPD Diagnosis

Weihing, J., Chermak, G.D., Musiek, F.E. 2015. Auditory training for central auditory processing disorder. *Semin Hear*, 36(4), 199-215. Thieme Medical Publishers.

Weihing, J., Guenette, L., Chermak, G.D., Brown, M., Ceruti, J. et al. 2015. Characteristics of pediatric performance on a test battery commonly used in the diagnosis of central auditory processing disorder (CAPD). *J Am Acad Audiol*, 26, 652-669.

Wiig, E., Semel, E., Secord, W. 2013. *Clinical Evaluation of Language Fundamentals (CELF-5)*.  
Bloomington, MN: NCS Pearson.

Willott, J.F. 1991. *Aging and the auditory system: Anatomy, physiology and psychophysics*.  
Whurr Pub Ltd.

Wilson, W. J., Jackson, A., Pender, A., Rose, C., Wilson, J. et al. 2011. The CHAPS, SIFTER, and TAPS-R as Predictors of (C) AP Skills and (C) APD. *J Speech Lang Hear Res*, 54(1), 278-291.

Zatorre, R.J., Belin, P. 2001. Spectral and temporal processing in human auditory cortex. *Cereb Cortex*, 11(10), 946-953.