

1 **Psychotherapy**

2 Introduction: Eric M. Plakun, MD

3 **Bias Toward Psychodynamic Therapy:**  
4 **Framing the Problem and Working Toward a Solution**

5 Allan Abbass MD, FRCPC, Patrick Luyten PhD, Christiane Steinert, PhD and Falk Leichsenring,  
6 DSc

7 *In press: Journal of Psychiatric Practice*

8 ABBASS: Centre for Emotions and Health, Dalhousie University, Canada; LUYTEN: University  
9 College London, United Kingdom, and University of Leuven, Belgium; STEINERT: University  
10 of Giessen, Germany, and Medical School, Berlin, Germany; LEICHSENDRING: University of  
11 Giessen, Germany; PLAKUN: Associate Medical Director, Director of Biopsychosocial  
12 Advocacy, Austen Riggs Center, Stockbridge, MA, and Leader, American Psychiatric  
13 Association Psychotherapy Caucus, Arlington, VA

14 Please send correspondence to: Allan Abbass, Professor and Director, Centre for Emotions and  
15 Health, Dalhousie University, Rm 7507, 5909 Veterans Memorial Lane, Halifax, Nova Scotia,  
16 Canada (email: [allan.abbass@dal.ca](mailto:allan.abbass@dal.ca)).

17

18 The authors declare no conflicts of interest.

19 **Running head: PSYCHOTHERAPY**

20

21

## 1 **Abstract**

2 Although psychodynamic therapy (PDT) is an evidence-based intervention for a broad spectrum  
3 of psychiatric conditions, there is often notable bias in the way PDT is depicted both in the  
4 popular media and in the scientific literature. This has contributed to a negative view of PDT,  
5 which hampers both patient access to this treatment and researcher access to funding for further  
6 research on PDT. The adverse effects of these distortions and biases are detrimental not only to  
7 PDT but also to the overall field of psychotherapy, raising questions about its credibility. Here we  
8 summarize current evidence for PDT, describe existing biases, and formulate a set of  
9 recommendations to foster a more balanced perspective on PDT.

10 **KEY WORDS:** psychotherapy, psychodynamic therapy, psychotherapy research

11

## 1 **Introduction to Guest Column**

2 **Eric M. Plakun, MD**

3 **Psychotherapy Section Editor**

4

5           This column had its origins on the 161st birthday of Sigmund Freud on May 6, 2017,  
6 while I was listening to Garrison Keillor's *Writer's Almanac* piece for that day on National  
7 Public Radio. Keillor cited Freud's birthday in his piece, but after describing Freud's seminal  
8 work on dreams and his focus on the centrality of unconscious factors in determining  
9 human behavior, Keillor concluded his comments by stating that science had by now largely  
10 debunked Freud's theories. I was dumbstruck by this offhanded and completely inaccurate  
11 statement from a usually knowledgeable radio commentator. It reminded me of other  
12 examples I had come across of gross misinformation and bias with respect to  
13 psychoanalysis and psychodynamic therapy. For example, among general psychiatrist  
14 colleagues with whom I work in the American Psychiatric Association (APA) Assembly, no  
15 one would ever utter a joke about a minority group, but periodic jokes poking fun at  
16 psychoanalysts still turn up now and then. About a month later, I saw the PsychiatryOnline  
17 version of a paper by Christiane Steinert<sup>1</sup> and her group that was in press in the American  
18 Journal of Psychiatry, reporting "equivalence" of psychodynamic therapy with other forms  
19 of therapy. I knew then that I wanted to do a column on the problem of implicit bias toward  
20 psychodynamic therapy and psychoanalysis.

21           As I hope has been clear in all of these columns to date, they are about  
22 psychotherapy in general and do not favor one school of psychotherapy over another. We  
23 have too much work to do together to persuade a field that tends toward biological

1 reductionism in its thinking about etiology and treatment of disorders for this column or  
2 this columnist to join the proverbial “circular firing squad,” in which psychotherapists from  
3 competing schools argue with one another about who has the best therapy or the best  
4 research. This implicit bias, as the guest columnists show, goes beyond needless  
5 competition between schools and it reflects poorly on the credibility of psychotherapy as a  
6 form of treatment and on the quality of psychotherapy research of all kinds. It also runs  
7 counter to the recommendation from the Institute of Medicine's *Report on Psychosocial*  
8 *Interventions in Mental Health and Substance Use Disorders* that we increase efforts to  
9 identify shared elements across forms of therapy that are associated with change.<sup>2</sup> We  
10 cannot allow implicit bias to lead us to ignore science. I am delighted and grateful that these 4  
11 leading psychodynamic therapy researchers agreed to take on the task of describing these biases,  
12 citing evidence that unmasks them, and making recommendations to move forward.

13

## 1 **Bias Toward Psychodynamic Therapy: Framing the Problem and Working Toward a** 2 **Solution**

3 Guest Columnists: Allan Abbass, Patrick Luyten, Christiane Steinert and Falk Leichsenring

4 Psychodynamic therapy (PDT) as a family of treatments is an evidence-based intervention  
5 for a broad spectrum of psychiatric conditions.<sup>3-5</sup> PDT has been shown to be as effective as other  
6 psychosocial interventions, including the family of treatments known under the rubric of  
7 cognitive behavioral therapy (CBT).<sup>6</sup> Despite this, PDT continues to receive what appears to be  
8 biased treatment in treatment guidelines, reviews, and related publications, and in media that  
9 inform the public's perception and ultimately patient access to this effective treatment modality.

10

## 11 **EVIDENCE FOR PDT**

12 The efficacy and effectiveness of PDT for common mental disorders have been supported  
13 by several systematic reviews and meta-analyses. A Cochrane review investigating the efficacy  
14 of brief (under 40 sessions) PDT for common mental disorders, for instance, found that PDT  
15 outperformed wait-list, treatment as usual, and minimal contact comparisons at both short- and  
16 long- term follow-up.<sup>3</sup> Longer-term psychodynamic therapy has been found to be effective in  
17 complex mental disorders, including in patients with personality disorders, chronic mental  
18 disorders, or multiple mental disorders.<sup>7-9</sup> In complex mental disorders, the longer term versions  
19 of PDT appear more effective than short-term therapies. According to the Chambless and Hollon  
20 criteria<sup>10</sup> for empirically supported therapies, PDT is "efficacious" or "probably efficacious" in  
21 most common mental disorders.<sup>11</sup> In addition, meta-analyses have found no statistically  
22 significant differences in outcome between individual PDT and other forms of individual  
23 psychotherapy in patients with anxiety or depressive disorders,<sup>12,13</sup> and in patients with more

1 complex mental disorders.<sup>14,15</sup> A recent meta-analysis specifically designed to test for  
2 equivalence in outcomes found PDT to be as efficacious as treatments with established efficacy,  
3 such as CBT, across various mental disorders.<sup>6</sup>

4

## 5 **BIAS IN THE DEPICTION OF PDT**

6 Despite the evidence for PDT, biases in the depiction of PDT and of PDT research remain,  
7 and threaten to reduce the further development of and thereby limit patient access to PDT. Many  
8 of these biases appear to be due to 5 general biases that have been documented in scientific  
9 research, most notably researcher allegiance and the application of double standards.<sup>16,17</sup>

10

### 11 **Bias 1. Distorted Depictions of Psychodynamic Therapy as a Science**

12 Many textbooks of clinical psychology and basic psychology describe psychodynamic  
13 approaches, at best, as historically important in psychology's development, but as currently out-  
14 dated and obsolete. At worst, psychodynamic approaches are depicted as unscientific or even  
15 pseudoscientific.<sup>18</sup> What these depictions have in common is that they are typically based on  
16 caricatured versions of early psychoanalytic assumptions (example: repressed libido as the only  
17 dynamic force) while ignoring contemporary psychodynamic approaches and the considerable  
18 empirical evidence for these views that has emerged over the past few decades.<sup>19</sup> Unfortunately,  
19 this distorted image of current PDT has penetrated popular media and university curricula,  
20 damaging the perspectives of both mental health professionals and prospective patients.<sup>20,21</sup>

21

### 22 **Bias 2. Exclusion or Distortion of Evidence Related to PDT in Treatment Guidelines**

1           Several treatment guidelines exclude or downgrade PDT, often under the guise of the  
2           purportedly smaller evidence base for PDT, even though a higher number studies does not in and  
3           of itself provide evidence for superiority. For instance, in the treatment of anxiety disorders, a  
4           recent meta-analysis<sup>22</sup> showed that more than 80% of 121 trials of CBT focusing on anxiety  
5           disorders used wait-list control groups; only 17% of studies were of high quality. But even when  
6           there are a large number of trials available for PDT of a specific condition, this does not  
7           necessarily lead to comparable and unbiased presentation of PDT. For instance, the recently  
8           published Canadian Guidelines for the Management of Adults with Major Depressive Disorder  
9           (Canadian Network for Mood & Anxiety Disorders or CANMAT)<sup>23</sup> placed brief PDT as a  
10          second- rather than a first-line treatment for depression, despite citing a 54 study meta-analysis  
11          showing large persistent effects and equal effects between individual PDT and other individual  
12          treatment modalities.<sup>12</sup> Furthermore, even when this inconsistency was pointed out, the guideline  
13          committee did not revise its conclusions and also neglected to consider the outcomes of large  
14          studies showing non-inferiority of PDT to CBT.<sup>24</sup> A similar struggle occurred recently in Sweden  
15          where the National Board of Health and Welfare recruited a skewed mix of professionals to  
16          develop treatment guidance: the opinions of the few PDT professionals were outvoted leading to  
17          guidance undervaluing PDT.<sup>25</sup>

18          Research bodies such as the U.S. National Institute of Mental Health similarly appear to  
19          perpetuate distorted or biased information about PDT.<sup>26</sup> For example, under the categories of  
20          anxiety disorders, borderline personality disorder, depression, and eating disorders, there is no  
21          mention of PDT as a valid treatment option. In a separate section describing psychotherapies,<sup>27</sup>  
22          there is a notable absence of discussion of psychodynamic as well as other contributions to  
23          psychotherapy. The focus in these descriptions is on cognitive and behavioral processes, while

1 the language used is typical of CBT, which could mislead the public to believe that the only  
2 relevant psychotherapy approach is CBT.

3 As another example, a recent comprehensive review of psychosocial interventions in  
4 anxiety disorders<sup>28</sup> completely downplayed the evidence for PDT in the treatment of these  
5 disorders, even when confronted with evidence from meta-analyses showing similar effects of  
6 PDT compared with other treatments in these conditions.<sup>13,29</sup>

### 7

### 8 **Bias 3. Exclusion of Psychodynamic Researchers From Funding and Guidelines**

### 9 **Committees**

10 Although there are considerable regional differences, psychodynamic researchers are  
11 often excluded from committees responsible for developing treatment guidelines or for reviewing  
12 research and making decisions about research funding. For example, the CANMAT group  
13 mentioned above initially included PDT researchers, but they were subsequently removed from  
14 the group without explanation, leaving PDT data to be interpreted by researchers with allegiances  
15 to different schools of therapy.<sup>30</sup>

### 16

### 17 **Bias 4. Use of Neutered Versions of “Psychodynamic Therapy” in Randomized Clinical**

### 18 **Trials: The “Straw Man” Bias**

19 A particularly pernicious problem has been the use of neutered versions of  
20 “psychodynamic therapy” in some trials. For example, in a study of posttraumatic stress disorder,  
21 therapists delivering the PDT model were restricted from speaking about the trauma itself, a  
22 withholding that patients must have found both unusual and frustrating.<sup>31</sup> The use of diluted PDT



1 methods as “straw man” controls that are intended to fail was described for the first time more  
2 than three decades ago by Smith et al<sup>32</sup>: “A comparison condition might be set up as a kind of  
3 straw man over which the favored therapy would prevail. The comparison condition (often an  
4 ‘insight therapy’) would be treated with fairly obvious disdain, and would not be given as much  
5 opportunity for success” (p. 119). Surprisingly, this bias still exists, as has been amply  
6 documented in a number of recent reviews.<sup>16,33,34</sup> This bias among others affects the replicability,  
7 validity, and credibility of all psychotherapy research.<sup>35</sup>

8

## 9 **Bias 5. Biased Study Selection in Meta-Analyses**

10 The selection of studies in meta-analyses is frequently biased against PDT. It has been  
11 easy to demonstrate that, in some meta-analyses, typically including researchers with allegiance  
12 to a single and different form of therapy, study selection is performed in ways that exclude valid  
13 PDT studies, on the one hand, and that include flawed PDT studies, on the other. A meta-analysis  
14 by Marcus et al.,<sup>36</sup> for example, which purportedly claimed to investigate the effectiveness of  
15 CBT versus other treatment modalities (including PDT) included only three questionable studies  
16 of PDT, but omitted a large number of RCTs comparing PDT with other bona fide  
17 psychotherapies. Baardseth et al.<sup>37</sup> showed that several studies of bona fide psychotherapies,  
18 including PDT, were excluded in a similar way for unclear reasons in another meta-analysis  
19 purporting to find a consistent advantage for a particular family of treatments.<sup>38</sup> Table 1 lists the  
20 varieties of bias toward PDT.

21

## 22 **A WAY FORWARD**

1           If this problem of bias is not addressed, we not only risk that patients will be denied  
2 access to effective treatments, but we miss the opportunity for dialogue and collaboration that  
3 could enhance the credibility of scientific psychosocial interventions— at present, the credibility  
4 of research in psychology is severely questioned.<sup>39</sup> We propose the following steps to help move  
5 the field in a more sensible and healthy direction. These recommendations are in accordance with  
6 Chambers’ manifesto<sup>39</sup> for reforming the culture of scientific practice.

7           Clearly, for the public to have balanced information, researchers and clinicians who are  
8 knowledgeable about the current literature on PDT should be routinely included in committees  
9 charged with guideline development, funding decisions, webpage publications, and organizations  
10 furthering psychotherapy as a collective treatment approach. Furthermore, researcher allegiances  
11 and other conflicts of interest should be collected and consistently disclosed, so readers have a  
12 context for the materials.

13           Given the shared objective of all psychotherapy proponents to increase the effectiveness  
14 and scope of psychosocial interventions, collaborative research should be done using what has  
15 been called “adversarial collaboration” to further develop psychotherapy as a collective. Those  
16 conducting meta-analyses and review groups should consult with researchers from other models  
17 beyond the allegiance of the core group to yield a richer synthesis and contextualization of  
18 findings. Finally, shared research should continue on key therapy elements versus overall therapy  
19 models toward identifying which interventions work best for whom. This may be especially  
20 relevant in relation to therapeutic factors such as emotional experience/exposure<sup>40</sup> or change in  
21 person-environment exchanges<sup>41</sup> as presumed key ingredients across treatment modalities. Table  
22 2 lists recommendations for corrective action.

23

## 1 REFERENCES

- 2 1. Steinert C, Munder T, Rabung S, Hoyer J, Leichsenring F. 2017. Psychodynamic  
3 Therapy: As Efficacious as Other Empirically Supported Treatments? A Meta-Analysis  
4 Testing Equivalence of Outcomes. *Am. J. Psychiatry*. [Published ahead of print May 25,  
5 2017]. Available at: <https://doi.org/10.1176/appi.ajp.2017.17010057>.
- 6 2. IOM (Institute of Medicine). *Psychosocial interventions for mental health and substance*  
7 *use disorders: A framework for establishing evidence-based standards*. 2015. Washington,  
8 DC: The National Academies Press.
- 9 3. Abbass A. *Reaching Through Resistance: Advanced Psychotherapy Techniques*. Kansas  
10 City, MO: Seven Leaves Press; 2015.
- 11 4. Leichsenring F, Luyten P, Hilsenroth MJ, et al. Psychodynamic therapy meets evidence-  
12 based medicine: a systematic review using updated criteria. *Lancet Psychiatry*. 2015;2  
13 648–660.
- 14 5. Fonagy P. The effectiveness of psychodynamic psychotherapies: an update. *World*  
15 *Psychiatry*. 2015;14:137–150.
- 16 6. Steinert C, Munder T, Rabung S, et al.. Psychodynamic therapy: as efficacious as other  
17 empirically supported treatments? a meta-analysis testing equivalence of outcomes. *Am. J.*  
18 *Psychiatry*. 2017 [Published ahead of print May 25, 2017]. Available at:  
19 <https://doi.org/10.1176/appi.ajp.2017.17010057>.
- 20 7. Leichsenring F, Abbass A, Luyten P, et al. The emerging evidence for long-term  
21 psychodynamic therapy. *Psychodyn Psychiatry*. 2013;41:361–384.
- 22 8. Leichsenring F, Rabung S. Effectiveness of long-term psychodynamic psychotherapy: a  
23 meta-analysis. *JAMA*. 2008;300:1551–1565.

- 1 9. Leichsenring F, Rabung S. Long-term psychodynamic psychotherapy in complex mental  
2 disorders: update of a meta-analysis. *Br J Psychiatry*. 2011;199:15–22.
- 3 10. Chambless DL, Hollon SD. Defining empirically supported therapies. *J Consult Clin*  
4 *Psychol*. 1998;66:7–18.
- 5 11. Leichsenring F, Leweke F, Klein S, et al. The empirical status of psychodynamic  
6 psychotherapy—an update: Bambi's alive and kicking. *Psychother Psychosom*.  
7 2015;84:129–148.
- 8 12. Driessen E, Hegelmaier LM, Abbass AA, et al. The efficacy of short-term psychodynamic  
9 psychotherapy for depression: a meta-analysis update. *Clin Psychol Rev*. 2015;42:1–15.
- 10 13. Keefe JR, McCarthy KS, Dinger U, et al. A meta-analytic review of psychodynamic  
11 therapies for anxiety disorders. *Clin Psychol Rev*. 2014;34:309–323.
- 12 14. Cristea IA, Gentili C, Cotet CD, et al. Psychotherapy for borderline personality disorder: a  
13 systematic review and meta-analysis. *JAMA Psychiatry*. 2017;74:319–328.
- 14 15. Leichsenring F, Leibing E. The effectiveness of psychodynamic therapy and cognitive  
15 behavior therapy in the treatment of personality disorders: a meta-analysis. *Am J*  
16 *Psychiatry*. 2003;160:1223–1232.
- 17 16. Leichsenring F, Rabung S. Double standards in psychotherapy research. *Psychother*  
18 *Psychosom*. 2011;80:48–51; author reply 53–54.
- 19 17. Munder T, Brusch O, Leonhart R, et al. Researcher allegiance in psychotherapy outcome  
20 research: an overview of reviews. *Clin Psychol Rev*. 2013;33:501–511.
- 21 18. Luyten P, Blatt SJ, Corveleyn J. Minding the gap between positivism and hermeneutics in  
22 psychoanalytic research. *J Am Psychoanal Assoc*. 2006;54:571–610.
- 23 19. Luyten P, Mayes LC, Fonagy P, et al. *Handbook of Psychodynamic Approaches to*  
24 *Psychopathology*. New York: The Guilford Press; 2015.

- 1 20. McWilliams N. On teaching psychoanalysis in antianalytic times: a polemic. *Am J*  
2 *Psychoanal.* 2000;60:371–390.
- 3 21. Redmond J, Shulman M. Access to psychoanalytic ideas in American undergraduate  
4 institutions. *J Am Psychoanal Assoc.* 2008;56:391–408.
- 5 22. Cuijpers P, Cristea IA, Karyotaki E, et al. How effective are cognitive behavior therapies  
6 for major depression and anxiety disorders? A meta-analytic update of the evidence.  
7 *World Psychiatry.* 2016;15:245–258.
- 8 23. Parikh SV, Quilty LC, Ravitz P, et al; CANMAT Depression Work Group. Canadian  
9 Network for Mood and Anxiety Treatments (CANMAT). 2016 Clinical Guidelines for the  
10 Management of Adults with Major Depressive Disorder: Section 2. Psychological  
11 Treatments. *Can J Psychiatry.* 2016;61:524–539.
- 12 24. Town JM, Abbass A, Driessen E, et al. Updating the evidence and recommendations for  
13 short-term psychodynamic psychotherapy in the treatment of major depressive disorder in  
14 adults. *Can J Psychiatry.* 2017;62:73–74.
- 15 25. Philips B, Lilliengren P, Klingströ A. Socialstyrelsens riktlinjer är ett haveri [The  
16 guidelines from the National Board of Health and Welfare are a wreck]. *Svenska*  
17 *Dagbladet.* published online February 17, 2017.
- 18 26. National Institute of Mental Health. Health Topics. Available at:  
19 <https://www.nimh.nih.gov/index.shtml>. Accessed June 9, 2017.
- 20 27. National Institute of Mental Health. Psychotherapies. Available at:  
21 <https://www.nimh.nih.gov/health/topics/psychotherapies/index.shtml>. Accessed June 9,  
22 2017.
- 23 28. Craske MG, Stein MB. Anxiety. *Lancet.* 2016; 388:3048–3059.

- 1 29. Steinert C, Leichsenring F. No psychotherapy monoculture for anxiety disorders. *Lancet*.  
2 2017;389:1882–1883.
- 3 30. Leichsenring F, Steinert C. Further evidence for short-term psychodynamic therapy in  
4 major depressive disorder. *Can J Psychiatry*. 2017;62:75–76.
- 5 31. Gilboa-Schechtman E, Foa EB, Shafran N, et al. Prolonged exposure versus dynamic  
6 therapy for adolescent PTSD: a pilot randomized controlled trial. *J Am Acad Child*  
7 *Adolesc Psychiatry*. 2010;49:1034–1042.
- 8 32. Smith ML, Glass GV, Miller TI. *The Benefits of Psychotherapy*. Baltimore, MD: Johns  
9 Hopkins University Press; 1980.
- 10 33. Wampold BE, Fluckiger C, Del Re AC, et al. In pursuit of truth: a critical examination of  
11 meta-analyses of cognitive behavior therapy. *Psychother Res*. 2017;27:14–32.
- 12 34. Wampold BE, Imel ZE. *The Great Psychotherapy Debate: The Evidence for What Makes*  
13 *Psychotherapy Work*. New York/London: Taylor & Francis; 2015.
- 14 35. Leichsenring F, Abbass A, Hilsenroth MJ, et al. Biases in research: risk factors for non-  
15 replicability in psychotherapy and pharmacotherapy research. *Psychol Med*.  
16 2017;47:1000–1011.
- 17 36. Marcus DK, O'Connell D, Norris AL, et al. Is the Dodo bird endangered in the 21st  
18 century? A meta-analysis of treatment comparison studies. *Clin Psychol Rev*. 2014;34:  
19 519–530.
- 20 37. Baardseth TP, Goldberg SB, Pace BT, et al. Cognitive-behavioral therapy versus other  
21 therapies: redux. *Clin Psychol Rev*. 2013;33:395–405.
- 22 38. Tolin DF. Is cognitive-behavioral therapy more effective than other therapies? A meta-  
23 analytic review. *Clin Psychol Rev*. 2010;30:710–720.

- 1 39. Chambers C. *The 7 Deadly Sins of Psychology, A Manifesto for Reforming the Culture*  
2 *of Scientific Practice*. Princeton, NJ: Princeton University Press 2017.
- 3 40. Abbass AA, Kisely SR, Town JM, et al. Short-term psychodynamic psychotherapies for  
4 common mental disorders. *Cochrane Database Syst Rev*. 2014;(7):CD004687.
- 5 41. Fonagy P, Luyten P, Allison E, et al. What we have changed our minds about: part 2.  
6 borderline personality disorder, epistemic trust and the developmental significance of  
7 social communication. *Borderline Personal Disord Emot Dysregul*. 2017;4: 9.
- 8