



MURDOCH RESEARCH REPOSITORY

<http://researchrepository.murdoch.edu.au>

This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination.

Haig, S.M., Beaver, E.A., Chambers, S.M., Draheim, H.M., Dugger, B.D., Dunham, S., Elliott-Smith, E., Fontaine, J.B., Kesler, D.C., Knaus, B.J., Lopes, I.F., Loschl, P., Mullins, T.D. and Sheffield, L.M. (2006) Taxonomic Considerations in Listing Subspecies Under the U.S. Endangered Species Act. Conservation Biology, 20 (6). pp. 1584-1594.

<http://researchrepository.murdoch.edu.au/2579>

Copyright © Society for Conservation Biology.
It is posted here for your personal use. No further distribution is permitted.

REVIEW:

TAXONOMIC CONSIDERATIONS IN LISTING
SUBSPECIES UNDER THE U.S. ENDANGERED
SPECIES ACT

12 June 2006

Susan M. Haig¹, Erik A. Beever^{1,2}, Steven M. Chambers³, Hope M. Draheim^{1,4}, Bruce D. Dugger⁴, Susie Dunham⁵, Elise Elliott-Smith¹, Joseph B. Fontaine⁴, Dylan C. Kesler^{1,4}, Brian J. Knaus⁵, Iara F. Lopes^{1,4,6}, Pete Loschl⁴, Thomas D. Mullins¹, and Lisa M. Sheffield^{4,7}.

¹USGS Forest and Rangeland Ecosystem Science Center, 3200 SW Jefferson Way, Corvallis, OR 97331; susan_haig@usgs.gov; 541-750-7482, 541-758-8806.

²Current Address: U.S. National Park Service Great Lakes Network, 2800 Lake Shore Dr. E., Ashland, WI 54806.

³Division of Ecological Services, U.S. Fish and Wildlife Service, P.O. Box 1306, Albuquerque NM 87103.

⁴Department of Fisheries and Wildlife, Oregon State University, Corvallis OR 97331.

⁵Department of Botany and Plant Pathology, Oregon State University, Corvallis OR 97331.

⁶Current Address: Departamento de Genética e Evolução, Universidade Federal de São Carlos, Rodovia Washington Luís Km 235, São Carlos SP 13565905 Brazil.

⁷Second and subsequent authors listed alphabetically.

Key words: Subspecies, U.S. Endangered Species Act, IUCN Red List, Nomenclature, Taxonomy, Evolutionarily Significant Unit, Distinct Population Segment

Running Head: Taxonomy and the ESA

Word Count: 6,936

ABSTRACT

The U.S. Endangered Species Act (ESA) allows listing of subspecies and other groupings below the rank of species. This provides the U.S. Fish and Wildlife Service and the National Marine Fisheries Service with a means to target the most critical unit in need of conservation. While roughly one-quarter of listed taxa are subspecies, these management agencies are hindered by uncertainties about taxonomic standards during listing or delisting activities. In a review of taxonomic publications and societies, we found few subspecies lists and none that stated standardized criteria for determining subspecific taxa. Lack of criteria is attributed to a centuries-old debate over species and subspecies concepts. However, the critical need to resolve this debate for ESA listings lead us to propose that minimal biological criteria to define disjunct subspecies (legally or taxonomically) should include the discreteness and significance criteria of Distinct Population Segments (as defined under the ESA). Our subspecies criteria are in stark contrast to that proposed by supporters of the Phylogenetic Species Concept and provide a clear distinction between species and subspecies. Efforts to eliminate or reduce ambiguity associated with subspecies-level classifications will assist with ESA listing decisions. Thus, we urge professional taxonomic societies to publish and periodically update peer-reviewed species and subspecies lists. This effort must be paralleled throughout the world for efficient taxonomic conservation to take place.

INTRODUCTION

Intraspecific taxa are important in discussions of biodiversity because they represent evolutionary potential within a species. Lists of taxonomic groupings are key to conservation efforts because they provide a foundation for identifying species, subspecies (i.e., taxa below the species level), and evolutionarily unique populations. Recognizing this, subspecific taxa are included in the International Union for the Conservation of Nature (IUCN) Red List of Threatened Species, appendices in the Convention on International Trade in Endangered Species of Wild Flora and Fauna (CITES), TRAFFIC (wildlife trade monitoring network), Brazil's Lista Nacional das Espécies da Fauna Brasileira Ameaçadas de Extinção, Canada's Species at Risk Act, Australia's Environmental Protection and Biodiversity Act, South Africa's Biodiversity Act, and others (<http://eelink.net/~asilwildlife/legislat.shtml>).

The U.S. Endangered Species Act of 1973 (ESA) also protects intraspecific taxa. Originally, the ESA definition of *species* included "any subspecies of fish or wildlife or plants. Also in 1978, the ESA was amended so "species" encompassed "any distinct population segment [i.e., DPS] of any species of vertebrate fish or wildlife which interbreeds when mature" (USFWS & National Marine Fisheries Service [NMFS] 1996). Currently, one-quarter of ESA-listed taxa have subspecific rank (Tables 1, 2; Fig. 1).

Listing subspecific taxa is becoming increasingly controversial (Isaac et al. 2004; Zink 2004; Harris & Froufe 2005; Mallet et al. 2005) due to incongruencies between biological and legal criteria for recognizing and protecting subspecific taxa and because of the continuing debate among taxonomists regarding the validity of subspecies as a taxonomic unit (e.g., Zink et al. 2000; Ramey et al. 2005). This has been exacerbated by

careless taxonomy in some cases (reviewed in Mayr & Ashlock 1991) and overapplication of the subspecies concept for species that attract human interest. Thus, the concept further fell out of favor when subspecies of notable species like the leopard (*Panthera onca*; Larson 1997; Eizirik et al. 2001), seaside sparrow (*Ammodramus nigrescens*; Avise & Nelson 1989), and redwing blackbird (*Agelaius phoeniceus*; Williams et al. 2004) did not hold up under scrutiny.

Among taxonomists, definitions of subspecies are a source of considerable disagreement. This uncertainty is compounded in a conservation context when the specialized taxonomic expertise required to evaluate conflicting interpretations does not exist within management agencies responsible for listing species. Thus, management agencies need to be made aware of the best available science and policies that can relate biological criteria to legal requirements under the ESA.

To illustrate and understand the taxonomic information available to agencies evaluating subspecies for ESA listing, we: 1) reviewed subspecific definitions and concepts across taxonomic groups, 2) summarized how subspecific taxa have been used in ESA listings, 3) reviewed variations in how taxonomists and management agencies use subspecific taxonomic groupings, 4) carried out discussions of issues related to subspecific listings with agency personnel responsible for ESA-related matters, and 4) proposed recommendations on how taxonomists can better contribute to conservation under the ESA and other similar legislative entities around the world.

SUBSPECIFIC DEFINITIONS AND CONCEPTS

Species are generally recognized as the fundamental units of taxonomy, but species concepts provide little assistance in recognizing taxa below the species level. For example, reliance of Mayr's (1963) Biological Species Concept (BSC) on reproductive isolation between species might lead one to assume that partial reproductive isolation would be an appropriate criterion for subspecies recognition. However, there is little evidence outside of *Drosophila* (e.g., Ayala et al. 1974) that this criterion has been routinely employed. Thus, in recent decades, taxonomists have applied subspecies names to geographic races without evidence of or reference to partial isolation. Under most phylogenetic species concepts (PSC, Cracraft 1983), species are the smallest, irreducible, monophyletic units as measured by molecular markers. Any groupings within such species do not warrant taxonomic standing, so subspecies are not recognized.

The concept of infraspecific taxa has been used at least since Linnaeus' time (1753). Qualitative definitions include: Darwin (1896) who considered varieties to be incipient species, potentially evolving into full species; Mayr (1963) defined subspecies as "geographically defined aggregates of local populations which differ taxonomically from other such subdivisions of the species"; and more recently Frankham et al. (2002) stated they were "...populations partway through the evolutionary process of divergence towards full speciation". Qualitative definitions have been criticized as arbitrary because some groups classified qualitatively as subspecies are not differentiated based on multiple characters (Wilson & Brown 1953; Mallet 2001).

Traditionally, subspecies have been defined by morphological traits or color variations, but recent critics are concerned that these traits may not reflect underlying

genetic structure and phylogenies. This concern stems from recent work where phylogenetic patterns of genetic variation were not concordant with some subspecies classifications defined by morphology (Zink 1989; Ball & Avise 1992; Zink et al. 2000; Zink 2004).

The only quantitative subspecies definition we found was the 75% rule (Amadon 1949; Patten & Unitt 2002) that states a subspecies is valid if 75% or more of a population is separable from all (or >99% of) members of the overlapping population. Although the 75% rule is more quantitative than other definitions, there is disagreement about the 75% threshold and the number of characters that should be used when comparing populations (Patten & Unitt 2002).

Despite all the criticisms, recent studies using multiple criteria (e.g., morphological, behavioral, and genetic characters) have confirmed that many subspecies are evolutionarily definable entities (e.g., Gavin et al. 1999; Pasquet 1999; Haig et al. 2004). Thus, while subspecies definitions may have been too liberally applied by some early taxonomists, this does not invalidate the concept of subspecies as meaningful biological entities. Taxonomists continue to recognize and sometimes describe subspecies. However, more work is needed to clarify this taxonomic concept to assist management agencies with identifying which entities are appropriate for providing regulatory protection. These efforts should focus on developing guidelines for how to reconcile multiple lines of evidence when evaluating the validity of a subspecies. One good example of this attempt is the recent British Ornithologists Union guidelines for defining subspecies in birds (Helbig et al. 2005) that considered PSCs and the BSC.

VARIATION IN SUBSPECIFIC CLASSIFICATIONS ACROSS TAXA

In an extensive literature review, we found no universally accepted subspecies definition within or across taxa (Table 3). Further, we found use of subspecies in modern taxonomy differs by taxonomic group. In general, more subspecies have been described for vertebrates and plants than the less studied invertebrates and fungi, where most taxonomic studies remain focused at the species level. Defining subspecies is complicated by the biology of an organism as well as by a paucity of knowledge about the diversity in some groups. For example, some groups (e.g., birds) are better dispersers than others; hence there may be fewer, less defined, or genetically differentiated avian subspecies than in less vagile taxa (e.g., amphibians). These factors and others result in an imbalance in how ESA protection can be applied across taxa.

Every taxonomic group has followed a similar evolution of ideas to resolve taxonomic questions. Historically, morphology and geography were used to separate taxa. Subsequently, BSC stimulated simplifications (i.e., lumping) at the species level and an enthusiasm for the use of subspecies level classifications to describe morphological variation within resulting polytypic species. Currently, taxonomists are struggling with how to incorporate results of modern molecular methods into their assessments using various phylogenetic species concepts.

In the following sections, we summarize taxon-specific use of subspecies rank, organizations that provide taxonomic standards or official lists of taxa, and the prevalence of subspecies listings under the ESA. Current species and subspecies numbers are presented in Table 3 for all taxa.

Animals (Kingdom Animalia)

The Zoological Record, implemented via the International Council on Zoological Nomenclature (ICZN), is the most comprehensive list of animal names in the world (Table 3; www.biosis.org/products/zr/). The ICZN has no role in determining which species concepts are applied; however, it provides clear instructions on the formation of specific and subspecific names and has published nomenclatural changes since 1905. Changes in recognized names of taxa are printed in the *Quarterly Bulletin of Zoological Nomenclature* with commentary from the ICZN. Additionally, its *Official List of Names in Zoology and Works in Zoological Nomenclature* includes names of taxa and titles of works that have been the subject of their rulings. ICZN recently proposed a web-based register (“ZooBank”) that identifies animal names and facilitates communication regarding their nomenclature (Polaszek 2005).

Birds (Class Aves)

Ornithologists have spent considerable effort refining and debating subspecies concepts (Wiens 1982). In the early 20th century, Walter Rothschild and Ernst Hartert made extensive use of their geographic-based subspecies concept in numerous avian taxonomic publications (Mayr 1976; Rothschild 1983). However, following introduction of the BSC, 315 of the 607 North American bird species were reclassified as subspecies (AOU 1957; Mayr 1982). The American Ornithologists’ Union (AOU) Committee on Classification and Nomenclature is the scientific body responsible for standardizing avian taxonomy in North America. The committee has published seven editions of its *Checklist*

of North American Birds since 1896. Subspecies were included in the first four editions, but have not been included since 1957 due to committee time constraints (AOU 1998). Currently, 43% of ESA-listed birds are listed at the subspecific rank, emphasizing the importance of reliable avian subspecific taxonomy (Table 1).

Recent avian subspecies debates have been motivated by examples of genetic population structure that differed from morphology-based subspecies delineations (Zink 1989; Ball & Avise 1992; Zink et al. 2000; Zink 2004). Some ornithologists argue that historic classifications may not accurately reflect phylogeny because morphological differences do not always have a phylogenetically significant basis (Zink 2004). Others debate whether the subspecies concept has intrinsic value to avian classification (Smith & White 1956; Barrowclough 1982; Wiens 1982; Zink 2004). Yet a new consideration of avian subspecies world-wide indicates that 36% are phylogenetically distinct and that lack of distinctness is most pronounced in North American and European taxa, where much of the debate has taken place (Phillimore & Owens 2006).

Mammals (Class Mammalia)

Mammalian classification is listed in *Mammal Species of the World* (Wilson & Reeder 2005) which is updated every 10 years in a cooperative effort between the Association of Systematics Collections and the American Society of Mammalogists. Editors assume that systematic and nomenclatural decisions are the province of the professional research community, but they address conflicting opinions regarding taxonomy. Modern systematists suggest that some traditional mammalian subspecific designations based on minor geographic variations in size and/or color may not necessarily represent actual

genetic differences. Conversely, cryptic taxa have been overlooked and denied status (Hershkovitz 1983; Smith & Patton 1988). Over two thirds of ESA mammal listings are for subspecies (Table 1). Although this trend toward subspecies listing has been consistent over time, more recent mammal listings have focused on DPSs (Figure 1).

Fishes (Classes Cephalaspidomorphi, Chondrichthyes, Myxini, and Osteichthyes)

Fishes comprise the most diverse vertebrate group at the species level, with approximately 24,600 formally recognized species globally (Helfman et al. 1997; Table 3). Subspecies classifications in fish are based most commonly on allopatry (Echelle 1991; Duvernell & Turner 1999). In this context, Linnaen trinomials applied to fishes are largely synonymous with geographic races. However, there is a wide disparity in levels of molecular genetic divergence among conspecific taxa representing different subspecies. Some taxonomists argue against use of subspecies designations for fish because under a strict subspecies definition (e.g., a population in a particular region that is genetically distinguishable from other such populations and is capable of interbreeding with them), every isolated creek and pond could have a unique subspecies or species of fish (Mayden 1999).

The American Fisheries Society and the American Society of Ichthyologists and Herpetologists are the scientific bodies responsible for reviewing published taxonomic descriptions of fish and for compiling a standardized list of fish species, subspecies, and other infraspecific names for North American taxa (Nelson et al. 2004). Their joint Name of Fishes Committee *List of Species* does not list subspecies but acknowledges them in

footnotes (e.g., subspecies of *O. mykiss*). Thus, because subspecific classification has been used unevenly and often sparingly with fishes, it is correspondingly underrepresented in ESA listings. Only 15% of U.S. fish listings are targeted at the subspecific rank (Table 1). USFWS and NMFS have more often listed fish DPSs (NMFS uses an ESU concept to define DPSs of Pacific salmon; NMFS 1991) than subspecies. Consequently, 31% (44/142) of domestic fish listings are at the level of DPS.

Reptiles (Class Reptilia) and Amphibians (Orders Anura, Caudata, and Gymnophiona)

In North America, the Society for the Study of Amphibians and Reptiles (SSAR) and the Center for North American Herpetology publish lists of scientific and standard herpetological names (Crother et al. 2001, 2003; Collins & Taggart 2002; Table 3). Both lists include subspecies. The SSAR list has been sanctioned by the major herpetological societies in North America (Stuart 2002). A review of the SSAR list (Crother et al. 2001) indicates that a similar number of species and subspecies have been identified in North America. The proportion of subdivided species within each genera ranges from 0% (crocodiles) to 56% (snakes) with up to 12 subspecies described per species.

Variation in the ratio of described species/subspecies under herpetofauna reflects the fact that the subspecies concept has fallen from favor in modern herpetofauna systematics (David Wake, UC-Berkeley, personal communication). Currently, 29% of ESA herpetofaunal listings are for subspecies (Table 1). Whether this disproportionate listing of species reflects a greater need for conservation at the species level in herpetofauna or the influence of other factors (e.g., a disproportionate listing of taxa with relatively few recognized subspecies) is unclear.

Invertebrates (Phyla Annelida, Arthropoda, Brachiopoda, Bryozoa, Cnidaria, Ctenophora, Echinodermata, Loricifera, Mollusca, Nematoda, Phoronida, Platyhelminthes, Porifera, and Rotifera)

Invertebrates make up a vast paraphyletic group that includes approximately 96% of all described animal species (Table 3). Thus, multiple professional groups oversee their nomenclature. In North America, these groups include: the Acarological Society of America (mites), American Malacological Society (molluscs), International Union for the Study of Social Insects (primarily Hymenoptera and some Isoptera), Orthopterists Society (grasshoppers), Crustacean Society (all crustaceans), Xerces Society (primarily butterflies), and Lepidopterists Societies (butterflies), and Coleopterists Society (beetles).

Among invertebrates, fine-scale variation in morphological traits has most often been ascribed at the species rather than subspecies level. Combined with a lower overall amount of taxonomic knowledge, relatively few species or subspecies of invertebrates have been described compared to most vertebrate groups, except among butterflies (Wilson & Brown 1953; Brusca & Brusca 2003). Subspecific ESA listings have largely been restricted to more fully described taxa such as butterflies and freshwater mollusks.

Plants (Kingdom Plantae)

Botanical species concepts are numerous (Stuessy 1990; McDade 1995; Mayden 1997; Bachmann 1998; Rieseberg & Burke 2001). However, most plant taxa have been described using a morphological (or Linnaean) species concept. Although subspecific concepts in plants have received less discussion than species concepts (but see Stuessy

1990; McDade 1995), subspecific taxa are frequently described. McDade (1995) found that of 104 contemporary monographs, 56% included infraspecific taxa. Authors generally utilized the rank of subspecies or variety (roughly equally) but not both. This suggests that although these concepts are technically different (Greuter et al. 2000), in practice they are being treated synonymously (McDade 1995; Table 3). Consequently, extension of the ESA's definition of species to include "variety" (USFWS 1978) appears congruent with current nomenclatural practice.

The naming of plants (and historically, fungi) is dictated by the International Code of Botanical Nomenclature (Greuter et al. 2000). The code provides a hierarchy (species, subspecies, variety, form) but no definitions to clarify ranks. There is no internationally recognized list of valid plant taxa, but the U.S. Department of Agriculture's PLANTS database (USDA NRCS 2006) is used to provide guidance for federal agencies. Many states have floras compiled that, although frequently out of date, provide a local authority on plants. The *Flora of North America* (1993+) is an ongoing, multi-volume project that has been partially published and is considered an authority on plant names for the United States.

In the United States, ESA authority to list DPSs and the ESA "take" prohibitions do not apply to plants (Table 2). The protections that do apply to plants cover only those that occur on federally owned land or are affected by a federal action. Thus, the ESA provides protection for plants to a more limited extent than to animals. Of 745 plants listed by USFWS, 123 are for subspecies (including varieties; Table 1).

Kingdom Fungi

Fungus taxonomy is focused at the species level. Approximately 800 new fungus species names are catalogued in the *Index Fungorum* each year, but many are never reassessed after their original description. It is estimated that mycologists inadvertently redescribe known species at a rate of 2.5:1 (Hawksworth 1991). Given this error rate, the estimated number of described fungal species ranges from 80,060 to 150,000. It is widely accepted that there are 1.5 million species of fungi worldwide (Hawksworth 1991), which suggests that < 10% of fungal species have been described.

The authoritative reference on fungal taxonomy is *The Dictionary of the Fungi* (Kirk et al. 2001), which is in its fifty-ninth year and ninth edition. This dictionary summarizes accumulated knowledge on all organisms studied by mycologists including lichens, mushrooms, slime molds, water molds, and yeasts. Criteria for delineating the three subspecific categories most commonly used in fungal taxonomy (subspecies, variety, form) are defined by individual monographers and vary among families and genera. The terms *subspecies* and *variety* appear to be used interchangeably (see *Index Fungorum*, www.indexfungorum.org). In many situations, however, the same infraspecific name is listed under the rank of subspecies and variety. Often the rank of subspecies is converted to the rank of variety in later taxonomic treatments, indicating either that most fungal taxonomists consider the ranks biologically equivalent or have not settled on criteria to define them. The uncertainty surrounding the number of described and actual fungi species suggests standardized criteria for designating subspecific ranks are unlikely to be defined in the near future.

There is no world or North American checklist for fungi, although establishment of MycoBank in 2004 (Hawksworth 2005; www.mycobank.org) is quickly filling that niche by providing online accession numbers to newly described taxa. Currently only two species of lichenized fungi (*Cladonia perforata*; *Gymnoderma lineare*) are listed under the ESA (USFWS 1993, 1995a). Future efforts related to listing fungi under the ESA should pay special attention to verifying names at the taxonomic rank of species, if not subspecies or variety.

PERSPECTIVES AND RECOMMENDATIONS

Listing Subspecies

Listing subspecies under the ESA has increasingly become a source of conflict in science and policy. The complex processes involved in speciation (Dobzhansky 1937; Grant 1981; Coyne & Orr 2004) can make it difficult to define species or subspecies relationships as simply as stated in the applicable legislation. While the scientific community has some level of comfort with the subjective nature of subspecies classification (Hey et al. 2003), agencies and the general public want subspecies criteria to be more quantitative or better defined so that conservation designations are applied more predictably. One potential criterion would favor adopting a phylogenetic species concept. However, they generally do not recognize subspecies and it would result in species-level recognition of smaller units. That is, subspecies would either be elevated to full species or not be recognized. This would lead to a proliferation of described species, most of which would occupy more restricted ranges and thereby be more vulnerable to extinction and listing under the ESA. In addition, phylogenetic taxonomy presents its

own methodological problems and is not universally accepted among biologists (Hudson & Coyne 2002; Avise 2004; Coyne & Orr 2004).

The lack of rigid definitions does not mean that currently described subspecies are not useful for defining populations worthy of ESA listings. For example, listings have included well known and accepted subspecies such as Florida Panther (*Felis concolor coryi*; USFWS 1967), Northern Spotted Owl (*Strix occidentalis caurina*; USFWS 1990), and Marbled Murrelet (*Brachyramphus marmoratus marmoratus*; USFWS 1992). Nevertheless, listing a poorly-defined or invalid subspecies could have unwarranted economic impact on private landowners, developers, and other interests (Zink et al. 2000; Ramey et al. 2005). Clearly, the best possible methods must be used to assess taxonomy to avoid this problem. These assessments may have to be re-evaluated periodically as techniques evolve, similar to adaptive management efforts in other types of biological management.

The ESA provides for protection of groups or populations to allow for the conservation of evolutionary potential within a species. This helps focus management efforts on vulnerable areas so that the entire species does not become endangered or threatened. Thus, subspecies and DPS listings will result in a far less long-term impact on stakeholders than listing entire species as they insure evolutionary potential is preserved while not invoking protections beyond what is needed for a specific population.

Resolution of Subspecies Taxonomy and ESA Listings

In view of the legal need to bring together taxonomy and the ESA, as well as provide greater consistency for describing subspecies within each of the major taxonomic groups, we need a unified set of criteria that are biologically and legally defensible under the ESA. Certainly, we need to assure these decisions occur unidirectionally (Bowen and Karl 1999). That is, conservation strategies should be influenced by taxonomy, but taxonomy cannot be influenced by conservation priorities.

We propose that, as a starting point for discussion about subspecies criteria, *minimal* biological requirements in most situations include two criteria based on discreteness of the population in relation to the remainder of the species to which it belongs and the biological significance of the population to the species. This is in contrast to "evolutionary significance" which implies more species-like relationships. In other words, if a taxonomic subspecies does not satisfy these criteria, then the biological legitimacy of the particular subspecies classification should be questioned. Modification of the discreteness criterion would be necessary to allow for cases where subspecies contact one another along a stable hybrid zone but maintain their taxonomic integrity elsewhere. Adopting these criteria would provide for consideration of multiple types of biological data, not just molecular genetic data, yet would still carry the requirement of defining differences. However, satisfaction of these criteria would not necessarily warrant subspecies recognition via a Linnaean trinomial.

The proposed minimum criteria are somewhat similar to the DPS criteria in that the DPS policy provides useful discussion and insight into a reasonable set of minimum criteria for evaluating potential designations for groups lacking prior serious taxonomic

consideration. However, there are distinct differences between DPS's and subspecies; in particular the caveat that DPS's are not listed for plants and invertebrates whereas subspecies of these taxa can be listed. Further, subspecies are a taxonomic grouping whereas DPS's are a legal classification.

Our proposal takes a distinctly different tact than that proposed by supporters of the phylogenetic species concept. The requirement that a taxonomic subspecies should satisfy the phylogenetic species rids taxonomy of the term "subspecies" and uses only molecular data in assessment of taxonomy. This emphasis on establishing evolutionary distinctness does not resolve the subspecies issue; hence it provides a narrow perspective not useful for ESA considerations of units below the species level. Further, factors other than genetics need to be considered in understanding relationships below the species level.

New Subspecific Descriptions and Taxonomic Revisions Involving Subspecies

When describing new subspecies or revising subspecies taxonomy, detailed descriptions of concepts and criteria used to determine taxonomic relationships should accompany each proposed classification or reclassification published in the scientific literature. We recommend explicit description of the geographic distributions of taxa to clearly define their limits for agency management. Experts in each field are better able to judge the legitimacy of subspecies if the methodology, including the species concept used to determine classification, is stated explicitly in taxonomic publications.

Molecular genetic techniques will continue to be useful for evaluating subspecies designations. However, the level and magnitude of genetic variation among and within

well-documented closely-related species should be described for comparison's sake (Barrowclough & Flesness 1996; Fraser & Bernatchez 2001). It is important to recognize that while these tools excel at exploring historic reproductive isolation, they usually do not directly address adaptive divergence. Therefore, all else being equal, species with high dispersal rates will have fewer subspecies identified via molecular markers than species with lower rates of dispersal. Consequently, they will generally require additional information beyond molecular markers to justify designation of subspecies, such as evidence of local adaptation in spite of ongoing gene flow.

Recent work has emphasized that adaptive divergence can take place despite gene flow (Wu 2001; Beaumont & Balding 2004). It is therefore important to use multiple sources of information when evaluating a taxon's status including tools that address the questions of reproductive isolation, adaptive divergence, and spatial patterns of local adaptation (Crandall et al. 2001; Fraser & Bernatchez 2001). Since it will often be impossible to clearly show adaptive divergence among populations (using translocation, common garden experiments, or population genomic approaches), significant differences in phenotypes (e.g., morphology, behavior, life history, or ecology) and environments may reflect local adaptation, and should be used in listing decisions when harder scientific evidence for local adaptation would be too difficult or expensive to acquire. Thus, higher levels of confidence can be obtained in classifications based on the concurrence of multiple morphological, molecular, ecological, behavioral and/or physiological characters.

Whatever criteria are accepted, professional taxonomic societies and journal editors need to adopt statements describing the range of currently acceptable taxonomic

standards and concepts so that agencies preparing ESA listings have the proper peer-reviewed criteria to judge professional taxonomic decisions. We further recommend expert nomenclatural and taxonomic committees to create and update accurate species and subspecies lists. The resulting lists could then be globally standardized and made accessible on the Internet through hosts such as the Global Biodiversity Information Facility (www.gbif.org/) or the National Biological Inventory Substructure (www.nbi.gov/index.html).

New ESA Subspecific Listings

The published final rules for endangered and threatened subspecies demonstrate that USFWS and NMFS use (and need) peer-reviewed publications, taxonomic checklists provided by professional societies, and comments from independent scientists and expert panels when evaluating taxonomic status. Therefore, USFWS and NMFS biologists actively seek professional comments when rules are under review, yet sometimes find limited published information, or few taxonomists who specialize in that species, groups of species, or techniques used to classify subspecies. Consequently, numerous listings require extended internal reviews of taxonomic status by USFWS or NMFS because published information was either contradictory or not available. In some cases, these agencies must evaluate highly esoteric disagreements among respected scientific experts with little expertise of their own (e.g., USFWS 2005). An example is the final rule for the Southwestern Willow Flycatcher (*Empidonax traillii extimus*). It stated that “the Service reviewed the information and found a majority opinion that *E. t. extimus* is a valid subspecies...”(USFWS 1995b). Although some degree of interpretation of what is

the “best available” scientific information will always be required of management agencies when making listing decisions, the scientific community can help to ensure that interpretation of taxonomy is solely a scientific endeavor.

Taxonomists can make a more significant contribution to conservation by developing an understanding of ESA policy (or the appropriate policies of their country) and regulatory requirements for listing species under the ESA. This will help taxonomists understand the nature of taxonomic challenges faced by agency biologists and provide directions for future research. Thus, the key is to provide scientists with enough practical background so they can understand consequences of different biological conclusions without making them advocates or encouraging them to blur boundaries between science and policy. Of course, these professionals must maintain an independence from the conservation implications of their work. Thus, we re-emphasize the point that the relationship between conservation and taxonomy must be unidirectional (Bowen and Karl 1999).

Professional societies can contribute to reducing taxonomic uncertainty by working with USFWS and NMFS to identify old listings that may need taxonomic updating. This will facilitate research that can clarify these questions using modern tools. An adaptive management strategy for dealing with taxonomy will keep taxonomy and listings up to date in view of the latest technology. Furthermore, it would be a major contribution to conservation if professional societies maintained a list of members who are qualified to make taxonomic evaluations or to participate in panels to evaluate special taxonomic cases for subspecies that are subjects of listing action or likely to become so. Peer reviewed assessments of particular issues conducted by independent professional

societies or their members will lend credibility to conclusions about taxonomy for the management agencies.

CONCLUSIONS

ESA protection of biodiversity through listing at the level of taxonomic species and subspecies provides taxonomists with a unique and challenging opportunity. Efforts to eliminate or reduce ambiguity associated with subspecies-level classifications will assist with ESA listing decisions and facilitate better identification of relationships among taxa by taxonomists. While their general application extends worldwide, our recommendations may appear fairly USA-centric. To our knowledge, these taxonomic issues have weighed more heavily in the U.S. than in countries without the legislative ability to protect subspecies, with newer legislation, and those with fewer penalties for violation. However, efforts to list subspecies under (for example) CITES or the IUCN Red-Data List already call for better clarification of taxa. Our hope is that professional societies throughout the world can see this as a global issue and participate in this important effort toward taxonomic clarification.

ACKNOWLEDGEMENTS

We thank our Oregon State University colleagues and others for ideas related to this topic: R. Bellinger, J. Boland, D. Markle, J. Scordino, F. Allendorf (University of Montana), R. Banks (USGS/NMNH), D. Campton (USFWS), C. Funk (USGS FRESC), A. Hecht (USFWS), D. Olson (U.S. Forest Service), J. Pagel (USFWS), D. Wake (University of California-Berkeley), R. Waples (NMFS), and D. Wilson (National

Museum of Natural History). Views expressed in this paper are the authors and do not necessarily reflect the view of USFWS, USGS, or OSU.

LITERATURE CITED

- Amadon, D. 1949. The seventy-five percent rule for subspecies. *Condor* 51:250–258.
- American Ornithologists' Union. 1957. Check-list of North American birds. 5th edition.
American Ornithologists Union, Washington, D.C.
- American Ornithologists' Union. 1998. Check-list of North American birds. 7th edition.
American Ornithologists Union, Washington, D.C.
- Avise, J.C. 2004. Molecular markers, natural history, and evolution, 2nd Edition.
Sinauer, Sunderland, Massachusetts.
- Avise, J.C., and W.S. Nelson. 1989. Molecular genetic relationships of the extinct dusky
seaside sparrow. *Science* 243: 646-648.
- Ayala, F.J. M.L. Tracey, D. Hedgecock, and R.C. Richmond. 1974. Genetic
differentiation during the speciation process in *Drosophila*. *Evolution* 28:576-
592.
- Bachmann, K. 1998. Species as units of diversity; an outdated concept. *Theory in
Biosciences* 17:213-230.
- Ball, R.M., and J.C. Avise. 1992. Mitochondrial DNA phylogeographic differentiation
among avian populations and the evolutionary significance of subspecies. *Auk*
109:626-636.
- Barrowclough, G.F. 1982. Geographic variation, predictiveness, and subspecies. *Auk*
99:601-603.
- Barrowclough, G.F., and N. Flesness. 1996. Species, subspecies, and races: the problem
of units of management in conservation. Pages 247–254 in D.G. Kleiman, M.

- Allen, K. Thompson, and S. Lumpkin, editors. Wild mammals in captivity: principles and techniques. University of Chicago Press, Chicago, Illinois.
- Beaumont, M.A., and D.J. Balding. 2004. Identifying adaptive genetic divergence among populations from genome scans. *Molecular Ecology* 13: 969-980.
- Bowen, B.W., and S.A. Karl. 1999. In war, truth is the first casualty. *Conservation Biology* 13:1013-1016.
- Brusca, R.C., and G.J. Brusca. 2003. *Invertebrates*. 2nd ed. Sinauer, Sunderland, Massachusetts.
- Collins, J. T., and T. W. Taggart. 2002. Standard common and current scientific names for North American amphibians, turtles, reptiles, and crocodylians. The Center for North American Herpetology, Lawrence, Kansas.
- Coyne, J.A., and H.A. Orr. 2004. *Speciation*. Sinauer, Sunderland, Massachusetts.
- Cracraft, J. 1983. Species concepts and speciation analysis. *Current Ornithology* 1:159-187.
- Crandall, K.A., O.R.P. Bininda-Emonds, G.M. Mace, and R.K. Wayne. 2001. Considering evolutionary processes in conservation biology: an alternative to 'evolutionary significant units'. *Trends in Ecology & Evolution*. 15:290-295.
- Crother, B.I., et al. 2000. Scientific and standard English names of amphibians and reptiles of North America North of Mexico, with comments regarding confidence in our understanding. *Herpetological circular* 29. Society for the Study of Amphibians and Reptiles, Lawrence, Kansas.

- Crother, B.I. et al. 2003. Scientific and standard English names of amphibians and reptiles of North America north of Mexico: update. *Herpetological Review* 34:196-203.
- Darwin, C. 1896. *The variation of animals and plants under domestication*. D. Appleton, New York.
- Dobzhansky, T. 1937. *Genetics and the origin of species*. Columbia University Press, New York.
- Duvernell, D.D., and B.J. Turner. 1999. Variation and divergence of Death Valley pupfish populations at retrotransposon-defined loci. *Molecular Biology and Evolution* 16:363–371.
- Echelle, A.A. 1991. Conservation genetics and genic diversity in freshwater fishes of western North America. Pages 141-153 in W.L. Minckley and J.E. Deacon, editors. *Battle against extinction*. University of Arizona Press, Tucson, Arizona.
- Eizirik, E., K. Jae-Heup, M. Menotti-Raymond, P.G. Crawshaw, Jr., S.J. O'Brien, and W.E. Johnson. 2001. Phylogeography, population history and conservation genetics of jaguars (*Panthera onca*, Mammalia, Felidae). *Molecular Ecology* 10: 65-79.
- Flora of North America Editorial Committee, eds. 1993+. *Flora of North America North of Mexico*. 7+ vols. New York and Oxford (also available from www.fna.org/FNA/).
- Frankham, R., J.D. Ballou, and D.A. Briscoe. 2002. Resolving taxonomic uncertainties and defining management units. Pages 365-392 in R. Frankham, J.D. Ballou, and

D.A. Briscoe, editors. Introduction to conservation genetics. Cambridge University Press, United Kingdom.

Fraser, D.J., and L. Bernatchez. 2001. Adaptive evolutionary conservation: towards a unified concept for defining conservation units. *Molecular Ecology* 10:2741-2752.

Gavin, T.A., P.W. Sherman, E. Yensen, and B. May. 1999. Population genetic structure of the northern Idaho ground squirrel (*Spermophilus brunneus brunneus*). *Journal of Mammalogy* 80:156-168.

Grant, V. 1981. Plant speciation. 2nd edition. Columbia University Press, New York.

Greuter, W., et al. 2000. International Code of Botanical Nomenclature (St Louis Code).

Regnum Vegetabile 138. Koeltz Scientific Books, Königstein, Denmark (Also

available from [\[berlin.de/iapt/nomenclature/code/SaintLouis/0000St.Luistitle.htm\]\(http://berlin.de/iapt/nomenclature/code/SaintLouis/0000St.Luistitle.htm\)\).](http://www.bgbm.fu-</u></p></div><div data-bbox=)

Haig, S.M., E.D. Forsman, and T.D. Mullins. 2004. Subspecies relationships and genetic structure in the Spotted Owl. *Conservation Genetics* 5: 683-705.

Harris, J.D., and E. Froufe. 2005. Taxonomic inflation: species concept or historical geopolitical bias? *Trends in Ecology & Evolution* 20:6-8.

Hawksworth, D.L. 1991. The fungal dimension of biodiversity: its magnitude and significance. *Mycological Research* 95:441-456.

Hawksworth, D.L. 2005. Universal fungal register offers pattern for zoology. *Nature* 438:24-34.

Helbig, A.J., A.J. Knox, D.T. Parkin, G. Sangster, and M. Collinson. 2005. Guidelines for assigning species rank. *Ibis* 144:518-522.

- Helfman, G.S., B.B. Collette, and D.E. Facey. 1997. *The diversity of fishes*. Blackwell Science, Malden, Massachusetts.
- HersHKovitz, P. 1983. Two new species of night monkeys, genus *Aotus* (Cebidae, Platyrrhini): a preliminary report of *Aotus* taxonomy. *American Journal of Primatology* 4:209-243.
- Hey, J., R.S. Waples, M.L. Arnold, R.K. Butlin, and R.G. Harrison. 2003. Understanding and confronting species uncertainty in biology and conservation. *Trends in Ecology & Evolution* 18: 597-603.
- Hudson, R.R., and J.A. Coyne. 2002. Mathematical consequences of the genealogical species concept. *Evolution* 56: 1557-1565.
- Isaac, N.J.B., J. Mallet, and G.M. Mace. 2004. Taxonomic inflation: it's influence on macroecology and conservation. *Trends in Ecology & Evolution* 19:464-469.
- Kirk, P.M., P.F. Cannon, J.C. David, J.A. Stalpers (eds). 2001. *Dictionary of the Fungi*. 9th CAB International Bioscience, Oxfordshire, United Kingdom.
- Larson, S.E. 1997. Taxonomic re-evaluation of the jaguar. *Zoo Biology* 16: 107–120.
- Linnaeus, C. 1753. *Species plantarum*. Facsimile of the 1st edition with introduction by W.T. Stearn. The Ray Society, London, 1957.
- Mallet, J. 2001. Subspecies, semispecies. Pages 523-526 in S. Levin, editor. *Encyclopedia of Biodiversity*. Vol. 5. Academic Press, San Diego, California.
- Mallet, J., N.J.B. Isaac, and G.M. Mace. 2005. Response to Harris and Froufe, and Knapp et al.: Taxonomic inflation. *Trends in Ecology & Evolution* 20:385-386.

- Mayden, R.L. 1997. A hierarchy of species concepts: the denouement in the saga of the species problem. Pages 381-424 in M.F. Claridge, H.A. Dawah, and M.R. Wilson, editors. *Species: the units of biodiversity*. Chapman and Hall, New York.
- Mayden, R.L. 1999. Consilience and a hierarchy of species concepts: advances towards closure on the species puzzle. *Journal of Nematology* 31:95-116.
- Mayr, E. 1963. *Animal species and evolution*. Belknap Press, Cambridge, Massachusetts.
- Mayr, E. 1976. *Evolution and the diversity of life*. Belknap Press, Cambridge, Massachusetts.
- Mayr, E. 1982. Of what use are subspecies? *Auk* 99:593-595.
- Mayr, E., and P.D. Ashlock. 1991. *Principles of systematic zoology*, second edition. McGraw-Hill, Inc., New York.
- Mayr, E., E.G. Linsley, and R.L. Usinger. 1953. *Methods and principles of systematic zoology*. McGraw-Hill, New York.
- McDade, L.A. 1995. Species concepts and problems in practice: insights from botanical monographs. *Systematic Botany* 20:606-622.
- NMFS (National Marine Fisheries Service). 1991. Policy on applying the definition of species under the Endangered Species Act to Pacific Salmon. *Federal Register* 56:58612-58618.
- Nelson, J.S., E.J. Crossman, H. Espinosa-Pérez, L.T. Findley, C.R. Gilbert, R.N. Lea, and J.D. Williams. 2004. *Common and scientific names of fishes from the United States, Canada, and Mexico*, 6th Edition. American Fisheries Society, Bethesda, Maryland.

- Pasquet, R.S. 1999. Genetic relationships among subspecies of *Vigna uguiculata* (L.) Walp. based on allozyme variation. *Theoretical & Applied Genetics* 98:1104-1119.
- Patten, M.A., and P. Unitt. 2002. Diagnosability versus mean differences of sage sparrow subspecies. *Auk* 119:26-35.
- Phillimore, A.B., and I.P.F. Owens. 2006. Are subspecies useful in evolutionary and conservation biology? *Proceedings of the Royal Society of London B* 273: 1049-1053.
- Polaszek, A. 2005. A universal register for animal names. *Nature* 437: 477.
- Ramey, R.R., II, H-P. Liu, C.W. Epps, L.M. Carpenter, and J.D. Wehausen. 2005. Genetic relatedness of the Preble's meadow jumping mouse (*Zapus hudsonius preblei*) to nearby subspecies of *Z. hudsonius* as inferred from variation in cranial morphology, mitochondrial DNA and microsatellite DNA: implication for taxonomy and conservation. *Animal Conservation* 8:329-346.
- Rieseberg, L.H. and J.M. Burke. 2001. The biological reality of species: gene flow, selection, and collective evolution. *Taxon* 50:47-67.
- Robins, C.R. et al. 1991. Common and scientific names of fishes from the United States and Canada. American Fisheries Society, Bethesda, Maryland.
- Rothschild, M. 1983. Dear Lord Rothschild; Birds, Butterflies and History. 1983. Balaban, Philadelphia, Pennsylvania.
- Smith, H.M., and F.N. White. 1956. A case for the trinomen. *Systematic Zoology* 5:183-190.

- Smith, M.F., and J.L. Patton. 1988. Subspecies of pocket gophers: causal bases for geographic differentiation in *Thomomys bottae*. *Systematic Zoology* 37:163-178.
- Stuart, J.N. 2002. Book reviews: Scientific and standard English names of amphibians and reptiles of North America North of Mexico, with comments regarding confidence in our understanding, by the committee on standard English and scientific names (B.I. Crother, chair), and standard common and current scientific names for North American amphibians, turtles, reptiles and crocodylians, 5th Edition, by J.T. Collins and T.W. Taggart. *Bulletin of the Chicago Herpetological Society* 37:197-199.
- Stuessy, T.F. 1990. *Plant taxonomy; the systematic evaluation of comparative data*. Columbia University Press, New York
- USDA, NRCS. 2006. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). Data compiled from various sources by M.W. Skinner. National Plant Data Center, Baton Rouge, LA 70874-4490 USA
- USFWS (U.S. Fish and Wildlife Service). 1967. U.S. Endangered Species List. *Federal Register* 32: 4001.
- USFWS (U.S. Fish and Wildlife Service). 1978. Determination that 11 plant taxa are endangered species and two plant taxa are threatened species. *Federal Register* 43:17910-17916.
- USFWS (U.S. Fish and Wildlife Service). 1990. ETWP; Determination of threatened status for the Northern spotted owl. *Federal Register* 55:26114-26194.

- USFWS (U.S. Fish and Wildlife Service). 1992. ETWP; Determination of threatened status for the Washington, Oregon and California population of the Marbled Murrelet. Federal Register 57:45328-45333.
- USFWS (U.S. Fish and Wildlife Service). 1993. Endangered or threatened status for seven central Florida plants. Federal Register 58:25746-25755.
- USFWS (U.S. Fish and Wildlife Service). 1995a. *Gymnoderma lineare* (Rock Gnome Lichen) determined to be endangered. Federal Register 60:3557-3562.
- USFWS (U.S. Fish and Wildlife Service). 1995b. Final rule determining endangered status for the Southwestern Willow Flycatcher. Federal Register 60:10693-10715.
- USFWS (U.S. Fish and Wildlife Service). 2005. 12-month finding on a petition to list *Cicurina cueva* (no common name) as an endangered species. Federal Register 70:75071-75074.
- USFWS (U.S. Fish and Wildlife Service) and NMFS (National Marine Fisheries Service). 1996. Policy regarding the recognition of distinct vertebrate population segments under the Endangered Species Act. Federal Register 61:4721-4725.
- Wiens, J.A. 1982. Forum: Avian subspecies in the 1980's. Auk 99:593-615.
- Williams, C.L., H.J. Homan, J.J. Johnston, and G.M. Linz. 2004. Microsatellite variation in Red-Winged Blackbirds (*Agelaius phoeniceus*). Biochemical Genetics 42: 35-41.
- Wilson, D.E., and D.M. Reeder (editors). 2005. Mammal species of the world: a taxonomic and geographic reference, 3rd ed. Smithsonian Institution Press, Washington, D.C.

- Wilson, E.O., and W.L. Brown, Jr., 1953. The subspecies concept and its taxonomic application. *Systematic Zoology* 2: 97-111.
- Wu, Chung-I. 2001. The genic view of the process of speciation. *Journal of Evolutionary Biology* 14: 851-865.
- Zink, R.M. 1989. The study of geographic variation. *Auk* 106:157-160.
- Zink, R.M. 2004. The role of subspecies in obscuring avian biological diversity and misleading conservation policy. *Proceedings of The Royal Society of London B* 271:561-564.
- Zink, R.M., G.F. Barrowclough, J.L. Atwood and R.C. Blackwell-Rago. 2000. Genetics, taxonomy, and conservation of the threatened California gnatcatcher. *Conservation Biology* 14:1394-1405.

Table 1. Species and subspecies listings by taxa under the U.S. Endangered Species Act as of October 2005. Data summarized from <http://www.fws.gov/endangered>. Trinomials are considered subspecies, whereas those listed as binomials are considered species.

Taxon	U.S. listings ^a			Species listed outside U.S. ^b			All listings
	Species	Subspecies	Total	Species	Subspecies	Total	
Birds	53	40	93	145	39	184	277
Mammals	28	56	84	209	68	277	361
Fish	121	21	142	12	0	12	154
Herps	40	16	56	74	25	99	155
Invertebrates	141	15	156	2	1	3	159
Plants	617	125	742	3	0	3	745
Fungi (lichens)	2	0	2	0	0	0	2

^a Species were considered to be U.S. listings if the reported historic range included the United States and the listing included their entire range.

^b Listings outside of the U.S. are principally CITES listings.

Table 2. Potential listing categories for various taxa under the U.S. Endangered Species Act.

Taxa	Species	Subspecies/variety	DPS ^a	ESU ^b
Birds	●	●	●	
Mammals	●	●	●	
Fish	●	●	●	● ^c
Herps	●	●	●	
Invertebrates	●	●		
Plants	●	●		
Lichens/fungi	●	●		

^aDistinct Population Segment

^bEvolutionary Significant Unit

^cESU is used for DPSs of Pacific Salmon

Table 3. Estimated numbers of worldwide and U.S. species per taxon and use of infraspecific categories in taxonomy.

Taxon	Species		Infraspecific categories	Authority Reference
	Worldwide	U.S.		
Birds	9,688	650	mostly subspecies	AOU 1998, globalforestwatch.org , bsc-eoc.org/avibase/avibase.jsp
Mammals	5,416	432	mostly subspecies	Wilson & Reeder 2005, globalforestwatch.org
Fish	24,600	2,428	subspecies but not often	Robins et al. 1991, Helfman et al. 1997
Amphibians	5,743	263	subspecies	globalamphibians.org , globalforestwatch.org
Reptiles	8,240	287	subspecies	reptile-database.org , globalforestwatch.org
Invertebrates	1,288,518	~93,000	subspecies, varieties	Brusca & Brusca 2003, www.iczn.org
Plants	300,000	?	subspecies, varieties, forms	Greuter et al. 2000
Fungi	80,000-150,000 described	5-10,000	subspecies, varieties, forms	Kirk et al. 2001

FIGURE LEGEND

Figure 1. Number of species and subspecies listed by year under the U.S. Endangered Species Act (ESA) as of October 2005.

