



**Murdoch**  
UNIVERSITY

## MURDOCH RESEARCH REPOSITORY

*This is the author's final version of the work, as accepted for publication following peer review but without the publisher's layout or pagination.*

*The definitive version is available at*

<http://dx.doi.org/10.1016/j.fishres.2011.02.008>

**Wakefield, C.B., Fairclough, D.V., Lenanton, R.C.J. and Potter, I.C. (2011) *Spawning and nursery habitat partitioning and movement patterns of *Pagrus auratus* (Sparidae) on the lower west coast of Australia. Fisheries Research, 109 (2-3). pp. 243-251.***

<http://researchrepository.murdoch.edu.au/4346/>

Copyright: © 2011 Elsevier B.V.

It is posted here for your personal use. No further distribution is permitted.

## Accepted Manuscript

Title: Spawning and nursery habitat partitioning and movement patterns of *Pagrus auratus* (Sparidae) on the lower west coast of Australia

Authors: Corey B. Wakefield, David V. Fairclough, Rod C.J. Lenanton, Ian C. Potter



PII: S0165-7836(11)00079-8  
DOI: doi:10.1016/j.fishres.2011.02.008  
Reference: FISH 3159

To appear in: *Fisheries Research*

Received date: 21-12-2010  
Revised date: 11-2-2011  
Accepted date: 11-2-2011

Please cite this article as: Wakefield, C.B., Fairclough, D.V., Lenanton, R.C.J., Potter, I.C., Spawning and nursery habitat partitioning and movement patterns of *Pagrus auratus* (Sparidae) on the lower west coast of Australia, *Fisheries Research* (2010), doi:10.1016/j.fishres.2011.02.008

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Spawning and nursery habitat partitioning and movement patterns of**  
2 ***Pagrus auratus* (Sparidae) on the lower west coast of Australia**

3

4

5 **Corey B. Wakefield<sup>a,b</sup>** (corresponding author)

6 **David V. Fairclough<sup>a</sup>**

7 **Rod C. J. Lenanton<sup>a</sup>**

8 **Ian C. Potter<sup>b</sup>**

9

10

11 <sup>a</sup>Western Australian Fisheries and Marine Research Laboratories

12 Department of Fisheries, Government of Western Australia

13 P.O. Box 20, North Beach, Western Australia, 6920, Australia.

14

15 <sup>b</sup>Centre for Fish and Fisheries Research

16 School of Biological Sciences and Biotechnology

17 Murdoch University

18 South Street, Perth, Western Australia, 6150, Australia.

19

20 Phone: +61 (08) 9203 0111; fax: +61 (08) 9203 0199

21 Email: corey.wakefield@fish.wa.gov.au

22

23

24

25 **Abstract**

26 The ages and lengths of *Pagrus auratus* caught by line fishing in three marine  
27 embayments (Owen Anchorage, Cockburn Sound and Warnrbo Sound) and inshore  
28 (< 80 m depth) and offshore waters (> 80 m depth) on the lower west coast of Australia  
29 (31°45' to 32°45' S) were used to infer the movement patterns and habitats occupied by  
30 this species at different stages in its life cycle on this coast. These data were  
31 supplemented by results obtained by tagging individuals in spawning aggregations in  
32 the embayments. 0+ *P. auratus* < 200 mm FL were caught exclusively in the three  
33 adjacent embayments. The ages and lengths of immature *P. auratus*, ranging from 1+  
34 (ca 200 mm FL) to 5+ years (ca 400 mm FL), increased progressively with distance  
35 from these embayments. During the spawning period (from September to January), the  
36 relative abundances of *P. auratus* with either developing, developed or recently-spent  
37 gonads were far greater in the three embayments (91 %) than in either inshore (12 %) or  
38 offshore waters (30 %). Some tagged *P. auratus* were recaptured among spawning  
39 aggregations in the same embayment during subsequent spawning seasons, while others  
40 were recaptured in these embayments outside the spawning period. However, some  
41 other tagged individuals were recaptured up to 92 km north, 33 km west and 134 km  
42 south outside the spawning period and up to five years after tagging. The results of this  
43 study emphasise that the above three adjacent marine embayments constitute important  
44 spawning and nursery areas for *P. auratus* and are thus potentially critical for sustaining  
45 the stocks of this recreationally and commercially important species on the lower west  
46 coast of Australia.

47

48

49 **Keywords:** *Pagrus auratus*, Sparidae, tagging, movement, spawning aggregation,  
50 nursery, embayment.

51

52

53 **Running heading:** Distributions and movement of *P. auratus* in WA

54

55

56 **Introduction**

57 The development of sound management policies for sustaining heavily-fished stocks of  
58 a species requires reliable data on various aspects of the life cycle of that species,  
59 including their habitats. It is important to determine, however, not only the habitats  
60 occupied by such species at different stages in their life cycles, but also the ages, lengths  
61 and life cycle stages and times of the year at which any movements occur between  
62 habitats. Thus, for example, if the stock of a species is shown to congregate predictably  
63 in restricted locations at a certain time of the year, it will be identified as potentially  
64 prone to particularly heavy exploitation by fishers at that time. A thorough knowledge  
65 of habitat use and migrations of the individuals of a heavily-fished stock is thus required  
66 to identify which life cycle stages of that stock and their habitats may require special  
67 protection. Finally, information on habitat use and movements is important for  
68 developing an ecosystem-based approach to fisheries management and thereby to ensure  
69 that ecosystem function in the different habitats is maintained.

70

71 Habitat partitioning between life stages, e.g. juveniles and adults, has been reported for  
72 many sparids (e.g. Bennett, 1993; Gillanders, 2002; Hesp et al., 2004a). Evidence for  
73 connectivity between juvenile and adult habitats is most commonly obtained from  
74 differences in the abundance of size and/or age classes in these habitats (Gillanders et  
75 al., 2003). The movement patterns of juveniles (0+ and 1+) inferred from this approach  
76 would otherwise be difficult to obtain from direct methods, due, for example, to the  
77 high levels of mortality and low recapture rates that are associated with traditional dart  
78 and anchor tagging of small fish. Mature individuals of many fish species also undergo  
79 seasonal movements each year to and from specific locations at the commencement and  
80 completion of the spawning period, respectively (e.g. Colin, 2010; Domeier and Colin,

81 1997; Heyman and Kjerfve, 2008). Seasonal accumulations of mature conspecific fish  
82 at specific locations and in significantly higher densities than are found outside the  
83 spawning period have been termed spawning aggregations (Domeier and Colin, 1997).  
84 The locations where spawning aggregations occur may also act as important nursery  
85 areas (Fowler et al., 2005; Hamer et al., 2005).  
86  
87 *Pagrus auratus* (Forster 1801) is distributed throughout the temperate Indo-Pacific  
88 coastal waters of Australia and New Zealand between 18° and 38° S (Paulin, 1990).  
89 Throughout its distribution, this species forms a few spawning aggregations (as defined  
90 by Domeier and Colin, 1997; Society for the Conservation of Reef Fish Aggregations,  
91 www.scrfa.org) in sheltered marine embayments (Coutin et al., 2003; Crossland, 1977;  
92 Jackson and Cheng, 2001; McGlennon, 2004; Scott et al., 1993). In Western Australia,  
93 these spawning aggregations are typically found in embayments where the  
94 geomorphology, hydrology and habitat characteristics facilitate the retention of eggs,  
95 larvae and, to a certain extent, juveniles (Doak, 2004 unpublished honours thesis;  
96 Lenanton, 1974; Nahas et al., 2003; Wakefield, 2006). The areas where spawning  
97 aggregations of *P. auratus* form in South Australia and Victoria also constitute  
98 important nursery areas (Fowler et al., 2005; Hamer et al., 2005). For example, Fowler  
99 et al. (2005) showed that, in South Australia, the progenies of the spawning  
100 aggregations of *P. auratus* in the northern areas of Gulf St Vincent and Spencer Gulf in  
101 1991 remained in those areas for up to the first three years. This strong year class  
102 derived from spawning in those gulfs dominated the catches taken along more than  
103 2,000 km of the South Australian coast in at least the year 2000 (Fowler et al., 2005),  
104 which demonstrated the importance of such discrete spawning/nursery areas for  
105 recruitment and sustainability of broader adult stocks.

106

107 Tagging studies have shown that the extent to which adult *P. auratus* move varies  
108 among populations. For example, individuals of this species can move up to 1,650 km  
109 in a northerly direction along the lower east coast of Australia (Sanders, 1974), whereas,  
110 those of the three stocks of *P. auratus* in the inner gulfs of Shark Bay in Western  
111 Australia (ca 26° S) remain in the same area, with the majority moving < 20 km from  
112 their location of release (Moran et al., 2003). The extent of the movements undertaken  
113 by the individuals of a given population of *P. auratus* thus presumably represents the  
114 degree to which the various life cycle stages are adapted to a given environment (Moran  
115 et al., 2003).

116

117 *Pagrus auratus* forms spawning aggregations during the austral spring/summer in three  
118 adjacent marine embayments on the lower west of Australia, i.e. Owen Anchorage,  
119 Cockburn Sound and Warnbro Sound (Fig. 1), but do not apparently spawn in the  
120 waters immediately to the west (Wakefield, 2010). The geomorphology of Cockburn  
121 and Warnbro Sounds and the prevailing south-westerly winds during spring/summer  
122 result in a counter-clockwise gyre, which coincides with the spawning period of  
123 *P. auratus* and facilitates the retention of eggs and larvae in these embayments during  
124 this period (Wakefield, 2010). This suggests that these relatively small, discrete  
125 embayments are potentially an important source of recruitment for nearby adult stocks  
126 along the lower west coast of Australia.

127

128 This study compares the length and age compositions of *P. auratus* in the three marine  
129 embayments (and particularly Cockburn and Warnbro Sounds, in which the spawning  
130 aggregations are by far the largest) with those in the nearshore shallow (< 80 m depth)

131 and offshore deeper (> 80 m) areas. These comparisons were used to determine whether  
132 habitat partitioning by *P. auratus* occurs in this region during the year and at different  
133 stages of their life cycle. Particular attention was paid to ascertaining the relative  
134 abundance of juvenile and mature *P. auratus* in all areas to gain an understanding of the  
135 importance of the three embayments as spawning and nursery areas for *P. auratus*.  
136 Finally, a mark-recapture tagging program was undertaken at known spawning  
137 aggregation locations (Cockburn and Warnbro Sounds) during the spawning period, in  
138 an attempt to determine the directions and distances that adult *P. auratus* might move  
139 from such locations.

140

## 141 **Methods**

### 142 **Sample collection and measurements**

143 Samples of *Pagrus auratus* were collected from 2002 to 2006 from three areas between  
144 *ca* 31°45' and 32°45' S on the lower west coast of Western Australia (Fig. 1). The  
145 offshore and inshore areas were located at depths > and < 80 m, respectively, and were  
146 situated immediately west of the marine embayments area which comprised Owen  
147 Anchorage, Cockburn Sound and Warnbro Sound (Fig. 1). These marine embayments  
148 have been identified previously as locations where spawning aggregations of *P. auratus*  
149 (Wakefield, 2006) and assemblages of 0+ juveniles (Lenanton, 1974) occur each year.

150

151 *Pagrus auratus* was caught by line fishing from either research vessels or recreational  
152 charter vessels with research staff onboard who were permitted to keep fish less than the  
153 minimum legal length (410 mm total length at that time). This sampling, which was  
154 undertaken at least monthly from April 2003 to March 2005, was not accompanied by  
155 the tagging of fish (described later). Although fishing effort was not quantified, the



156 sample sizes of *P. auratus* were sufficient to determine the proportions of the different  
157 life history stages in each area in all months. The same range of hook sizes and variety  
158 of rig types, which were known collectively to catch a large size range of *P. auratus*  
159 (Otway and Craig, 1993), were used on each sampling occasion. *Pagrus auratus* caught  
160 from research vessels were later processed in the laboratory, while those caught from  
161 recreational charter vessels were processed onboard during each trip. The fork length  
162 (FL) of each *P. auratus* was measured to the nearest 1 mm. The two sagittal otoliths  
163 were removed from each fish, cleaned and stored in paper envelopes, and the  
164 macroscopic appearance of the gonads were used to sex each fish and to determine its  
165 stage of development (Table 1).

166

#### 167 **Treatment of otoliths**

168 The right otolith of each fish was embedded in epoxy resin and, using a slow speed saw  
169 (Buehler Ltd.) with a diamond tipped saw blade, sectioned transversely through its  
170 primordium, perpendicular to the sulcus acusticus. The sections were mounted on a  
171 glass microscope slide with a cover slip using casting resin.

172

173 The opaque zones in each otolith section were counted under reflected light at 20 to 40  
174 times magnification, without any knowledge of the length of the fish or its date of  
175 capture. The first opaque zone was easily distinguished, as its formation resulted in the  
176 development of an inflection point in the Subcupular Meshwork Fibre zone (Francis et  
177 al., 1992).

178

179 A single opaque zone has previously been shown to form annually in the otoliths of *P.*  
180 *auratus* from the lower west coast of Australia (Wakefield, 2006). Thus, the age of each

181 *P. auratus* on its date of capture was estimated using a combination of an average birth  
 182 date and the number of opaque zones in its otolith. An average birth date of 1 November  
 183 was chosen because it represented the approximate peak time of spawning derived  
 184 previously from the trends exhibited throughout the year by mean monthly values for  
 185 gonadosomatic indices and the proportions of mature fish in samples collected from the  
 186 lower west coast of Australia (Wakefield, 2006). These trends in gonadal variables  
 187 demonstrated that *P. auratus* spawn from spring to mid-summer, *i.e.* from September to  
 188 January each year (Wakefield, 2006).

189

### 190 **Juvenile habitat partitioning**

191 The differences between the abundances of juvenile *P. auratus* in the marine  
 192 embayments and the inshore and offshore areas were described using length and age  
 193 distributions up to the minimum length and age at which a fish was recorded with  
 194 mature gonads (stages II-V, Table 1), *i.e.* 320 mm FL and 3.74 yr. The proportions of  
 195 juveniles at a given fork length in the marine embayments, *i.e.*  $P_{L,embayments}$ , compared  
 196 with those in areas outside were calculated using a reparameterised form of the logistic  
 197 equation (Hesp et al., 2004b; Punt and Kennedy, 1997),

$$198 \quad P_{L,embayments} = 1 - \left\{ 1 + \exp \left[ -\ln(19) \frac{(L - L_{50})}{(L_{95} - L_{50})} \right] \right\}^{-1},$$

199 where the parameters  $L_{50}$  and  $L_{95}$  represent the estimated lengths at which 50 and 95 %  
 200 of *P. auratus* were present in the marine embayments, respectively. The  $L_{50}$  and  $L_{95}$   
 201 values and their 95 % confidence intervals were determined by bootstrapping, where  
 202 estimates were obtained from the analysis of data sets produced by random resampling,  
 203 with replacement, of each data assemblage to generate 1000 estimates of the parameters  
 204 of the logistic equation. The parameters for the reparameterised logistic equation were

205 calculated as the median of the 1000 bootstrap estimates of each length class. Estimates  
206 of the proportions of juvenile *P. auratus* at a given age in the marine embayments, i.e.  
207  $P_{A,embayments}$ , compared to outside areas were calculated using the same equation, but  
208 with  $A_{50}$  and  $A_{95}$  substituted for  $L_{50}$  and  $L_{95}$ , respectively.

209

### 210 **Tagging study**

211 *Pagrus auratus* caught in spawning aggregations in Cockburn Sound and, to a lesser  
212 extent, Warnbro Sound were tagged in each year between 2003 and 2008. This study is  
213 restricted to recapture data recorded up to September 2009. Each tag had a unique  
214 identification number, the name of the responsible governing organisation, 'FISHERIES  
215 WA', and a free-call phone number to report recaptures. A reward was offered for  
216 reporting recaptures and was stated on the tag as 'REWARD – MEASURE'. The  
217 tagging program was advertised through multi-media sources. Tags were 9 cm in length  
218 and constructed of yellow plastic with a large dart tip (type PDA, Hallprint Australia  
219 Pty Ltd). Tags were inserted with a hollow needle through the dorsal musculature and  
220 locked behind the pterygiophore bones approximately 1 to 2 cm below the base of the  
221 dorsal fin. Two tags were inserted in the majority of *P. auratus* to increase the  
222 likelihood of being able to identify recaptured individuals in case of tag loss, and to  
223 estimate rates of such losses. Previous studies using dart tags and identical tag insertion  
224 methods have demonstrated that tagging did not have a detrimental influence on the  
225 survival and growth of *P. auratus* elsewhere (McGlennon and Partington, 1997;  
226 Quartararo and Kearney, 1996).

227

### 228 **Results**

#### 229 **Length and age compositions**

230 A total of 837 *P. auratus* were collected (not including tagged fish) ranging from 68 to  
231 980 mm FL (Table 2). The majority of *P. auratus* caught in the offshore area were  
232 between 300 and 450 mm FL, with a prominent modal length class at 375-399 mm FL  
233 (Fig. 2). In contrast, the samples caught in the inshore area contained a substantial  
234 number of fish with FL < 300 mm, with no conspicuous mode in the length-frequency  
235 composition. The marine embayments essentially contained fish encompassing the  
236 entire length range of *P. auratus* (Fig. 2) and was the only area where fish < 200 mm FL  
237 were caught. However, in contrast to the situation in offshore and inshore waters, the  
238 fish between 300 and 400 mm FL, which represented fish approaching maturity, were  
239 poorly represented in this area (Fig. 2). Further, the length-frequency distribution in the  
240 marine embayments contained a prominent mode at *ca* 600 to 800 mm FL, representing  
241 the majority of mature individuals caught during this study (Fig. 2).

242  
243 The majority of *P. auratus* in the marine embayments belonged to the 0-1+ or 5-12+  
244 age classes (Fig. 3). In contrast, the majority of fish in both inshore and offshore areas  
245 belonged to the 2-6+ age classes, producing modal age classes of 3+ and 4+ years,  
246 respectively (Fig. 3).

247  
248 All *P. auratus* < 200 mm FL were caught in the marine embayments (Figs 2 & 4). The  
249 percentage contribution of individuals caught in the marine embayments to the total  
250 catch from all areas declined progressively from 100 % in the 150-199 mm FL class to  
251 5.4 % in the 300-349 mm FL class (Fig. 4). This marked decline in the proportions of  
252 juveniles in the marine embayments was reflected in the logistic parameter  $L_{50}$  and  $L_{95}$   
253 values of 251 and 322 mm FL, respectively (Fig. 4). The proportions of *P. auratus* in

254 the marine embayments then increased progressively to > 90 % in all length classes  
255 from 600 to 849 mm FL (Fig. 4).

256

257 The above trends exhibited by the prevalence of successive length classes of *P. auratus*  
258 in the marine embayments were paralleled by those of the age classes. Thus, the  
259 proportion of juveniles caught in the marine embayments declined from 100 % in the 0+  
260 age class to ca 50 % at two years of age and 5 % by three years of age, i.e.  $A_{50}$  and  $A_{95}$   
261 of 1.9 and 2.9 yr, respectively (Fig. 4). All 1+ *P. auratus* were either caught in the  
262 marine embayments or inshore areas (Fig. 4), with the youngest *P. auratus* caught in the  
263 offshore area being 2.2 yr. The lowest percentage contribution to the catches of *P.*  
264 *auratus* in the embayments was 5.4 % in the 4+ age class, with ca 70 % of this age class  
265 being recorded in the offshore area (Fig. 4). The percentage contribution to the catches  
266 in the embayments increased progressively after the 4+ age class, with fish from this  
267 area representing  $\geq 80$  % in the majority of age classes above 7+ (Fig. 4).

268

269 The trends exhibited by monthly length-frequency distributions emphasised that very  
270 few *P. auratus* > 200 mm FL were caught in the marine embayments between March  
271 and July, but that the numbers of such fish increased markedly after August when  
272 spawning commenced, and declined in January when spawning ceased (Fig. 5).

273 Juveniles < 250 mm FL were caught in the marine embayments in several months.

274 Substantial numbers of *P. auratus* between 250 and 500 mm FL were caught outside the  
275 embayments in all months (Fig. 5).

276

277 The proportions of *P. auratus* in spawning condition (i.e. with gonads at stages II to V,  
278 Table 1) between September and January, when spawning occurs, was greatest in the

279 three adjacent embayments (91 %) than in offshore waters immediately west at depths  
280 > 80 m, i.e. 12 %, and in inshore waters < 80 m depth (, i.e. 30 %, Fig. 6). Thus,  
281 although *P. auratus* in spawning condition were caught outside the embayments,  
282 catches in that location were dominated by smaller, immature individuals (Fig. 6). In  
283 contrast, essentially all *P. auratus* > 325 mm FL in catches from the embayments during  
284 the spawning months of September to January possessed gonads that were developing  
285 (stage II), developed (stages III), spawning (stage IV) or had recently spawned (stage V,  
286 Fig. 6). The lengths of the majority of fish caught in the embayments during the  
287 spawning period in each year from 2003 to 2008 were between 550 and 800 mm FL  
288 (Fig. 6), a length range that was poorly represented in the samples from the other two  
289 areas at this time of year (Fig. 6).

290

### 291 **Fish recaptures**

292 The lengths of the 777 *P. auratus* that were tagged and released in Cockburn and  
293 Warnbro Sounds between September and December of 2003 to 2009, i.e. during seven  
294 successive spawning seasons, ranged from 355 to 920 mm FL. Forty-nine of these  
295 tagged fish were reported to have been recaptured (6.3 %, Table 3), with their times at  
296 liberty ranging from 4 to 1,827 days (ca 5 yr). All but 11 of these fish were recaptured  
297 in Cockburn Sound. The longest distances moved by these 11 exceptions were 92 km to  
298 the north, ca 35 km to the west and 125 and 134 km to the south and southwest,  
299 respectively (Fig. 1). All but one of these eleven *P. auratus* were re-caught during non-  
300 spawning months, with the single exception originally tagged in October 2006 and re-  
301 caught 51 days later, i.e. in December, south-east of Rottnest Island. The greatest  
302 movement northwards (92 km) was that exhibited by a mature fish of 832 mm FL, when  
303 tagged in October 2003 in Cockburn Sound and recaptured 454 days later in January

304 2005. The greatest movement southwards (134 km) was that exhibited by a mature fish  
305 of 783 mm FL, which was tagged in October 2008 in Cockburn Sound and recaptured  
306 231 days later in June 2009.

307

308 The 38 *P. auratus*, which were recaptured in Cockburn Sound, comprised 12 that were  
309 caught in the spawning season in which they were tagged and 12 that were caught in  
310 spawning periods subsequent to that of their tagging and release. The remaining 14  
311 were caught in Cockburn Sound during non-spawning months. A total of 530 of the 777  
312 tagged *P. auratus* had two tags inserted. A total of 31 of these fish were recaptured, 11  
313 of which had lost one tag. Due to the low number of recaptures of double-tagged  
314 *P. auratus*, the tag shedding rates with respect to time at liberty and a taggers ability to  
315 insert tags was uncertain.

316

### 317 **Discussion**

318 The data collected during this study showed that the distribution of *P. auratus* between  
319 *ca* 31°45' and 32°45' S on the lower west coast of Western Australia varied with life  
320 cycle stage. Within these latitudes, the coastal marine embayments of Cockburn Sound,  
321 Warnbro Sound and Owen Anchorage were found clearly to constitute nursery and  
322 spawning areas for this species. The monthly length- and age-frequency distributions  
323 demonstrated that *P. auratus* use these embayments as a nursery area during the first  
324 two years of life. This finding is consistent with the results of Lenanton (1974), who  
325 collected small (< 150 mm FL) and young (< *ca* 15 months old) *P. auratus* by trawling  
326 over soft substrates in Cockburn Sound. Studies of *P. auratus* in New Zealand and  
327 south-eastern Australia also found that the abundance of juveniles was relatively high in  
328 sheltered inshore areas (e.g. Francis, 1995; Hamer and Jenkins, 2004; Paul and Tarring,

329 1980). In addition, two recent studies on the age-related elemental profiles of otoliths  
330 showed that *P. auratus* collected over a large stretch of coast, i.e. 700 km in Victoria  
331 and > 2000 km in South Australia, could be linked to one or two points of origin  
332 (nursery areas) in coastal marine embayments (Fowler et al., 2005; Hamer et al., 2005).  
333 This highlights the importance of discrete nursery areas for *P. auratus*, and in Western  
334 Australia could apply to these three adjacent marine embayments (between 32°15' and  
335 32°40'S). In contrast, two tagging studies of *P. auratus* that targeted stocks in large sub-  
336 tropical marine embayments on the east and west coasts of Australia found that there  
337 was limited or no exchange of juveniles and adults between these bays and outside  
338 waters (Moran et al., 2003; Sumpton et al., 2003). This suggests that these two  
339 subtropical embayments provide the necessary resources (e.g. habitat and prey) to  
340 support the full life cycle of *P. auratus* (see Ross, 1986).

341

342 In the case of immature *P. auratus* on the lower west coast of Australia, i.e. those with  
343 lengths <  $L_{50}$  of 488-505 mm FL and  $A_{50}$  of 5.6-5.7 years at maturity (Wakefield, 2006),  
344 their lengths and ages increased progressively from within to immediately outside the  
345 embayments (< 80 m) and then to the area further offshore (> 80 m). This increase in  
346 length and age with distance west of the embayments and prior to maturation indicates  
347 dispersal from a nursery habitat. It is thus concluded that the three adjacent marine  
348 embayments constitute important nursery areas, at least between 32°15' and 32°40'S on  
349 the lower west coast of Australia. The increase in size and age from shallow, inshore to  
350 deeper, offshore waters parallels the trends exhibited by other sparids, including *P.*  
351 *auratus* in the East Cape region of New Zealand (Paul and Tarring, 1980),  
352 *Rhabdosargus sarba* in Western Australia (Hesp et al., 2004a) and *Argyrozona*  
353 *argyrozona* in South Africa (Griffiths and Wilke, 2002).



354

355 As the water circulation in Cockburn and Warnbro Sounds during the spawning period  
356 of *P. auratus* facilitates the retention of eggs and larvae in those embayments  
357 (Wakefield, 2010) and 0+ fish < 200 mm FL were found exclusively in these  
358 embayments, the period and mode of dispersal for this species on the lower west coast  
359 of Australia appears to be that derived from movement of individuals from these  
360 embayments at ca 2-4 years of age and thus prior to reaching sexual maturity  
361 (Wakefield, 2006). Fowler et al. (2005) found that, in South Australia, *P. auratus*  
362 moved considerable distances between the ages of 2-5 years, which also represents its  
363 sub-adult stage in that region and is consistent with the movements inferred from length  
364 and age compositions in this study.

365

366 During the spawning season, the relative abundances of *P. auratus*  $\geq L_{50}$  at maturity of  
367 505 mm FL females and 488 mm FL for males (Wakefield, 2006), were far greater in  
368 the marine embayments than in surrounding oceanic waters. As large numbers of large  
369 fish were not present, however, in these embayments after spawning had been  
370 completed, the majority of *P. auratus* that had aggregated and spawned in marine  
371 embayments had apparently moved back into inshore and offshore waters. This  
372 conclusion is consistent with tag-recapture data, which demonstrated that, at least some  
373 but not all *P. auratus*, underwent such a movement. The relatively low numbers of large  
374 fish (> 600 mm FL) in all areas during non-spawning months (February to August),  
375 suggest that fish belonging to the spawning aggregations in the marine embayments  
376 undergo a wide dispersal to surrounding waters and become less vulnerable to capture  
377 throughout these locations and months. Such movement patterns associated with  
378 spawning and non-spawning periods parallel those of *P. auratus* off the mid-west coast

379 of Australia (near Koks, Bernier and Dorre Islands in the waters outside of Shark Bay),  
380 approximately 620 km to the north. At those latitudes, adults move from surrounding  
381 waters of the continental shelf to form a few spawning aggregations on the inshore reefs  
382 during the spawning season (Moran et al., 2003). Tag-recapture results from the current  
383 study demonstrated that the home range of *P. auratus* during non-spawning months  
384 extended ca 100 km north to 31°25' S (south of Lancelin, Fig. 1), ca 135 km south to  
385 33°20' S (west of Bunbury) and ca 35 km west from Cockburn and Warnbro Sounds.  
386

387 Fowler et al. (2005) found that, after ca five years of age, the individuals of *P. auratus*  
388 in South Australia remained in the regions to which they had moved. However, *P.*  
389 *auratus* from that study were collected over more than 2000 km of coast, whereas, in  
390 the present study, the distance from the coastal marine embayments to the western  
391 border of the study area was only ca 65 km and the latitudinal range sampled was only  
392 ca 110 km. Thus, the proportion of young *P. auratus* that move beyond the study area,  
393 after leaving the coastal marine embayments where they were spawned, and the distance  
394 along the Western Australian coast they travel is not well understood.

395  
396 The movements inferred by the length and age distributions and tagging results would  
397 benefit from further investigation using the elemental profiles of otoliths of *P. auratus*  
398 collected over a larger range of coast, similar to the approaches used by Fowler et al.  
399 (2005) and Hamer et al. (2005). This would help identify the extent to which *P. auratus*  
400 from the three adjacent marine embayments recruit to adult stocks along the west coast  
401 of Australia. Nonetheless, the present study has demonstrated that the three adjacent  
402 marine embayments of Owen Anchorage, Cockburn Sound and Warnbro Sound  
403 constitute important spawning and nursery areas for *P. auratus*. Based on available data,

404 the importance of these embayments as a recruitment source for *P. auratus* stocks  
405 should not be underestimated, considering there is a paucity of similar nearshore marine  
406 embayments and hence potential recruitment sources along the essentially linear lower  
407 west coast of Australia. It is thus important for the sustainability of at least nearby adult  
408 stocks to maintain adequate biomass levels associated with spawning aggregations in  
409 those embayments. Furthermore, knowledge of the current status of those stocks would  
410 benefit from annual monitoring of both spawning aggregation biomass and subsequent  
411 juvenile recruitment strength. This should be undertaken in conjunction with existing  
412 monitoring of age structures and rates of fishing mortality at the broader scale, which  
413 are currently the main method used to assess stock status (see Wise et al., 2007).  
414 However, the present study has revealed that such monitoring would need to consider  
415 both temporal and spatial effects of the movement of individuals of this species on  
416 sampled age structures, to ensure that they are representative of the populations along  
417 the lower west coast of Australia. It is also vital that habitats within these nursery  
418 embayments are protected in order to maintain adequate recruitment success to nearby  
419 exploited adult fisheries for this species. This is particularly the case with Cockburn  
420 Sound, as this embayment has a history of ecosystem changes resulting from intense  
421 human use (e.g. Kendrick et al., 2002), with further large-scale development and  
422 dredging proposed, given that larvae of *P. auratus* are susceptible to gill-fouling caused  
423 by increased sedimentation in the water column (Partridge and Michael, 2010).  
424  
425 In summary, this study has demonstrated that three embayments on the lower west coast  
426 of Australia constitute very important spawning grounds and nursery areas for  
427 *P. auratus* in this region. From a fisheries perspective, this makes the stock of this  
428 species particularly susceptible to the effects of recreational and commercial fishing and

429 habitat degradation in the embayments. This point was recognised by the Department of  
430 Fisheries Western Australia and led to the closure of the embayments to fishing for  
431 *P. auratus* from October to January, which encompasses the majority of its spawning  
432 period. Furthermore, the data produced during the study has led to the development of a  
433 more sophisticated and ongoing approach to determining the status of the stock within  
434 and outside the embayments. Our results have also highlighted the need to guard against  
435 deleterious anthropogenic changes to the three embayments as these habitats are so  
436 important to *P. auratus* and thus apparently required to sustain the stocks of this species  
437 at a level that will maintain the structure of the ecosystem of which this species is a part.

438

#### 439 **Acknowledgements**

440 We extend our gratitude to all scientists, charter operators and fishers who helped with  
441 the collection of samples and tagging. Thanks also to Brett Molony, Dan Gaughan and  
442 Gary Jackson whose comments on earlier versions of the manuscript were greatly  
443 appreciated. Financial support was provided by the Department of Fisheries, Western  
444 Australia, Murdoch University and Fisheries Research and Development Corporation  
445 (FRDC, Project No. 2003/052).

446

#### 447 **References**

- 448 Bennett, B.A., 1993. Aspects of the biology and life history of white steenbras  
449 *Lithognathus lithognathus* in southern Africa. S. Afr. J. Mar. Sci. 13, 83-96.
- 450 Colin, P.L., 2010. Aggregation and spawning of the humphead wrasse *Cheilinus*  
451 *undulatus* (Pisces: Labridae): general aspects of spawning behaviour. J. Fish  
452 Biol. 76, 987-1007.

- 453 Coutin, P., Cashmore, S., Sivakumuran, K.P., 2003. Assessment of the snapper fishery  
454 in Victoria. FRDC (Project No. 97/127). Marine and Freshwater Resources  
455 Institute, Melbourne, 205 pp.
- 456 Crossland, J., 1977. Seasonal reproductive cycle of snapper *Chrysophrys auratus*  
457 (Forster) in the Hauraki Gulf. N. Z. J. Mar. Freshwater Res. 11, 37-60.
- 458 Doak, C., 2004. Hydrodynamic modeling of snapper (*Pagrus auratus*) eggs and larvae  
459 in Cockburn Sound. Centre for Water Research. University of Western  
460 Australia, 81 pp.
- 461 Domeier, M.L., Colin, P.L., 1997. Tropical reef fish spawning aggregations: defined  
462 and reviewed. Bull. Mar. Sci. 60, 698-726.
- 463 Fowler, A.J., Gillanders, B.M., Hall, K.C., 2005. Relationships between elemental  
464 concentration and age from otoliths of adult snapper (*Pagrus auratus*, Sparidae):  
465 implications for movement and stock structure. Mar. Freshwater Res. 56, 661-  
466 676.
- 467 Francis, M.P., 1995. Spatial and seasonal variation in the abundance of juvenile snapper  
468 (*Pagrus auratus*) in the north-western Hauraki Gulf. N. Z. J. Mar. Freshwater  
469 Res. 29, 565-579.
- 470 Francis, M.P., Williams, M.W., Pryce, A.C., Pollard, S., Scott, S.G., 1992. Daily  
471 increments in otoliths of juvenile snapper, *Pagrus auratus* (Sparidae). Aust. J.  
472 Mar. Freshwater Res. 43, 1015-1032.
- 473 Gillanders, B.M., 2002. Connectivity between juvenile and adult fish populations: do  
474 adults remain near their recruitment estuaries? Mar. Ecol. Prog. Ser. 240, 215-  
475 223.

- 476 Gillanders, B.M., Able, K.W., Brown, J.A., Eggleston, D.B., Sheridan, P.F., 2003.  
477 Evidence of connectivity between juvenile and adult habitats for mobile marine  
478 fauna: an important component of nurseries. *Mar. Ecol. Prog. Ser.* 247, 281-295.
- 479 Griffiths, M.H., Wilke, C.G., 2002. Long-term movement patterns of five temperate-  
480 reef fishes (Pisces: Sparidae): implications for marine reserves. *Mar. Freshwater*  
481 *Res.* 53, 233-244.
- 482 Hamer, P.A., Jenkins, G.P., 2004. High levels of spatial and temporal recruitment  
483 variability in the temperate sparid *Pagrus auratus*. *Mar. Freshwater Res.* 55,  
484 663-673.
- 485 Hamer, P.A., Jenkins, G.P., Gillanders, B.M., 2005. Chemical tags in otoliths indicate  
486 the importance of local and distant settlement areas to populations of a temperate  
487 sparid, *Pagrus auratus*. *Can. J. Fish. Aquat. Sci.* 62, 623-630.
- 488 Hesp, S.A., Hall, N.G., Potter, I.C., 2004a. Size-related movements of *Rhabdosargus*  
489 *sarba* in three different environments and their influence on estimates of von  
490 Bertalanffy growth parameters. *Mar. Biol.* 144, 449-462.
- 491 Hesp, S.A., Potter, I.C., Hall, N.G., 2004b. Reproductive biology and protandrous  
492 hermaphroditism in *Acanthopagrus latus*. *Environ. Biol. Fishes* 70, 257-272.
- 493 Heyman, W.D., Kjerfve, B., 2008. Characterization of transient multi-species reef fish  
494 spawning aggregations at Gladden Spit, Belize. *Bull. Mar. Sci.* 83, 531-551.
- 495 Jackson, G., Cheng, Y.W., 2001. Parameter estimation with egg production surveys to  
496 estimate snapper, *Pagrus auratus*, biomass in Shark Bay, Western Australia. *J.*  
497 *Agric. Biol. Environ. Stat.* 6, 243-257.
- 498 Kendrick, G.A., Aylwarda, M.J., Hegge, B.J., Cambridge, M.L., Hillman, K., Wyllie,  
499 A., Lord, D.A., 2002. Changes in seagrass coverage in Cockburn Sound,  
500 Western Australia between 1967 and 1999. *Aquat. Bot.* 73, 75-87.

- 501 Lenanton, R.C.J., 1974. The abundance and size composition of trawled juvenile  
502 snapper *Chrysophrys unicolor* (Quoy and Gaimard) from Cockburn Sound,  
503 Western Australia. Aust. J. Mar. Freshwater Res. 25, 281-285.
- 504 McGlennon, D., 2004. The fisheries biology and population dynamics of snapper  
505 *Pagrus auratus* in northern Spencer Gulf, South Australia. Department of  
506 Environmental Biology. University of Adelaide, Australia, 219 pp.
- 507 McGlennon, D., Partington, D., 1997. Mortality and tag loss in dart and loop-tagged  
508 captive snapper, *Pagrus auratus* (Sparidae), with comparisons to relative  
509 recapture rates from a field study. N. Z. J. Mar. Freshwater Res. 31, 39-49.
- 510 Moran, M., Burton, C., Jenke, J., 2003. Long-term movement patterns of continental  
511 shelf and inner gulf snapper (*Pagrus auratus*, Sparidae) from tagging in the  
512 Shark Bay region of Western Australia. Mar. Freshwater Res. 54, 913-922.
- 513 Nahas, E.L., Jackson, G., Pattiaratchi, C.B., Ivey, G.N., 2003. Hydrodynamic modelling  
514 of snapper *Pagrus auratus* egg and larval dispersal in Shark Bay, Western  
515 Australia: reproductive isolation at a fine spatial scale. Mar. Ecol. Prog. Ser.  
516 265, 213-226.
- 517 Otway, N.M., Craig, J.R., 1993. Effects of hook size on the catches of undersized  
518 snapper *Pagrus auratus*. Mar. Ecol. Prog. Ser. 93, 9-15.
- 519 Partridge, G., Michael, R.J., 2010. Direct and indirect effects of simulated calcareous  
520 dredge material on eggs and larvae of pink snapper *Pagrus auratus*. J. Fish Biol.  
521 77, 227-240.
- 522 Paul, L.J., Tarring, S.C., 1980. Growth rate and population structure of snapper,  
523 *Chrysophrys auratus*, in the East Cape region, New Zealand. N. Z. J. Mar.  
524 Freshwater Res. 14, 237-247.

- 525 Paulin, C.D., 1990. *Pagrus auratus*, a new combination for the species known as  
526 'snapper' in Australasian waters (Pisces: Sparidae). N. Z. J. Mar. Freshwater Res.  
527 24, 259-265.
- 528 Punt, A.E., Kennedy, R.B., 1997. Population modelling of Tasmanian rock lobster,  
529 *Jasus edwardsii*, resources. Mar. Freshwater Res. 48, 967-980.
- 530 Quartararo, N., Kearney, R.E., 1996. Effects of dart tags on the growth and survival of  
531 captive snapper, *Pagrus auratus* (Sparidae). Fish. Res. 25, 231-238.
- 532 Ross, S.T., 1986. Resource partitioning in fish assemblages: A review of field studies.  
533 Copeia 1986, 352-388.
- 534 Sanders, M.J., 1974. Tagging indicates at least two stocks of snapper *Chrysophrys*  
535 *auratus* in south-east Australian waters. N. Z. J. Mar. Freshwater Res. 8, 371-  
536 374.
- 537 Scott, S.G., Zeldis, J.R., Pankhurst, N.W., 1993. Evidence of daily spawning in natural  
538 populations of the New Zealand snapper *Pagrus auratus* (Sparidae). Environ.  
539 Biol. Fish. 36, 149-156.
- 540 Sumpton, W.D., Sawynok, B., Carstens, N., 2003. Localised movement of snapper  
541 (*Pagrus auratus*, Sparidae) in a large subtropical marine embayment. Mar.  
542 Freshwater Res. 54, 923-930.
- 543 Wakefield, C.B., 2006. Latitudinal and temporal comparisons of the reproductive  
544 biology and growth of snapper, *Pagrus auratus* (Sparidae), in Western Australia.  
545 Centre for Fish and Fisheries Research. Murdoch University, Western Australia,  
546 162 pp.
- 547 Wakefield, C.B., 2010. Annual, lunar and diel reproductive periodicity of a spawning  
548 aggregation of snapper *Pagrus auratus* (Sparidae) in a marine embayment on the  
549 lower west coast of Australia. J. Fish Biol. 77, 1359-1378.



550 Wise, B.S., St John, J., Lenanton, R.C.J., 2007. Spatial scales of exploitation among  
551 populations of demersal scalefish: implications for management. Part 1: Stock  
552 status of the key indicator species for the demersal scalefish fishery in the West  
553 Coast Bioregion. Department of Fisheries, Western Australia, 130 pp.

554

### 555 **Table and Figure captions**

556 **Table 1.** Macroscopic characteristics in the appearance of ovaries and testes of *Pagrus*  
557 *auratus*.

558

559 **Table 2.** Sample sizes and fork lengths (FL, mm) of *Pagrus auratus* caught by hook and  
560 line in each of the three areas.

561

562 **Table 3.** Numbers of *Pagrus auratus* tagged and recaptured in each year.

563

564 **Figure 1.** Map showing the three sampling areas, *i.e.* (1) offshore (depth > 80 m), (2)  
565 inshore (depth < 80 m) and (3) marine embayments (black areas, *i.e.* Owen Anchorage,  
566 Cockburn Sound & Warnbro Sound) and the recapture locations of snapper that had  
567 been tagged in Cockburn Sound (dashed lines with circles).

568

569 **Figure 2.** Length-frequency distributions of *Pagrus auratus* that were retained (grey  
570 bars) or tagged and released (white bars, embayments only) in sequential 25 mm length  
571 classes from the three areas (sample sizes given).

572

573 **Figure 3.** Age-frequency distributions for *Pagrus auratus* in successive age classes up  
574 to 15+ years caught in each of the three areas. The solid line represents the age at 50 %  
575 maturity ( $A_{50}$ )  $\pm$  95 % CI (dashed lines) for females and males (Wakefield, 2006).

576

577 **Figure 4.** Cumulative frequency (%) contributions made by the numbers of *Pagrus*  
578 *auratus* in successive length (excludes tagged fish) and age classes in samples obtained  
579 from the embayments (dark grey bars), inshore (light grey bars) and offshore (white  
580 bars) areas. Sample sizes are given for each length and age class (above). Lines  
581 represent the expected percentage ( $\pm$  95 % CI) of juvenile *P. auratus* in the marine  
582 embayments (solid) as determined from logistic regression analysis.

583

584 **Figure 5.** Monthly length-frequency distributions for *Pagrus auratus* in marine  
585 embayments (black bars, excludes tagged fish) and the other areas combined (grey  
586 bars).

587

588 **Figure 6.** Length-frequency distributions of immature (gonad stage I, grey bars) and  
589 mature (gonad stages II-V, black bars) *Pagrus auratus* in sequential 25 mm length  
590 classes caught during the spawning period, *i.e.* September to January, in each of the  
591 three areas. Lines represent the length at 50 % maturity ( $L_{50}$ ) for females (dashed) and  
592 males (solid) of 505 and 488 mm FL, respectively (Wakefield, 2006).

1 **Table 1.**

Stage	Ovaries	Testes
I. Immature / Resting	Occupy up to one half of length of ventral cavity. Cylindrical, blood capillaries visible and pink to orange.	Occupy up to one half of ventral cavity. Flat and white.
II. Developing	Occupy up to two thirds of ventral cavity. Blood capillaries and oocytes visible.	Occupy up to two thirds of ventral cavity and white colour more apparent.
III. Developed	Occupy full length of ventral cavity. Oocytes clearly visible but not hydrated. Blood capillaries more conspicuous.	Testis much larger occupying up to full length of ventral cavity. No milt discharged when slight pressure is applied to lobes or abdomen.
IV. Ripe / Spawning	Similar in size to stage III. Hydrated oocytes (translucent) visible throughout ovarian lobes or concentrated in oviduct.	Similar in size to stage III. Milt discharged when slight pressure is applied to lobes or abdomen.
V. Spent	Ovaries reduced in size, flaccid and red in areas.	Testes reduced in size, flaccid and red in areas.

2

3

4

5

6

7

**Table 2.**

Area	<i>n</i>	Fork length (mm)			<i>n</i>	Age (years)		
		Range	Mean	SD		Range	Mean	SD
Offshore								
Retained	337	245 - 869	402	89	274	2.2 - 28.8	5.18	2.98
Inshore								
Retained	227	202 - 980	414	124	193	1.5 - 11.5	4.43	1.81
Marine embayments								
Retained	273	68 - 901	444	245	273	0.3 - 24.0	6.13	4.31
Tagged	777	355 - 920	664	104				

8

*n* = Sample size, SD = Standard deviation


9

10

11 **Table 3.**

		<b>Tagged in</b>							
		<i>Year</i>	<i>2003</i>	<i>2004</i>	<i>2005</i>	<i>2006</i>	<i>2007</i>	<i>2008</i>	<i>Total</i>
		<i>(from Oct)</i>							
<b>Recaptured in</b>	<i>Year</i>	<b>Total</b>	<b>54</b>	<b>277</b>	<b>143</b>	<b>120</b>	<b>22</b>	<b>161</b>	<b>777</b>
	<i>2003</i>	<b>2</b>	2						
	<i>(from Oct)</i>								
	<i>2004</i>	<b>8</b>	4	4					
	<i>2005</i>	<b>9</b>	3	5	1				
	<i>2006</i>	<b>17</b>	1	9	4	3			
	<i>2007</i>	<b>5</b>	0	1	2	2	0		
	<i>2008</i>	<b>5</b>	0	2	1	0	0	2	
	<i>2009</i>	<b>3</b>	0	1	0	0	0	2	
	<i>(to Sep)</i>								
	<b>Total</b>	<b>49</b>							
	Recapture rate (%)		18.52	7.94	5.59	4.17	0	2.48	


12



13


14

15 **Figure 1.**



16  
17


**Figure 2.**



18

19

20 **Figure 3.**




21

22


23 **Figure 4.**





24

25 **Figure 5.**



26

27

28 **Figure 6.**

**Highlights**

- Habitat partitioning is reported among major life cycle stages of *Pagrus auratus*
- Three metropolitan embayments are crucial for spawning aggregations and juveniles
- Sub-adults occupy deeper offshore waters adjacent to the embayments
- A closure to fishing has been implemented to protect spawning aggregations
- Industrial development in the embayments may influence spawning and recruitment success

Accepted Manuscript