

electronic reprint

Acta Crystallographica Section C

**Crystal Structure
Communications**

ISSN 0108-2701

Ethyl and isopropyl 4-ferrocenylbenzoate

Frankie P. Anderson, John F. Gallagher, Peter T. M. Kenny, Clodagh Ryan and David Savage

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

Ethyl and isopropyl 4-ferrocenylbenzoate

Frankie P. Anderson,^a John F. Gallagher,^{a*} Peter T. M. Kenny,^{a*} Clodagh Ryan^b and David Savage^b^aSchool of Chemical Sciences, National Institute for Cellular Biotechnology, Dublin City University, Dublin 9, Ireland, and ^bSchool of Chemical Sciences, Dublin City University, Dublin 9, Ireland

Correspondence e-mail: john.gallagher@dcu.ie

Received 18 November 2002

Accepted 26 November 2002

Online 10 December 2002


The title compounds, $[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{14}\text{H}_{13}\text{O}_2)]$ and $[\text{Fe}(\text{C}_5\text{H}_5)(\text{C}_{15}\text{H}_{15}\text{O}_2)]$, respectively, contain the ferrocenyl $\eta^5(\text{C}_5\text{H}_4)$ and phenylene $-\text{C}_6\text{H}_4-$ rings in a nearly coplanar arrangement, with interplanar angles of 6.88 (12) and 10.5 (2)°, respectively. Molecules of the ethyl ester form dimers through $\eta^5(\text{C}_5\text{H}_5)\text{C}-\text{H}\cdots\text{O}=\text{C}$ hydrogen bonds, with graph set $R_2^2(20)$, and, together with $\text{Csp}^3-\text{H}\cdots\pi(\text{C}_5\text{H}_5)$ interactions, generate a one-dimensional column (irregular ladder). Molecules of the isopropyl ester aggregate through $\eta^5(\text{C}_5\text{H}_5)\text{C}-\text{H}\cdots\pi(\text{C}_6\text{H}_4)$ interactions.

Comment

The design of new redox-active ligands for application in diverse research areas, such as medicinal chemistry and materials science, has engrossed scientists in recent years. Ferrocene (Fc) derivatives, which are efficient redox systems, have been studied extensively in charge-transfer chemistry, hydrogen bonding and molecular-recognition science, peptide chemistry, non-linear optical materials, and liquid crystal research (Chesney *et al.*, 1998; Glidewell *et al.*, 1997; Zakaria *et al.*, 2002; Kraatz *et al.*, 1999; Gallagher *et al.*, 1999*a,b*; Hudson, Asselsbergh *et al.*, 2001; Hudson, Manning *et al.*, 2001; Even *et al.*, 2001; Seo *et al.*, 2001). Our interest in ferrocenylbenzoyl derivatives stems from their use as precursors to ferrocenylbenzoyl amino acid ester and dipeptide derivatives. We have recently reported the crystal structure of methyl 4-ferrocenylbenzoate, (I) (Savage *et al.*, 2002).

An understanding of the interactions present in a given crystal structure can provide valuable information on the hydrogen-bonding and aggregation modes not just in the solid state but also in the liquid-crystalline state. The structures of ethyl 4-ferrocenylbenzoate, (II), and the isopropyl analogue, (III), are reported herein for comparison with both the methyl analogue, (I), and our ongoing research on longer chain alkyl derivatives.

The Fe1—C bond lengths for the $\eta^5(\text{C}_5\text{H}_4)$ ring of (II) are in the range 2.0341 (16)–2.0452 (14) Å, similar to the $\eta^5(\text{C}_5\text{H}_5)$ ring, with a range of 2.0315 (18)–2.0407 (17) Å. For (III), these values are in the ranges 2.024 (3)–2.044 (3) and 2.023 (4)–2.033 (4) Å, respectively. In (II), the Fe1 \cdots Cg1 and Fe1 \cdots Cg2 distances are 1.6425 (8) and 1.6463 (9) Å, respectively, and the Cg1 \cdots Fe1 \cdots Cg2 angle is 179.49 (5)°, where Cg1 and Cg2 are the centroids of the $\eta^5(\text{C}_5\text{H}_4)$ and $\eta^5(\text{C}_5\text{H}_5)$ rings, respectively. In (III), these values are 1.6413 (15) and 1.643 (2) Å, and 178.88 (9)°, respectively. In (II), the cyclopentadienyl C—C bond-length ranges are small, being 1.413 (3)–1.435 (2) and 1.398 (3)–1.420 (3) Å for the $\eta^5(\text{C}_5\text{H}_4)$ and $\eta^5(\text{C}_5\text{H}_5)$ rings, respectively. In (III), these ranges are 1.407 (5)–1.428 (4) and 1.389 (6)–1.410 (6) Å, respectively. These results are as expected and highlight the similarity in the ferrocenyl bond lengths and angles in the esters (I) (Savage *et al.*, 2002), and (II) and (III), described herein.



(II) R = ethyl

(III) R = isopropyl

The cyclopentadienyl rings deviate slightly from an eclipsed geometry in (II), as evidenced by the C1 $n\cdots$ Cg1 \cdots Cg2 \cdots C2 n ($n = 1-5$) torsion angles ranging from -2.87 (14) to -3.61 (14)°. In (III), the angles are in the range 6.1 (3)– 7.0 (3)°, similar to the eclipsed geometry in (I), where the range is 0.8 (4)– 2.3 (4)°. In contrast, this range of angles is 13.7 (2)– 15.4 (3)° in *para*-ferrocenylbenzoyl-L-alanine methyl ester (Savage *et al.*, 2002).

The essentially linear molecular conformations adopted by (II) and (III) are comparable, with interplanar angles of 6.88 (12)° between the $\eta^5(\text{C}_5\text{H}_4)$ and $-\text{C}_6\text{H}_4-$ rings in (II), 10.5 (2)° in (III) and 9.35 (13)° in (I). The major differences are in the terminal O1—C1—C34—C33 torsion angles, the value of which is -0.8 (2)° in (II), -18.2 (5)° in (III) and 171.2 (3)° in (I). However, the disposition of the terminal alkoxy group, which gives a C1—O1—C2—C3 torsion angle of




Figure 1

A view of the molecule of (II) with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.




Figure 2
A stereoview of the interactions in the crystal structure of (II).

179.76 (16)° in (II) and 112.0 (5)° in (III), means that the methyl groups of the isopropyl moiety in (III) are oriented almost orthogonal to the ester CO₂ plane (Figs. 1 and 3).

Analysis of the hydrogen bonding in (II) shows only two interactions of note, involving the substituted ring and the ester O=C group as $\eta^5(\text{C}_5\text{H}_4)\text{C}-\text{H}\cdots\text{O}=\text{C}$ interactions (Table 2 and Fig. 2). Molecules of the ethyl ester form dimers through $\eta^5(\text{C}_5\text{H}_5)\text{C}-\text{H}\cdots\text{O}=\text{C}$ hydrogen bonds, graph set $R_2^2(20)$ and, together with $\text{Csp}^3-\text{H}\cdots\pi(\text{C}_5\text{H}_5)$ interactions, these generate a one-dimensional column (irregular ladder). The molecules of the isopropyl ester, (III), aggregate through $\eta^5(\text{C}_5\text{H}_5)\text{C}-\text{H}\cdots\pi(\text{C}_6\text{H}_4)$ interactions (Table 4 and Fig. 4).

Analysis of the April 2002 Version of the Cambridge Structural Database using *ConQuest* Version 1.4 (Allen, 2002)




Figure 3
A view of the molecule of (III) with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.




Figure 4
A stereoview of the interactions in the crystal structure of (III). For the sake of clarity, the unit-cell box has been omitted.

for the mono-substituted *para*-Fc-C₆H₄-X group (X is any atom) was undertaken for structures which fulfil the three-dimensional coordinates and $R < 0.10$ criteria. A total of 13 relevant structures were found. The interplanar angle between the C₅H₄ and C₆ rings varies between 2.2 and 29.1° for 17 examples, with a median of 12.8° (11 examples are within the range 7.7–19.3°). Our results above compare favourably with these values.

Experimental

Compounds (II) and (III) were prepared according to standard literature procedures. Analytical data for (II): m.p. 358–359 K (uncorrected); IR (KBr, ν , cm⁻¹): 1678 (C=O); UV-vis, λ_{max} (CH₃CN): 360 (880), 445 (290) nm; ¹H NMR (400 MHz, CDCl₃, δ , p.p.m.): 7.86 (2H, *d*, $J = 8.4$ Hz, Ar-H), 7.66 (2H, *d*, $J = 8.4$ Hz, Ar-H), 4.89 [2H, *t*, $J = 1.8$ Hz, *o*- $\eta^5(\text{C}_5\text{H}_4)$], 4.44 [2H, *t*, $J = 1.8$ Hz, *m*- $\eta^5(\text{C}_5\text{H}_4)$], 4.32 (2H, *q*, $J = 7$ Hz, -OCH₂CH₃), 4.03 [5H, *s*, $\eta^5(\text{C}_5\text{H}_5)$], 1.33 (3H, *t*, -OCH₂CH₃); ¹³C NMR (CDCl₃, δ , p.p.m.): 166.1, 145.5, 129.6, 127.2, 126.0, 83.0, 70.2, 69.9, 67.2, 60.9, 14.6. Analytical data for (III): m.p. 351–352 K (uncorrected); IR (KBr, ν , cm⁻¹): 1710 (C=O); UV-vis, λ_{max} (CH₃CN): 360 (1120), 458 (302) nm; ¹H NMR (400 MHz, CDCl₃, δ , p.p.m.): 7.87 (2H, *d*, $J = 8.4$ Hz, Ar-H), 7.44 (2H, *d*, $J = 8.4$ Hz, Ar-H), 5.18 [1H, *m*, OCH(CH₃)₂], 4.64 [2H, *s*, *o*- $\eta^5(\text{C}_5\text{H}_4)$], 4.32 [2H, *s*, *m*- $\eta^5(\text{C}_5\text{H}_4)$], 3.96 [5H, *s*, $\eta^5(\text{C}_5\text{H}_5)$], 1.30 [6H, *t*, OCH(CH₃)₂]; ¹³C NMR (CDCl₃, δ , p.p.m.): 166.6, 145.2, 130.0, 128.5, 126.0, 83.9, 70.2, 70.1, 68.5, 67.3, 22.4.

Compound (II)

Crystal data

[Fe(C₅H₅)(C₁₄H₁₃O₂)]
 $M_r = 334.18$
Monoclinic, $P2_1/c$
 $a = 7.9563$ (5) Å
 $b = 16.3464$ (11) Å
 $c = 12.0088$ (10) Å
 $\beta = 94.128$ (5)°
 $V = 1557.78$ (19) Å³
 $Z = 4$

$D_x = 1.425$ Mg m⁻³
Mo $K\alpha$ radiation
Cell parameters from 80 reflections
 $\theta = 5.5$ –20.7°
 $\mu = 0.97$ mm⁻¹
 $T = 294$ (1) K
Block, orange
0.50 × 0.45 × 0.45 mm

Data collection

Siemens P4 diffractometer
 $\omega/2\theta$ scans
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\text{min}} = 0.629$, $T_{\text{max}} = 0.646$
4875 measured reflections
3780 independent reflections
3276 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$
 $\theta_{\text{max}} = 28^\circ$
 $h = -10 \rightarrow 1$
 $k = -21 \rightarrow 1$
 $l = -15 \rightarrow 15$
4 standard reflections
every 296 reflections
intensity variation: $\pm 1\%$

Table 1

Selected geometric parameters (Å, °) for (II).

O1—C1	1.200 (2)	C1—C34	1.489 (2)
O2—C1	1.3282 (19)	C2—C3	1.486 (3)
O2—C2	1.4503 (19)	C11—C31	1.470 (2)
C1—O2—C2	115.63 (13)	O2—C2—C3	107.37 (15)
O1—C1—O2	123.17 (16)	C1—C34—C33	122.85 (14)
O1—C1—C34	124.07 (15)	C1—C34—C35	118.07 (14)
O2—C1—C34	112.76 (13)		
C2—O2—C1—C34	-178.95 (13)	C12—C11—C31—C36	7.9 (2)
C1—O2—C2—C3	179.76 (16)	O1—C1—C34—C33	178.40 (17)

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.079$
 $S = 1.05$
 3780 reflections
 200 parameters
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.039P)^2 + 0.319P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$

Table 2

Hydrogen-bonding and short-contact geometry (\AA , $^\circ$) for (II).

Cg1 is the centroid of the substituted cyclopentadienyl ring.

D—H...A	D—H	H...A	D...A	D—H...A
C22—H22...O1 ⁱ	0.93	2.54	3.360 (3)	147
C2—H2A...Cg1 ⁱⁱ	0.97	2.87	3.770 (2)	155

Symmetry codes: (i) $1 - x, 1 - y, -z$; (ii) $-x, 1 - y, -z$.

Compound (III)

Crystal data

[Fe(C₅H₅)(C₁₅H₁₅O₂)]
 $M_r = 348.21$
 Monoclinic, $P2_1/a$
 $a = 9.3406$ (9) \AA
 $b = 10.1663$ (6) \AA
 $c = 17.9072$ (10) \AA
 $\beta = 90.136$ (6) $^\circ$
 $V = 1700.5$ (2) \AA^3
 $Z = 4$

$D_x = 1.360 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation
 Cell parameters from 72 reflections
 $\theta = 6.1\text{--}15.3^\circ$
 $\mu = 0.89 \text{ mm}^{-1}$
 $T = 294$ (1) K
 Block, red
 $0.39 \times 0.26 \times 0.15 \text{ mm}$

Data collection

Siemens P4 diffractometer
 $\omega/2\theta$ scans
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.769, T_{\max} = 0.875$
 4440 measured reflections
 3355 independent reflections
 2439 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.006$
 $\theta_{\max} = 26^\circ$
 $h = -11 \rightarrow 1$
 $k = -1 \rightarrow 12$
 $l = -22 \rightarrow 22$
 4 standard reflections
 every 296 reflections
 intensity variation: $\pm 1\%$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.109$
 $S = 1.04$
 3355 reflections
 210 parameters
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0468P)^2 + 0.7638P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$

Table 3

Selected geometric parameters (\AA , $^\circ$) for (III).

O1—C1	1.186 (4)	C1—C34	1.485 (4)
O2—C1	1.321 (4)	C2—C3	1.440 (7)
O2—C2	1.481 (4)	C2—C4	1.448 (6)
C1—O2—C2	117.7 (2)	O2—C2—C4	108.2 (3)
O1—C1—O2	124.1 (3)	C3—C2—C4	114.8 (4)
O1—C1—C34	123.8 (3)	C1—C34—C33	118.2 (3)
O2—C1—C34	112.1 (3)	C1—C34—C35	123.7 (3)
O2—C2—C3	107.9 (4)		
C2—O2—C1—O1	5.9 (5)	C1—O2—C2—C3	112.0 (5)

Table 4

Hydrogen-bonding and short-contact geometry (\AA , $^\circ$) for (III).

Cg3 is the centroid of the phenylene ring system.

D—H...A	D—H	H...A	D...A	D—H...A
C2—H2...O1	0.98	2.25	2.696 (4)	106
C12—H12...Cg3 ⁱ	0.93	2.75	3.658 (3)	167

Symmetry code: (i) $x - \frac{3}{2}, -\frac{1}{2} - y, z$.

For compounds (II) and (III), space groups $P2_1/c$ and $P2_1/a$, respectively, were uniquely assigned from the systematic absences and confirmed by the analyses. H atoms were treated as riding atoms, with C—H distances in the range 0.93–0.98 \AA .

For both compounds, data collection: XSCANS (Siemens, 1994); cell refinement: XSCANS; data reduction: XSCANS; program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: ORTEPIII (Burnett & Johnson, 1996) and PLATON (Spek, 1998); software used to prepare material for publication: SHELXL97 and PREP8 (Ferguson, 1998).

FPA thanks the National Institute for Cellular Biotechnology, for a studentship under the Irish Government PRTLI#3 funding scheme. JFG thanks Dublin City University for the purchase of a P4 diffractometer and computer system. DS thanks the School of Chemical Sciences and the Irish–American Partnership for funding to undertake chemical research.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: GD1238). Services for accessing these data are described at the back of the journal.

References

Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
 Burnett, M. N. & Johnson, C. K. (1996). ORTEPIII. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
 Chesney, A., Bryce, M. R., Batsanov, A. S., Howard, J. A. K. & Goldenberg, L. M. (1998). *J. Chem. Soc. Chem. Commun.* pp. 677–679.
 Even, M., Heinrich, B., Guillon, D., Guldi, D. M., Prato, M. & Deschenaux, R. (2001). *Chem. Eur. J.* **7**, 2595–2604.
 Ferguson, G. (1998). PREP8. University of Guelph, Canada.
 Gallagher, J. F., Kenny, P. T. M. & Sheehy, M. J. (1999a). *Inorg. Chem. Commun.* **2**, 200–204.
 Gallagher, J. F., Kenny, P. T. M. & Sheehy, M. J. (1999b). *Inorg. Chem. Commun.* **2**, 327–330.
 Glidewell, C., Ahmed, S. Z., Gallagher, J. F. & Ferguson, G. (1997). *Acta Cryst.* **C53**, 1775–1778.
 Hudson, R. D. A., Asselsbergh, I., Clays, K., Cuffe, L. P., Gallagher, J. F., Manning, A. R., Persoons, A. & Wostyn, K. (2001). *J. Organomet. Chem.* **637–639**, 435–444.
 Hudson, R. D. A., Manning, A. R., Nolan, D. F., Asselsbergh, I., Van Boxel, R., Persoons, A. & Gallagher, J. F. (2001). *J. Organomet. Chem.* **619**, 141–151.
 Kraatz, H.-B., Leek, D. M., Houmam, A., Enright, G. D., Lustyk, J. & Wayner, D. D. M. (1999). *J. Organomet. Chem.* **589**, 38–49.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Savage, D., Gallagher, J. F., Ida, Y. & Kenny, P. T. M. (2002). *Inorg. Chem. Commun.* **5**, 1034–1040.
 Seo, J. S., Yoo, Y. S. & Choi, M. G. (2001). *J. Mater. Chem.* **11**, 1332–1338.
 Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.
 Siemens (1994). XSCANS. Version 2.2. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Spek, A. L. (1998). PLATON. University of Utrecht, The Netherlands.
 Zakaria, C. M., Ferguson, G., Lough, A. J. & Glidewell, C. (2002). *Acta Cryst.* **C58**, m1–m4.