

Coherent Spin Ratchets

Matthias Scheid

Institut für Theoretische Physik, Universität Regensburg, D-93040, Germany

Andreas Pfund

*Institut für Theoretische Physik, Universität Regensburg, D-93040, Germany and
ETH Hönggerberg, Solid State Physics Lab, HPF E6, Zürich, CH-8093, Switzerland.*

Dario Bercioux*

*Institut für Theoretische Physik, Universität Regensburg, D-93040, Germany and
Physikalisches Institut, Albert-Ludwigs-Universität, D-79104 Freiburg, Germany*

Klaus Richter

Institut für Theoretische Physik, Universität Regensburg, D-93040, Germany

(Dated: February 5, 2008)

We demonstrate that the combined effect of a spatially periodic potential, lateral confinement and spin-orbit interaction gives rise to a quantum ratchet mechanism for spin-polarized currents in two-dimensional coherent conductors. Upon adiabatic ac-driving, in the absence of a net static bias, the system generates a directed spin current while the total charge current is zero. We analyze the underlying mechanism by employing symmetry properties of the scattering matrix and numerically verify the effect for different setups of ballistic conductors. The spin current direction can be changed upon tuning the Fermi energy or the strength of the Rashba spin-orbit coupling.

PACS numbers: 05.60.Gg, 73.23.-b, 72.25.-b

I. INTRODUCTION

Charge transport is usually studied by considering current in response to an externally applied bias. However, there has been growing interest throughout the last decade in mechanisms enabling directed particle motion in nanosystems without applying a net dc-bias. In this respect, ratchets, periodic structures with broken spatial symmetry, *e.g.* saw tooth-type potentials, represent a prominent class. Ratchets in the original sense are devices operating far from equilibrium by converting thermal fluctuations into directed particle transport in the presence of unbiased time-periodic driving¹. First discovered in the context of (overdamped) classical Brownian motion^{2,3}, the concept of dissipative ratchets was later generalized to the quantum realm⁴. More recently, coherent ratchets and rectifiers have gained increasing attention. They are characterized by coherent quantum dynamics in the central periodic system in between leads where dissipation takes place. Proposals comprise molecular wires⁵ and cold atoms in optical lattices⁶, besides Hamiltonian ratchets⁷. Experimentally, ratchet-induced charge flow in the coherent regime was first observed in a chain of triangular-shaped lateral quantum dots⁸ and later in lateral superlattices⁹.

Here we propose a different class of ratchet devices, namely *spin ratchets* which act as sources for spin currents with simultaneously vanishing charge, respectively particle currents. To be definite we consider coherent transport through ballistic mesoscopic conductors in the presence of spin-orbit (SO) interaction¹⁰. Contrary to particle ratchets, which rely on asymmetries in either the

spatially periodic modulation or the time-periodic driving, a SO-based ratchet works even for symmetric periodic potentials. As possible realizations we have in mind semiconductor heterostructures with Rashba SO interaction¹¹ that can be tuned in strength by an external gate voltage allowing to control the spin evolution.

Among other features it is this property which is triggering recent broad interest in semiconductor-based spin electronics¹². Also since direct spin injection from a ferromagnet into a semiconductor remains problematic¹³, alternatively, several suggestions have been made for generating spin-polarized charge carriers without using magnets. In this respect, spin pumping appears promising, *i.e.* the generation of spin-polarized currents at zero bias via cyclic variation of at least two parameters. Different theoretical proposals based on SO¹⁴ and Zeeman¹⁵ mediated spin pumping in non-magnetic semiconductors have been put forward¹⁶ and, in the latter case, experimentally observed in mesoscopic cavities¹⁷.

While pumps and ratchets share the appealing property of generating directed flow without net bias, ratchet transport requires only a single driving parameter, the periodic ratchet potential has a strong collective effect on the spin current and gives rise to distinct features such as spin current reversals upon parameter changes.

II. MODEL AND SYMMETRY CONSIDERATIONS

We consider a two-dimensional coherent ballistic conductor in the plane (x, z) connected to two nonmagnetic

leads. The Hamiltonian of the central system in presence of Rashba SO interaction reads

$$\mathcal{H}_c = \frac{\hat{p}^2}{2m^*} + \frac{\hbar k_{\text{SO}}}{m^*} (\hat{\sigma}_x \hat{p}_z - \hat{\sigma}_z \hat{p}_x) + U(x, z). \quad (1)$$

Here m^* is the effective electron mass, $U(x, z)$ includes the ratchet potential in x - and a lateral transverse confinement in z -direction, and $\hat{\sigma}_i$ denote Pauli spin matrices. The effect of the SO coupling with strength k_{SO} is twofold: it is leading to spin precession and it is coupling transversal modes in the confining potential¹⁸.

In view of a ratchet setup we consider an additional time-periodic driving term $\mathcal{H}_V(t)$ due to an external bias potential $V(t)$ with zero net bias (rocking ratchet). We study adiabatic driving (such that the system can adjust to the instantaneous equilibrium state), assuming that the driving period t_0 is large compared to the relevant time scales for transmission. This is the case in related experiments⁸. The entire Hamiltonian then reads

$$\mathcal{H} = \mathcal{H}_c + \mathcal{H}_V(t) \quad ; \quad \mathcal{H}_V(t) = V(t)g(x, z; V), \quad (2)$$

where $g(x, z; V)$ describes the spatial distribution of the voltage drop and should in principle be obtained self-consistently from the particle density.

We model spin-dependent transport within a scattering approach assuming that inelastic processes take place only in the reservoirs. Then the probability amplitude for an electron to pass through the conductor is given by the scattering matrix $\mathcal{S}_{n\sigma; n'\sigma'}(E, V)$, where n', n denote transverse modes and $\sigma', \sigma = \pm 1$ the spin directions in the incoming and outgoing lead, respectively. Making use of the unitarity of the scattering matrix, $\mathcal{S}\mathcal{S}^\dagger = \mathcal{S}^\dagger\mathcal{S} = \mathbf{1}$, and summing over all open channels in the left (L) and right (R) lead, respectively, we find the relations

$$\sum_{\substack{n, \sigma \in \text{R} \\ n', \sigma' \in \text{R} \cup \text{L}}} |\mathcal{S}_{n, \sigma; n', \sigma'}|^2 = \sum_{n, \sigma \in \text{R}} 1, \quad \sum_{\substack{n, \sigma \in \text{R} \\ n', \sigma' \in \text{R} \cup \text{L}}} \sigma |\mathcal{S}_{n, \sigma; n', \sigma'}|^2 = 0. \quad (3)$$

For the further analysis, we consider an unbiased square wave driving $V(t) = V_0 \text{sign}[\sin(2\pi t/t_0)]$, restricted to the values $\pm V_0$ ($V_0 > 0$); generalizations to, *e.g.*, harmonic driving are straight forward. The ratchet current is then given by the average of the steady-state currents in the two opposite rocking situations, $\langle I(V_0) \rangle = [I(+V_0) + I(-V_0)]/2$, which we compute within the Landauer formalism relating conductance to transmission.

Contrary to charge current, spin current is usually not conserved. Thus it is crucial to fix the measuring point, which we choose to be inside the right lead. Then, in view of Eq. (3), the ratchet charge $\langle I_C \rangle$ and spin $\langle I_S \rangle$ currents can be expressed as

$$\langle I_{C/S}(V_0) \rangle = G_{C/S} \int_{E_C}^{\infty} dE \Delta f(E, V_0) \Delta T_{C/S}(E, V_0). \quad (4)$$

Here, the prefactor $G_{C/S}$ is equal to $e/2h$ for the charge current and $1/8\pi$ for the spin current. E_C denotes




FIG. 1: (*Color online*) Spin-dependent transmissions as a function of the injection energy $E = (kL)^2$ in the presence of Rashba spin-orbit interaction ($k_{\text{SO}}L = 1.5$) for a short periodic chain of five symmetrical potential barriers (see inset, barrier height $U_0 = 22$) and moderate rocking amplitude $V_0 = 2$. The dashed (red) and dotted (blue) lines indicate T_S , Eq. (7), in the two rocking situations. The solid (black) line depicts the ratchet spin transmission, Eq. (5), the sign indicating the flow direction. For reference, the dashed-dotted (green) curve shows T_C , Eq. (6), and the staircase function T_C for a wire without potential barriers and SO interaction.

the energy of the conduction band edge, $\Delta f(E, V_0) = [f(E, E_F + V_0/2) - f(E, E_F - V_0/2)]$ is the difference between the Fermi functions in the leads, and

$$\Delta T_{C/S}(E, V_0) = T_{C/S}(E, +V_0) - T_{C/S}(E, -V_0). \quad (5)$$

With $T_{\sigma, \sigma'} = \sum_{n \in \text{R}, n' \in \text{L}} |\mathcal{S}_{n, \sigma; n', \sigma'}|^2$, the transmission probabilities for charge and spin in (5) are defined as

$$T_C(E, V) = \sum_{\substack{\sigma' = \pm 1 \in \text{L} \\ \sigma = \pm 1 \in \text{R}}} T_{\sigma, \sigma'}(E, V), \quad (6)$$

$$T_S(E, V) = \sum_{\sigma' = \pm 1 \in \text{L}} [T_{+, \sigma'}(E, V) - T_{-, \sigma'}(E, V)]. \quad (7)$$

The latter is given by the difference between the transmission of spin-up and spin-down electrons upon exit, with the spin measured with respect to the z -axis.

Equation (5) indicates that $\Delta T_{C/S}(E, V_0)$, and thereby the average conductance, vanishes in the linear response limit $V_0 \rightarrow 0$. In the following we consider the nonlinear regime and devise a minimum model for a spin ratchet mechanism by assuming identical leads and a spatially symmetric potential $U(x, z)$ in Eq. (1). The total Hamiltonian (2) is then invariant under the symmetry operation $\hat{\mathcal{P}} = \hat{\mathcal{C}}\hat{R}_x\hat{R}_V\hat{\sigma}_z$, where $\hat{\mathcal{C}}$ is the operator of complex conjugation, \hat{R}_x inverts the x -coordinate and \hat{R}_V changes the sign of the applied voltage ($\pm V \leftrightarrow \mp V$). The action of $\hat{\mathcal{P}}$ on the scattering states is to switch between the two rocking situations and to exchange the leads, *i.e.*, a mode index n is replaced by its corresponding mode \tilde{n} . Moreover, incoming (outgoing) states are




FIG. 2: (*Color online*) (a) Ratchet spin transmission as a function of the number of barriers N for $k_{\text{SO}}L = 1.5$, $U_0 = 22$, $V_0 = 2$, and energies $E = (kL)^2 = 24$ (black symbols, lower line), 33 (red, middle line) and 35.5 (green, upper green). (b) Ratchet spin conductance $\langle I_S \rangle (e/V_0)$ at zero temperature in units of eG_S as a function of applied voltage V_0 for $N = 20$, $k_{\text{SO}}L = 1.5$, $U_0 = 22$ and $E = 24$ (black solid line), 33 (red dashed line) and 35.5 (green dash-dotted line).

transformed into outgoing (incoming) states with complex conjugated amplitude. It is then straightforward to show that

$$S_{n,\sigma;n',\sigma'}(E, \mp V_0) = \sigma\sigma' S_{\bar{n},\sigma';\bar{n},\sigma}(E, \pm V_0), \quad (8)$$

leading to a vanishing charge current $\langle I_C(V_0) \rangle$ and a simplified expression for the ratchet spin transmission (5):

$$\Delta T_S(E, V_0) = 2 [T_{+,-}(E, +V_0) - T_{-,+}(E, +V_0)]. \quad (9)$$

III. RATCHET MECHANISM: NUMERICAL RESULTS

We illustrate the prediction for a ratchet spin current (Eq. (4) with (9)) by performing numerical calculations for the Hamiltonian (1,2). The amplitudes $\mathcal{S}_{n\sigma';m\sigma}(E, V)$ are obtained by projecting the Green function of the open ratchet system onto an appropriate set of asymptotic spinors defining incoming and outgoing channels. For the efficient calculation of the \mathcal{S} -matrix elements a real-space discretization of the Schrödinger equation combined with a recursive algorithm for the Green functions was implemented for spin-dependent transport^{21,22}.

As a model for a spin ratchet we consider a ballistic two-dimensional quantum wire of width W with Rashba SO strength k_{SO} and a one-dimensional periodic modulation (period L) composed of a set of N symmetric potential barriers $U(x) = U_0[1 - \cos(2\pi x/L)]$. We assume a linear voltage drop¹⁹ across the system, $g(x, z) = 1/2 - x/(NL)$ in Eq. (2). To simplify the assessment of the rich parameter space $(E_F, U(x), V, k_{\text{SO}}, N)$ of the problem (L can be scaled out and W is fixed to $1.5L$) and to analyze the mechanisms for spin currents, we first consider a strip with $N=5$ potential barriers (see inset in Fig. 1) and few open transverse modes. Figure 1 shows the numerically obtained spin transmission probabilities $T_S(E, V)$, Eq. (7), for $k_{\text{SO}}L = 1.5$ in the two rocking situations $\pm V_0$ (dashed and dotted line, respectively). The solid line represents the resulting ratchet spin transmission ΔT_S , Eq. (5). For comparison, the dashed-dotted curve shows $T_C(+V_0) = T_C(-V_0)$, Eq. (6), and the staircase function the successive opening of transverse modes

$n = 1, 2, 3$ in the overall transmission of the conductor without potential barriers and SO interaction.

At energies below U_0 and within the first conducting transverse mode the spin transmissions $T_S(\pm V_0)$ are zero, while the total transmission $T_C(\pm V_0)$ is suppressed up to a sequence of four peaks representing resonant tunneling through states which can be viewed as precursors of the lowest Bloch band in the limit of an infinite periodic potential. When the second mode is opened spin polarization is possible (see model below) and takes different values in the two rocking situations leading to a finite ratchet spin transmission. Two transmission peak sequences, related to the lowest one, reappear at higher energies (around $E=24$ and 45), both for $T_C(\pm V_0)$ and for $T_S(\pm V_0)$, owing to corresponding resonant Bloch states involving the second and third transverse mode. The enhanced ratchet spin transmission at the opening of the third mode (at $E = 38$) can be associated to a 'classical' rectification effect resulting from a different number of open modes in one lead in the two rocking situations.

Figure 1 demonstrates moreover that the associated spin current changes sign several times upon variation of the energy, opening up the experimental possibility to control the spin current direction through the carrier density via an external gate. This energy dependence




FIG. 3: (*Color online*) Ratchet spin transmission as a function of energy $E = (kL)^2$ and SO interaction $k_{\text{SO}}L$ for $N = 20$, $V_0 = 2$ and $U_0 = 22$. The dashed lines are a guide to the eye for the shift of the first Bloch band.

of the spin current implies also current inversion as a function of temperature²¹. Such behavior is considered as typical for quantum (particle) ratchets^{4,8}.

In Fig. 2(a) we present the ratchet spin transmission ΔT_S as a function of the barrier number N . Obviously, ΔT_S approaches different asymptotic values depending on the Fermi energy: For energies in resonance with the first Bloch band (lowest trace), ΔT_S exhibits a long- and a short-scale frequency oscillation owing to commensurability between the spin precession length $L_{SO} = \pi/k_{SO}$ and the geometry of the periodic system. For off-resonant injection energies two characteristic, distinct behaviors are shown: a large-scale oscillation (upper curve) and a nearly constant behavior (middle trace), respectively. It is remarkable that in all cases the periodic structure enhances considerably the absolute value of ΔT_S .

In Fig. 2(b) we show the ratchet spin conductance, $\langle I_S \rangle (e/V_0)$, as a function of the applied driving voltage for a system with 20 barriers. For energies within the first Bloch band (solid line), the ratchet spin conductance exhibits a non-monotonic behavior. For the off-resonant cases (dashed and dashed-dotted line) it is monotonically increasing in the voltage window considered.

In Fig. 3 we present the ratchet spin transmission as a function of injection energy E and Rashba SO interaction k_{SO} . We find a rich structure in the explored parameter space, where both large positive and negative values of the ratchet spin transmission can be observed. In the whole energy range peaks due to resonant tunneling are visible, which are shifted to lower energies for increasing SO coupling (*e.g.*, region between dashed lines). Furthermore, we observe discontinuities in the spin transmission at energies where an additional transversal mode in one of the leads opens up (marked by arrows).

For InAs quantum wells L_{SO} is of the order of $0.2 \mu\text{m}$ ²³, in InGaAs it has been tuned from 0.7 to $1.6 \mu\text{m}$ ²⁴ and in GaAs from 2.3 to $5.6 \mu\text{m}$ ²⁵; the range of SO coupling $k_{SO}L = \pi L/L_{SO}$ given in Fig. 3 can be achieved in experiments for period L on scales of μm . Spin-polarized currents as predicted here exceed those observed with experimental detection schemes, reported, *e.g.*, in Ref. 17.

IV. RATCHET MECHANISM: SIMPLIFIED MODEL

Finally we present a simplified model providing additional insight into the underlying mechanism for the occurrence of a finite ratchet spin current. We consider a wire with two open transverse modes ($n = 1, 2$) and a smooth symmetric potential barrier $U(x)$ in the two rocking situations, see Fig. 4. Upon adiabatically traversing the barrier from A via B to C, the spin-orbit split energy spectrum $E_n(k_x)$ for electrons is shifted up and down. For fixed Fermi energy E_F , the initial shift causes a depopulation of the upper levels ($n=2$) and a spin-dependent repopulation while moving from B to C. When E_F is traversing an anti-crossing between succes-




FIG. 4: (*Color online*) Illustration of the spin polarization mechanism for transmission through a strip with a single adiabatic symmetric potential barrier $U(x)$ (solid line) in the two rocking situations (dashed and dotted line). At points A, B and C the position-dependent energy dispersion relation $E_n(k_x)$ is sketched with respect to the Fermi energy E_F (horizontal line) for two transverse modes and SO-induced spin splitting of each mode.

sive modes (see the region indicated by the dashed window in Fig. 4), there is a certain probability P for the electrons to change their spin state. This causes an asymmetry between spin-up and -down states for the repopulated levels²⁰. The related transition probability can be computed in a Landau-Zener picture and reads, for a transverse parabolic confinement of frequency ω_0 ,

$$P(\pm V_0) = 1 - \exp \left\{ \frac{-\pi k_{SO} \omega_0 / \Sigma_z}{(\partial/\partial x)[U(x, z) \pm V_0 g(x, z)]} \right\}. \quad (10)$$

Here Σ_z denotes the difference in the polarizations of the two modes involved. The spin transmission is proportional to $P(V)$ and thus different in the two rocking situations. Hence, the ratchet spin current $\langle I_S(V_0) \rangle$ is nonzero, even in the case of a symmetric barrier. Expanding Eq. (10) for small V_0 allows to qualitatively understand the linear dependence of the ratchet spin conductance for small V_0 in Fig. 2. However, a quantitative explanation of the spin ratchet effect for a periodic, non-necessarily adiabatic potential is beyond this model.

V. CONCLUSIONS

The overall analysis indicates that the ratchet setup, carrying features of a spin rectifier, differs from the proposals^{14,15,17} for spin pumps, since it operates with a single driving parameter, invokes quantum tunneling effects, and the spin transmission is governed by the spatial periodicity of the underlying potential. Further calculations²¹ for combined Rashba- and Dresselhaus²⁶ SO coupling do not alter the overall picture but show that the spin current direction can be changed upon tuning the relative strength of the two coupling mechanisms.

To summarize, we showed that ratchets built from mesoscopic conductors with SO interaction generate spin currents in an experimentally accessible parame-

ter regime. Many further interesting questions open up within this new concept, including the exploration of spin ratchet effects for non-adiabatic driving and for dissipative and non-equilibrium particle and spin dynamics.

Acknowledgements: We thank P. Hänggi, M. Grifoni and M. Strehl for useful discussions and acknowledge support from the German Science Foundation (DFG) within SFB 689.

-
- * Electronic address: dario.bercioux@physik.uni-freiburg.de
- ¹ For recent overviews see the special issue in Appl. Phys. A **75** (2002) and P. Reimann, Phys. Rep. **361**, 57 (2002).
 - ² P. Hänggi and R. Bartussek in: *Lecture Notes in Physics* **476**, edited by J. Parisi, S. C. Müller, and W. W. Zimmermann (Springer, Berlin, 1996), p. 294.
 - ³ F. Jülicher, A. Adjari, and J. Prost, Rev. Mod. Phys. **69**, 1269 (1997); R. D. Astumian, Science **276**, 917 (1997).
 - ⁴ P. Reimann, M. Grifoni, and P. Hänggi, Phys. Rev. Lett. **79**, 10 (1997).
 - ⁵ J. Lehmann, S. Kohler, P. Hänggi, and A. Nitzan, Phys. Rev. Lett. **88**, 228305 (2002).
 - ⁶ E. Lundh and M. Wallin, Phys. Rev. Lett. **94**, 110603 (2005).
 - ⁷ S. Flach, O. Yevtushenko, and Y. Zolotaryuk, Phys. Rev. Lett. **84**, 2358 (2000); H. Schanz, M.-F. Otto, R. Ketzmerick, and T. Dittrich, *ibid.* **87**, 60701 (2001).
 - ⁸ H. Linke, T. E. Humphrey, A. Löfgren, A. O. Sushkov, R. Newbury, R. P. Taylor, and P. Omling, Science **286**, 2314 (1999).
 - ⁹ E. M. Höhberger, A. Lorke, W. Wegscheider, and M. Bichler, Appl. Phys. Lett. **78**, 2905 (2001).
 - ¹⁰ For recent proposals of spin ratchets based on the Zeeman effect see: M. Scheid, M. Wimmer, D. Bercioux, and K. Richter, phys. stat. sol. (c) **3**, 4235 (2006); M. Scheid, D. Bercioux, and K. Richter, arXiv:0707.2478v1 [cond-mat.mes-hall].
 - ¹¹ E. Rashba, Fiz. Tverd. Tela (Leningrad) **2**, 1224 (1960) [Sov. Phys. Solid State **2**, 1109 (1960)].
 - ¹² I. Žutić, J. Fabian, and S. Das Sarma, Rev. Mod. Phys. **76**, 323 (2004).
 - ¹³ G. Schmidt, D. Ferrand, L.W. Molenkamp, A.T. Filip, and B.J. van Wees Phys. Rev. B **62**, R4790 (2000).
 - ¹⁴ P. Sharma and P. W. Brouwer, Phys. Rev. Lett. **91**, 166801 (2003); M. Governale, F. Taddei, and R. Fazio, Phys. Rev. B **68**, 155324 (2003).
 - ¹⁵ E. R. Mucciolo, C. Chamon, and C. M. Marcus, Phys. Rev. Lett. **89**, 146802 (2002); M. Blaauboer and C. M. L. Fricot, Phys. Rev. B **71**, 041303(R) (2005).
 - ¹⁶ Corresponding settings, where a precessing magnetization in a ferromagnet emits spin currents, were reviewed by Y. Tserkovnyak, A. Brataas, G. E. W. Bauer, and B. I. Halperin, Rev. Mod. Phys. **77**, 1375 (2005).
 - ¹⁷ S. K. Watson, R.M. Potok, C.M. Marcus, and V. Umansky, Phys. Rev. Lett. **91**, 258301 (2003).
 - ¹⁸ F. Mireles and G. Kirczenow, Phys. Rev. B **64**, 24426 (2001); M. Governale and U. Zülicke, *ibid.* **66**, 073311 (2002); C.A. Perroni, D. Bercioux, V. M. Ramaglia, and V. Cataudella, J. Phys.: Condens. Matter **19**, 186227 (2007).
 - ¹⁹ An analysis²¹ of different models for the voltage drop shows that the results for the spin current, up to slight quantitative changes, are not altered qualitatively.
 - ²⁰ M. Eto, T. Hayashi, and Y. Kurotani, J. Phys. Soc. Jpn. **74**, 1934 (2005).
 - ²¹ D. Bercioux, M. Scheid, A. Pfund, and K. Richter, unpublished.
 - ²² D. Frustaglia, M. Hentschel, and K. Richter, Phys. Rev. Lett. **87**, 256602 (2001); Phys. Rev. B **69**, 155327 (2004).
 - ²³ D. Grundler, Phys. Rev. Lett. **84**, 6074 (2000);
 - ²⁴ T. Bergsten, T. Kobayashi, Y. Sekine, and J. Nitta, Phys. Rev. Lett. **97**, 196803 (2006).
 - ²⁵ D.M. Zumbühl, J. B. Miller, C.M. Marcus, K. Campman, and A. C. Gossard, Phys. Rev. Lett. **89**, 276803 (2002).
 - ²⁶ G. Dresselhaus, Phys. Rev. **100**, 580 (1955).