

Luminescence Characterizations of Cyclometalated Rhenium(I) Carbonyl Complexes

Peter Spellane[†] and Richard J. Watts*

Department of Chemistry, University of California,
Santa Barbara, California 93106

Arnd Vogler


Institut für Anorganische Chemie, Universität Regensburg,
Postfach 397, Universitätsstrasse 31,
8400 Regensburg, Germany

Received March 25, 1993

Introduction

The high level of current interest in the photochemistry of tricarbonylrhenium(I) complexes of the type $\text{Re}(\text{CO})_3(\text{LL}')\text{X}$ (LL' is a bidentate N,N' -chelating ligand such as 1,10-phenanthroline or 2,2'-bipyridine, and X is a monodentate anionic ligand such as Cl^- or Br^-) stems from reports of the activity of these species in a variety of processes including photochemical reduction of CO_2 ,¹⁻⁵ chemiluminescence,^{6,7} electrochemiluminescence,⁸⁻¹⁰ exciplex emission,¹¹ and as luminescent probes in curing of epoxy resins.^{12,13} The luminescent excited state of this class of complexes, which is generally the excited state responsible for photochemical activity, is often metal-to-ligand charge transfer (MLCT or $d-\pi^*$) with substantial triplet character.¹⁴ However, for a set of complexes $\text{Re}(\text{CO})_3(\text{L})_2\text{X}$ ($\text{L} = 4$ -phenylpyridine, 4,4'-bipyridine; $\text{X} = \text{Cl}^-$, Br^- , I^-), luminescence derives from either a ligand-centered (LC or $\pi-\pi^*$) state or the MLCT state, depending upon the identity of X .¹⁵ A variety of related Re(I) carbonyl complexes have been found to display multiple emissions due to the presence of close-lying MLCT and LC states.¹⁶⁻²⁰

Neutral tetracarbonylrhenium(I) complexes $\text{XRe}(\text{CO})_4\text{L}$ ($\text{X} = \text{Cl}$, I ; $\text{L} = \text{CO}$, piperidine, triphenylphosphine) were reported to display metal-centered (MC or dd) emission properties,^{21,22} and quenching of emission due to low-lying charge-transfer states



$\text{L} = \text{CO}, \text{PPh}_3$

Figure 1. Structural representation of cyclometalated benzo[*h*]quinolin-10-ylrhenium(I) carbonyl complexes.

between donor and acceptor ligand sites (ligand-to-ligand charge transfer or LLCT) has been reported in several instances.²³⁻²⁷ Studies of the photoproperties of neutral as well as cationic Re(I) complexes have greatly expanded since the first work in this area. These studies encompass a variety of diverse topics including monochromatic and dichromatic photolysis,^{28,29} photoelectrochemistry,^{30,31} and surfactant photochemistry.^{32,33}

A few photochemical studies of cationic tetracarbonylrhenium(I) complexes have been reported, including those of *cis*- $[\text{Re}(\text{CO})_4(\text{phen})]^+$,³⁴ *cis*- $[\text{Re}(\text{CO})_4(\text{diphos})]^+$,³⁵ and *trans*- $[\text{Re}(\text{CO})_4(\text{PPh}_3)_2]^+$.³⁶ Recent reports indicate that the cationic tetracarbonyl species $[\text{Re}(\text{CO})_4(\text{bpm})]^+$ ($\text{bpm} = 2,2'$ -bipyrimidine) has a substantially longer lifetime and higher emission energy than many related tricarbonyl species³⁷ and that several species of the type $[\text{Re}(\text{CO})_4(\text{LL}')]^+$ ($\text{LL}' = \text{phen}$, bpy , bpm , dmb (4,4'-dimethyl-2,2'-bipyridine)) are strong photooxidants.³⁸

We report here results of a study of related neutral tetracarbonyl and tricarbonyl complexes $\text{Re}(\text{CO})_4(\text{bzq})$ ($\text{bzq} = \text{benzo}[h]\text{-quinolin-10-yl anion}$) and $\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$ ($\text{PPh}_3 = \text{triphenylphosphine}$) (Figure 1). Although the tetracarbonyl complex was first prepared some 20 years ago,³⁹ the present work is the first report of the emission properties of orthometalated Re(I) complexes.

Experimental Section

Synthetic Procedures. $\text{Re}(\text{CO})_4(\text{bzq})$. Two methods of preparation were employed; the first of these was a modification of the method of Bruce *et al.*³⁹ $\text{Re}(\text{CO})_5\text{Cl}$ (Pressure Chemical Co., Pittsburgh, PA) and benzo[*h*]quinoline (20% molar excess, Fluka) were combined in freshly distilled toluene and refluxed for several hours. The product was extracted into warm hexanes from the solid mixture which remained after toluene had been evaporated, and this product was recrystallized several times

[†] Current address: Akzo Research Laboratory Dobbs Ferry, 1 Livingston Ave., Dobbs Ferry, NY 10522.

- (1) Hawecker, J.; Lehn, J.-M.; Ziessel, R. *J. Chem. Soc., Chem. Commun.* **1983**, 536-538.
- (2) Hawecker, J.; Lehn, J.-M.; Ziessel, R. *J. Chem. Soc., Chem. Commun.* **1984**, 328.
- (3) Kutal, C.; Weber, M. A.; Ferraudi, G.; Geiger, D. *Organometallics* **1985**, *4*, 2161-2166.
- (4) Kutal, C.; Corbin, A. J.; Ferraudi, G. *Organometallics* **1987**, *6*, 553-557.
- (5) Ziessel, R. *NATO ASI Ser., Ser. C* **1987**, *206*, 113-138 (Carbon Dioxide Source Carbon: Biochem. Chem. Uses).
- (6) Vogler, A.; Kunkely, H. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 469.
- (7) Vogler, A.; El-Sayed, L.; Jones, R. G.; Namnath, J.; Adamson, A. W. *Inorg. Chim. Acta* **1981**, *53*, L35-L37.
- (8) Luong, J. C.; Nadjo, L.; Wrighton, M. S. *J. Am. Chem. Soc.* **1978**, *100*, 5790-5795.
- (9) Vogler, A.; Kunkely, H. *ACS Symp. Ser.* **1987**, *No. 333*, 155-168 (High Energy Processes Organomet. Chem.).
- (10) Kunkely, H.; Merz, A.; Vogler, A. *J. Am. Chem. Soc.* **1983**, *105*, 7241.
- (11) Vogler, A.; Kunkely, H. *Inorg. Chim. Acta* **1980**, *45*, L265.
- (12) Kotch, T. G.; Lees, A. J.; Fuerniss, S. J.; Papatthomas, K. I. *Chem. Mater.* **1991**, *3*, 25-27.
- (13) Kotch, T. G.; Lees, A. J.; Fuerniss, S. J.; Papatthomas, K. I. *Chem. Mater.* **1992**, *4*, 675-683.
- (14) Wrighton, M.; Morse, D. L. *J. Am. Chem. Soc.* **1974**, *96*, 998-1003.
- (15) Giordano, P. J.; Wrighton, M. *J. Am. Chem. Soc.* **1979**, *101*, 2888-2897.
- (16) Fredericks, S. M.; Luong, J. C.; Wrighton, M. S. *J. Am. Chem. Soc.* **1979**, *101*, 7415-7417.
- (17) Shaw, J. R.; Schmehl, R. H. *J. Am. Chem. Soc.* **1991**, *113*, 389-394.
- (18) Glezen, M. M.; Lees, A. J. *J. Am. Chem. Soc.* **1988**, *110*, 6243-6244.
- (19) Glezen, M. M.; Lees, A. J. *J. Am. Chem. Soc.* **1989**, *111*, 6602-6610.
- (20) Sacksteder, L.; Zipp, A.; Brown, E. A.; Streich, J.; Demas, J. N. *Inorg. Chem.* **1990**, *29*, 4335.
- (21) Glezen, M. M.; Lees, A. J. *J. Chem. Soc., Chem. Commun.* **1987**, 1752.
- (22) Glezen, M. M.; Lees, A. J. *J. Am. Chem. Soc.* **1988**, *110*, 3892-3897.

- (23) Westmoreland, T. D.; Schanze, K. S.; Neveux, P. E.; Danielson, E.; Sullivan, B. P.; Chen, P.; Meyer, T. *J. Inorg. Chem.* **1985**, *24*, 2596-2597.
- (24) Schanze, K. S.; Cabana, L. S. *J. Phys. Chem.* **1990**, *94*, 2740.
- (25) Perkins, T. A.; Humer, W.; Netzel, T. L.; Schanze, K. S. *J. Phys. Chem.* **1990**, *94*, 2229.
- (26) MacQueen, D. B.; Schanze, K. S. *J. Am. Chem. Soc.* **1991**, *113*, 6108-6110.
- (27) MacQueen, D. B.; Schanze, K. S. *J. Am. Chem. Soc.* **1991**, *113*, 7470-7479.
- (28) Feliz, M.; Ferraudi, G.; Altmiller, H. *J. Phys. Chem.* **1992**, *96*, 257-264.
- (29) Feliz, M.; Ferraudi, G. *J. Phys. Chem.* **1992**, *96*, 3059-3062.
- (30) Shu, C.-F.; Wrighton, M. S. *Inorg. Chem.* **1988**, *27*, 4326-4329.
- (31) Juris, A.; Campagna, S.; Bidd, I.; Lehn, J.-M.; Ziessel, R. *Inorg. Chem.* **1988**, *27*, 4007-4011.
- (32) Reitz, G. A.; Dressick, W. J.; Demas, J. N. *J. Am. Chem. Soc.* **1986**, *108*, 5344-5345.
- (33) Reitz, G. A.; Demas, J. N.; DeGraff, B. A.; Stephens, E. M. *J. Am. Chem. Soc.* **1988**, *110*, 5051-5059.
- (34) Angelici, R. J.; Brink, R. W. *Inorg. Chem.* **1973**, *12*, 1067.
- (35) Anglin, J. R.; Graham, W. A. G. *J. Am. Chem. Soc.* **1978**, *100*, 5790.
- (36) Kruck, R.; Hofler, M. *Chem. Ber.* **1963**, *96*, 3035.
- (37) Shaver, R. J.; Rillema, D. P.; Woods, C. *J. Chem. Soc., Chem. Commun.* **1990**, 179.
- (38) Shaver, R. J.; Rillema, D. P. *Inorg. Chem.* **1992**, *31*, 4101.
- (39) Bruce, M. I.; Goodall, B. L.; Stone, F. G. A. *J. Organomet. Chem.* **1973**, *60*, 343.

from hexanes to remove the unreacted benzo[*h*]quinoline and yield the light yellow product. An alternative preparation involved the photochemical oxidative addition of benzo[*h*]quinoline to $\text{Re}_2(\text{CO})_{10}$ (Alfa) in toluene. The reagents were combined in freshly distilled toluene under N_2 and stirred under Pyrex-filtered irradiation from a 100-W medium-pressure Hg arc lamp. Heat from the lamp was sufficient to cause the reaction solution to reflux during the 24-h reaction period. After completion of the reaction the toluene was removed by flash evaporation, and the product mixture was dissolved in boiling hexanes. The crystalline product precipitated from this solution upon cooling. Anal. Calcd for $\text{ReC}_{17}\text{H}_8\text{NO}_4$: C, 42.85; H, 1.69; N, 2.94. Found: C, 43.16; H, 1.57; N, 3.06.

$\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$. $\text{Re}(\text{CO})_4(\text{bzq})$ was combined with excess triphenylphosphine (PPh_3) in tetrahydrofuran (freshly distilled from Na/benzophenone). This mixture was irradiated with the Pyrex glass filtered output of a 100-W medium-pressure Hg arc lamp for 21 h. During this period, the mixture refluxed due to heat from the lamp. The mixture was sampled by thin-layer chromatography before, and periodically during, the irradiation period. A new product could be detected after 3 h of irradiation, and the irradiation period was continued until no further increase in the product could be detected by TLC. After completion of the reaction, the THF solvent was flash evaporated, and the orange/yellow liquid was dissolved in CH_2Cl_2 . This solution was spotted in a line on a preparative silica chromatography plate, and the product was eluted with hexanes/ CH_2Cl_2 (3:2 by volume). Several bands could be seen to be resolved due to their luminescence under ultraviolet irradiation. The unreacted starting material moved to the top of the plate, while the primary product could be seen as a yellow band with an R_f value of 0.37. Other luminescent products were evident under ultraviolet irradiation with R_f values of 0.23 (orange), 0.14 (orange), 0.05 (blue), and 0 (orange). The band with $R_f = 0.37$ was scraped from the plate, and the product was removed from the silica by washing with CH_2Cl_2 and filtration. Evaporation of the resulting CH_2Cl_2 solvent under a stream of dry nitrogen gas yielded bright yellow crystals of the product. Anal. Calcd for $\text{ReC}_{34}\text{H}_{23}\text{NO}_3\text{P}$: C, 57.38; H, 3.26; N, 1.97. Found: C, 57.02; H, 3.33; N, 1.94.

Measurements. ^1H and ^{13}C NMR measurements were performed on samples dissolved in CD_2Cl_2 with a Nicolet NT-300 spectrometer operated at 300 MHz for protons and at 75 Mz for ^{13}C . Low-temperature luminescence spectra and luminescence lifetimes were determined with instrumentation described in prior publications.⁴⁰ For these measurements, samples of the complexes were dissolved in a mixture of ethanol and methanol (4:1 by volume) and cooled to 77 K in liquid nitrogen. Room-temperature emission measurements were performed using deoxygenated benzene solutions of the complexes. In order to minimize exposure of the photoreactive samples to light, emission was monitored with a 512-element Oriol Instaspec diode array after passage of the emitted light through an Oriol $1/8$ -m Multispec monochromator. Excitation for these measurements was at 337 nm with the pulsed output of a Laser Photonics UV-22 nitrogen laser. Complete spectral analysis was accomplished by averaging of exposures of the diode array during approximately 100 laser pulses. Mass spectral analysis was performed with a ZAB 2F double-focusing reverse-geometry mass spectrometer.

Results

$\text{Re}(\text{CO})_4(\text{bzq})$. The low-resolution positive ion chemical ionization mass spectrum of $\text{Re}(\text{CO})_4(\text{bzq})$ includes a parent peak at m/e 477 ($\text{C}_{17}\text{H}_8\text{NO}_4\text{Re}$) and clusters of peaks for species resulting from the loss of one, two, and three CO ligands. Each peak cluster shows the expected rhenium isotopic distribution. Metalation of the benzo[*h*]quinoline ligand is confirmed by ^{13}C NMR spectroscopy. Whereas the uncoordinated ligand shows nine carbon atoms strongly coupled to protons (as indicated in comparison of fully proton-decoupled to off-resonance-decoupled spectra), the rhenium compound shows only eight. The resonance at 162.6 ppm shows no strong coupling to a proton and differs greatly from any resonance of the uncoordinated ligand; this resonance is therefore assigned to the metalated C atom. The carbonyl resonances at 189.2, 192.9, and 193.6 ppm show intensities comparable to those of the protonated carbons. Data were acquired with a 1.5-s delay between scans; the intensity of




Figure 2. Luminescence, absorption, and excitation spectra of $\text{Re}(\text{CO})_4(\text{bzq})$ (a) and $\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$ (b). Key: Luminescence spectra in EtOH/MeOH (4:1 by volume) glass at 77 K (—); luminescence in nitrogen-saturated benzene at 298 K (---); absorption in EtOH/MeOH at 298 K (- - -); excitation monitored at 520 nm in EtOH/MeOH at 77 K (· · ·). (All luminescence spectra have been corrected for the wavelength dependence of the optical train and detector to yield corrected spectra in which the wavelength dependence of the intensity is proportional to the actual number of photons emitted per unit time. Excitation spectra were monitored as the ratio of sample intensity/lamp intensity to give the corrected excitation result.)

the resonances suggests that these carbons have short T_1 relaxation times, possibly due to the electric quadrupole of rhenium.

$\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$. Parent lines at m/e 709–713 were observed in the low-resolution positive ion chemical ionization mass spectrum of the product ($\text{C}_{34}\text{H}_{23}\text{NO}_3\text{PRE}$); additional clusters of lines corresponding to the loss of each of the three carbonyl ligands were also observed.

The entire ^1H NMR spectrum of this compound appears at ca. 0.5 ppm higher field than the spectrum of either uncoordinated benzo[*h*]quinoline or $\text{Re}(\text{CO})_4(\text{bzq})$; however, the bzq protons of the phosphine complex do show a pattern of resonances similar to that of $\text{Re}(\text{CO})_4(\text{bzq})$. The phosphine ligand appears to be less effective at deshielding the bzq protons than is CO, an effect consistent with the lower π -acidity of PPh_3 compared to CO.

The aromatic region of the ^{13}C NMR spectrum of $\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$ is much like that of $\text{Re}(\text{CO})_4(\text{bzq})$, with added resonances arising from the carbon atoms of triphenylphosphine (identified by their constants for coupling to phosphorous). Eight resonances of bzq carbon atoms bonded to hydrogen are seen in the range 121–153 ppm, and a ninth resonance at 157.2 ppm corresponds to the metalated aromatic carbon (seen at 162.0 ppm in $\text{Re}(\text{CO})_4(\text{bzq})$; vide supra). However, the carbonyl region of the phosphine complex differs greatly from that of the tetracarbonyl: no strong resonances are evident in the room-temperature spectrum. At lower temperature (223 K), broader, low-intensity lines are seen at 171.5, 192.6, and 200.6 ppm.

Luminescence. Emission spectra of $\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$ and $\text{Re}(\text{CO})_4(\text{bzq})$ in EtOH/MeOH glasses (4:1 by volume) at 77 K each consist of a series of three well-resolved peaks and two shoulders in the visible region (Figure 2). Both excitation spectra

(40) Wilde, A. P.; King, K. A.; Watts, R. J. *J. Phys. Chem.* 1991, 95, 629.

Table I. Spectral Features and Luminescence Lifetimes (τ) of Re(I) and Ir(III) Complexes at 77 and 298 K^a

complex	emission (10^3-cm^{-1}), 77 K	excitation (10^3-cm^{-1}), 77 K	τ , 77 K (298 K)
Re(CO) ₄ (bzq)	20.66, 20.20 (sh), 19.16, 17.79, 16.39 (sh)	26.66, 25.45	4.7 ms (6.4 μ s) ^b
Re(CO) ₃ (bzq)(PPh ₃)	20.33, 19.88 (sh), 18.90, 17.45, 16.13 (sh)	25.97, 24.75	2.0 ms (12.9 μ s) ^b
ClRe(CO) ₃ (phen) ^{c,d}	18.94		9.6 μ s
ClRe(CO) ₃ (3-bzp) ₂ ^{e,f}	24.2, 22.6, 21.1, 19.5 ^g		1.4 ms ^h
ClIr(bzq) ₂ (CO) ⁱ	20.66, 19.05, 17.70		0.95 ms
ClIr(bzq) ₂ (PPh ₃) ⁱ	20.08, 18.69, 17.54		0.10 ms

^a 4:1 EtOH/MeOH glass solvent unless stated otherwise. ^b Nitrogen-saturated benzene solvent at 298 K. ^c Phen = 1,10-phenanthroline. ^d Data from ref 14; EPA glass solvent. ^e 3-bzp = 3-benzoylpyridine. ^f Data from ref 41; EPA glass solvent. ^g Long-lived emission component in dual emission measured with a phosphoroscope. ^h Long-lived component of dual emission; short-lived component due to MLCT reported to have 18.0- μ s lifetime. ⁱ Data measured in this laboratory; see ref 44.

monitored at 520 nm under the same conditions show a band with two well-resolved features in the region around 400 nm (Figure 2). The positions of these features are compiled in Table I along with lifetimes for the luminescence decays.

Although no emission could be seen from either complex in air-saturated solutions at room temperature, strong emissions were observed in deoxygenated solutions. The emissions were found to decrease rapidly with photolysis, and the decreases were found to be accompanied by easily measurable changes in the absorption spectra of the photolyzed solutions. Emission spectra of the complexes in deoxygenated benzene solutions are illustrated in Figure 2, and luminescence lifetimes under these conditions are compiled in Table I. Absorption features in room-temperature fluid EtOH/MeOH solutions are quite similar to those found in the excitation spectra at 77 K; however, these bands are slightly red-shifted and broadened as anticipated in the fluid solutions (Figure 2). Previously reported results for several tricarbonyl-rhenium(I) complexes as well as several bzq complexes of Ir(III) are included in Table I for purposes of comparison.

Discussion

Since the initial studies of Wrighton et al.,¹⁴⁻¹⁶ it has been clear that the low-energy excited states which give rise to luminescence in tricarbonylrhenium(I) complexes may be of either LC or MLCT origin. It has been further noted,^{20,38} that structured luminescences are generally indicative of emission from a LC excited state, while a broad, structureless emission generally suggests MLCT luminescence. In addition, low-temperature luminescence lifetimes associated with the structured LC emissions are generally found to be in the millisecond time domain, while the MLCT luminescence lifetimes tend to be in the microsecond region.^{15,20,41} Thus, characteristic MLCT behavior is illustrated in Table I by the single emission peak and microsecond lifetime of ClRe(CO)₃(phen); characteristics of LC behavior are illustrated by structured emissions and millisecond lifetimes for ClRe(CO)₃(3-benzoylpyridine),¹⁴ ClIr(bzq)₂(CO),⁴² and ClIr(bzq)₂(PPh₃).⁴² The luminescence spectra of Re(CO)₄(bzq) and Re(CO)₃(bzq)(PPh₃) are therefore assigned to LC excited states of the metalated bzq ligand. Further support for this conclusion is found in comparison of the emission spectra of Re(CO)₄(bzq) and Re(CO)₃(bzq)(PPh₃) with those of [Rh(bzq)₂Cl]₂⁴² and [Rh(bzq)₂(phen)]⁺.⁴³ These latter two complexes display well-characterized emissions from LC excited states of metalated bzq with lifetimes of 2.7 and 4.25 ms, respectively, at 77 K. The emission spectra of these complexes have origins of $20.8 \times 10^3 \text{ cm}^{-1}$ and $20.7 \times 10^3 \text{ cm}^{-1}$ —nearly identical to that of Re(CO)₄(bzq) ($20.7 \times 10^3 \text{ cm}^{-1}$) and only slightly above the origin of the Re(CO)₃(bzq)(PPh₃) ($20.3 \times 10^3 \text{ cm}^{-1}$). Finally, the vibrational progressions in these complexes are quite similar with a spacing between the first two features of about 1500 cm^{-1} .

While cyclometalating ligands such as benzo[h]quinoline are known to promote low-energy MLCT to other ligands by enrichment of electron density at the metal center through their strong σ -donor ability, these ligands are not nearly so good electron acceptors as N,N'-bidentate coordinating ligands such as bpy or phen.^{43,44} The net result in both Re(CO)₄(bzq) and Re(CO)₃(bzq)(PPh₃) is an LC emitting state, which is probably due in part to the relatively poor electron-accepting ability of metalated bzq and in part to the relatively low energy of the LC state of bzq. It is interesting to note that these two factors are probably both reversed in cationic tetracarbonyl complexes such as [Re(CO)₄(bpy)]⁺ and [Re(CO)₄(phen)]⁺, which are reported to display unstructured emissions due to MLCT excited states.³⁸ Re carbonyl complexes with less strongly electron-accepting N,N'-coordinating ligands such as bpm, on the other hand, do display a structured emission associated with a LC excited state.^{37,38} However, the red-shift in the emission spectrum and shortening of the emission lifetime of Re(CO)₃(bzq)(PPh₃) as compared to Re(CO)₄(bzq) are likely to be due to mixing of the LC state with a MLCT state; a similar trend has been noted⁴⁴ in comparison of the low-temperature luminescence lifetimes and emission spectra (see Table I) of ClIr(bzq)₂(PPh₃) and ClIr(bzq)₂CO. Thus, while the MLCT state is higher in energy than the LC states in rigid glass solutions of these complexes, it remains sufficiently close in energy to influence their luminescence characteristics.

Although these complexes show no emission in air-saturated solutions at room temperature, nitrogen-saturated benzene solutions are found to emit quite strongly with microsecond lifetimes (Table I). This is quite unusual for LC emissions, since most of these are completely quenched in fluid solutions due to their small radiative decay rates. The emission spectra of both complexes are red-shifted in fluid benzene relative to EtOH/MeOH glass, and this effect is somewhat larger in Re(CO)₃(bzq)(PPh₃) than it is in Re(CO)₄(bzq). The rather longer emission lifetime of the former complex in fluid benzene is probably another indication of enhanced mixing of MLCT character into the emissive state of this species. In fact, the relatively strong emissions and microsecond lifetimes for fluid solution luminescence of these species suggest that the MLCT state of each may have moved below the LC state in fluid solutions. Rigidochromic effects, which are well-known for MLCT excited states of other Re(I) complexes,^{14,15} may substantially lower the MLCT energies in fluid solutions. It was found that although these complexes emit rather strongly in freshly prepared nitrogen-saturated solutions, the emissions lose intensity rapidly under illumination and the appearance of a photoproduct is evident in enhanced absorption in the region around 350 nm as well as the appearance of a new shoulder which extends well into the visible region. However, it is not clear whether or not the two complexes yield a common photoproduct, and no attempt has been made to identify the photolysis products at this time.

The room-temperature absorption spectra of both complexes are quite similar to the excitation spectra, although the latter are

(41) Giordano, P. J.; Fredericks, S. M.; Wrighton, M. S.; Morse, D. L. *J. Am. Chem. Soc.* **1978**, *100*, 2257.

(42) Sprouse, S.; King, K. A.; Spellane, P. J.; Watts, R. J. *J. Am. Chem. Soc.* **1984**, *106*, 6647.

(43) Ohsawa, Y.; Sprouse, S.; King, K. A.; DeArmond, M. K.; Hanck, K. W.; Watts, R. J. *J. Phys. Chem.* **1987**, *91*, 1047.

(44) King, K. A. Ph.D. Dissertation, University of California, Santa Barbara, CA, 1986.

more sharply resolved due to the low-temperature rigid glass used for the excitation measurement (Figure 2). Absorption and excitation results show a major band in the region around 400 nm, and this band undergoes a large red-shift in absorption with benzene as a solvent as compared to EtOH/MeOH. This solvent sensitivity indicates that it is due to a spin-allowed MLCT transition.

The temperature dependence noted in the CO region of the ^{13}C NMR of $\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$ invites further investigation. While the infrared spectrum of this compound in CH_2Cl_2 solutions at room temperature shows the presence of carbonyl ligands, the absence of ^{13}C NMR resonances due to CO at room temperature contrasts to the rather intense carbonyl resonances seen in $\text{Re}(\text{CO})_4(\text{bzq})$ under similar conditions. Insertion of CO into metal-carbon bonds is known to be influenced by triphenylphosphine,^{45,46}

and a rapid CO insertion could scramble CO with acyl carbon to influence the ^{13}C NMR spectrum. The ^{13}C chemical shift of the acyl carbon in $\text{C}_6\text{H}_5\text{C}(\text{O})\text{Re}(\text{CO})_5$, for example, is reported to be 245.4 ppm.⁴⁷ Clarification of the origin of the temperature dependence noted in the ^{13}C NMR of $\text{Re}(\text{CO})_3(\text{bzq})(\text{PPh}_3)$ awaits further investigation.

Acknowledgment. This work was supported by the Office of Basic Energy Sciences, United States Department of Energy, Project DE-FG03-88ER13842.

Registry Numbers (supplied by author): $\text{Re}(\text{CO})_4(\text{bzq})$, 51466-82-1; $\text{Re}(\text{CO})_3(\text{Bzq})(\text{PPh}_3)$, not yet assigned.

(45) Cotton, J. D.; Markwell, R. D. *Inorg. Chim. Acta* **1982**, *63*, 13.

(46) Anderson, G. K.; Cross, R. J. *J. Chem. Soc., Dalton Trans.* **1980**, 712.

(47) Webb, M. J.; Graham, W. A. G. *J. Organomet. Chem.* **1975**, *93*, 119.