

Dirac eigenspinors for generic metrics



DISSERTATION ZUR ERLANGUNG DES DOKTORGRADES
DER NATURWISSENSCHAFTEN (DR. RER. NAT.)
AN DER FAKULTÄT FÜR MATHEMATIK DER UNIVERSITÄT REGENSBURG

vorgelegt von
Andreas Hermann
Regensburg, im Januar 2012

Promotionsgesuch eingereicht am 17.1.2012

Die Arbeit wurde angeleitet von Prof. Dr. Bernd Ammann

Prüfungsausschuss:

Vorsitzender:	Prof. Dr. Harald Garcke
1. Gutachter:	Prof. Dr. Bernd Ammann
2. Gutachter:	Prof. Dr. Marc Herzlich, Universität Montpellier
weiterer Prüfer:	Prof. Dr. Roman Sauer
Ersatzprüfer:	Prof. Dr. Felix Finster

Abstract

We consider a Riemannian spin manifold (M, g) with a fixed spin structure. The zero sets of solutions of generalized Dirac equations on M play an important role in some questions arising in conformal spin geometry and in mathematical physics. In this setting the mass endomorphism has been defined as the constant term in an expansion of Green's function for the Dirac operator. One is interested in obtaining metrics, for which it is not zero.

In this thesis we study the dependence of the zero sets of eigenspinors of the Dirac operator on the Riemannian metric. We prove that on closed spin manifolds of dimension 2 or 3 for a generic Riemannian metric the non-harmonic eigenspinors have no zeros. Furthermore we prove that on closed spin manifolds of dimension 3 the mass endomorphism is not zero for a generic Riemannian metric.

Zusammenfassung

Sei (M, g) eine Riemannsche Spin-Mannigfaltigkeit mit einer fixierten Spin-Struktur. In manchen Fragen aus der konformen Spin-Geometrie oder der mathematischen Physik spielen Nullstellenmengen von Lösungen verallgemeinerter Dirac-Gleichungen auf M eine wichtige Rolle. In diesem Zusammenhang wurde der Massen-Endomorphismus als der konstante Term in einer asymptotischen Entwicklung der Greenschen Funktion des Dirac-Operators definiert. Gesucht sind Riemannsche Metriken, für die er nicht Null ist.

In dieser Dissertation untersuchen wir die Abhängigkeit der Nullstellenmenge der Eigenspinoren des Dirac-Operators von der Riemannschen Metrik. Wir beweisen, dass auf einer geschlossenen Spin-Mannigfaltigkeit der Dimension 2 oder 3 für eine generische Riemannsche Metrik die nicht-harmonischen Eigenspinoren keine Nullstellen haben. Weiter zeigen wir, dass auf einer geschlossenen Spin-Mannigfaltigkeit der Dimension 3 für eine generische Riemannsche Metrik der Massen-Endomorphismus nicht Null ist.

Acknowledgements

I am grateful to many people, who have supported me during the preparation of this thesis. First of all I would like to thank my advisor Prof. Bernd Ammann for accepting me as his student and for proposing a very interesting research topic. I have profited very much from his explanations and numerous mathematical discussions with him. He has encouraged me a lot during the time that progress on my work was slow. Many thanks go to my colleagues for creating a nice working atmosphere and especially to Dr. Nadine Große and Dr. Nicolas Ginoux for many interesting discussions.

I am grateful to Dr. Mattias Dahl who invited me to Stockholm and from whom I could learn very much about techniques for demonstrating generic properties. Also I would like to thank Prof. Emmanuel Humbert for an invitation to Nancy.

Thanks are due to the DFG for supporting part of my work by the graduate program “Curvature, Cycles and Cohomology”. I have also profited very much from conferences and seminars within this program.

Finally I would like to thank my family for their emotional support during all the years of my studies.

Contents

1	Overview	7
2	Recapitulation of some facts	11
2.1	Review of spin geometry	11
2.2	Real and quaternionic structures	18
2.3	Spinors for different metrics	22
3	Motivation	31
3.1	Conformal bounds on Dirac eigenvalues	31
3.2	Witten spinors	33
3.3	Zero sets of eigenspinors	35
4	Examples of zero sets of eigenspinors	39
4.1	Harmonic spinors on Riemann surfaces	39
4.2	The sphere with the standard metric	41
4.3	Flat tori	44
5	Green's function for the Dirac operator	47
5.1	Trivialization of the spinor bundle	47
5.2	The Euclidean Dirac operator	50
5.3	Expansion of Green's function	52
5.4	Definition of the mass endomorphism	60
6	Eigenspinors for generic metrics	63
6.1	Transversality	63
6.2	Eigenspinors in dimensions 2 and 3	65
6.3	Mass endomorphism in dimension 3	78
A	Analytic perturbation theory	85

Chapter 1

Overview

Some questions arising in conformal spin geometry and in mathematical physics involve the study of the zero sets of solutions to generalized Dirac equations. For the first example let (M, g) be a compact Riemannian spin manifold with a fixed orientation and a fixed spin structure. One is interested in finding bounds on the eigenvalues of the Dirac operator D^g which are uniform in the conformal class $[g]$ of g . The two conformal invariants

$$\lambda_{\min}^{\pm} := \inf_{h \in [g]} |\lambda_1^{\pm}(h)| \operatorname{vol}(M, h)^{1/n}$$

have been studied by many authors. A natural question is whether the infimum is attained at a Riemannian metric. By a result of B. Ammann this is the case, if the nonlinear partial differential equation

$$D^g \psi = \lambda_{\min}^+ |\psi|_g^{2/(n-1)} \psi, \quad \|\psi\|_{2n/(n-1)} = 1 \quad (1.1)$$

has a solution ψ , which is nowhere zero on M (see [Am4]). It is not obvious that a solution without zeros exists.

A second example comes from general relativity, more precisely from a remarkable proof of the positive energy theorem obtained by E. Witten (see [Wi]). He uses harmonic spinors (i. e. spinors ψ satisfying $D^g \psi = 0$) on asymptotically flat manifolds, which are called Witten spinors. It has been suggested to use these spinors in order to construct special orthonormal frames of the tangent bundle of an asymptotically flat manifold of dimension 3 (see [N], [DM], [FNS]). It turns out that this is possible, if one can find a Witten spinor which is nowhere zero. However it is not clear that such a spinor exists.

In this thesis we first consider the zero sets of eigenspinors of the Dirac operator on closed spin manifolds. It is interesting for several reasons. First of all it is easier than the questions mentioned above, since the underlying

manifold is compact and all the operators involved are linear. Apart from that it is useful to have eigenspinors, which are nowhere zero. For example one obtains in this case a simple proof of Hijazi's inequality.

The spectrum of the Dirac operator has been computed explicitly for some Riemannian spin manifolds. The result on the round sphere (see e. g. [Bä2]) shows that the multiplicity of an eigenvalue can be greater than the rank of the spinor bundle, which implies that there exist eigenspinors with non-empty zero set for special choices of a Riemannian metric. However we can show that the situation is different for generic Riemannian metrics (see Section 6.2 for a precise definition of the term "generic"). More precisely let M be a closed spin manifold and denote by $R(M)$ the set of all smooth Riemannian metrics on M . For every $g \in R(M)$ denote by $[g] \subset R(M)$ the conformal class of g . Furthermore let $N(M)$ be the set of all $g \in R(M)$ such that all the non-harmonic eigenspinors of D^g are nowhere zero on M . Then we prove the following.

Theorem 1.0.1. *Let M be a closed connected spin manifold of dimension 2 or 3 with a fixed orientation and a fixed spin structure. Then the set $N(M) \cap [g]$ is residual in $[g]$ with respect to every C^k -topology, $k \geq 1$.*

Recall that a subset is residual, if it contains a countable intersection of open and dense sets. In Section 4.1 we will give an example showing that in dimension 2 an analogue of this theorem for harmonic spinors does not hold.

The main idea of the proof of Theorem 1.0.1 is as follows. If g, h are two Riemannian metrics on the spin manifold M , then a natural isomorphism between the two vector bundles $\Sigma^g M$ and $\Sigma^h M$ is well known. We construct a continuous map F defined on a suitable space of Riemannian metrics, which associates to every metric h an eigenspinor of the corresponding Dirac operator D^h viewed as a section of $\Sigma^g M$. Theorem 1.0.1 then follows from a transversality theorem. In order to apply this theorem we have to make sure that the evaluation map corresponding to F is transverse to the zero section of $\Sigma^g M$. Our assumption that this is not the case leads to an equation involving Green's function for the operator $D^g - \lambda$ with $\lambda \in \mathbb{R}$. From the expansion of this Green's function we obtain a contradiction using the unique continuation property of the Dirac operator.

In this thesis we also treat a certain aspect of the question on conformal bounds of Dirac eigenvalues mentioned above. The non-linear partial differential equation (1.1) cannot be solved by standard methods, since the corresponding Sobolev embedding is critical. However under some additional assumptions on the Riemannian spin manifold (M, g) the existence of a solution has been shown, if there is a point p on M such that a certain endomorphism of the fibre $\Sigma_p^g M$ of the spinor bundle does not vanish (see

[AHM]). This endomorphism can be regarded as the constant term in an expansion of Green's function for the Dirac operator around p . It is called the mass endomorphism at p because of an analogy in conformal geometry: the constant term of Green's function $\Gamma(\cdot, p)$ for the conformal Laplacian at p is related to the mass of the asymptotically flat Riemannian manifold $(M \setminus \{p\}, \Gamma(\cdot, p)^{4/(n-2)}g)$. Unfortunately the mass endomorphism is known explicitly only for very few spin manifolds. However in dimension 3 we can show that in the generic case it is not zero. More precisely, for a fixed point $p \in M$ we denote by $R_p(M)$ the set of all Riemannian metrics on M , such that the mass endomorphism at p can be defined and we denote by $S_p(M)$ the set of all such Riemannian metrics, for which the mass endomorphism does not vanish. Then we prove the following.

Theorem 1.0.2. *Let M be a closed spin manifold of dimension 3 with a fixed spin structure and let $p \in M$. Then $S_p(M)$ is dense in $R_p(M)$.*

The structure of this thesis is as follows. In Chapter 2 we review the basic definitions and results in spin geometry. We begin with the definition of spin groups and the Dirac operator and then proceed with an overview on real and quaternionic structures on spinor modules. Since for two different Riemannian metrics the spinor bundles are two different vector bundles, we need a natural way of identifying spinors for different metrics. This is known in the literature and is described in Section 2.3. Results from perturbation theory, which apply in our situation, are collected in the Appendix.

In Chapter 3 we explain the questions mentioned above in more detail.

Chapter 4 contains some examples of zero sets of Dirac eigenspinors. This should serve as an illustration. The reader who is only interested in the main results may skip Sections 4.2, 4.3.

In Chapter 5 we introduce Green's function for the Dirac operator, which will be one of our tools in the proofs of our results. We describe a method for the explicit calculation of some terms in the expansion of Green's function around the singularity. After that we give the definition of the mass endomorphism from the literature.

In Chapter 6 we state and prove our main results. First let M be a closed spin manifold of dimension 2 or 3. We prove that for a generic Riemannian metric on M the non-harmonic eigenspinors of the Dirac operator are nowhere zero. The proof is based on a well known transversality theorem, which we state in Section 6.1 including its proof. After that we prove that on every closed spin manifold of dimension 3 for a generic Riemannian metric the mass endomorphism is not zero.

Chapter 2

Recapitulation of some facts

2.1 Review of spin geometry

We review some basic definitions and results in spin geometry in order to fix the notation. Brief and nicely written introductions to spin geometry can be found in [Hij3], [BGM05]. The reader can find a detailed treatment of this subject in the books [LM], [F4]. Our notation will be similar to some of the notation in these texts.

Let V be a real vector space with a scalar product g . Then the real Clifford algebra $\text{Cl}(V, g)$ for (V, g) is the unital \mathbb{R} -algebra generated by V with the relation

$$v \cdot w + w \cdot v = -2g(v, w)1, \quad v, w \in V. \quad (2.1)$$

We denote by $\mathbb{C}\text{Cl}(V, g) := \text{Cl}(V, g) \otimes_{\mathbb{R}} \mathbb{C}$ its complexification and call it the Clifford algebra for (V, g) . In the case $V = \mathbb{R}^n$, $n \in \mathbb{N} \setminus \{0\}$, $g = g_{\text{eucl}}$ we use the notation

$$\text{Cl}(n) := \text{Cl}(\mathbb{R}^n, g_{\text{eucl}}), \quad \mathbb{C}\text{Cl}(n) := \mathbb{C}\text{Cl}(\mathbb{R}^n, g_{\text{eucl}}).$$

Every $v \in \mathbb{R}^n \setminus \{0\}$ is invertible in $\text{Cl}(n)$ and the map

$$\text{Ad}_v : \text{Cl}(n) \rightarrow \text{Cl}(n), \quad w \mapsto v \cdot w \cdot v^{-1}$$

preserves the subspace $\mathbb{R}^n \subset \text{Cl}(n)$. The restriction acts on \mathbb{R}^n as the reflection at the line generated by v and thus is in $O(n)$.

Let $\text{Cl}(n)^*$ be the multiplicative group of invertible elements of $\text{Cl}(n)$. We define the spin group $\text{Spin}(n)$ as the subgroup of $\text{Cl}(n)^*$ generated by elements of the form $v_1 \cdot v_2$, where $v_1, v_2 \in \mathbb{R}^n$, $|v_1| = |v_2| = 1$. We obtain a group homomorphism

$$\vartheta : \text{Spin}(n) \rightarrow \text{SO}(n), \quad q \mapsto \text{Ad}_q$$

which is a two-fold covering. The covering is nontrivial for $n \geq 2$ and it is the universal covering for $n \geq 3$.

If $(E_i)_{i=1}^n$ denotes the standard basis of \mathbb{R}^n , then

$$\omega_{\mathbb{C}} := i^{[(n+1)/2]} E_1 \cdot \dots \cdot E_n \in \mathbb{C}\text{I}(n)$$

is called the complex volume element. Here $[\cdot]$ denotes the integer part. We have $\omega_{\mathbb{C}}^2 = 1$ and

$$\omega_{\mathbb{C}} \cdot v = (-1)^{n-1} v \cdot \omega_{\mathbb{C}}$$

for all $v \in \mathbb{R}^n$.

For even n there is exactly one irreducible complex representation of $\mathbb{C}\text{I}(n)$. The module has complex dimension $2^{n/2}$ and is denoted by Σ_n . It is the direct sum of the eigenspaces Σ_n^{\pm} of $\omega_{\mathbb{C}}$ for the values ± 1 :

$$\Sigma_n = \Sigma_n^+ \oplus \Sigma_n^-.$$

Since $\omega_{\mathbb{C}}$ anticommutes with elements of \mathbb{R}^n , the two eigenspaces have the same dimension.

For odd n the complex volume element commutes with all elements of $\mathbb{C}\text{I}(n)$ and thus by Schur's lemma acts as a multiple of the identity on every irreducible module. There exist exactly two inequivalent irreducible complex representations of $\mathbb{C}\text{I}(n)$, both of dimension $2^{(n-1)/2}$, and they are distinguished by the action of $\omega_{\mathbb{C}}$ as Id or $-\text{Id}$ respectively. In this thesis we will use the representation for which $\omega_{\mathbb{C}}$ acts as Id . The module is again denoted by Σ_n .

Thus for every n we have $\Sigma_n \cong \mathbb{C}^N$, where $N := 2^{\lfloor n/2 \rfloor}$. The representation will be denoted by ρ . The action of $\mathbb{C}\text{I}(n)$ on Σ_n via ρ is called the Clifford multiplication on Σ_n . It will be denoted by $v \cdot \sigma := \rho(v)\sigma$ for $v \in \mathbb{C}\text{I}(n)$, $\sigma \in \Sigma_n$.

On Σ_n there exists a positive definite hermitian inner product $\langle \cdot, \cdot \rangle$, such that Clifford multiplication with all elements of \mathbb{R}^n is antisymmetric with respect to $\langle \cdot, \cdot \rangle$, i. e. such that

$$\langle v \cdot \psi, \varphi \rangle + \langle \psi, v \cdot \varphi \rangle = 0$$

for all $\varphi, \psi \in \Sigma_n$ and all $v \in \mathbb{R}^n$.

Let M be an n -dimensional oriented manifold. We denote by $\text{GL}^+(n, \mathbb{R})$ the group of all real $n \times n$ -matrices with positive determinant and we write $A: \widetilde{\text{GL}}^+(n, \mathbb{R}) \rightarrow \text{GL}^+(n, \mathbb{R})$ for its connected two-fold covering. Let

$$\pi: \text{P}_{\text{GL}^+}(M) \rightarrow M$$

be the principal $\text{GL}^+(n, \mathbb{R})$ -bundle over M whose fibre over $x \in M$ consists of all positively oriented bases of $T_x M$. It is called the bundle of positively oriented frames of the tangent bundle.

Definition 2.1.1. A spin structure on M is a principal $\widetilde{\text{GL}}^+(n, \mathbb{R})$ -bundle

$$\pi' : P_{\widetilde{\text{GL}}^+}(M) \rightarrow M$$

over M together with a two-fold covering $\Theta: P_{\widetilde{\text{GL}}^+}(M) \rightarrow P_{\text{GL}^+}(M)$, such that the following diagram commutes

$$\begin{array}{ccc} P_{\widetilde{\text{GL}}^+}(M) \times \widetilde{\text{GL}}^+(n, \mathbb{R}) & \longrightarrow & P_{\widetilde{\text{GL}}^+}(M) \\ \Theta \times A \downarrow & & \downarrow \Theta \\ P_{\text{GL}^+}(M) \times \text{GL}(n, \mathbb{R}) & \longrightarrow & P_{\text{GL}^+}(M) \end{array} \quad \begin{array}{c} \nearrow \pi' \\ \\ \searrow \pi \end{array} \quad \begin{array}{c} \\ \\ M \end{array}$$

where the horizontal arrows denote the group actions. M is called a spin manifold, if there exists a spin structure on M .

Not every oriented manifold has a spin structure and some oriented manifolds have more than one spin structure. An oriented manifold M is a spin manifold if and only if the second Stiefel-Whitney class

$$w_2(TM) \in H^2(M, \mathbb{Z}/2\mathbb{Z})$$

vanishes. If $w_2(TM) = 0$, then the distinct spin structures on M are in one-to-one correspondence with the elements of $H^1(M, \mathbb{Z}/2\mathbb{Z})$. For example every orientable manifold M of dimension $n \leq 3$ is a spin manifold.

For every Riemannian metric g on an oriented spin manifold M we denote by $P_{\text{SO}}(M, g) \subset P_{\text{GL}^+}(M)$ the principal $\text{SO}(n)$ -bundle over M whose fibre over $x \in M$ consists of all positively oriented g -orthonormal bases of $T_x M$. It is called the bundle of positively oriented g -orthonormal frames of the tangent bundle. The restriction of $A: \widetilde{\text{GL}}^+(n, \mathbb{R}) \rightarrow \text{GL}^+(n, \mathbb{R})$ to the preimage of $\text{SO}(n) \subset \text{GL}^+(n, \mathbb{R})$ coincides with $\vartheta: \text{Spin}(n) \rightarrow \text{SO}(n)$. Furthermore

$$P_{\text{Spin}}(M, g) := \Theta^{-1}(P_{\text{SO}}(M, g))$$

is a principal $\text{Spin}(n)$ -bundle over M and the maps in the above commutative diagram restrict to the following commutative diagram

$$\begin{array}{ccc} P_{\text{Spin}}(M, g) \times \text{Spin}(n) & \longrightarrow & P_{\text{Spin}}(M, g) \\ \Theta \times \vartheta \downarrow & & \downarrow \Theta \\ P_{\text{SO}}(M, g) \times \text{SO}(n) & \longrightarrow & P_{\text{SO}}(M, g) \end{array} \quad \begin{array}{c} \nearrow \pi' \\ \\ \searrow \pi \end{array} \quad \begin{array}{c} \\ \\ M \end{array}$$

By a Riemannian spin manifold (M, g, Θ) we will always mean an orientable Riemannian spin manifold (M, g) together with a fixed orientation and a fixed spin structure Θ .

To every principal bundle over M we can associate a vector bundle in the following way. Let G be a Lie group and let $\pi: P \rightarrow M$ be a principal G -bundle over a manifold M . Furthermore let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , let V be a vector space over \mathbb{K} and let $\rho: G \rightarrow \text{Aut}_{\mathbb{K}}(V)$ be a representation of G . On $P \times V$ we define an equivalence relation by

$$(p, v) \sim (p', v') \iff \exists g \in G \text{ such that } p' = pg^{-1} \text{ and } v' = \rho(g)v.$$

We denote by $[p, v]$ the equivalence class of (p, v) and by $P \times_{\rho} V$ the set of all equivalence classes. Then $P \times_{\rho} V$ is a vector bundle over M with fibre V . It is called the associated vector bundle to P via ρ .

As an example let $\tau: \text{SO}(n) \rightarrow \text{Aut}_{\mathbb{R}}(\mathbb{R}^n)$ be the standard representation. Then there exists a canonical isomorphism of vector bundles

$$TM \cong P_{\text{SO}}(M, g) \times_{\tau} \mathbb{R}^n.$$

Now let (M, g, Θ) be an n -dimensional Riemannian spin manifold. The restriction of the complex spinor representation ρ to $\text{Spin}(n) \subset \text{Cl}(n)$ is again denoted by ρ , i. e.

$$\rho: \text{Spin}(n) \rightarrow \text{Aut}_{\mathbb{C}}(\Sigma_n).$$

Definition 2.1.2. *The complex spinor bundle $\Sigma^g M$ over M for the metric g and the spin structure Θ is the associated vector bundle*

$$\Sigma^g M := P_{\text{Spin}}(M, g) \times_{\rho} \Sigma_n.$$

The complex spinor bundle $\Sigma^g M$ is a vector bundle with fibre $\Sigma_n \cong \mathbb{C}^N$. For $(M, g) = (\mathbb{R}^n, g_{\text{eucl}})$ equipped with the unique spin structure we will write

$$\Sigma \mathbb{R}^n := \Sigma^{g_{\text{eucl}}} \mathbb{R}^n.$$

We define the Clifford multiplication on $\Sigma_x^g M$, $x \in M$, by

$$T_x M \otimes \Sigma_x^g M \rightarrow \Sigma_x^g M, \quad [\Theta(s), v] \otimes [s, \sigma] \mapsto [s, v \cdot \sigma].$$

The inner product $\langle \cdot, \cdot \rangle$ on Σ_n yields a hermitian metric on $\Sigma^g M$ which we also denote by $\langle \cdot, \cdot \rangle$. It is defined by

$$\langle [s, \sigma_1], [s, \sigma_2] \rangle := \langle \sigma_1, \sigma_2 \rangle.$$

We will denote the induced norm by

$$|\psi|_g := \langle \psi, \psi \rangle^{1/2}.$$

Clifford multiplication with elements of TM is antisymmetric with respect to $\langle \cdot, \cdot \rangle$, i. e. we have

$$\langle v \cdot \psi, \varphi \rangle + \langle \psi, v \cdot \varphi \rangle = 0$$

for all $v \in T_x M$, $\psi, \varphi \in \Sigma_x^g M$, $x \in M$.

Let $(e_i)_{i=1}^n$ be a positively oriented local orthonormal frame of TM . Then

$$\omega_{\mathbb{C}} := i^{[(n+1)/2]} e_1 \cdot \dots \cdot e_n$$

is independent of the choice of the e_i and thus can be defined on all of M . Fibrewise Clifford multiplication by $\omega_{\mathbb{C}}$ is an endomorphism of $\Sigma^g M$ which is again denoted by $\omega_{\mathbb{C}}$. We find $\omega_{\mathbb{C}}^2 = 1$ and

$$\omega_{\mathbb{C}} \cdot v = (-1)^{n-1} v \cdot \omega_{\mathbb{C}}$$

for all $v \in TM$. If n is odd, we have by convention $\omega_{\mathbb{C}} = \text{Id}$. If n is even, then, since $\omega_{\mathbb{C}}$ commutes with $\text{Spin}(n)$, the above splitting of the spinor module induces a splitting of the spinor bundle

$$\Sigma^g M = \Sigma^+ M \oplus \Sigma^- M.$$

A local section of the spinor bundle $\Sigma^g M$ is called a spinor. If n is even the local sections of $\Sigma^{\pm} M$ are called positive respectively negative spinors. We denote by $C^r(\Sigma^g M)$, $r \in \mathbb{N}$, (resp. $C^\infty(\Sigma^g M)$) the space of all r times continuously differentiable (resp. smooth) spinors.

In order to define a covariant derivative on the spinor bundle we recall the following general fact. Let G be a Lie group, $P \rightarrow M$ a principal G -bundle and let $\rho: G \rightarrow \text{Aut}_{\mathbb{C}}(V)$ be a representation of G on a complex vector space V . Let ω be a connection one-form on P . Every section ψ of the associated vector bundle $P \times_{\rho} V$ is locally given by $\psi = [s, \sigma]$, where s is a locally defined section of P on an open subset $U \subset M$ and σ is a function on U with values in V . Then for $X \in TM|_U$ we define the spinor $\nabla_X \psi$ on U by

$$\nabla_X \psi := [s, X(\sigma) + d\rho(\omega(ds(X)))\sigma]. \quad (2.2)$$

Here and henceforth for any differentiable function f defined on an open subset $U \subset M$ with values in a real or complex vector space and any vector field X on U we denote by $X(f)$ the derivative of f in the direction X . One can show that ∇ is well-defined and yields a covariant derivative on $P \times_{\rho} V$. Furthermore if V carries a G -invariant hermitian scalar product, then one obtains a hermitian metric on $P \times_{\rho} V$ and ∇ is compatible with this metric.

The Levi Civita connection ∇^g for the Riemannian metric g on M induces a covariant derivative on $\Sigma^g M$ also denoted by ∇^g as follows. The connection

one-form of the Levi Civita connection lifts to a connection one-form ω on $P_{\text{Spin}}(M, g)$. Then we apply the formula (2.2) to ω . The result may locally be written as follows. Let $(e_i)_{i=1}^n$ be a positively oriented local orthonormal frame of TM on an open subset $U \subset M$. There exists a locally defined section $s \in \Gamma(U, P_{\text{Spin}}(M, g)|_U)$ such that $(e_i)_{i=1}^n = \Theta \circ s$ on U . Let $(E_i)_{i=1}^n$ be the standard basis of \mathbb{C}^n . The section s determines a local orthonormal frame $(\psi_i)_{i=1}^n$ of $\Sigma^g M|_U$ via $\psi_i = [s, E_i]$ for $1 \leq i \leq n$. We denote by ∂ the locally defined flat connection with respect to the local frame $(\psi_i)_{i=1}^n$, i. e. for $h_1, \dots, h_n \in C^\infty(U, \mathbb{C})$ and $X \in TM|_U$ we define

$$\partial_X \left(\sum_{i=1}^n h_i \psi_i \right) := \sum_{i=1}^n X(h_i) \psi_i.$$

We define the symbols $\tilde{\Gamma}_{ij}^k$ by the equation

$$\tilde{\Gamma}_{ij}^k := g(\nabla_{e_i}^g e_j, e_k)$$

and write locally $\psi = \sum_{i=1}^n h_i \psi_i$. Then by (2.2) for all $i \in \{1, \dots, n\}$ we have on U

$$\nabla_{e_i}^g \psi = \partial_{e_i} \psi + \frac{1}{4} \sum_{j,k=1}^n \tilde{\Gamma}_{ij}^k e_j \cdot e_k \cdot \psi = \partial_{e_i} \psi + \frac{1}{4} \sum_{j=1}^n e_j \cdot (\nabla_{e_i}^g e_j) \cdot \psi \quad (2.3)$$

(see [LM], p. 103, 110). One finds that ∇^g is a metric connection with respect to $\langle \cdot, \cdot \rangle$ and that it satisfies

$$\nabla_X^g (Y \cdot \psi) = (\nabla_X^g Y) \cdot \psi + Y \cdot \nabla_X^g \psi$$

for all $X \in TM|_U$, $Y \in C^\infty(TM|_U)$, $\psi \in C^\infty(\Sigma^g M|_U)$.

Let $(e_i)_{i=1}^n$ be a local orthonormal frame of TM . The Dirac operator is defined as

$$D^g : C^\infty(\Sigma^g M) \rightarrow C^\infty(\Sigma^g M), \quad D^g \psi := \sum_{i=1}^n e_i \cdot \nabla_{e_i}^g \psi.$$

It is easily seen that the definition does not depend on the choice of the local frame $(e_i)_{i=1}^n$. If n is even, then with respect to the above splitting of the spinor bundle the Dirac operator has the form

$$D^g = \begin{pmatrix} 0 & D^- \\ D^+ & 0 \end{pmatrix} \quad (2.4)$$

with $D^\pm : C^\infty(\Sigma^\pm M) \rightarrow C^\infty(\Sigma^\mp M)$.

If X, Y are smooth vector fields on M we define the second covariant derivative operator $\nabla_{X,Y}^2: C^\infty(\Sigma^g M) \rightarrow C^\infty(\Sigma^g M)$ by

$$\nabla_{X,Y}^2 \psi := \nabla_X^g \nabla_Y^g \psi - \nabla_{\nabla_X^g Y}^g \psi$$

and the connection Laplacian $\nabla^* \nabla: C^\infty(\Sigma^g M) \rightarrow C^\infty(\Sigma^g M)$ by

$$\nabla^* \nabla \psi := -\text{tr}(\nabla_{\dots}^2 \psi),$$

i. e. if $(e_i)_{i=1}^n$ is a local orthonormal frame on M we have

$$\nabla^* \nabla \psi = - \sum_{i=1}^n \nabla_{e_i}^g \nabla_{e_i}^g \psi + \sum_{i=1}^n \nabla_{\nabla_{e_i}^g e_i}^g \psi. \quad (2.5)$$

A very important result is the Schrödinger-Lichnerowicz formula

Theorem 2.1.3. *For all $\psi \in C^\infty(\Sigma^g M)$ the formula*

$$(D^g)^2 \psi = \nabla^* \nabla \psi + \frac{\text{scal}^g}{4} \psi. \quad (2.6)$$

holds, where scal^g is the scalar curvature of (M, g) .

Proof. see [LM] p. 160. □

Let $\psi \in C^\infty(\Sigma^g M)$ and $f \in C^\infty(M, \mathbb{C})$. Then the following Leibniz rule holds on M :

$$D^g(f\psi) = \text{grad}^g(f) \cdot \psi + f D^g \psi,$$

where $\text{grad}^g(f)$ is the gradient of f with respect to g . For a proof see [LM], p. 116.

Let $\mu \in \mathbb{C}$. A spinor $\psi \in C^\infty(\Sigma^g M)$ is called a Killing spinor for μ , if we have

$$\nabla_X^g \psi = \mu X \cdot \psi$$

for all $X \in TM$.

A spinor $\psi \in C^\infty(\Sigma^g M)$ is called parallel, if we have $\nabla_X^g \psi = 0$ for all $X \in TM$.

Definition 2.1.4. *The elements of $\ker D^g$ are called harmonic spinors. If n is even, then the elements of $\ker D^\pm$ are called positive respectively negative harmonic spinors.*

Let M be compact. The hermitian metric $\langle \cdot, \cdot \rangle$ on $\Sigma^g M$ induces a positive definite scalar product $(\cdot, \cdot)_2$ on smooth spinors by

$$(\psi, \varphi)_2 := \int_M \langle \psi, \varphi \rangle \mathrm{d}v^g,$$

where $\mathrm{d}v^g$ denotes the volume form induced by the Riemannian metric g . Let $p \in \mathbb{R}$ be positive. The L^p -norm of a smooth spinor ψ is by definition

$$\|\psi\|_p := \left(\int_M |\psi|_g^p \mathrm{d}v^g \right)^{1/p}.$$

The completion of $C^\infty(\Sigma^g M)$ with respect to $\|\cdot\|_p$ is called $L^p(\Sigma^g M)$. For $k \in \mathbb{N}$ the Sobolev k -norm of a smooth spinor ψ is defined by

$$\|\psi\|_{H^k} := \sum_{i=0}^k \|\nabla^i \psi\|_2$$

and the Sobolev space $H^k(\Sigma^g M)$ is the completion of $C^\infty(\Sigma^g M)$ with respect to $\|\cdot\|_{H^k}$. If (M, g) is complete, then with respect to the scalar product $(\cdot, \cdot)_2$ the Dirac operator D^g is essentially self-adjoint, i. e. its closure in $L^2(\Sigma^g M)$ is self-adjoint. If (M, g, Θ) is a closed Riemannian spin manifold, then the spectrum $\mathrm{spec}(D^g)$ of the Dirac operator consists of a sequence of isolated real eigenvalues, which is neither bounded from above nor bounded from below.

2.2 Real and quaternionic structures

In certain dimensions n there exist real or quaternionic structures on the modules Σ_n of the complex spinor representation which are $\mathrm{Spin}(n)$ -equivariant. Given a Riemannian spin manifold (M, g, Θ) of dimension n these structures then induce conjugate linear endomorphisms of $\Sigma^g M$.

Definition 2.2.1. *Let W be a complex vector space.*

1. *A real structure on W is a \mathbb{R} -linear map $J: W \rightarrow W$ such that $J^2 = \mathrm{Id}$ and $J(iw) = -iJ(w)$ for all $w \in W$.*
2. *A quaternionic structure on W is a \mathbb{R} -linear map $J: W \rightarrow W$ such that $J^2 = -\mathrm{Id}$ and $J(iw) = -iJ(w)$ for all $w \in W$.*

Let $\rho: \mathbb{Cl}(n) \rightarrow \text{End}_{\mathbb{C}}(\Sigma_n)$ be the complex spinor representation. A real or quaternionic structure J on Σ_n is called commuting, if it commutes with Clifford multiplication by elements of \mathbb{R}^n , i. e. if

$$J(x \cdot \sigma) = x \cdot J(\sigma) \text{ for all } x \in \mathbb{R}^n \subset \mathbb{Cl}(n), \sigma \in \Sigma_n.$$

It is called anti-commuting, if it anti-commutes with Clifford multiplication by elements of \mathbb{R}^n , i. e. if

$$J(x \cdot \sigma) = -x \cdot J(\sigma) \text{ for all } x \in \mathbb{R}^n \subset \mathbb{Cl}(n), \sigma \in \Sigma_n.$$

The existence of real or quaternionic structures on Σ_n for certain n is proved in [F4]. In the following theorem we state the result and mention some further structures.

Theorem 2.2.2. *On Σ_n the following structures exist*

- a) *If $n \equiv 0 \pmod{8}$ there exist a commuting real structure and an anti-commuting real structure.*
- b) *If $n \equiv 1 \pmod{8}$ there exists an anti-commuting real structure. There exists no commuting real structure.*
- c) *If $n \equiv 2 \pmod{8}$ there exist an anti-commuting real structure and a commuting quaternionic structure. There exists no commuting real structure.*
- d) *If $n \equiv 3 \pmod{8}$ there exists a commuting quaternionic structure. There exists no commuting real structure.*
- e) *If $n \equiv 4 \pmod{8}$ there exist a commuting quaternionic structure and an anti-commuting quaternionic structure. There exists no commuting real structure.*
- f) *If $n \equiv 5 \pmod{8}$ there exists an anti-commuting quaternionic structure. There exists no commuting real structure.*
- g) *If $n \equiv 6 \pmod{8}$ there exist a commuting real structure and an anti-commuting quaternionic structure.*
- h) *If $n \equiv 7 \pmod{8}$ there exists a commuting real structure.*

Proof. a) $n \equiv 0 \pmod{8}$: By [F4], p. 33 there exists an anti-commuting real structure J . Then $\omega_{\mathbb{C}} \circ J$ is a commuting real structure.

- b) $n \equiv 1 \pmod{8}$: By [F4] there exists an anti-commuting real structure. Suppose that there was a commuting real structure J . We restrict the spinor representation to $\text{Cl}(n) \subset \text{Cl}(n)$. If $n = 8k + 1$ then the eigenspace of J corresponding to 1 is a real subspace of Σ_n of real dimension 2^{4k} and is an invariant subspace of this restriction. This is a contradiction, since every irreducible real module for $\text{Cl}(n)$ has real dimension 2^{4k+1} (see [LM], p. 33).
- c) $n \equiv 2 \pmod{8}$: By [F4] there exists a commuting quaternionic structure J . Then $\omega_{\mathbb{C}} \circ J$ is an anti-commuting real structure. As in the case $n \equiv 1 \pmod{8}$ one sees that there is no commuting real structure.
- d) $n \equiv 3 \pmod{8}$: By [F4] there exists a commuting quaternionic structure. As in the case $n \equiv 1 \pmod{8}$ one sees that there is no commuting real structure.
- e) $n \equiv 4 \pmod{8}$: By [F4] there exists an anti-commuting quaternionic structure J . Then $\omega_{\mathbb{C}} \circ J$ is a commuting quaternionic structure. As in the case $n \equiv 1 \pmod{8}$ one sees that there is no commuting real structure.
- f) $n \equiv 5 \pmod{8}$: By [F4] there exists an anti-commuting quaternionic structure J . As in the case $n \equiv 1 \pmod{8}$ one sees that there is no commuting real structure.
- g) $n \equiv 6 \pmod{8}$: By [F4] there exists a commuting real structure J . Then $\omega_{\mathbb{C}} \circ J$ is an anti-commuting quaternionic structure.
- h) $n \equiv 7 \pmod{8}$: By [F4] there exists a commuting real structure. The assertion follows. \square

As an example we consider the case $n = 2$. The spinor representation of $\text{Cl}(2) = \text{End}_{\mathbb{C}}(\mathbb{C}^2)$ is the standard representation of $\text{Mat}(2, \mathbb{C})$ on \mathbb{C}^2 . We define the action of E_1 and E_2 by

$$E_1 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad E_2 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

We define the map $J: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ by $J(z, w) = (-\bar{w}, \bar{z})$. Then J is a commuting quaternionic structure. A motivation for this definition comes from considering the Hamilton quaternions

$$\mathbb{H} := \{a + ib + cj + dk \mid a, b, c, d \in \mathbb{R}, i^2 = j^2 = k^2 = -1, ij = -ji = k\}$$

as a complex vector space, \mathbb{C} acting by quaternion multiplication from the right, and from using the \mathbb{C} -linear isomorphism

$$\mathbb{C}^2 \rightarrow \mathbb{H}, \quad (z, w) \mapsto z + jw.$$

Under this isomorphism the actions of E_1, E_2 correspond to quaternion multiplication with $j, -k$ from the left and J corresponds to quaternion multiplication with j from the right. We define the map $K: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ by $K(z, w) := (-\bar{w}, -\bar{z})$. Then K is an anticommuting real structure.

Next let $n = 3$. The Clifford algebra $\mathbb{Cl}(3) = \text{End}_{\mathbb{C}}(\mathbb{C}^2) \oplus \text{End}_{\mathbb{C}}(\mathbb{C}^2)$ has two inequivalent irreducible complex representations ρ_1, ρ_2 given by

$$\rho_1(A, B)(x) := Ax, \quad \rho_2(A, B)(x) := Bx$$

for $x \in \mathbb{C}^2$. We define the action of E_1, E_2, E_3 under ρ_1 by

$$E_1 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad E_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad E_3 = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}.$$

Under the isomorphism

$$\mathbb{C}^2 \rightarrow \mathbb{H}, \quad (z, w) \mapsto z + jw.$$

this corresponds to quaternion multiplication with i, j, k from the left. The map $J: \mathbb{C}^2 \rightarrow \mathbb{C}^2$ by $J(z, w) = (-\bar{w}, \bar{z})$ is a commuting quaternionic structure.

Now let (M, g, Θ) be a Riemannian spin manifold of dimension n . Let J be one of the structures mentioned above. Then J commutes with the action of $\text{Spin}(n)$ on Σ_n and thus induces a map $\Sigma^g M \rightarrow \Sigma^g M$, $[s, \sigma] \mapsto [s, J\sigma]$ which will again be denoted by J . This map is fibre-preserving and \mathbb{R} -linear on the fibres and it satisfies $Ji = -iJ$ and $J^2 = \pm \text{Id}$. Furthermore for any $g \in \text{Spin}(n)$ we have

$$J \circ \rho(g) = \rho(g) \circ J.$$

Taking the derivative of this equation we obtain $J \circ d\rho = d\rho \circ J$. It follows from the formula (2.2) that $\nabla_X^g J\psi = J\nabla_X^g \psi$ for all $X \in TM$ and for all $\psi \in C^\infty(\Sigma^g M)$, i. e. J is parallel with respect to ∇^g .

We note the following important consequences for the spectrum of the Dirac operator.

Remark 2.2.3. *If J is an anti-commuting real structure, it anti-commutes with the Dirac operator. It follows that if λ is an eigenvalue of the Dirac operator, then also $-\lambda$ is an eigenvalue. By Theorem 2.2.2 this occurs in dimensions $n \equiv 0, 1, 2 \pmod{8}$.*

If J is a commuting quaternionic structure, it commutes with the Dirac operator. Since for every nonzero spinor ψ the system $\{\psi, J\psi\}$ is linearly independent over \mathbb{C} , every eigenspace of D^g has even complex dimension. By Theorem 2.2.2 this is the case in dimensions $n \equiv 2, 3, 4 \pmod{8}$.

If J is an anti-commuting quaternionic structure, it anti-commutes with the Dirac operator. In this case the kernel of D^g has even complex dimension and, if λ is an eigenvalue of D^g , then also $-\lambda$ is an eigenvalue of D^g . By Theorem 2.2.2 this occurs in dimensions $n \equiv 4, 5, 6 \pmod{8}$.

2.3 Spinors for different metrics

Let (M, g, Θ) be a Riemannian spin manifold and let h be another Riemannian metric on M . Since the spinor bundles $\Sigma^g M$ and $\Sigma^h M$ are two different vector bundles, the question arises how one can identify spinors on (M, g, Θ) with spinors on (M, h, Θ) in a natural way. The case of conformally related metrics g and h has been treated in [Hit], [Hij1]. For general Riemannian metrics g and h Bourguignon and Gauduchon [BG] have solved this problem. The question when such an identification can be obtained in the case of semi-Riemannian metrics has been treated in [BGM05]. For our purpose we will recall the method of [BG] and use some remarks from [Ma].

Given the metrics g and h there exists a unique endomorphism $a_{g,h}$ of TM such that for all $x \in M$ and for all $v, w \in T_x M$ we have

$$g(a_{g,h}v, w) = h(v, w).$$

For each $x \in M$ the endomorphism $a_{g,h}(x) \in \text{End}(T_x M)$ is g -self-adjoint and positive definite. Thus there exists a unique endomorphism $b_{g,h}(x)$ of $T_x M$ which is positive definite and satisfies $b_{g,h}(x)^2 = a_{g,h}(x)^{-1}$. In this way we obtain an endomorphism $b_{g,h}$ of TM . The compositions of the endomorphisms $a_{g,h}$ and $b_{g,h}$ have the following properties.

Lemma 2.3.1. *Let g, h, k be three Riemannian metrics on M . Then we have $a_{g,h}a_{h,k} = a_{g,k}$. The equation $b_{g,h} \circ b_{h,k} = b_{g,k}$ holds if and only if $a_{g,h}$ and $a_{h,k}$ commute.*

Proof. This follows immediately from the definitions of $a_{g,h}$ and $b_{g,h}$. \square

For example if g, h, k are conformally related, then $a_{g,h}$ and $a_{h,k}$ commute.

Lemma 2.3.2. *The endomorphism $b_{g,h}$ induces an isomorphism of principal $SO(n)$ -bundles*

$$c_{g,h} : \text{P}_{\text{SO}}(M, g) \rightarrow \text{P}_{\text{SO}}(M, h), \quad (e_i)_{i=1}^n \mapsto (b_{g,h}e_i)_{i=1}^n$$

Proof. One easily finds that $b_{g,h}$ maps positively oriented g -orthonormal bases to positively oriented h -orthonormal bases. Obviously the map $c_{g,h}$ is $SO(n)$ -equivariant and an isomorphism with $c_{h,g} = c_{g,h}^{-1}$. \square

Lemma 2.3.3. *The isomorphism $c_{g,h}$ lifts to an isomorphism of principal $\text{Spin}(n)$ -bundles*

$$\gamma_{g,h} : \text{P}_{\text{Spin}}(M, g) \rightarrow \text{P}_{\text{Spin}}(M, h).$$

Proof. Let $\Theta: \text{P}_{\text{Spin}}(M, g) \rightarrow \text{P}_{\text{SO}}(M, g)$ be the covering map. We define a family of Riemannian metrics $(g_t)_{t \in [0,1]}$ on M by $g_t := (1-t)g + th$ and we define the map

$$F : \text{P}_{\text{Spin}}(M, g) \times [0, 1] \rightarrow \text{P}_{\text{GL}^+}(M), \quad (s, t) \mapsto c_{g,g_t}(\Theta(s)).$$

We consider the following commutative diagram

$$\begin{array}{ccc} \text{P}_{\text{Spin}}(M, g) \times \{0\} & \xrightarrow{j} & \text{P}_{\widetilde{\text{GL}}^+}(M) \\ \downarrow i & & \downarrow \Theta \\ \text{P}_{\text{Spin}}(M, g) \times [0, 1] & \xrightarrow{F} & \text{P}_{\text{GL}^+}(M) \end{array}$$

where i and j are inclusions. Since the map Θ has the homotopy lifting property, there exists a unique map

$$G : \text{P}_{\text{Spin}}(M, g) \times [0, 1] \rightarrow \text{P}_{\widetilde{\text{GL}}^+}(M)$$

such that $\Theta \circ G = F$ and $G(s, 0) = s$ for all $s \in \text{P}_{\text{Spin}}(M, g)$. We define $\gamma_{g,h}(s) := G(s, 1)$. The definition of $\gamma_{g,h}$ does not depend on the family of Riemannian metrics g_t chosen above. Namely any two paths between g and h are homotopic and therefore yield the same result $G(s, 1)$. We find that $\gamma_{g,h}$ is $\text{Spin}(n)$ -equivariant, since c_{g,g_t} is $\text{SO}(n)$ -equivariant. Using the uniqueness of lifts one can show that $\gamma_{g,h}$ is an isomorphism with $\gamma_{h,g} = \gamma_{g,h}^{-1}$. \square

Lemma 2.3.4. *The isomorphism $\gamma_{g,h}$ induces an isomorphism of vector bundles*

$$\beta_{g,h} : \Sigma^g M \rightarrow \Sigma^h M, \quad [s, \sigma] \mapsto [\gamma_{g,h}(s), \sigma]$$

which is a fibrewise isometry with respect to the hermitian metrics on $\Sigma^g M$ and $\Sigma^h M$. Furthermore for all $v \in TM$ and for all $\psi \in \Sigma^g M$ we have $\beta_{g,h}(v \cdot \psi) = b_{g,h}v \cdot \beta_{g,h}\psi$, where \cdot denotes both Clifford multiplications on $\Sigma^g M$ and on $\Sigma^h M$.

Proof. The map $\beta_{g,h}$ is well defined, since $\gamma_{g,h}$ is $\text{Spin}(n)$ -equivariant. We find that $\beta_{h,g} = \beta_{g,h}^{-1}$. By the definition of the hermitian metrics on $\Sigma^g M$ and $\Sigma^h M$ the map $\beta_{g,h}$ is a fibrewise isometry. For the last assertion observe that under the isomorphism $TM \cong \text{P}_{\text{SO}}(M, g) \times_{\tau} \mathbb{R}^n$ for all $s \in \text{P}_{\text{Spin}}(M, g)$, $w \in \mathbb{R}^n$ we have

$$b_{g,h}([\Theta(s), w]) = [c_{g,h}(\Theta(s)), w] = [\Theta(\gamma_{g,h}(s)), w]$$

and therefore for all $s \in \text{P}_{\text{Spin}}(M, g)$, $w \in \mathbb{R}^n$, $\sigma \in \Sigma_n$

$$b_{g,h}([\Theta(s), w]) \cdot \beta_{g,h}([s, \sigma]) = [\gamma_{g,h}(s), w \cdot \sigma] = \beta_{g,h}([\Theta(s), w] \cdot [s, \sigma])$$

which completes the proof. \square

Next we want to compare the Dirac operators D^g and D^h . The map $\beta_{g,h}$ does not induce an isometry of Hilbert spaces $L^2(\Sigma^g M) \rightarrow L^2(\Sigma^h M)$, since the volume forms $\text{d}v^g$, $\text{d}v^h$ induced by the metrics g , h are different (see [Ma]). In order to compensate this we note, that there exists a smooth positive function $f_{g,h}$ on M such that $\text{d}v^h = f_{g,h}^2 \text{d}v^g$. We define

$$\bar{\beta}_{g,h} := \frac{1}{f_{g,h}} \beta_{g,h} : \Sigma^g M \rightarrow \Sigma^h M.$$

The maps $\beta_{g,h}$, $\bar{\beta}_{g,h}$ induce isomorphisms $C^\infty(\Sigma^g M) \rightarrow C^\infty(\Sigma^h M)$, which will also be denoted by $\beta_{g,h}$, $\bar{\beta}_{g,h}$. We use the map $\bar{\beta}_{g,h}$ to pull back the Dirac operator on $\Sigma^h M$ to spinors for the metric g .

$$D^{g,h} := \bar{\beta}_{h,g} D^h \bar{\beta}_{g,h}.$$

We see that these operators have the following properties.

Lemma 2.3.5. *The map $\bar{\beta}_{g,h}$ induces an isometry of Hilbert spaces*

$$\bar{\beta}_{g,h} : L^2(\Sigma^g M) \rightarrow L^2(\Sigma^h M).$$

If D^h has self-adjoint closure on $L^2(\Sigma^h M)$, then $D^{g,h}$ has self-adjoint closure on $L^2(\Sigma^g M)$.

Proof. This is clear from the definitions of $\bar{\beta}_{g,h}$ and $D^{g,h}$. \square

Next we want to express the operator $D^{g,h}$ in terms of a local orthonormal frame. For a Riemannian metric g on M we denote both the Levi Civita connection on TM and the induced connection on $\Sigma^g M$ by ∇^g .

Theorem 2.3.6. *Let $(e_i)_{i=1}^n$ be a local g -orthonormal frame defined on an open subset $U \subset M$. Then for all $\psi \in C^\infty(\Sigma^g M)$ we have on U :*

$$\begin{aligned}
D^{g,h}\psi &= \sum_{i=1}^n e_i \cdot \nabla_{b_{g,h}(e_i)}^g \psi \\
&\quad + \frac{1}{4} \sum_{i,j=1}^n e_i \cdot e_j \cdot (b_{h,g} \nabla_{b_{g,h}(e_i)}^h (b_{g,h} e_j) - \nabla_{b_{g,h}(e_i)}^g e_j) \cdot \psi \\
&\quad - \frac{1}{f_{g,h}} b_{g,h}(\text{grad}^g(f_{g,h})) \cdot \psi.
\end{aligned} \tag{2.7}$$

Here $\text{grad}^g(f)$ denotes the gradient of a smooth function f with respect to the metric g .

Proof. We find

$$\begin{aligned}
D^{g,h}\psi &= \bar{\beta}_{h,g} \left(\frac{1}{f_{g,h}} D^h \beta_{g,h} \psi - \frac{1}{f_{g,h}^2} \text{grad}^h(f_{g,h}) \cdot \beta_{g,h} \psi \right) \\
&= \beta_{h,g} D^h \beta_{g,h} \psi - \frac{1}{f_{g,h}} b_{h,g}(\text{grad}^h(f_{g,h})) \cdot \psi \\
&= \beta_{h,g} D^h \beta_{g,h} \psi - \frac{1}{f_{g,h}} b_{g,h}(\text{grad}^g(f_{g,h})) \cdot \psi.
\end{aligned}$$

For the first summand we obtain

$$\begin{aligned}
\beta_{h,g} D^h \beta_{g,h} \psi &= \beta_{h,g} \left(\sum_{i=1}^n b_{g,h} e_i \cdot \nabla_{b_{g,h}(e_i)}^h \beta_{g,h} \psi \right) \\
&= \sum_{i=1}^n e_i \cdot \beta_{h,g} \nabla_{b_{g,h}(e_i)}^h \beta_{g,h} \psi \\
&= \sum_{i=1}^n e_i \cdot \nabla_{b_{g,h}(e_i)}^g \psi + \sum_{i=1}^n e_i \cdot (\beta_{h,g} \nabla_{b_{g,h}(e_i)}^h \beta_{g,h} \psi - \nabla_{b_{g,h}(e_i)}^g \psi).
\end{aligned}$$

Using the formula (2.3) we get

$$\begin{aligned}
&\beta_{h,g} \nabla_{b_{g,h}(e_i)}^h \beta_{g,h} \psi - \nabla_{b_{g,h}(e_i)}^g \psi \\
&= \frac{1}{4} \sum_{j=1}^n \beta_{h,g} (b_{g,h} e_j \cdot (\nabla_{b_{g,h}(e_i)}^h (b_{g,h} e_j)) \cdot \beta_{g,h} \psi) - \frac{1}{4} \sum_{j=1}^n e_j \cdot (\nabla_{b_{g,h}(e_i)}^g e_j) \cdot \psi \\
&= \frac{1}{4} \sum_{j=1}^n e_j \cdot (b_{h,g} \nabla_{b_{g,h}(e_i)}^h (b_{g,h} e_j) - \nabla_{b_{g,h}(e_i)}^g e_j) \cdot \psi.
\end{aligned}$$

This gives the formula of the assertion. \square

For $r \in \mathbb{N}$ denote by $C^r(\text{Sym}^2(T^*M))$ (resp. $C^\infty(\text{Sym}^2(T^*M))$) the space of r times continuously differentiable (resp. smooth) symmetric $(2, 0)$ tensor fields on M . In order to compute the derivative of $D^{g,h}$ with respect to the metric h we let $k \in C^\infty(\text{Sym}^2(T^*M))$. Then there exists an open neighborhood $I \subset \mathbb{R}$ of 0 such that for every $t \in I$ the tensor field $g_t := g + tk$ is a Riemannian metric on M .

Theorem 2.3.7. *Let $(e_i)_{i=1}^n$ be a local g -orthonormal frame defined on an open subset $U \subset M$. Then for all $\psi \in C^\infty(\Sigma^g M)$ we have on U :*

$$\frac{d}{dt} D^{g,g_t} \Big|_{t=0} \psi = -\frac{1}{2} \sum_{i=1}^n e_i \cdot \nabla_{a_{g,k}(e_i)}^g \psi - \frac{1}{4} \sum_{i=1}^n \text{div}^g(k)(e_i) e_i \cdot \psi. \quad (2.8)$$

Here $\text{div}^g(k)$ denotes the divergence of k with respect to the metric g . If $(e_i)_{i=1}^n$ is a local orthonormal frame of TM , it is the one-form on M which is locally defined by

$$\text{div}^g(k)(X) := \sum_{i=1}^n (\nabla_{e_i}^g k)(X, e_i)$$

for all $X \in TM$. By $\text{tr}^g(k) \in C^\infty(M, \mathbb{R})$ we will denote the g -trace of k , which is locally defined by

$$\text{tr}^g(k) := \sum_{i=1}^n k(e_i, e_i).$$

Of course these definitions are independent of the choice of orthonormal frame.

Proof. From $g_t = g + tk$ we obtain $b_{g,g_t} = (\text{Id} + ta_{g,k})^{-1/2}$. It follows that

$$\frac{d}{dt} b_{g,g_t} \Big|_{t=0} = -\frac{1}{2} a_{g,k}.$$

The first summand is now obtained from the first summand of (2.7). By [Be], p. 62, we have for all $X, Y, Z \in TM$

$$2g\left(\frac{d}{dt} \nabla_X^{g_t} Y \Big|_{t=0}, Z\right) = (\nabla_X^g k)(Y, Z) + (\nabla_Y^g k)(X, Z) - (\nabla_Z^g k)(X, Y).$$

We calculate

$$\begin{aligned} & \frac{d}{dt} (b_{g_t,g} \nabla_{b_{g_t,g}(e_i)}^{g_t} (b_{g_t,g} e_j) - \nabla_{b_{g_t,g}(e_i)}^g e_j) \Big|_{t=0} \\ &= \frac{1}{2} a_{g,k} (\nabla_{e_i}^g e_j) + \frac{d}{dt} \nabla_{e_i}^{g_t} e_j \Big|_{t=0} - \frac{1}{2} \nabla_{a_{g,k}(e_i)}^g e_j - \frac{1}{2} \nabla_{e_i}^g (a_{g,k} e_j) + \frac{1}{2} \nabla_{a_{g,k}(e_i)}^g e_j \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \sum_{m=1}^n k(\nabla_{e_i}^g e_j, e_m) e_m + \frac{d}{dt} \nabla_{e_i}^{g_t} e_j|_{t=0} - \frac{1}{2} \sum_{m=1}^n g(\nabla_{e_i}^g (a_{g,k} e_j), e_m) e_m \\
&= \frac{1}{2} \sum_{m=1}^n (k(\nabla_{e_i}^g e_j, e_m) + k(e_j, \nabla_{e_i}^g e_m) - \partial_{e_i} k(e_j, e_m)) e_m + \frac{d}{dt} \nabla_{e_i}^{g_t} e_j|_{t=0} \\
&= -\frac{1}{2} \sum_{m=1}^n (\nabla_{e_i}^g k)(e_j, e_m) e_m + \frac{d}{dt} \nabla_{e_i}^{g_t} e_j|_{t=0} \\
&= \frac{1}{2} \sum_{m=1}^n ((\nabla_{e_j}^g k)(e_i, e_m) - (\nabla_{e_m}^g k)(e_i, e_j)) e_m.
\end{aligned}$$

It follows that

$$\begin{aligned}
&\frac{d}{dt} \sum_{i,j=1}^n e_i \cdot e_j \cdot (b_{g_t, g} \nabla_{b_{g_t, g_t}(e_i)}^{g_t} (b_{g_t, g_t} e_j) - \nabla_{b_{g_t, g_t}(e_i)}^g e_j)|_{t=0} \\
&= \frac{1}{2} \sum_{i,j,m=1}^n (\nabla_{e_j}^g k)(e_i, e_m) e_i \cdot e_j \cdot e_m - \frac{1}{2} \sum_{i,j,m=1}^n (\nabla_{e_m}^g k)(e_i, e_j) e_i \cdot e_j \cdot e_m \\
&= -\sum_{i,j=1}^n (\nabla_{e_j}^g k)(e_i, e_j) e_i - \sum_{i,j,m=1}^n (\nabla_{e_m}^g k)(e_i, e_j) e_i \cdot e_j \cdot e_m \\
&= -\sum_{i=1}^n \operatorname{div}^g(k)(e_i) e_i + \operatorname{grad}^g(\operatorname{tr}^g(k)).
\end{aligned}$$

From $\operatorname{dv}^{g_t} = \det(\operatorname{Id} + t a_{g,k})^{1/2} \operatorname{dv}^g$ it follows that $f_{g,g_t} = \det(\operatorname{Id} + t a_{g,k})^{1/4}$. Since $f_{g,g} \equiv 1$ we obtain

$$\frac{d}{dt} \frac{1}{f_{g,g_t}} b_{g,g_t}(\operatorname{grad}^g(f_{g,g_t}))|_{t=0} = \operatorname{grad}^g\left(\frac{d}{dt} f_{g,g_t}|_{t=0}\right) = \frac{1}{4} \operatorname{grad}^g(\operatorname{tr}^g(k)).$$

The assertion follows. \square

We define the following $(2, 0)$ tensor field on M .

Definition 2.3.8. *Let $\psi \in C^\infty(\Sigma^g M)$. The energy momentum tensor for ψ is a symmetric $(2, 0)$ tensor field on M defined by*

$$Q_\psi(X, Y) := \frac{1}{2} \operatorname{Re} \langle X \cdot \nabla_Y^g \psi + Y \cdot \nabla_X^g \psi, \psi \rangle$$

for $X, Y \in TM$.

Again let $k \in C^\infty(\text{Sym}^2(T^*M))$ and let $I \subset \mathbb{R}$ be an open interval containing 0 such that for all $t \in I$ the tensor field $g_t = g + tk$ is a Riemannian metric on M . Let λ be an eigenvalue of D^g with $d := \dim_{\mathbb{C}} \ker(D^g - \lambda)$. By Theorem A.0.13 and Lemma A.0.16 there exist real-analytic functions $\lambda_1, \dots, \lambda_d$ on I such that $\lambda_j(t)$ is an eigenvalue of D^{g_t} for all j and all t and $\lambda_j(0) = \lambda$ for all j . Furthermore there exist spinors $\psi_{j,t}$, $1 \leq j \leq d$, $t \in I$, which are real-analytic in t , such that $\psi_{j,t}$ is an eigenspinor of D^{g_t} corresponding to $\lambda_j(t)$. We can choose these spinors such that $\|\psi_{j,t}\|_2 = 1$ for all j and all t . Then we have

$$\lambda_j(t) = (\psi_{j,t}, D^{g_t} \psi_{j,t})_2$$

for all t . If we take the derivative with respect to t at $t = 0$, then, since D^g is self-adjoint and $\psi_{j,t}$ is normalized, the only contribution comes from the derivative of the Dirac operator.

$$\frac{d\lambda_j(t)}{dt} \Big|_{t=0} = (\psi_{j,0}, \frac{d}{dt} D^{g_t} \Big|_{t=0} \psi_{j,0})_2 \quad (2.9)$$

Recall that if $(e_i)_{i=1}^n$ is a local orthonormal frame, then by (2.8) we have locally

$$\frac{d}{dt} D^{g_t} \Big|_{t=0} \psi_{j,0} = -\frac{1}{2} \sum_{i=1}^n e_i \cdot \nabla_{a_{g,k}(e_i)}^g \psi_{j,0} - \frac{1}{4} \sum_{i=1}^n \text{div}^g(k)(e_i) e_i \cdot \psi_{j,0}.$$

The scalar product of the first term with $\psi_{j,0}$ can locally be written as

$$\begin{aligned} & -\frac{1}{2} \sum_{i=1}^n \langle \psi_{j,0}, e_i \cdot \nabla_{a_{g,k}(e_i)}^g \psi_{j,0} \rangle \\ &= -\frac{1}{4} \sum_{i,m=1}^n k(e_i, e_m) \langle \psi_{j,0}, e_i \cdot \nabla_{e_m}^g \psi_{j,0} + e_m \cdot \nabla_{e_i}^g \psi_{j,0} \rangle \end{aligned}$$

The scalar product of the second term with $\psi_{j,0}$ is purely imaginary. Since the left hand side of (2.9) is real, it is sufficient to consider the real part. Then the second term gives no contribution. If (\cdot, \cdot) denotes the standard pointwise inner product of $(2,0)$ tensor fields, then we obtain

$$\frac{d\lambda_j(t)}{dt} \Big|_{t=0} = -\frac{1}{2} \int_M (k, Q_{\psi_{j,0}}) dv^g. \quad (2.10)$$

We consider now the special case of conformally related Riemannian metrics on M . Let $u \in C^\infty(M, \mathbb{R})$ and let g, h be two Riemannian metrics on M such that $h = e^{2u}g$. The connections on the spinor bundles $\Sigma^g M$ and $\Sigma^h M$ are then related in the following way.

Theorem 2.3.9. *For all $\psi \in C^\infty(\Sigma^g M)$ and for all $X \in TM$ we have*

$$\begin{aligned}\nabla_X^h \beta_{g,h} \psi &= \beta_{g,h} (\nabla_X^g \psi - \frac{1}{2} X \cdot \text{grad}^g(u) \cdot \psi - \frac{1}{2} X(u) \psi) \\ \nabla_X^h \bar{\beta}_{g,h} \psi &= \bar{\beta}_{g,h} (\nabla_X^g \psi - \frac{1}{2} X \cdot \text{grad}^g(u) \cdot \psi - \frac{n+1}{2} X(u) \psi).\end{aligned}\quad (2.11)$$

Proof. By [Be], p. 58 we have for all $X, Y \in TM$

$$\nabla_X^h Y = \nabla_X^g Y + X(u)Y + Y(u)X - g(X, Y) \text{grad}^g(u).$$

Note that $b_{g,h} = e^{-u} \text{Id}$. Let $(e_i)_{i=1}^n$ be a local g -orthonormal frame on an open subset $U \subset M$. Then $(b_{g,h} e_i)_{i=1}^n$ is a local h -orthonormal frame on U . We obtain by the formula (2.3)

$$\begin{aligned}\nabla_X^h \beta_{g,h} \psi &= \partial_X \beta_{g,h} \psi + \frac{1}{4} \sum_{j=1}^n b_{g,h} e_j \cdot (\nabla_X^h b_{g,h} e_j) \cdot \beta_{g,h} \psi \\ &= \partial_X \beta_{g,h} \psi + \frac{1}{4} \sum_{j=1}^n b_{g,h} e_j \cdot b_{g,h} (\nabla_X^h e_j - X(u) e_j) \cdot \beta_{g,h} \psi \\ &= \partial_X \beta_{g,h} \psi + \frac{1}{4} \sum_{j=1}^n \beta_{g,h} (e_j \cdot (\nabla_X^g e_j + e_j(u) X) \cdot \psi) \\ &\quad - \frac{1}{4} \beta_{g,h} (X \cdot \text{grad}^g(u) \cdot \psi) \\ &= \beta_{g,h} (\nabla_X^g \psi + \frac{1}{4} \text{grad}^g(u) \cdot X \cdot \psi - \frac{1}{4} X \cdot \text{grad}^g(u) \cdot \psi).\end{aligned}$$

The first formula in the assertion now follows from (2.1). Furthermore using that $f_{g,h} = e^{nu/2}$ we get

$$\nabla_X^h \bar{\beta}_{g,h} \psi = X(e^{-nu/2}) \beta_{g,h} \psi + e^{-nu/2} \nabla_X^h \beta_{g,h} \psi.$$

The second formula now follows. \square

The Dirac operators on $\Sigma^g M$ and $\Sigma^h M$ are related by the following formulas.

Theorem 2.3.10. *For all $\psi \in C^\infty(\Sigma^g M)$ we have*

$$\begin{aligned}D^h(e^{-(n-1)u/2} \beta_{g,h} \psi) &= e^{-(n+1)u/2} \beta_{g,h} D^g \psi \\ D^h(e^{u/2} \bar{\beta}_{g,h} \psi) &= e^{-u/2} \bar{\beta}_{g,h} D^g \psi.\end{aligned}\quad (2.12)$$

Proof. Let $(e_i)_{i=1}^n$ be a local orthonormal frame. Using the formula (2.11) we find

$$\begin{aligned} D^h \beta_{g,h} \psi &= \sum_{i=1}^n b_{g,h}(e_i) \cdot \nabla_{b_{g,h}(e_i)}^h \beta_{g,h} \psi \\ &= \sum_{i=1}^n \beta_{g,h}(e_i \cdot e^{-u} (\nabla_{e_i}^g \psi - \frac{1}{2} e_i(u) \psi - \frac{1}{2} e_i \cdot \text{grad}^g(u) \cdot \psi)) \\ &= \beta_{g,h}(e^{-u} D^g \psi + \frac{n-1}{2} e^{-u} \text{grad}^g(u) \cdot \psi). \end{aligned}$$

The first formula follows from this calculation. Using $\bar{\beta}_{g,h} = e^{-nu/2} \beta_{g,h}$ one obtains the second formula. \square

We specialise the above formulas for the derivative of the Dirac operator and the derivative of the eigenvalue to the case of conformal changes of the Riemannian metric. Let $f \in C^\infty(M, \mathbb{R})$ and let $I \subset \mathbb{R}$ be an open interval around 0 such that for all $t \in I$ the tensor field $g_t := g + tfg$ is a Riemannian metric on M . Then we immediately obtain the following.

Theorem 2.3.11. *For all $\psi \in C^\infty(\Sigma^g M)$ we have*

$$\frac{d}{dt} D^{g_t} \psi \Big|_{t=0} = -\frac{f}{2} D^g \psi - \frac{1}{4} \text{grad}^g(f) \cdot \psi \quad (2.13)$$

and

$$\frac{d\lambda_j(t)}{dt} \Big|_{t=0} = -\frac{\lambda}{2} \int_M f |\psi_{j,0}|_g^2 dv^g. \quad (2.14)$$

Chapter 3

Motivation

3.1 Conformal bounds on Dirac eigenvalues

Let (M, g, Θ) be a compact Riemannian spin manifold of dimension $n \geq 2$. We denote by $\lambda_1^+(g)$ the smallest positive eigenvalue of D^g and by $\lambda_1^-(g)$ the largest negative eigenvalue of D^g . The question arises how the values $\lambda_1^\pm(g)$ depend on the geometry of the underlying manifold. Lower bounds on $|\lambda_1^\pm(g)|$ in terms of the scalar curvature of (M, g) have been obtained in [Li], [F1], [Ki1], [Ki2], [KSW1], [KSW2]. Searching for lower bounds which are uniform on the conformal class

$$[g] := \{e^{2u}g \mid u \in C^\infty(M, \mathbb{R})\}$$

of the metric g one observes first that for every constant $a > 0$ one has $\lambda_1^\pm(a^2g) = a^{-1}\lambda_1^\pm(g)$ by (2.12). Thus the expressions $|\lambda_1^\pm(g)|\text{vol}(M, g)^{1/n}$ are invariant under constant rescalings of the metric. We define the two invariants

$$\begin{aligned}\lambda_{\min}^+(M, [g], \Theta) &:= \inf_{h \in [g]} \lambda_1^+(h)\text{vol}(M, h)^{1/n} \\ \lambda_{\min}^-(M, [g], \Theta) &:= \inf_{h \in [g]} |\lambda_1^-(h)|\text{vol}(M, h)^{1/n}\end{aligned}$$

of the conformal class $[g]$. It was proven in [Am3] that

$$\lambda_{\min}^\pm(M, [g], \Theta) > 0.$$

These invariants have been treated by many authors (see e. g. [Hij1], [Hij2], [Lo], [Bä1], [Am2], [Am4]).

Explicit values are known only for very few spin manifolds. For example if g_{can} denotes the standard metric on S^n and σ the unique spin structure on

S^n , we use the notation $\mathbb{S}^n := (S^n, [g_{\text{can}}], \sigma)$. Then we have

$$\lambda_{\min}^+(\mathbb{S}^n) = \lambda_{\min}^-(\mathbb{S}^n) = \frac{n}{2} \text{vol}(S^n, g_{\text{can}})^{1/n}$$

(see e. g. [Am3], [Am4]). Furthermore one has the following inequalities.

Theorem 3.1.1 ([Am3], [AGHM]). *For every compact Riemannian spin manifold (M, g, Θ) we have*

$$\lambda_{\min}^+(M, [g], \Theta) \leq \lambda_{\min}^+(\mathbb{S}^n), \quad \lambda_{\min}^-(M, [g], \Theta) \leq \lambda_{\min}^-(\mathbb{S}^n).$$

It is a natural question whether the infimum in the definition of the invariants $\lambda_{\min}^\pm(M, [g], \Theta)$ is attained at a Riemannian metric $h \in [g]$. This has been investigated in [Am4], where the following result is obtained.

Theorem 3.1.2. *Let (M, g, Θ) be a compact Riemannian spin manifold of dimension $n \geq 2$. Assume that we have the strict inequality*

$$\lambda_{\min}^+ := \lambda_{\min}^+(M, [g], \Theta) < \lambda_{\min}^+(\mathbb{S}^n).$$

Then there exists a spinor $\psi \in C^2(\Sigma^g M)$, which is smooth on $M \setminus \psi^{-1}(0)$, such that

$$D^g \psi = \lambda_{\min}^+ |\psi|_g^{2/(n-1)} \psi, \quad \|\psi\|_{2n/(n-1)} = 1. \quad (3.1)$$

If there is such a spinor ψ , which is nowhere zero on M , then $h := |\psi|_g^{4/(n-1)} g$ is a Riemannian metric on M such that

$$\lambda_{\min}^+ = \lambda_1^+(h) \text{vol}(M, h)^{1/n}.$$

An analogous assertion holds for $\lambda_{\min}^-(M, [g], \Theta)$.

Therefore one is interested in finding solutions to (3.1) which are nowhere zero. It is not clear, whether this is possible. An important result by C. Bär on the zero sets of solutions of generalized Dirac equations is the following.

Theorem 3.1.3 ([Bä4]). *Let (M, g, Θ) be a connected Riemannian spin manifold of dimension $n \geq 2$, not necessarily compact and possibly with boundary. Let h be a smooth endomorphism field of $\Sigma^g M$ and let ψ be a nontrivial solution of*

$$(D^g + h)\psi = 0.$$

Then the zero set of ψ is a countably $(n - 2)$ -rectifiable set and thus has Hausdorff dimension $n - 2$ at most. If $n = 2$, then the zero set of ψ is a discrete subset of M .

One can apply this theorem for $n = 2$. However, if ψ is zero somewhere and $n \geq 3$, then $|\psi|_g^{2/(n-1)}$ is not smooth on the zero set of ψ and thus the theorem cannot be applied. In this case one can still prove that the zero set of a nontrivial solution does not contain any open subset of M and that its complement is connected (see [AMM]).

In this thesis we will not solve this problem. However our result on the zero sets of eigenspinors for generic metrics suggests that for generic metrics any solution to (3.1) should be nowhere zero.

3.2 Witten spinors

In his proof of the positive energy theorem for spin manifolds Witten introduces asymptotically constant harmonic spinors on certain non-compact Riemannian manifolds (see [Wi]). A rigorous proof of his ideas is given in [PT]. Following the latter article we will state existence and uniqueness of these so called Witten spinors. Then we will see that the zero set of certain Witten spinors is not empty. For $x \in \mathbb{R}^n$ we denote by r its euclidean distance from 0. We assume that $n \geq 3$.

Definition 3.2.1. For $s \in \mathbb{R}$ we define

$$f \in O''(r^s) : \iff \nabla^j f \in O(r^{s-j}) \text{ for } j = 0, 1, 2.$$

Definition 3.2.2. A complete Riemannian manifold (M, g) of dimension n is called asymptotically flat, if there exists a compact subset $K \subset M$ such that $M \setminus K$ is the disjoint union of a finite number of subsets M_1, \dots, M_d with the property that for every $i \in \{1, \dots, d\}$ there exists a contractible and compact subset $K_i \subset \mathbb{R}^n$ and a diffeomorphism $\Phi_i: \mathbb{R}^n \setminus K_i \rightarrow M_i$, such that in the standard coordinates of \mathbb{R}^n for all j, k we have:

$$(\Phi_i^* g)_{jk} - \delta_{jk} \in O''(r^{2-n})$$

as $r \rightarrow \infty$. The subsets M_i are called the ends of M .

The simplest example of an asymptotically flat Riemannian manifold is the Euclidean space $(\mathbb{R}^n, g_{\text{eucl}})$. A more interesting example is $M = \mathbb{R}^3 \setminus \{0\}$ with the Schwarzschild metric $g = (1 + \frac{1}{r})^4 g_{\text{eucl}}$, where $m > 0$ is a constant. It has two asymptotically flat ends, one at infinity and one at zero. In order to see the end at zero note that the map $x \mapsto \frac{1}{r^2} x$ is an isometry of (M, g) (see [BN]). Using this map one obtains a diffeomorphism, which defines the end at zero.

Since the different spin structures over $\mathbb{R}^n \setminus K_i$ are classified by the elements of $H^1(\mathbb{R}^n \setminus K_i, \mathbb{Z}/2\mathbb{Z}) = 0$ we see that the pullback bundle $\Phi_i^* \Sigma^g M$ is equal to the trivial spinor bundle over $\mathbb{R}^n \setminus K_i$ and thus extends to the trivial bundle $\mathbb{R}^n \times \Sigma_n$ over all of \mathbb{R}^n . The pullbacks under Φ_i^{-1} of the constant sections of this bundle will be called the constant spinors on the end M_i . From now we assume $n = 3$.

Theorem 3.2.3 ([PT]). *Let (M, g, Θ) be a 3-dimensional asymptotically flat Riemannian spin manifold. For all $i \in \{1, \dots, d\}$ let γ_i be a constant spinor on the end M_i . Then there exists a unique spinor ψ on M such that $D^g \psi = 0$ and such that for every $\varepsilon > 0$ we have $\lim_{r \rightarrow \infty} r^{1-\varepsilon} |\psi - \gamma_i| = 0$ on each end M_i .*

The spinor ψ is called the Witten spinor for the constant spinors $\gamma_1, \dots, \gamma_d$.

Next we describe the relation between nowhere vanishing spinors on manifolds of dimension 3 and orthonormal frames of its tangent bundle. It has long been known that every orientable manifold M of dimension 3 is parallelizable, i. e. there exists a global frame of the tangent bundle, which can of course be orthonormalized (see [St], [Wh]). More recently it has been suggested by people working in general relativity (see [N], [DM], [FNS]) to use Witten spinors in order to construct special orthonormal frames of TM . This is possible if the Witten spinor is nowhere zero on M . In order to explain this, we first recall that $\text{Spin}(3) = \text{SU}(2)$ and therefore

$$\text{Spin}(3) = \left\{ \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \in \text{GL}(2, \mathbb{C}) \mid |a|^2 + |b|^2 = 1 \right\}.$$

It follows that for any $v, w \in S^3 \subset \mathbb{C}^2$ there exists exactly one $q \in \text{SU}(2)$ such that $\rho(q)v = w$. Using this fact we can show:

Theorem 3.2.4. *Let (M, g, Θ) be a Riemannian spin manifold of dimension 3. Then every smooth spinor on M which is nowhere zero leads to a global orthonormal frame of TM .*

Proof. Let $\psi \in C^\infty(\Sigma^g M)$ be nowhere zero on M . We define $\chi \in C^\infty(\Sigma^g M)$ by $\chi(x) := \frac{\psi(x)}{|\psi(x)|_g}$. Let $U \subset M$ be open such that there exists a smooth section s of $P_{\text{Spin}}(M, g)|_U$ and a smooth map $\sigma: U \rightarrow S^3 \subset \mathbb{C}^2$ such that on U we have $\chi|_U = [s, \sigma]$. For $x \in U$ let $q(x) \in \text{Spin}(3)$ be the unique element such that $\rho(q(x))\sigma(x)$ is equal to the first vector E_1 of the standard basis of \mathbb{C}^2 . We obtain $q \in C^\infty(U, \text{Spin}(3))$ such that for all $x \in U$ we have $\chi|_U(x) = [s(x)q(x)^{-1}, E_1]$. It follows that $s(x)q(x)^{-1}$ is independent of the choices of s, σ and thus can be defined on all of M . Then $e: M \rightarrow P_{\text{SO}}(M, g)$ with

$$e(x) := \Theta(s(x)q(x)^{-1})$$

is a smooth section of $P_{\text{SO}}(M, g)$. \square

However it is easy to see that Witten spinors can have zeros. More precisely if W denotes the space of all Witten spinors on an asymptotically flat manifold M with d ends, then by Theorem 3.2.3 there exists a \mathbb{C} -linear isomorphism $(\Sigma_3)^d \rightarrow W$. Let $x \in M$. The evaluation map

$$ev_x : W \rightarrow \Sigma_x^g M, \quad \psi \mapsto \psi(x)$$

is a \mathbb{C} -linear map between complex vector spaces of dimensions $2d$ and 2 . Thus in the case $d > 1$ for every $x \in M$ there exists a Witten spinor on M , which is zero in x .

In the case $d = 1$ it is not clear whether Witten spinors can have zeros. If ψ is a Witten spinor, then $W = \text{span}_{\mathbb{C}}\{\psi, J\psi\}$, where J is a quaternionic structure. Thus if there exists $x \in M$ and a Witten spinor ψ with $\psi(x) = 0$, then all Witten spinors on (M, g, Θ) are zero in x .

In this thesis we will not solve this problem. However one can hope to carry over some of our techniques in order to prove that in the generic case Witten spinors are nowhere zero. This would mean that the zero sets of Witten spinors have no physical significance.

3.3 Zero sets of eigenspinors

Let (M, g, Θ) be a closed Riemannian spin manifold of dimension $n \geq 2$. We observe that eigenspinors of the Dirac operator can have non empty zero set. Namely there are explicit computations of eigenvalues whose multiplicities are greater than the rank of the spinor bundle (see e. g. Theorem 4.2.1). Thus for these eigenvalues there exist eigenspinors with zeros. Harmonic spinors on Riemann surfaces provide another example, which we will treat in Section 4.1.

It would be useful to have eigenspinors, which are nowhere zero, since then one would have a simple proof of Hijazi's inequality, which we now explain. First let

$$\Delta^g : C^\infty(M, \mathbb{R}) \rightarrow C^\infty(M, \mathbb{R})$$

denote the Laplace operator for the metric g acting on functions on M . If x_1, \dots, x_n are local coordinates on M with coordinate vector fields $\partial_1, \dots, \partial_n$ and the matrices (g_{ij}) and (g^{ij}) are defined by

$$g_{ij} := g(\partial_i, \partial_j), \quad g := (g_{ij}), \quad (g^{ij}) := g^{-1},$$

then we have for all locally defined smooth functions f :

$$\Delta^g f = -\det(g)^{-1/2} \sum_{i,j=1}^n \partial_i(g^{ij} \det(g)^{1/2} \partial_j f). \quad (3.2)$$

The conformal Laplace operator is a linear second order differential operator

$$L^g : C^\infty(M, \mathbb{R}) \rightarrow C^\infty(M, \mathbb{R}),$$

which is defined by

$$L^g f := \frac{4(n-1)}{n-2} \Delta^g f + \text{scal}^g f,$$

where scal^g denotes the scalar curvature for the metric g . The operator L^g is called the conformal Laplace operator because it satisfies a nice transformation law under conformal changes of the metric. More precisely let g and $h = e^{2u}g$ be two conformally related metrics, where u is a smooth function on M . Then we have for all $f \in C^\infty(M, \mathbb{R})$ (see [LP], p. 43)

$$L^g(e^{(n-2)u/2} f) = e^{(n+2)u/2} L^h f. \quad (3.3)$$

The spectrum of L^g consists of a sequence of isolated real eigenvalues, which is bounded from below by the smallest eigenvalue μ and is not bounded from above.

$$\mu := \mu_0 \leq \mu_1 \leq \dots \rightarrow \infty.$$

Hijazi's inequality gives a lower bound on the eigenvalues of the Dirac operator in terms of the smallest eigenvalue of the conformal Laplace operator.

Theorem 3.3.1 ([Hij1]). *Let (M, g, Θ) be a closed Riemannian spin manifold of dimension $n \geq 3$ and let μ be the smallest eigenvalue of the conformal Laplace operator. Then any eigenvalue λ of the Dirac operator satisfies*

$$\lambda^2 \geq \frac{n}{4(n-1)} \mu.$$

For the new simple proof we need the following assumption

- (A) There exists an eigenspinor ψ corresponding to λ , which is nowhere zero on M .

Proof of Theorem 3.3.1 using Assumption (A). We define a smooth function u on M by $u := \frac{2}{n-1} \ln |\psi|_g$. The metric $h := e^{2u}g$ is conformal to g and the spinor

$$\varphi := e^{-(n-1)u/2} \beta_{g,h} \psi \in C^\infty(\Sigma^h M)$$

satisfies $|\varphi|_h \equiv 1$ on M and $D^h\varphi = e^{-u}\lambda\varphi$ by (2.12). By the Schrödinger-Lichnerowicz formula (2.6) we find

$$\int_M |D^h\varphi|_h^2 dv^h = \int_M |\nabla^h\varphi|_h^2 dv^h + \frac{1}{4} \int_M \text{scal}^h dv^h.$$

We use the estimate

$$\int_M |\nabla^h\varphi|_h^2 dv^h \geq \frac{1}{n} \int_M |D^h\varphi|_h^2 dv^h.$$

Since the volume elements are related by $dv^h = e^{nu}dv^g$ we find

$$\int_M |D^h\varphi|_h^2 dv^h = \lambda^2 \int_M e^{(n-2)u} dv^g$$

and thus altogether

$$\int_M \text{scal}^h dv^h \leq \frac{4(n-1)}{n} \lambda^2 \int_M e^{(n-2)u} dv^g.$$

By the transformation formula (3.3) we find

$$\text{scal}^h = L^h 1 = e^{-(n+2)u/2} L^g(e^{(n-2)u/2})$$

and thus by setting $f := e^{(n-2)u/2}$ we obtain

$$\int_M f L^g f dv^g \leq \frac{4(n-1)}{n} \lambda^2 \int_M f^2 dv^g.$$

Since L^g is an elliptic self-adjoint differential operator on a compact manifold, there exists an L^2 -orthonormal Hilbert basis of $L^2(M, \mathbb{R})$ consisting of eigenfunctions of L^g (see e. g. [LM], p. 196). We write f as a possibly infinite linear combination of these eigenfunctions and conclude that the left hand side is bounded from below by $\mu \int_M f^2 dv^g$. \square

The eigenvalues of D^g depend continuously on g with respect to the C^k -topology for all $k \geq 1$ (see Proposition 6.2.3). For $k \in \mathbb{N}$ the k -th eigenvalue $\mu_k(g)$ of the conformal Laplacian L^g can be characterized as follows. For every $(k+1)$ -dimensional subspace V_{k+1} of $C^\infty(M, \mathbb{R})$ we define

$$\Lambda_g(V_{k+1}) := \sup\{(f, L^g f)_2 / \|f\|_2^2 \mid f \in V_{k+1} \setminus \{0\}\}.$$

Then we have

$$\mu_k(g) := \inf_{V_{k+1}} \Lambda_g(V_{k+1}),$$

where the infimum is taken over all $(k + 1)$ -dimensional subspaces. This can be proven similarly to Proposition 2.1 in [BU]. Since the scalar curvature contains second derivatives of the metric, it follows that $\mu_k(g)$ depends continuously on g with respect to the C^2 -topology.

We will show in Section 6.2 that for $n \in \{2, 3\}$ the subset of all Riemannian metrics on M , for which Assumption (A) holds, is dense in the space of all smooth Riemannian metrics on M with respect to any C^k -topology, $k \geq 1$. By continuity of the eigenvalues of D^g and L^g we obtain a new proof of Theorem 3.3.1 for $n = 3$.

Chapter 4

Examples of zero sets of eigenspinors

4.1 Harmonic spinors on Riemann surfaces

First we recall a way of identifying positive harmonic spinors on closed Riemann surfaces with holomorphic sections of certain line bundles (see [Hit]). Let (M, g) be a compact Kähler manifold of complex dimension m . We denote by $K := \Lambda^m T^{1,0}M$ the canonical line bundle over M . Then M is a spin manifold if and only if there exists a holomorphic line bundle L such that $L \otimes_{\mathbb{R}} L = K$. In this case the different spin structures on M are in one-to-one-correspondence with such holomorphic line bundles. The spinor bundle is isomorphic to $\Lambda^* T^{0,1}M \otimes_{\mathbb{R}} L$ and the $\mathbb{Z}/2\mathbb{Z}$ -grading is given by the decomposition of $\Lambda^* T^{0,1}M$ into even and odd forms. Let

$$\bar{\partial} : \Lambda^p T^{0,1}M \rightarrow \Lambda^{p+1} T^{0,1}M$$

be the exterior derivative in the Dolbeault complex. If s is a local holomorphic section of L and φ is a smooth local section of $\Lambda^* T^{0,1}M$, we define $\bar{\partial}(\varphi \otimes s) := \bar{\partial}\varphi \otimes s$. Then the Dirac operator corresponds to $\sqrt{2}(\bar{\partial} + \bar{\partial}^*)$. Now let M be a closed orientable surface with a Riemannian metric g . Then M is a spin manifold and since $\Lambda^0 T^{0,1}M$ is the trivial line bundle, there is a canonical isomorphism $\Sigma^+ M \cong L$. Thus positive harmonic spinors can be identified with holomorphic sections of L .

Next let (M, g, Θ) be a two-dimensional Riemannian spin manifold. We recall a way of associating a vector field to a positive or negative spinor on M (see [Am1]). First we define $\tau_{\pm}: \text{SO}(2) \rightarrow \mathbb{C}$ by

$$\begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} \mapsto \exp(\pm it).$$

We define a complex structure J on M such that for every $p \in M$ and for every unit vector $X \in T_p M$ the system (X, JX) is a positively oriented orthonormal basis of $T_p M$. Then the map $P_{\text{SO}}(M, g) \times_{\tau_+} \mathbb{C} \rightarrow (TM, J)$ which sends $[(e_1, e_2), 1]$ to e_1 is an isomorphism of complex line bundles. In the same way we obtain an isomorphism $P_{\text{SO}}(M, g) \times_{\tau_-} \mathbb{C} \rightarrow (TM, -J)$. We quote the following Lemma from [Am1] including the proof. It enables us to associate a vector field on M to a given section of $\Sigma^\pm M$.

Lemma 4.1.1. *Let Θ be a spin structure on (M, g) . Then the map*

$$\begin{aligned} \Phi_\pm : \quad \Sigma^\pm M = P_{\text{Spin}}(M, g) \times_\rho \Sigma_2^\pm &\rightarrow P_{\text{SO}}(M, g) \times_{\tau_\mp} \mathbb{C} = (TM, \mp J) \\ [s, \sigma] &\mapsto [\Theta(s), \sigma^2] \end{aligned}$$

is well defined.

Proof. For any n the multiplicative group $\text{Cl}(n)^*$ of invertible elements in the Clifford algebra $\text{Cl}(n)$ is an open subset. Thus one may identify the Lie algebra of the spin group $\text{Spin}(n)$ with a subspace of $\text{Cl}(n)$. By [LM], Proposition I.6.1 the Lie algebra of $\text{Spin}(2)$ is spanned by $E_1 \cdot E_2$. Therefore every element of $\text{Spin}(2)$ can be written in the form $\exp(tE_1 \cdot E_2)$. Since $E_1 \cdot E_2$ acts on Σ_2^\pm as $\mp i\text{Id}$, we find that $\exp(tE_1 \cdot E_2)$ acts as $\exp(\mp it)$. Furthermore by [LM], Proposition I.6.2 we have

$$\vartheta_*(E_1 \cdot E_2) = \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix} = 2J.$$

Thus for every $g \in \text{Spin}(n)$ the element g^2 acts on Σ_2^\pm as $\tau_\mp(\vartheta(g))$. We conclude that

$$\begin{aligned} [\Theta(sg), \sigma^2] &= [\Theta(s)\vartheta(g), \sigma^2] \\ &= [\Theta(s), \tau_\mp(\vartheta(g)) \cdot \sigma^2] \\ &= [\Theta(s), (g \cdot \sigma)^2] \end{aligned}$$

and therefore Φ_\pm is well defined. \square

From now assume that M is a closed oriented surface of genus γ with a Riemannian metric g . Assume that ψ is a nontrivial positive harmonic spinor on M and that $p \in M$ is a point with $\psi(p) = 0$. Since $\Sigma^+ M$ can be canonically identified with a holomorphic line bundle, there exists a local chart of M mapping p to $0 \in \mathbb{C}$ and a local trivialization of $\Sigma^+ M$ around p , such that in this chart and in this trivialization ψ corresponds to a holomorphic function $f: U \rightarrow \mathbb{C}$ on an open subset $U \subset \mathbb{C}$ containing 0. Then in a neighborhood of 0 the function f is given by a convergent power series

$f(z) = \sum_{k=1}^{\infty} a_k z^k$ with $a_k \in \mathbb{C}$. Define m_p as the smallest integer k such that $a_k \neq 0$. Let X be the vector field on M associated to ψ via Lemma 4.1.1. It follows that X has an isolated zero at p with index equal to $-2m_p$. Let $\chi(M) = 2 - 2\gamma$ denote the Euler characteristic of M . Denote by N the zero set of ψ . Since M is compact, the set N is finite. By the Poincaré-Hopf Theorem (see e.g. [Mi], p. 35) we obtain $-2 \sum_{p \in N} m_p = \chi(M)$. Thus we find the following result.

Theorem 4.1.2. *Assume that ψ is a positive harmonic spinor on a closed oriented surface M and let $N \subset M$ be its zero set. Then N is finite and we have*

$$\sum_{p \in N} m_p = -\frac{1}{2}\chi(M) = \gamma - 1.$$

From this we can see the well known fact, that for $\gamma = 0$ there are no harmonic spinors for any Riemannian metric (see e.g. [Hit]). In the case $\gamma = 1$ it has been proven in [F2] that for the flat Riemannian metric g there exists exactly one spin structure with $\dim \ker(D^g) = 2$ and exactly three spin structures with no harmonic spinors. Since the dimension of the space of harmonic spinors is conformally invariant and since any two Riemannian metrics on M are conformally related, it follows that this result holds for all Riemannian metrics on M . If ψ is a positive harmonic spinor and J is a parallel quaternionic structure on the spinor bundle, then $J\psi$ is a negative harmonic spinor and it has the same zero set as ψ . By Theorem 4.1.2 all the harmonic spinors are nowhere zero on M . For $\gamma = 2$ it follows from [Hit], Proposition 2.3, that there exist 6 distinct spin structures on M , such that for every Riemannian metric g on M we have $\dim \ker(D^g) = 2$. We take one of these spin structures and fix a Riemannian metric g on M . Using again a parallel quaternionic structure and using Theorem 4.1.2 we find that the zero set of every harmonic spinor consists of exactly one point and it is the same point for all harmonic spinors.

4.2 The sphere with the standard metric

We will explicitly calculate some zero sets of eigenspinors on the sphere with the standard metric. The reader, who is only interested in the main results may skip this section and continue in Chapter 5.

Theorem 4.2.1 ([Bä2]). *Let $M = S^n$, $n \geq 2$, be equipped with the standard metric and the unique spin structure. The eigenvalues of the Dirac operator*

are $\lambda = \pm(\frac{n}{2} + m)$, $m \in \mathbb{N}$, with multiplicities

$$2^{\lfloor n/2 \rfloor} \binom{m+n-1}{m}.$$

We will describe the method used in [Bä2] for the computation of the spectrum. In this way we will obtain the eigenspinors and their zero sets explicitly.

Let Δ^g be the Laplace operator acting on smooth functions on S^n defined as in (3.2). It is a classical result that the eigenvalues of Δ^g are given by $k(n+k-1)$, $k = 0, 1, 2, \dots$ with multiplicities

$$\frac{n+2k-1}{n+k-1} \binom{n+k-1}{k}.$$

The eigenfunctions for $k(n+k-1)$ are exactly the restrictions to S^n of polynomial functions defined on \mathbb{R}^{n+1} which are homogeneous of degree k and harmonic with respect to the Laplace operator on \mathbb{R}^{n+1} (see [BGM]).

If n is even, let α_j , $1 \leq j \leq N$, be constant and pointwise orthonormal sections of $\Sigma\mathbb{R}^{n+1}$. If n is odd, then we have a decomposition

$$\Sigma\mathbb{R}^{n+1} = \Sigma^+\mathbb{R}^{n+1} \oplus \Sigma^-\mathbb{R}^{n+1},$$

and we define α_j , $1 \leq j \leq N$ to be constant and pointwise orthonormal sections of $\Sigma^+\mathbb{R}^{n+1}$. In each case we define spinors β_j on \mathbb{R}^{n+1} by $x \mapsto x \cdot \alpha_j$, $1 \leq j \leq N$. One finds that the restrictions of the α_j to the hypersurface (S^n, g) are Killing spinors on S^n for the value $1/2$ and that the restrictions of the β_j to S^n are Killing spinors for the value $-\frac{1}{2}$. We denote the restrictions of the α_j by $\varphi_{1,j}$ and the restrictions of the β_j by $\varphi_{-1,j}$. Using this one can show that for every eigenfunction f_k of Δ^g corresponding to $k(n+k-1)$, for all $j \in \{1, \dots, N\}$ and for all $\mu \in \{\pm 1\}$ the equation

$$(D^g + \frac{\mu}{2})^2(f_k\varphi_{\mu,j}) = (k + \frac{n-1}{2})^2 f_k\varphi_{\mu,j}.$$

holds. Therefore if $\{f_i\}_{i \in \mathbb{N}}$ is an orthonormal Hilbert basis of $L^2(S^n, \mathbb{R})$ consisting of eigenfunctions of Δ^g on S^n then the $f_i\varphi_{\mu,j}$ form an orthonormal Hilbert basis of $L^2(\Sigma^g S^n)$ consisting of eigenspinors of $(D^g + \frac{\mu}{2})^2$. In general, if A is an endomorphism of a complex vector space and ν is a nonzero complex number, then we have an isomorphism

$$\ker(A^2 - \nu^2) \rightarrow \ker(A - \nu) \oplus \ker(A + \nu), \quad u \mapsto (Au + \nu u) \oplus (Au - \nu u).$$

Therefore the eigenvectors of A corresponding to $\pm\nu$ are exactly the nonzero vectors $Au \pm \nu u$, where $u \in \ker(A^2 - \nu^2)$. In this way the eigenvalues of D^g and their multiplicities are obtained in [Bä2].

We put $A = D^g + \frac{\mu}{2}$, $\lambda = k + \frac{n-1}{2}$ and compute

$$\begin{aligned}\psi_{k,j}^{\varepsilon,\mu} &:= (D^g + \frac{\mu}{2})(f_k \varphi_{\mu,j}) + \varepsilon(k + \frac{n-1}{2})f_k \varphi_{\mu,j} \\ &= (\frac{\mu(1-n)}{2} + \varepsilon(k + \frac{n-1}{2}))f_k \varphi_{\mu,j} + \text{grad}^{S^n}(f_k) \cdot \varphi_{\mu,j},\end{aligned}$$

where $\varepsilon \in \{\pm 1\}$. If $\psi_{k,j}^{\varepsilon,\mu}$ is not identically zero, then it is an eigenspinor of D^g corresponding to the value $\varepsilon(k + \frac{n-1}{2}) - \frac{\mu}{2}$. We find

$$\begin{aligned}\psi_{k,j}^{-} &= -kf_k \varphi_{-1,j} + \text{grad}^{S^n}(f_k) \cdot \varphi_{-1,j}, & \lambda &= -\frac{n}{2} - k + 1, \\ \psi_{k,j}^{+-} &= (k+n-1)f_k \varphi_{-1,j} + \text{grad}^{S^n}(f_k) \cdot \varphi_{-1,j}, & \lambda &= \frac{n}{2} + k, \\ \psi_{k,j}^{-+} &= (-k+1-n)f_k \varphi_{1,j} + \text{grad}^{S^n}(f_k) \cdot \varphi_{1,j}, & \lambda &= -\frac{n}{2} - k, \\ \psi_{k,j}^{++} &= kf_k \varphi_{1,j} + \text{grad}^{S^n}(f_k) \cdot \varphi_{1,j}, & \lambda &= \frac{n}{2} + k - 1.\end{aligned}$$

Therefore all the $\psi_{k,j}^{\varepsilon,\mu}$ except $\psi_{0,j}^{-}$ and $\psi_{0,j}^{++}$ are not identically zero.

In order to investigate the zero sets of the eigenspinors we first note that the spinors $\psi_{0,j}^{+-}$, $1 \leq j \leq 2^{\lfloor n/2 \rfloor}$, are eigenspinors for $\lambda = \frac{n}{2}$ and are non-zero multiples of $\varphi_{-1,j}$. Thus any nontrivial linear combination of them is nowhere zero on S^n . Similarly the spinors $\psi_{0,j}^{-+}$, $1 \leq j \leq 2^{\lfloor n/2 \rfloor}$, are eigenspinors for $\lambda = -\frac{n}{2}$ and are non-zero multiples of $\varphi_{1,j}$. Thus any nontrivial linear combination of them is nowhere zero on S^n . In the case $k \geq 1$ we determine the zero set of $\psi_{k,j}^{\varepsilon,\mu}$ as follows: We have

$$\psi_{k,j}^{\varepsilon,\mu}(x) = 0 \iff \text{grad}^{S^n}(f_k)(x) = 0 \text{ and } f_k(x) = 0.$$

Since f_k is homogeneous of degree k , we obtain for the normal component of $\text{grad}^{\mathbb{R}^{n+1}}(f_k)$ in $x \in S^n$ that

$$g_{\text{eucl}}(\text{grad}^{\mathbb{R}^{n+1}}(f_k)(x), x) = kf_k(x).$$

If $k \geq 1$, it follows that

$$\psi_{k,j}^{\varepsilon,\mu}(x) = 0 \iff \text{grad}^{\mathbb{R}^{n+1}}(f_k)(x) = 0.$$

For example one finds eigenspinors which vanish to high order at some point, i. e. the spinor and all of its derivatives up to a finite order are zero at this point. Namely given $m \in \mathbb{N} \setminus \{0\}$ we take

$$f_{m+1}(x_1, \dots, x_{n+1}) := \text{Re}((x_1 + ix_2)^{m+1}).$$

Then the spinor $\psi_{m+1,1}^{--}$ on S^n defined by

$$\psi_{m+1,1}^{--} := -(m+1)f_{m+1}\varphi_{-1,1} + \text{grad}^{S^n}(f_{m+1}) \cdot \varphi_{-1,1}$$

is an eigenspinor for the value $-\frac{n}{2} - m$ and with $p = e_{n+1} \in S^n$ we have $\nabla^r \psi_{m+1,1}^{--}(p) = 0$ for $0 \leq r \leq m-1$.

We also find spinors whose zero set is not a submanifold of S^n . As an example consider $n = 3$ and take $f_3(x_1, \dots, x_4) := x_1x_2x_3$. Then we have

$$\text{grad}^{\mathbb{R}^4}(f_3) = (x_2x_3, x_1x_3, x_1x_2, 0).$$

Therefore $\psi_{3,1}^{--}$ has the zero set

$$\{x \in S^3 \mid x_1 = x_2 = 0 \text{ or } x_1 = x_3 = 0 \text{ or } x_2 = x_3 = 0\}$$

which consists of three copies of S^1 intersecting in $\pm e_4$.

4.3 Flat tori

We will give some explicit computations of zero sets of eigenspinors on flat tori. The reader, who is only interested in the main results, may skip this section.

Let $\Gamma \subset \mathbb{R}^n$ and $M = \mathbb{R}^n/\Gamma$, $n \geq 2$ and let g be the metric on M induced by the Euclidean metric on \mathbb{R}^n . The spin structures on (M, g) are classified by

$$H^1(M, \mathbb{Z}/2\mathbb{Z}) \cong \text{Hom}(\Gamma, \mathbb{Z}/2\mathbb{Z})$$

and thus there are 2^n distinct spin structures. The calculation of the eigenvalues of the Dirac operator on (M, g) for all spin structures has been carried out in [F2]. It is also explained e. g. in [G]. Denote by Γ^* the dual lattice. Let $(\gamma_j)_{j=1}^n$ be a basis of Γ and let $(\gamma_j^*)_{j=1}^n$ be the dual basis. The spin structures will be denoted by the tuples $(\delta_1, \dots, \delta_n)$, where $\delta_j \in \{0, 1\}$ is the image of γ_j under a homomorphism from Γ to $\mathbb{Z}/2\mathbb{Z}$, $j \in \{1, \dots, n\}$. For a fixed spin structure $(\delta_1, \dots, \delta_n)$ we define the one-form δ on \mathbb{R}^n by $\delta = \frac{1}{2} \sum_{j=1}^n \delta_j \gamma_j^*$. Then the result is the following:

Theorem 4.3.1. *The eigenvalues of the Dirac operator on $(\mathbb{R}^n/\Gamma, g)$ with the spin structure $(\delta_1, \dots, \delta_n)$ are*

$$\{\pm 2\pi|\alpha + \delta| \mid \alpha \in \Gamma^*\}.$$

In case $\delta_1 = \dots = \delta_n = 0$ the multiplicity of the eigenvalue 0 is $2^{\lfloor n/2 \rfloor}$. For every other spin structure 0 is not an eigenvalue. For the spin structure $(\delta_1, \dots, \delta_n)$ the multiplicity of every non-zero eigenvalue λ is

$$2^{\lfloor n/2 \rfloor - 1} \#\{\alpha \in \Gamma^* \mid 2\pi|\alpha + \delta| = |\lambda|\}.$$

In order to obtain the eigenspinors explicitly we summarize the proof. It is well known that the spinors on M equipped with the spin structure $(\delta_1, \dots, \delta_n)$ are in one-to-one-correspondence with the spinors ψ on \mathbb{R}^n which satisfy the condition

$$\psi(x + \gamma_j) = (-1)^{\delta_j} \psi(x) \quad (4.1)$$

for all $x \in \mathbb{R}^n$ and all $j \in \{1, \dots, n\}$ (see e.g. [G]). We fix a spin structure $(\delta_1, \dots, \delta_n)$ on M . If $\delta = 0$, we see that every constant spinor on \mathbb{R}^n satisfies the equivariance condition (4.1) and thus is a harmonic spinor on (M, g, Θ) . Therefore we have $\dim \ker(D^g) = 2^{\lfloor n/2 \rfloor}$. Let $\alpha \in \Gamma^*$ and assume that $\alpha \neq 0$ or that $\delta \neq 0$. Then $\alpha + \delta \neq 0$ and the action of the one-form $i \frac{\alpha + \delta}{|\alpha + \delta|}$ on $\Sigma \mathbb{R}^n$ is parallel, unitary and an involution. Thus one obtains a parallel and orthogonal splitting of the spinor bundle

$$\Sigma \mathbb{R}^n = \Sigma_\alpha^+ \mathbb{R}^n \oplus \Sigma_\alpha^- \mathbb{R}^n,$$

where $\Sigma_\alpha^\pm \mathbb{R}^n$ are the eigenbundles corresponding to the eigenvalue ± 1 . They are isomorphic, since the Clifford multiplications with two orthogonal vectors anticommute. Note that

$$\Sigma_{-\alpha-2\delta}^\pm \mathbb{R}^n = \Sigma_\alpha^\mp \mathbb{R}^n.$$

For every $\mu \in \{\pm 1\}$ we choose sections $\varphi_{\alpha,j}^\mu$ of $\Sigma_\alpha^\mu \mathbb{R}^n$, $1 \leq j \leq 2^{\lfloor n/2 \rfloor - 1}$, which are constant and pointwise orthonormal. Then for all $\mu \in \{\pm 1\}$ the spinor $\psi_{\alpha,j}^\mu$ defined by

$$\psi_{\alpha,j}^\mu(x) := e^{2\pi i(\alpha + \delta)(x)} \varphi_{\alpha,j}^\mu$$

satisfies the equivariance condition (4.1) and is an eigenspinor of D^g for the value $2\pi\mu|\alpha + \delta|$. Since the functions $\{e^{2\pi i\alpha} | \alpha \in \Gamma^*\}$ form a Hilbert basis of $L^2(M, \mathbb{C})$, it follows that the spinors $\psi_{\alpha,j}^\mu$ form a Hilbert basis of $L^2(\Sigma^g M)$.

Concerning the zero set of eigenspinors consider first the so called trivial spin structure $\delta = 0$. It is the only spin structure which admits harmonic spinors. These are parallel spinors and they are nowhere zero. Next consider an arbitrary spin structure. In order to investigate the zero set of eigenspinors to nonzero eigenvalues first note that for any nonzero $\alpha \in \Gamma^*$ the spinors

$$\psi_{\alpha,1}^+, \dots, \psi_{\alpha,2^{\lfloor n/2 \rfloor - 1}}^+, \psi_{-\alpha-2\delta,1}^+, \dots, \psi_{-\alpha-2\delta,2^{\lfloor n/2 \rfloor - 1}}^+$$

are eigenspinors corresponding to $2\pi|\alpha + \delta|$ and they are pointwise linearly independent. Thus any non trivial linear combination of them is nowhere zero. These spinors are contributed by the elements $\alpha, -\alpha - 2\delta$ of Γ^* . Assume now that there is $\alpha' \notin \{\alpha, -\alpha - 2\delta\}$ with $|\alpha' + \delta| = |\alpha + \delta|$. Let φ be a

parallel spinor on \mathbb{R}^n such that $i \frac{\alpha'+\delta}{|\alpha'+\delta|} \cdot \varphi = \varphi$ and write $\varphi = \varphi^+ + \varphi^-$ where $i \frac{\alpha+\delta}{|\alpha+\delta|} \cdot \varphi^\pm = \pm \varphi^\pm$. Then the spinor ψ defined by

$$\begin{aligned} \psi(x) &= e^{2\pi i(\alpha+\delta)(x)} \varphi^+ + e^{-2\pi i(\alpha+\delta)(x)} \varphi^- - e^{2\pi i(\alpha'+\delta)(x)} \varphi \\ &= (e^{2\pi i(\alpha+\delta)(x)} - e^{2\pi i(\alpha'+\delta)(x)}) \varphi^+ + (e^{-2\pi i(\alpha+\delta)(x)} - e^{2\pi i(\alpha'+\delta)(x)}) \varphi^- \end{aligned}$$

is an eigenspinor corresponding to the value $2\pi|\alpha + \delta|$ and has the zero set

$$\{[x] \in M \mid (\alpha + \delta)(x), (\alpha' + \delta)(x) \in \mathbb{Z} \text{ or } (\alpha + \delta)(x), (\alpha' + \delta)(x) \in \frac{1}{2} + \mathbb{Z}\}.$$

This is a submanifold of codimension 2.

Chapter 5

Green's function for the Dirac operator

5.1 Trivialization of the spinor bundle

Let (M, g, Θ) be a Riemannian spin manifold of dimension n . In this section we explain a certain local trivialization of the spinor bundle $\Sigma^g M$, which we will use later. In the literature (e.g. [AGHM], [AH]) it is known as the Bourguignon-Gauduchon trivialization.

Let $p \in M$ and let $W \subset T_p M$ be an open neighborhood of 0 such that the restriction of the exponential map \exp_p to W is a diffeomorphism. Let $I: (\mathbb{R}^n, g_{\text{eucl}}) \rightarrow (T_p M, g)$ be a linear isometry, let $U := \exp_p(W) \subset M$ and let $V := I^{-1}(W)$. Then

$$\rho := \exp_p \circ I|_V : V \rightarrow U$$

defines a local parametrization of M by Riemannian normal coordinates. For $x \in V$ we use the canonical isomorphism $T_x V \cong \mathbb{R}^n$ and in this way we obtain the Euclidean scalar product g_{eucl} on $T_x V$. In addition we have the scalar product $\rho^* g$ on $T_x V$ defined by

$$(\rho^* g)(v, w) = g(d\rho|_x v, d\rho|_x w),$$

where $v, w \in T_x V$. There exists $G_x \in \text{End}(T_x V)$, such that for all vectors $v, w \in T_x V$ we have

$$(\rho^* g)(v, w) = g_{\text{eucl}}(G_x v, w)$$

and G_x is g_{eucl} -self-adjoint and positive definite. There is a unique positive definite endomorphism $B_x \in \text{End}(T_x V)$ such that we have $B_x^2 = G_x^{-1}$. If $(E_i)_{i=1}^n$ is any g_{eucl} -orthonormal basis of $T_x V$, then $(B_x E_i)_{i=1}^n$ is a $\rho^* g$ -orthonormal basis of $T_x V$. Therefore the vectors $d\rho|_x B_x E_i$, $1 \leq i \leq n$, form

a g -orthonormal basis of $T_{\rho(x)}M$. We assemble the maps B_x to obtain a vector bundle endomorphism B of TV and we define

$$b : TV \rightarrow TM|_U, \quad b = d\rho \circ B.$$

From this we obtain an isomorphism of principal $\mathrm{SO}(n)$ -bundles

$$P_{\mathrm{SO}}(V, g_{\mathrm{eucl}}) \rightarrow P_{\mathrm{SO}}(U, g), \quad (E_i)_{i=1}^n \mapsto (b(E_i))_{i=1}^n,$$

which lifts to an isomorphism

$$c : P_{\mathrm{Spin}}(V, g_{\mathrm{eucl}}) \rightarrow P_{\mathrm{Spin}}(U, g)$$

of principal $\mathrm{Spin}(n)$ -bundles. We define

$$\beta : \Sigma\mathbb{R}^n|_V \rightarrow \Sigma^g M|_U, \quad [s, \sigma] \mapsto [c(s), \sigma].$$

This gives an identification of the spinor bundles, which is a fibrewise isometry with respect to the bundle metrics on $\Sigma\mathbb{R}^n|_V$ and on $\Sigma^g M|_U$. Furthermore for all $X \in TV$ and for all $\varphi \in \Sigma\mathbb{R}^n|_V$ we have $\beta(X \cdot \varphi) = b(X) \cdot \beta(\varphi)$. We obtain an isomorphism

$$A : C^\infty(\Sigma^g M|_U) \rightarrow C^\infty(\Sigma\mathbb{R}^n|_V), \quad \psi \mapsto \beta^{-1} \circ \psi \circ \rho,$$

which sends a spinor on U to the corresponding spinor in the trivialization. Let ∇^g and ∇ denote the Levi Civita connections on (U, g) resp. on (V, g_{eucl}) as well as its lifts to $\Sigma^g M|_U$ and $\Sigma\mathbb{R}^n|_V$. Let $(e_i)_{i=1}^n$ be a positively oriented orthonormal frame of $TM|_U$. Then for all $\psi \in C^\infty(\Sigma^g M|_U)$ we have by the formula (2.3)

$$\nabla_{e_i}^g \psi = \partial_{e_i} \psi + \frac{1}{4} \sum_{j,k=1}^n \tilde{\Gamma}_{ij}^k e_j \cdot e_k \cdot \psi,$$

where

$$\tilde{\Gamma}_{ij}^k := g(\nabla_{e_i}^g e_j, e_k).$$

In particular we can take the standard basis $(E_i)_{i=1}^n$ of \mathbb{R}^n and put $e_i := b(E_i)$, $1 \leq i \leq n$. We define the matrix coefficients B_i^j by $B(E_i) = \sum_{j=1}^n B_i^j E_j$. It follows that

$$\begin{aligned} A\nabla_{e_i}^g \psi &= \nabla_{d\rho^{-1}(e_i)} A\psi + \frac{1}{4} \sum_{j,k=1}^n \tilde{\Gamma}_{ij}^k E_j \cdot E_k \cdot A\psi \\ &= \nabla_{B(E_i)} A\psi + \frac{1}{4} \sum_{j,k=1}^n \tilde{\Gamma}_{ij}^k E_j \cdot E_k \cdot A\psi \\ &= \nabla_{E_i} A\psi + \sum_{j=1}^n (B_i^j - \delta_i^j) \nabla_{E_j} A\psi + \frac{1}{4} \sum_{j,k=1}^n \tilde{\Gamma}_{ij}^k E_j \cdot E_k \cdot A\psi. \end{aligned} \quad (5.1)$$

Hence we obtain

$$\begin{aligned} AD^g\psi &= D^{g_{\text{eucl}}}A\psi + \sum_{i,j=1}^n (B_i^j - \delta_i^j)E_i \cdot \nabla_{E_j}A\psi \\ &\quad + \frac{1}{4} \sum_{i,j,k=1}^n \tilde{\Gamma}_{ij}^k E_i \cdot E_j \cdot E_k \cdot A\psi. \end{aligned} \quad (5.2)$$

Let $\partial_j := d\rho(E_j)$, $1 \leq j \leq n$, be the coordinate vector fields of the normal coordinates. It is well known that the Taylor expansion of

$$g_{ij} : V \rightarrow \mathbb{R}, \quad g_{ij}(x) := g(\partial_i|_{\rho(x)}, \partial_j|_{\rho(x)})$$

around 0 is given by

$$g_{ij}(x) = \delta_{ij} + \frac{1}{3} \sum_{a,b=1}^n R_{iabj}(p)x_a x_b + O(|x|^3),$$

where $R_{iabj} = g(R(\partial_b, \partial_j)\partial_a, \partial_i)$ denotes the components of the Riemann curvature tensor (see e. g. [LP], p. 61). Since we have $(B_i^j)_{ij} = (g_{ij})_{ij}^{-1/2}$ it follows that

$$B_i^j(x) = \delta_i^j - \frac{1}{6} \sum_{a,b=1}^n R_{iabj}(p)x_a x_b + O(|x|^3). \quad (5.3)$$

By definition we have

$$e_i = b(E_i) = d\rho(B(E_i)) = \sum_{j=1}^n B_i^j d\rho(E_j) = \sum_{j=1}^n B_i^j \partial_j.$$

We define the Christoffel symbols Γ_{kr}^m by

$$\nabla_{\partial_k}^g \partial_r = \sum_{m=1}^n \Gamma_{kr}^m \partial_m.$$

We obtain

$$\begin{aligned} \sum_{k=1}^n \tilde{\Gamma}_{ij}^k e_k &= \sum_{s,m=1}^n B_i^s \nabla_{\partial_s}^g (B_j^m \partial_m) \\ &= \sum_{s,m=1}^n B_i^s (\partial_s B_j^m + \sum_{r=1}^n B_j^r \Gamma_{sr}^m) \partial_m \\ &= \sum_{s,m,k=1}^n B_i^s (\partial_s B_j^m + \sum_{r=1}^n B_j^r \Gamma_{sr}^m) (B^{-1})_m^k e_k. \end{aligned}$$

From the formula

$$\Gamma_{kr}^m = \frac{1}{2} \sum_{s=1}^n g^{ms} (\partial_k g_{rs} + \partial_r g_{ks} - \partial_s g_{kr})$$

it follows that $\Gamma_{kr}^m(x) = O(|x|)$ and thus

$$\tilde{\Gamma}_{kr}^m(x) = O(|x|) \quad (5.4)$$

as $x \rightarrow 0$ for all m, k, r .

5.2 The Euclidean Dirac operator

The aim of this section is to calculate preimages under the Dirac operator of certain spinors on $\mathbb{R}^n \setminus \{0\}$ with the Euclidean metric. The results will be useful for obtaining the expansion of Green's function for the Dirac operator on a closed spin manifold.

Definition 5.2.1. For $k \in \mathbb{R}$, $m \in \mathbb{N}$ and $i \in \{0, 1\}$ we define the vector subspaces $P_{k,m,i}(\mathbb{R}^n)$ of $C^\infty(\Sigma\mathbb{R}^n|_{\mathbb{R}^n \setminus \{0\}})$ as follows. For $k \neq 0$ define

$$P_{k,m,0}(\mathbb{R}^n) := \text{span} \left\{ x \mapsto x_{i_1} \dots x_{i_m} |x|^k \gamma \mid \begin{array}{l} 1 \leq i_1, \dots, i_m \leq n, \\ \gamma \in \Sigma_n \text{ constant} \end{array} \right\}$$

$$P_{k,m,1}(\mathbb{R}^n) := \text{span} \left\{ x \mapsto x_{i_1} \dots x_{i_m} |x|^k x \cdot \gamma \mid \begin{array}{l} 1 \leq i_1, \dots, i_m \leq n, \\ \gamma \in \Sigma_n \text{ constant} \end{array} \right\}$$

and furthermore define

$$P_{0,m,0}(\mathbb{R}^n) := \text{span} \left\{ x \mapsto x_{i_1} \dots x_{i_m} \ln|x| \gamma \mid \begin{array}{l} 1 \leq i_1, \dots, i_m \leq n, \\ \gamma \in \Sigma_n \text{ constant} \end{array} \right\}$$

$$P_{0,m,1}(\mathbb{R}^n) := \text{span} \left\{ x \mapsto x_{i_1} \dots x_{i_m} (1 - n \ln|x|) x \cdot \gamma \mid \begin{array}{l} 1 \leq i_1, \dots, i_m \leq n, \\ \gamma \in \Sigma_n \text{ constant} \end{array} \right\}.$$

Note that there exist inclusions among these spaces. For example one has $P_{k+2,m,0}(\mathbb{R}^n) \subset P_{k,m+2,0}(\mathbb{R}^n)$ for all m and all $k \neq -2$. However in the following we will very often not use these inclusions, since exceptions like the case $k = -2$ in this example would make the following statements rather more complicated. The exception $k = -2$ in this example is due to our definition of $P_{0,m,0}(\mathbb{R}^n)$. The following proposition will show that this definition is nevertheless useful.

Proposition 5.2.2. *For all $m \in \mathbb{N}$, $k \in \mathbb{R}$ with $-n \leq k$ and $-n < k+m \leq 0$ we have*

$$P_{k,m,0}(\mathbb{R}^n) \subset D^{g_{\text{eucl}}} \left(\sum_{j=1}^{\lfloor (m+1)/2 \rfloor} P_{k+2j,m+1-2j,0}(\mathbb{R}^n) + \sum_{j=0}^{\lfloor m/2 \rfloor} P_{k+2j,m-2j,1}(\mathbb{R}^n) \right)$$

For all $m \in \mathbb{N}$, $k \in \mathbb{R}$ with $-n \leq k$ and $-n < k+m+1 \leq 0$ we have

$$P_{k,m,1}(\mathbb{R}^n) \subset D^{g_{\text{eucl}}} \left(\sum_{j=0}^{\lfloor m/2 \rfloor} P_{k+2+2j,m-2j,0}(\mathbb{R}^n) + \sum_{j=1}^{\lfloor (m+1)/2 \rfloor} P_{k+2j,m+1-2j,1}(\mathbb{R}^n) \right).$$

Proof. We use induction on m . Let $m = 0$ and let γ be a constant spinor on $(\mathbb{R}^n, g_{\text{eucl}})$. We want to prove that $P_{k,0,0}(\mathbb{R}^n) \subset D^{g_{\text{eucl}}}(P_{k,0,1}(\mathbb{R}^n))$ for all k with $-n < k \leq 0$ and $P_{k,0,1}(\mathbb{R}^n) \subset D^{g_{\text{eucl}}}(P_{k+2,0,0}(\mathbb{R}^n))$ for all k with $-n \leq k$ and $k+1 \leq 0$. One calculates easily that

$$\begin{aligned} D^{g_{\text{eucl}}} \left(-\frac{1}{n+k} |x|^k x \cdot \gamma \right) &= |x|^k \gamma, \quad k \neq -n \\ D^{g_{\text{eucl}}} \left(\frac{1-n \ln |x|}{n^2} x \cdot \gamma \right) &= \ln |x| \gamma, \\ D^{g_{\text{eucl}}} \left(\frac{1}{k+2} |x|^{k+2} \gamma \right) &= |x|^k x \cdot \gamma, \quad k \neq -2 \\ D^{g_{\text{eucl}}} (\ln |x| \gamma) &= |x|^{-2} x \cdot \gamma. \end{aligned}$$

Using the definition of $P_{k,0,i}(\mathbb{R}^n)$ one finds that the assertion holds for $m = 0$.

Let $m \geq 1$ and assume that all the inclusions in the assertion hold for $m-1$. Using the equation $E_i \cdot x = -2x_i - x \cdot E_i$ we find

$$\begin{aligned} &D^{g_{\text{eucl}}} (-x_{i_1} \dots x_{i_m} |x|^k x \cdot \gamma) \\ &= -\sum_{j=1}^m x_{i_1} \dots \widehat{x_{i_j}} \dots x_{i_m} |x|^k E_{i_j} \cdot x \cdot \gamma - x_{i_1} \dots x_{i_m} D^{g_{\text{eucl}}} (|x|^k x \cdot \gamma) \\ &= (2m+n+k) x_{i_1} \dots x_{i_m} |x|^k \gamma + \sum_{j=1}^m x_{i_1} \dots \widehat{x_{i_j}} \dots x_{i_m} |x|^k x \cdot E_{i_j} \cdot \gamma. \end{aligned}$$

Since $E_{i_j} \cdot \gamma$ is a parallel spinor the sum on the right hand side is contained in $P_{k,m-1,1}(\mathbb{R}^n)$. We apply the induction hypothesis and since $2m+n+k \neq 0$ we find that the assertion for $P_{k,m,0}(\mathbb{R}^n)$ holds. We define $f_k(x) := \frac{1}{k} |x|^k$ for $k \neq 0$ and $f_0(x) := \ln |x|$. Then we find

$$\begin{aligned} &D^{g_{\text{eucl}}} (x_{i_1} \dots x_{i_m} f_{k+2}(x) \gamma) \\ &= x_{i_1} \dots x_{i_m} |x|^k x \cdot \gamma + \sum_{j=1}^m x_{i_1} \dots \widehat{x_{i_j}} \dots x_{i_m} f_{k+2}(x) E_{i_j} \cdot \gamma. \end{aligned}$$

The sum on the right hand side is in $P_{k+2,m-1,0}(\mathbb{R}^n)$. Again we apply the induction hypothesis and we find that the assertion for $P_{k,m,1}(\mathbb{R}^n)$ holds. \square

5.3 Expansion of Green's function

Let (M, g, Θ) be a closed Riemannian spin manifold of dimension n and let $\lambda \in \mathbb{R}$. Since $D^g - \lambda$ is a self-adjoint elliptic differential operator, there is an L^2 -orthogonal decomposition

$$C^\infty(\Sigma^g M) = \ker(D^g - \lambda) \oplus \text{im}(D^g - \lambda).$$

Let $P: C^\infty(\Sigma^g M) \rightarrow \ker(D^g - \lambda)$ denote the L^2 -orthogonal projection. Then there is an operator $G: C^\infty(\Sigma^g M) \rightarrow C^\infty(\Sigma^g M)$, called Green's operator such that

$$G(D^g - \lambda) = (D^g - \lambda)G = \text{Id} - P$$

and it extends to a bounded linear operator $H^k(\Sigma^g M) \rightarrow H^{k+1}(\Sigma^g M)$ for every $k \in \mathbb{N}$ (see e.g. [LM], p. 195). In this section we will examine the integral kernel of Green's operator, which is called Green's function for $D^g - \lambda$. Let $\pi_i: M \times M \rightarrow M$, $i = 1, 2$ be the projections. We define

$$\Sigma^g M \boxtimes \Sigma^g M^* := \pi_1^* \Sigma^g M \otimes (\pi_2^* \Sigma^g M)^*$$

i. e. $\Sigma^g M \boxtimes \Sigma^g M^*$ is the vector bundle over $M \times M$ whose fibre over the point $(x, y) \in M \times M$ is given by $\text{Hom}_{\mathbb{C}}(\Sigma_y^g M, \Sigma_x^g M)$. Let $\Delta := \{(x, x) | x \in M\}$ be the diagonal. In the following we will abbreviate

$$\int_{M \setminus \{p\}} := \lim_{\varepsilon \rightarrow 0} \int_{M \setminus B_\varepsilon(p)}.$$

Definition 5.3.1. A smooth section $G_\lambda^g: M \times M \setminus \Delta \rightarrow \Sigma^g M \boxtimes \Sigma^g M^*$ which is locally integrable on $M \times M$ is called a Green's function for $D^g - \lambda$ if for all $p \in M$, for all $\varphi \in \Sigma_p^g M$ and for all $\psi \in \text{im}(D^g - \lambda)$ we have

$$\int_{M \setminus \{p\}} \langle (D^g - \lambda)\psi, G_\lambda^g(\cdot, p)\varphi \rangle dv^g = \langle \psi(p), \varphi \rangle, \quad (5.5)$$

and if for all $p \in M$, for all $\varphi \in \Sigma_p^g M$ and for all $\psi \in \ker(D^g - \lambda)$ we have

$$\int_{M \setminus \{p\}} \langle \psi, G_\lambda^g(\cdot, p)\varphi \rangle dv^g = 0. \quad (5.6)$$

In this section we will prove existence and uniqueness of Green's function for $D^g - \lambda$ in such a way that we also obtain the expansion of Green's function around the singularity. The smooth spinor $G_\lambda^g(\cdot, p)\varphi$ on $M \setminus \{p\}$ will sometimes also be called Green's function for φ . Thus we have for all $\psi \in C^\infty(\Sigma^g M)$ and for all $\varphi \in \Sigma_p^g M$

$$\int_{M \setminus \{p\}} \langle (D^g - \lambda)\psi, G_\lambda^g(\cdot, p)\varphi \rangle dv^g = \langle \psi(p) - P\psi(p), \varphi \rangle. \quad (5.7)$$

On Euclidean space we define a Green's function as follows.

Definition 5.3.2. *Let $(M, g) = (\mathbb{R}^n, g_{\text{eucl}})$ with the unique spin structure. A smooth section $G_\lambda^g: M \times M \setminus \Delta \rightarrow \Sigma^g M \boxtimes \Sigma^g M^*$ which is locally integrable on $M \times M$ is called a Green's function for $D^g - \lambda$ if for all $p \in M$, for all $\varphi \in \Sigma_p^g M$ and for all $\psi \in C^\infty(\Sigma^g M)$ with compact support the equation (5.5) holds.*

Of course a Green's function for $D^{g_{\text{eucl}}} - \lambda$ is not uniquely determined by this definition. We will explicitly write down a Green's function for $D^{g_{\text{eucl}}} - \lambda$ and use it later to find the expansion of Green's function for $D^g - \lambda$ on a closed Riemannian spin manifold (M, g, Θ) around the diagonal. First observe that for every spinor $\chi \in C^\infty(\Sigma \mathbb{R}^n|_{\mathbb{R}^n \setminus \{0\}})$ and for every $\lambda \in \mathbb{R}$ the equation

$$(D^{g_{\text{eucl}}} - \lambda)(D^{g_{\text{eucl}}} + \lambda)\chi = - \sum_{i=1}^n \nabla_{E_i} \nabla_{E_i} \chi - \lambda^2 \chi$$

holds on $\mathbb{R}^n \setminus \{0\}$. Therefore if γ is a constant spinor on $(\mathbb{R}^n, g_{\text{eucl}})$ and if $f \in C^\infty(\mathbb{R}^n \setminus \{0\}, \mathbb{R})$ satisfies

$$- \sum_{i=1}^n \frac{\partial^2 f}{\partial x_i^2} - \lambda^2 f = \delta_0,$$

then the spinor $G_\lambda^{g_{\text{eucl}}}(\cdot, 0)\gamma := (D^{g_{\text{eucl}}} + \lambda)(f\gamma)$ is a Green's function. Writing $f(x) = g(|x|)$ for a function g of one variable we get a solution if g solves the ordinary differential equation

$$g''(z) + \frac{n-1}{z}g'(z) + \lambda^2 g(z) = -\delta_0. \quad (5.8)$$

In the following let Γ denote the Gamma function and J_m, Y_m the Bessel functions of the first and second kind for the parameter $m \in \mathbb{R}$. In the

notation of [AS], p. 360 they are defined for $z \in (0, \infty)$ by

$$\begin{aligned} J_m(z) &= \frac{1}{2^m \Gamma(m+1)} z^m \left(1 + \sum_{k=1}^{\infty} a_k z^{2k}\right), \quad m \in \mathbb{R}, \\ Y_0(z) &= \frac{2}{\pi} \left(\ln\left(\frac{z}{2}\right) + c\right) J_0(z) + \sum_{k=1}^{\infty} b_k z^{2k}, \\ Y_m(z) &= -\frac{2^m}{\pi} \Gamma(m) z^{-m} \left(1 + \sum_{k=1}^{\infty} c_k z^{2k}\right), \quad m = \frac{1}{2} + k, k \in \mathbb{N}, \\ Y_m(z) &= -\frac{2^m}{\pi} \Gamma(m) z^{-m} \left(1 + \sum_{k=1}^{\infty} d_k z^{2k}\right) + \frac{2}{\pi} \ln\left(\frac{z}{2}\right) J_m(z), \quad m \in \mathbb{N} \setminus \{0\}, \end{aligned}$$

where c is a real constant, the a_k, b_k, c_k, d_k are real coefficients, the a_k, c_k, d_k depend on m and all the power series converge for all $z \in (0, \infty)$. Let $\omega_{n-1} = \text{vol}(S^{n-1}, g_{\text{can}})$ be the volume of the $(n-1)$ -dimensional unit sphere with the standard metric.

Theorem 5.3.3. *Let $m := \frac{n-2}{2}$. We define $f_\lambda: \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{R}$ as follows. For $\lambda \neq 0$ and $n = 2$*

$$f_\lambda(x) := -\frac{1}{4} Y_0(|\lambda x|) + \frac{\ln|\lambda| - \ln(2) + c}{2\pi} J_0(|\lambda x|),$$

for $\lambda \neq 0$ and odd $n \geq 3$

$$f_\lambda(x) := -\frac{\pi|\lambda|^m}{2^m \Gamma(m)(n-2)\omega_{n-1}} |x|^{-m} Y_m(|\lambda x|),$$

for $\lambda \neq 0$ and even $n \geq 4$

$$f_\lambda(x) := -\frac{\pi|\lambda|^m}{2^m \Gamma(m)(n-2)\omega_{n-1}} |x|^{-m} \left(Y_m(|\lambda x|) - \frac{2(\ln|\lambda| - \ln(2))}{\pi} J_m(|\lambda x|) \right)$$

and

$$f_0(x) := -\frac{1}{2\pi} \ln|x|, \quad n = 2, \quad f_0(x) := \frac{1}{(n-2)\omega_{n-1}|x|^{n-2}}, \quad n \geq 3.$$

Then for every constant spinor γ on \mathbb{R}^n a Green's function for $D^{g_{\text{eucl}}} - \lambda$ is given by

$$G_\lambda^{g_{\text{eucl}}}(x, 0)\gamma = (D^{g_{\text{eucl}}} + \lambda)(f_\lambda \gamma)(x).$$

Corollary 5.3.4. *For every constant spinor $\gamma \in \Sigma_n$ there exists a Green's function of $D^{g_{\text{eucl}}} - \lambda$, which has the following form. For $n = 2$*

$$G_\lambda^{g_{\text{eucl}}}(x, 0)\gamma = -\frac{1}{2\pi|x|} \frac{x}{|x|} \cdot \gamma - \frac{\lambda}{2\pi} \ln|x| \gamma + \ln|x| \vartheta_\lambda(x) + \zeta_\lambda(x),$$

for odd $n \geq 3$

$$G_\lambda^{g_{\text{eucl}}}(x, 0)\gamma = -\frac{1}{\omega_{n-1}|x|^{n-1}} \frac{x}{|x|} \cdot \gamma + \frac{\lambda}{(n-2)\omega_{n-1}|x|^{n-2}} \gamma + |x|^{2-n} \zeta_\lambda(x),$$

for even $n \geq 4$

$$\begin{aligned} G_\lambda^{g_{\text{eucl}}}(x, 0)\gamma &= -\frac{1}{\omega_{n-1}|x|^{n-1}} \frac{x}{|x|} \cdot \gamma + \frac{\lambda}{(n-2)\omega_{n-1}|x|^{n-2}} \gamma + |x|^{2-n} \zeta_\lambda(x) \\ &\quad - \frac{\lambda^{n-1}}{2^{n-2} \Gamma(\frac{n}{2})^2 \omega_{n-1}} \ln|x| \gamma + \ln|x| \vartheta_\lambda(x), \end{aligned}$$

where for every n and for every λ the spinors ϑ_λ , ζ_λ extend smoothly to \mathbb{R}^n and satisfy

$$|\zeta_\lambda(x)|_{g_{\text{eucl}}} = O(|x|), \quad |\vartheta_\lambda(x)|_{g_{\text{eucl}}} = O(|x|) \quad \text{as } x \rightarrow 0$$

and where for every n and for every x the spinors $\vartheta_\lambda(x), \zeta_\lambda(x) \in \Sigma_n$ are power series in λ with $\vartheta_0(x) = \zeta_0(x) = 0$.

Proof of Corollary 5.3.4. We find

$$Y_0(|\lambda x|) - \frac{2(\ln|\lambda| - \ln(2) + c)}{\pi} J_0(|\lambda x|) = \frac{2}{\pi} \ln|x| J_0(|\lambda x|) + \sum_{k=1}^{\infty} b_k |\lambda x|^{2k}$$

and for $m \in \mathbb{N} \setminus \{0\}$

$$\begin{aligned} Y_m(|\lambda x|) - \frac{2(\ln|\lambda| - \ln(2))}{\pi} J_m(|\lambda x|) &= -\frac{2^m \Gamma(m)}{\pi |\lambda x|^m} \left(1 + \sum_{k=1}^{\infty} d_k |\lambda x|^{2k}\right) \\ &\quad + \frac{2}{\pi} \ln|x| J_m(|\lambda x|). \end{aligned}$$

Thus we find for $n = 2$

$$f_\lambda(x) = -\frac{1}{2\pi} \ln|x| \left(1 + \sum_{k=1}^{\infty} a_k |\lambda x|^{2k}\right) - \frac{1}{4} \sum_{k=1}^{\infty} b_k |\lambda x|^{2k},$$

for odd $n \geq 3$

$$f_\lambda(x) = \frac{1}{(n-2)\omega_{n-1}|x|^{n-2}} \left(1 + \sum_{k=1}^{\infty} c_k |\lambda x|^{2k}\right)$$

and for even $n \geq 4$

$$\begin{aligned} f_\lambda(x) &= \frac{1}{(n-2)\omega_{n-1}|x|^{n-2}} \left(1 + \sum_{k=1}^{\infty} d_k |\lambda x|^{2k}\right) \\ &\quad - \frac{\lambda^{n-2}}{2^{n-2}(n!)^2\omega_{n-1}} \ln|x| \left(1 + \sum_{k=1}^{\infty} a_k |\lambda x|^{2k}\right). \end{aligned}$$

The assertion follows. \square

Proof of Theorem 5.3.3. Let f_λ be as in the assertion and write $f_\lambda(x) = g_\lambda(|x|)$ with $g_\lambda: (0, \infty) \rightarrow \mathbb{R}$. One finds that g_λ solves the equation (5.8). It remains to show that $(D^{g_{\text{eucl}}} + \lambda)(f_\lambda\gamma)$ satisfies (5.5). The calculation of the proof of the Corollary shows

$$(D^{g_{\text{eucl}}} + \lambda)(f_\lambda\gamma)(x) = -\frac{1}{\omega_{n-1}} \frac{x}{|x|^n} \cdot \gamma + \zeta(x),$$

where $|\zeta(x)|_{g_{\text{eucl}}} = o(|x|^{1-n})$ as $x \rightarrow 0$. It is well known (e. g. [LM], p. 115) that for a Riemannian spin manifold (M, g, Θ) with boundary $\partial M \neq \emptyset$ and ψ, φ compactly supported spinors we have

$$(D^g\psi, \varphi)_2 - (\psi, D^g\varphi)_2 = \int_{\partial M} \langle \nu \cdot \psi, \varphi \rangle dA,$$

where ν is the outer unit normal vector field on ∂M . We apply this to $(\mathbb{R}^n \setminus B_\varepsilon(0), g_{\text{eucl}})$ and $\nu(x) := -\frac{x}{|x|}$ and we find

$$\begin{aligned} &\int_{\mathbb{R}^n \setminus B_\varepsilon(0)} \langle (D^{g_{\text{eucl}}} - \lambda)\psi(x), (D^{g_{\text{eucl}}} + \lambda)(f_\lambda\gamma)(x) \rangle dv^g \\ &= - \int_{\partial B_\varepsilon(0)} \left\langle \frac{x}{|x|} \cdot \psi(x), (D^{g_{\text{eucl}}} + \lambda)(f_\lambda\gamma)(x) \right\rangle dA \\ &= \int_{\partial B_\varepsilon(0)} \left\langle \psi(x), \frac{1}{\omega_{n-1}|x|^{n-1}}\gamma + \frac{x}{|x|} \cdot \zeta(x) \right\rangle dA. \end{aligned}$$

With the substitution $x = \varepsilon y$ we find that

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^n \setminus B_\varepsilon(0)} \langle (D^{g_{\text{eucl}}} - \lambda)\psi(x), (D^{g_{\text{eucl}}} + \lambda)(f_\lambda\gamma)(x) \rangle dv^g = \langle \psi(0), \gamma \rangle.$$

The assertion follows. \square

Remark 5.3.5. In the case $n = 3$ we get more familiar expressions by using that

$$\sqrt{\frac{\pi}{2z}} Y_{1/2}(z) = -\frac{\cos(z)}{z}$$

(see [AS], p. 437f). Similar formulas exist for all odd n .

Definition 5.3.6. For $m \in \mathbb{R}$ we define

$$P_m(\mathbb{R}^n) := \sum_{\substack{r+s+t \geq m \\ r \geq -n}} P_{r,s,t}(\mathbb{R}^n) + (C^\infty(\Sigma \mathbb{R}^n|_{\mathbb{R}^n \setminus \{0\}}) \cap C^0(\Sigma \mathbb{R}^n)),$$

where the second space is the space of all spinors which are smooth on $\mathbb{R}^n \setminus \{0\}$ and have a continuous extension to \mathbb{R}^n .

Remark 5.3.7. Let $\vartheta \in P_m(\mathbb{R}^n)$. Then we have $E_i \cdot \vartheta \in P_m(\mathbb{R}^n)$ for all $i \in \{1, \dots, n\}$. If $f \in C^\infty(\mathbb{R}^n)$ then for every $s \in \mathbb{N}$ by Taylor's formula we may write

$$f(x) = \sum_{|\alpha| < s} \frac{1}{\alpha!} \frac{\partial^{|\alpha|} f(0)}{\partial x^\alpha} x^\alpha + R_s(x),$$

where $|R_s(x)| = O(|x|^s)$ as $x \rightarrow 0$. Thus by choosing $s \geq m + 1$ we find that the spinor $f\vartheta$ is in $P_m(\mathbb{R}^n)$.

Remark 5.3.8. A spinor $\vartheta \in P_m(\mathbb{R}^n)$ has a continuous extension to \mathbb{R}^n if and only if $m > 0$. Furthermore by Proposition 5.2.2 it follows that for all $m \in (-n, 0]$ we have

$$\begin{aligned} P_m(\mathbb{R}^n) &= \sum_{\substack{r+s+t=m \\ r \geq -n}} P_{r,s,t}(\mathbb{R}^n) + P_{m+1}(\mathbb{R}^n) \\ &\subset D^{g_{\text{eucl}}}(P_{m+1}(\mathbb{R}^n)) + P_{m+1}(\mathbb{R}^n). \end{aligned}$$

Lemma 5.3.9. Let (M, g, Θ) be a closed Riemannian spin manifold of dimension n and let $\lambda \in \mathbb{R}$. Let $\gamma \in \Sigma_n$ be a constant spinor on \mathbb{R}^n . Then the spinor $G_\lambda^{g_{\text{eucl}}}(\cdot, 0)\gamma$ is in $P_{1-n}(\mathbb{R}^n)$. Let the matrix coefficients B_i^j be defined as in (5.2). Then for all i the spinor

$$x \mapsto \sum_{j=1}^n (B_i^j(x) - \delta_i^j) \nabla_{E_j} G_\lambda^{g_{\text{eucl}}}(x, 0)\gamma$$

is in $P_{2-n}(\mathbb{R}^n)$.

Proof. The assertion for $G_\lambda^{g_{\text{encl}}}(\cdot, 0)\gamma$ can be seen immediately from Corollary 5.3.4. Let f_λ be as in Theorem 5.3.3 and write $f_\lambda(x) = g_\lambda(|x|)$ with a function $g_\lambda: (0, \infty) \rightarrow \mathbb{R}$. Then we have

$$G_\lambda^{g_{\text{encl}}}(x, 0)\gamma = \frac{g'_\lambda(|x|)}{|x|}x \cdot \gamma + \lambda g_\lambda(|x|)\gamma$$

and for every $j \in \{1, \dots, n\}$ we have

$$\nabla_{E_j} G_\lambda^{g_{\text{encl}}}(x, 0)\gamma = \frac{g''_\lambda(|x|)x_j}{|x|^2}x \cdot \gamma - \frac{g'_\lambda(|x|)x_j}{|x|^3}x \cdot \gamma + \frac{g'_\lambda(|x|)}{|x|}E_j \cdot \gamma + \lambda \frac{g'_\lambda(|x|)x_j}{|x|}\gamma.$$

Since the exponential map is a radial isometry, we have $\sum_{j=1}^n g_{ij}(x)x_j = x_i$ and thus $\sum_{j=1}^n B_i^j(x)x_j = x_i$ for every fixed i . Thus we find

$$\sum_{j=1}^n (B_i^j(x) - \delta_i^j) \nabla_{E_j} G_\lambda^{g_{\text{encl}}}(x, 0)\gamma = \sum_{j=1}^n (B_i^j(x) - \delta_i^j) \frac{g'_\lambda(|x|)}{|x|} E_j \cdot \gamma.$$

Since we have $g'_\lambda(|x|) = O(|x|^{1-n})$ as $x \rightarrow 0$ the assertion now follows from the Taylor expansion (5.3) of $B_i^j(x)$. \square

Next we prove existence and uniqueness of Green's function for $D^g - \lambda$ on a closed Riemannian spin manifold in such a way that we also obtain the expansion of Green's function around the singularity. The idea is to apply the equation (5.2) for the Dirac operator in the trivialization to a Euclidean Green's function and then determine the correction terms. This has been carried out in [AH], where for some technical steps Sobolev embeddings were used. We present a more simple argument using the preimages under the Dirac operator from Proposition 5.2.2.

In the following for a fixed point $p \in M$ let $\rho: V \rightarrow U$ be a local parametrization of M by Riemannian normal coordinates, where $U \subset M$ is an open neighborhood of p , $V \subset \mathbb{R}^n$ is an open neighborhood of 0 and $\rho(0) = p$. Furthermore let

$$\beta: \Sigma\mathbb{R}^n|_V \rightarrow \Sigma^g M|_U, \quad A: C^\infty(\Sigma^g M|_U) \rightarrow C^\infty(\Sigma\mathbb{R}^n|_V)$$

denote the maps which send a spinor to its corresponding spinor in the Bourguignon-Gauduchon trivialization defined in Section 5.1.

Theorem 5.3.10. *Let (M, g, Θ) be a closed n -dimensional Riemannian spin manifold, $p \in M$. For every $\varphi \in \Sigma_p^g M$ there exists a unique Green's function $G_\lambda^g(\cdot, p)\varphi$. If $\gamma := \beta^{-1}\varphi \in \Sigma_n$ is the constant spinor on \mathbb{R}^n corresponding to φ , then the first two terms of the expansion of $AG_\lambda^g(\cdot, p)\varphi$ at 0 coincide with the first two terms of the expansion of $G_\lambda^{g_{\text{encl}}}(\cdot, 0)\gamma$ given in Corollary 5.3.4.*

Proof. Let $\varepsilon > 0$ such that $B_{2\varepsilon}(0) \subset V$ and let $\eta : \mathbb{R}^n \rightarrow [0, 1]$ be a smooth function with $\text{supp}(\eta) \subset B_{2\varepsilon}(0)$ and $\eta \equiv 1$ on $B_\varepsilon(0)$. Then the spinor Θ_1 defined on $\mathbb{R}^n \setminus \{0\}$ by $\Theta_1(x) := \eta(x)G_\lambda^{\text{eucl}}(x, 0)\gamma$ is smooth on $\mathbb{R}^n \setminus \{0\}$. For $r \in \{1, \dots, n\}$ we define smooth spinors Φ_r on $M \setminus \{p\}$ and Θ_{r+1} on $\mathbb{R}^n \setminus \{0\}$ inductively as follows. For $r = 1$ we define

$$\Phi_1(q) := \begin{cases} A^{-1}\Theta_1(q), & q \in U \setminus \{p\} \\ 0, & q \in M \setminus U \end{cases}$$

and

$$\Theta_2(x) := \begin{cases} A(D^g - \lambda)\Phi_1(x), & x \in V \setminus \{0\} \\ 0, & x \in \mathbb{R}^n \setminus V \end{cases}.$$

By the formula (5.2) for the Dirac operator in the trivialization we have on $V \setminus \{0\}$

$$\begin{aligned} \Theta_2 &= (D^{\text{eucl}} - \lambda)\Theta_1 + \sum_{i,j=1}^n (B_i^j - \delta_i^j)E_i \cdot \nabla_{E_j}\Theta_1 \\ &\quad + \frac{1}{4} \sum_{i,j,k=1}^n \tilde{\Gamma}_{ij}^k E_i \cdot E_j \cdot E_k \cdot \Theta_1. \end{aligned}$$

The first term vanishes on $B_\varepsilon(0) \setminus \{0\}$. It follows from the expansions of $\tilde{\Gamma}_{ij}^k$ and $B_i^j - \delta_i^j$ in (5.3), (5.4) and from Lemma 5.3.9 that $\Theta_2 \in P_{2-n}(\mathbb{R}^n)$.

Next let $r \in \{2, \dots, n\}$ and assume that Φ_{r-1} and Θ_r have already been defined. We may assume that $\Theta_r \in P_{r-n}(\mathbb{R}^n)$. By Remark 5.3.8 there exists $\beta_{r+1} \in P_{r+1-n}(\mathbb{R}^n)$ such that $\Theta_r - (D^{\text{eucl}} - \lambda)\beta_{r+1} \in P_{r+1-n}(\mathbb{R}^n)$. We define Φ_r and Θ_{r+1} by

$$\Phi_r(q) := \begin{cases} \Phi_{r-1}(q) - A^{-1}(\eta\beta_{r+1})(q), & q \in U \setminus \{p\} \\ 0, & q \in M \setminus U \end{cases}$$

and

$$\Theta_{r+1}(x) := \begin{cases} A(D^g - \lambda)\Phi_r(x), & x \in V \setminus \{0\} \\ 0, & x \in \mathbb{R}^n \setminus V \end{cases}.$$

By the formula (5.2) for the Dirac operator in the trivialization we have on $B_\varepsilon(0) \setminus \{0\}$

$$\begin{aligned} \Theta_{r+1} &= A(D^g - \lambda)\Phi_{r-1} - A(D^g - \lambda)A^{-1}\beta_{r+1} \\ &= \Theta_r - (D^{\text{eucl}} - \lambda)\beta_{r+1} - \sum_{i,j=1}^n (B_i^j - \delta_i^j)E_i \cdot \nabla_{E_j}\beta_{r+1} \\ &\quad - \frac{1}{4} \sum_{i,j,k=1}^n \tilde{\Gamma}_{ij}^k E_i \cdot E_j \cdot E_k \cdot \beta_{r+1}. \end{aligned}$$

Using the expansions of $\tilde{\Gamma}_{ij}^k$ and $B_i^j - \delta_i^j$ in (5.3), (5.4) we conclude that $\Theta_{r+1} \in P_{r+1-n}(\mathbb{R}^n)$.

We see that Θ_{n+1} has a continuous extension to \mathbb{R}^n and we obtain a continuous extension Ψ of $(D^g - \lambda)\Phi_n$ to all of M . Thus there exists

$$\Psi' \in C^\infty(\Sigma^g M|_{M \setminus \{p\}}) \cap H^1(\Sigma^g M)$$

such that $(D^g - \lambda)\Psi' = P\Psi - \Psi$. Define

$$\Gamma := \Phi_n + \Psi', \quad \Theta := -\eta\beta_3 - \dots - \eta\beta_{n+1} + A\Psi'.$$

Then on $B_\varepsilon(0) \setminus \{0\}$ we have $A\Gamma = G_\lambda^{g_{\text{eucl}}}(\cdot, 0)\gamma + \Theta$.

If P is the L^2 -orthogonal projection onto $\ker(D^g - \lambda)$, then

$$G_\lambda^g(\cdot, p)\varphi := \Gamma - P\Gamma$$

satisfies (5.5), (5.6) and thus is a Green's function. Uniqueness also follows from (5.5), (5.6). The statement on the expansion of $AG_\lambda^g(\cdot, p)\varphi$ is obvious, since we have $\Theta \in P_{3-n}(\mathbb{R}^n)$. \square

5.4 Definition of the mass endomorphism

Let (M, g, Θ) be a closed Riemannian spin manifold and let $p \in M$. Assume that the metric g is flat on an open neighborhood of p . Let $U \subset M$ be an open neighborhood of p , let $V \subset \mathbb{R}^n$ be an open neighborhood of 0 and let $\rho: V \rightarrow U$ be a local parametrization of M by Riemannian normal coordinates sending 0 to p . Let

$$\beta^g : \Sigma\mathbb{R}^n|_V \rightarrow \Sigma^g M|_U.$$

be the identification of the spinor bundles defined in Section 5.1 and let

$$A^g : C^\infty(\Sigma^g M|_U) \rightarrow C^\infty(\Sigma\mathbb{R}^n|_V), \quad \psi \mapsto (\beta^g)^{-1} \circ \psi \circ \rho.$$

Let $\varphi \in \Sigma_p^g M$ and let $\gamma := (\beta^g)^{-1}\varphi$. Since g is flat on an open neighborhood of p , the terms $\tilde{\Gamma}_{ij}^k$ and $B_i^j - \delta_i^j$ in the formula (5.2) vanish on an open neighborhood of $0 \in \mathbb{R}^n$. Thus in the proof of Theorem 5.3.10 the spinor Θ_2 has a smooth extension to \mathbb{R}^n vanishing on an open neighborhood of 0. Then one obtains a smooth extension Ψ of $(D^g - \lambda)\Phi_1$ to all of M vanishing on an open neighborhood of p . Now there exists $\Psi' \in C^\infty(\Sigma^g M)$ such that $(D^g - \lambda)\Psi' = P\Psi - \Psi$. As above we define

$$\Gamma := \Phi_1 + \Psi', \quad G_\lambda^g(\cdot, p)\varphi := \Gamma - P\Gamma.$$

Thus the spinor

$$x \mapsto w^g(x, p)\varphi := \Psi'(x) - P\Gamma(x)$$

is smooth on all of M . It is independent of the choice of Ψ' and can be regarded as the difference between Green's function for $D^g - \lambda$ and the spinor $(A^g)^{-1}(\eta G_\lambda^{g_{\text{eucl}}}(\cdot, 0)\gamma)$.

In the following we are interested in the case $\lambda = 0$ and we denote Green's function for D^g by $G^g := G_0^g$. By Theorem 5.3.3 a Euclidean Green's function for $D^{g_{\text{eucl}}}$ is given by

$$G^{g_{\text{eucl}}}(x, y)\gamma := -\frac{1}{\omega_{n-1}|x-y|^{n-1}} \frac{x-y}{|x-y|} \cdot \gamma.$$

Using the definition of Φ_1 and $w^g(\cdot, p)\varphi$ we have for all $x \in M \setminus \{p\}$

$$G^g(x, p)\varphi = (A^g)^{-1}(\eta G^{g_{\text{eucl}}}(\cdot, 0)\gamma)(x) + w^g(x, p)\varphi,$$

where the first term on the right hand side is understood to be zero for $x \in M \setminus U$. Since $\eta \equiv 1$ on $B_\varepsilon(0)$, the spinor $w^g(p, p)\varphi \in \Sigma_p^g M$ is independent of the choice of η . It can be regarded as the constant term in an expansion of Green's function for D^g around p .

As in [AHM] we define the mass endomorphism.

Definition 5.4.1. *Let (M, g, Θ) be a closed Riemannian spin manifold with $\dim M \geq 2$, which is conformally flat on an open neighborhood of $p \in M$. Choose a metric $h \in [g]$, which is flat on an open neighborhood of p and such that $h_p = g_p$. Let G^h be Green's function for D^h . Then, we define the mass endomorphism in p as*

$$m_p^g : \Sigma_p^g M \rightarrow \Sigma_p^g M, \quad \varphi \mapsto \beta_{h,g} w^h(p, p) \beta_{g,h} \varphi,$$

where w^h is the term in the above expansion.

It is shown in [AHM] that this definition does not depend on the choice of $h \in [g]$ and that m_p^g is linear and self-adjoint. There is an analogy in conformal geometry: the constant term of Green's function $\Gamma(\cdot, p)$ for the Yamabe operator in p can be interpreted as the mass of the asymptotically flat manifold $(M \setminus \{p\}, \Gamma(\cdot, p)^{4/(n-2)}g)$ (see [LP]). Therefore, the endomorphism is called mass endomorphism.

The aim of introducing the mass endomorphism in [AHM] is to obtain at least one of the strict inequalities

$$\lambda_{\min}^+(M, [g], \Theta) < \lambda_{\min}^+(\mathbb{S}^n), \quad \lambda_{\min}^-(M, [g], \Theta) < \lambda_{\min}^+(\mathbb{S}^n)$$

from the hypothesis of Theorem 3.1.2. The result of this article then reads as follows.

Theorem 5.4.2. *Let (M, g, Θ) be a closed Riemannian spin manifold of dimension $n \geq 2$ with $\ker(D^g) = 0$. Assume that there is a point $p \in M$ which has a conformally flat neighborhood and that the mass endomorphism in p possesses a positive (resp. negative) eigenvalue. Then we have*

$$\lambda_{\min}^+(M, [g], \Theta) < \lambda_{\min}^+(\mathbb{S}^n) \quad \text{resp.} \quad \lambda_{\min}^-(M, [g], \Theta) < \lambda_{\min}^-(\mathbb{S}^n).$$

If $h \in [g]$ is a Riemannian metric such that $m_p^g = \beta_{h,g} w^h(p, p) \beta_{g,h}$, then the additional assumption $\ker(D^g) = 0$ implies that $D^h w^h(\cdot, p) \varphi$ vanishes on an open neighborhood of p and this fact is used in the proof. It is not clear how to obtain a proof without this assumption.

Unfortunately it is only known for very few Riemannian spin manifolds, whether there are points with nonzero mass endomorphism. For example, on the flat torus and on the sphere with the standard metric it always vanishes, whereas on the projective spaces $\mathbb{R}P^{4k+3}$ with the standard metric one has points with nonzero mass endomorphism (see [AHM]). We will prove in Section 6.3 that in dimension 3 the mass endomorphism is not zero in the generic case.

Chapter 6

Eigenspinors for generic metrics

6.1 Transversality

A nice introduction to finite dimensional transversality theory can be found in [Hir], while the infinite dimensional case is treated in [La]. Transversality theory has already been used in the literature to determine generic properties of eigenvalues and eigenfunctions of differential operators. Some examples are the scalar Laplacian [U] and the Hodge Laplacian [EP] for generic metrics and solutions of the Dirac equation in Minkowski spacetime for generic initial data [TT]. The goal of this section is to quote a transversality theorem which will be useful later.

Definition 6.1.1. *Let $f: Q \rightarrow N$ be a C^1 map between two smooth manifolds. Let $A \subset N$ be a submanifold. f is called transverse to A , if for all $x \in Q$ with $f(x) \in A$ we have*

$$T_{f(x)}A + \text{im}(df|_x) = T_{f(x)}N.$$

One reason for the importance of transversality is the following theorem.

Theorem 6.1.2. *Assume that Q, N are smooth manifolds without boundary. Let $f: Q \rightarrow N$ be a C^r map, $r \geq 1$ and $A \subset N$ a C^r submanifold. If f is transverse to A , then $f^{-1}(A)$ is a C^r submanifold of Q . The codimension of $f^{-1}(A)$ in Q is the same as the codimension of A in N .*

Proof. see [Hir], p. 22 □

Definition 6.1.3. *Let X be a topological space. A subset $E \subset X$ is called nowhere dense in X , if the interior of \overline{E} is empty. A subset of X is called of first category, if it is a countable union of sets which are nowhere dense in X . Otherwise it is called of second category. A subset $B \subset X$ is called*

residual in X , if it contains a countable intersection of sets which are open and dense in X .

If $A \subset X$ is of first category, it follows that the complement of A is residual in X . An important result is the following theorem, which is called the Baire category theorem.

Theorem 6.1.4. *If (X, d) is a complete metric space, then every residual subset of X is dense in X .*

Proof. see e. g. [Ru], p. 43. □

An important result is Sard's theorem.

Theorem 6.1.5. *Let $f: Q \rightarrow N$ be a C^r -map between smooth manifolds. If*

$$r > \max\{0, \dim Q - \dim N\},$$

then the set of all critical values of f has measure zero in N . The set of all regular values of f is residual and therefore dense.

Proof. see [Hir], p. 69. □

We quote the following transversality theorem from [Hir], [U] including the proof.

Theorem 6.1.6. *Let V, M, N be smooth manifolds and let $A \subset N$ be a smooth submanifold. Let $F: V \rightarrow C^r(M, N)$ be a map, such that the evaluation map $F^{ev}: V \times M \rightarrow N$, $(v, m) \mapsto F(v)(m)$ is C^r and transverse to A and furthermore*

$$r > \max\{0, \dim M + \dim A - \dim N\}.$$

Then the set of all $v \in V$, such that the map $F(v)$ is transverse to A , is residual and therefore dense in V .

Proof. Since F^{ev} is transverse to A , the set $P := (F^{ev})^{-1}(A)$ is a C^r submanifold of $V \times M$ of dimension $\dim V + \dim M + \dim A - \dim N$. Let $\pi_1: V \times M \rightarrow V$ be the projection and let $\alpha := \pi_1|_P$. Let $v \in V$, $m \in M$ and $F^{ev}(v, m) = a \in A$. Then

$$\operatorname{im}(d\alpha|_{(v,m)}) = d\pi_1|_{(v,m)}(dF^{ev}|_{(v,m)}^{-1}(T_a A))$$

and thus we find

$$\begin{aligned} \operatorname{codim} \operatorname{im}(d\alpha|_{(v,m)}) &= \operatorname{codim}(dF^{ev}|_{(v,m)}^{-1}(T_a A) + \ker d\pi_1|_{(v,m)}) \\ &= \operatorname{codim}(dF^{ev}|_{(v,m)}^{-1}(T_a A) + T_m M). \end{aligned}$$

By Theorem 6.1.5 the set of all regular values of α is residual in V . Fix a regular value v of α . If $\alpha^{-1}(v) = \emptyset$, then $\text{im}(F(v)) \cap A = \emptyset$ and thus $F(v)$ is transverse to A . Now assume that there exists $(v, m) \in \alpha^{-1}(v)$. Since $\text{codim im}(d\alpha|_{(v,m)}) = 0$ we obtain

$$dF^{ev}|_{(v,m)}^{-1}(T_a A) + T_m M = T_{(v,m)}(V \times M).$$

We apply $dF^{ev}|_{(v,m)}$ to both sides. Since F^{ev} is transverse to A , the right hand side contains a complement to $T_a A$. Thus $dF^{ev}|_{(v,m)}(T_m M)$ also contains a complement to $T_a A$, which means that $F(v)$ is transverse to A . The assertion follows. \square

Remark 6.1.7. *There exist infinite dimensional versions of this theorem and of Sard's theorem (see e. g. [U], [Sm], [Ab]). However for our purpose this theorem is sufficient.*

6.2 Eigenspinors in dimensions 2 and 3

Let M be a closed oriented spin manifold of dimension n with a fixed spin structure. We define $R(M)$ as the set of all smooth Riemannian metrics on M .

Definition 6.2.1. *A subset of $R(M)$ is called generic, if it is open in $R(M)$ with respect to the C^1 -topology and dense in $R(M)$ with respect to all C^k -topologies, $k \in \mathbb{N}$.*

Remark 6.2.2. *Let $k \in \mathbb{N}$ and let $A \subset R(M)$. If A is open in $R(M)$ with respect to the C^k -topology, then it is open in $R(M)$ with respect to the C^m -topology for all $m > k$. If A is dense in $R(M)$ with respect to the C^k -topology, then it is dense in $R(M)$ with respect to the C^m -topology for all $m < k$.*

It is an important result that the eigenvalues of D^g depend continuously on g with respect to the C^1 -topology. Namely Proposition 7.1 in [Bä3] states the following.

Proposition 6.2.3. *Let g be a smooth Riemannian metric on M and let $\varepsilon > 0$, $\Lambda > 0$ such that $-\Lambda, \Lambda \notin \text{spec}(D^g)$. Write*

$$\text{spec}(D^g) \cap (-\Lambda, \Lambda) = \{\lambda_1 \leq \dots \leq \lambda_k\}.$$

Then there exists an open neighborhood V of g in the set of all smooth Riemannian metrics with respect to the C^1 -topology such that for any $h \in V$ we have

$$\text{spec}(D^h) \cap (-\Lambda, \Lambda) = \{\mu_1 \leq \dots \leq \mu_k\}$$

and $|\lambda_i - \mu_i| < \varepsilon$ for all i . Here the eigenvalues λ_i and μ_i are repeated according to their complex multiplicities.

Assume that $n \equiv 2, 3, 4 \pmod{8}$. Then there exists a quaternionic structure on $\Sigma^g M$, which commutes with the Dirac operator D^g and thus the eigenspaces of D^g have even complex dimension (see Remark 2.2.3). Therefore the following notation introduced in [Da] is useful.

Definition 6.2.4. *A non-zero eigenvalue λ of D^g is called simple, if*

$$\dim_{\mathbb{C}} \ker(D^g - \lambda) = \begin{cases} 2, & n \equiv 2, 3, 4 \pmod{8} \\ 1, & \text{otherwise} \end{cases}$$

If $n \equiv 2, 3, 4 \pmod{8}$ and λ is simple, then we can always choose an L^2 -orthonormal basis of $\ker(D^g - \lambda)$ of the form $\{\psi, J\psi\}$.

For every $g \in R(M)$ we enumerate the nonzero eigenvalues of D^g in the following way

$$\dots \leq \lambda_2^- \leq \lambda_1^- < 0 < \lambda_1^+ \leq \lambda_2^+ \leq \dots$$

Here all the non-zero eigenvalues are repeated by their complex multiplicities, while $\dim \ker(D^g) \geq 0$ is arbitrary. For $m \in \mathbb{N} \setminus \{0\}$ we define

$$\begin{aligned} S_m(M) &:= \{g \in R(M) \mid \lambda_1^\pm, \dots, \lambda_m^\pm \text{ are simple}\} \\ N_m(M) &:= \{g \in R(M) \mid \text{all eigenspinors to } \lambda_1^\pm, \dots, \lambda_m^\pm \text{ are nowhere zero}\}. \end{aligned}$$

Using Proposition 6.2.3 together with a study of the change of the non-zero eigenvalues under conformal deformations of the metric M , Dahl could prove the following result.

Theorem 6.2.5 ([Da]). *Let M be a closed spin manifold of dimension 2 or 3 and let $m \in \mathbb{N} \setminus \{0\}$. Then for every $g \in R(M)$ the subset $S_m(M) \cap [g]$ is generic in $[g]$.*

In order to prove Theorem 1.0.1 it is sufficient to prove the following.

Theorem 6.2.6. *Let M be a closed connected spin manifold of dimension 2 or 3 and let $m \in \mathbb{N} \setminus \{0\}$. Then for every $g \in R(M)$ the subset $N_m(M) \cap [g]$ is generic in $[g]$.*

Remark 6.2.7. *On a closed oriented surface M of genus 2 there exist spin structures, such that for every Riemannian metric on M there exist harmonic spinors (see Proposition 2.3 in [Hit]). By the result on the zero set of positive harmonic spinors in Theorem 4.1.2 we find that Theorem 6.2.6 in the case $n = 2$ is not true for harmonic spinors.*

In the proof we will use the following unique continuation theorem due to Aronszajn ([Ar], quoted from [Bä4]).

Theorem 6.2.8. *Let M be a connected Riemannian manifold and let Σ be a vector bundle over M with a connection ∇ . Let P be an operator of the form $P = \nabla^* \nabla + P_1 + P_0$ acting on sections of Σ , where P_1, P_0 are differential operators of order 1 and 0 respectively. Let ψ be a solution to $P\psi = 0$. If there exists a point, at which ψ and all derivatives of ψ of any order vanish, then ψ vanishes identically.*

We will also use Theorem 6.1.6 for families of spinors parametrized by Riemannian metrics. A basic problem is that $[g]$ is not a smooth manifold. If one replaces $[g]$ by the space of all k times continuously differentiable metrics, $k \geq 1$, which are conformal to g , then we see from formula (2.3) that the coefficients of the Dirac operator are not smooth in general. In this case we cannot expect that the eigenspinors are smooth and that Theorem 6.2.8 remains valid. In order to get around this problem we will use finite dimensional manifolds of smooth Riemannian metrics of the form

$$V_{f_1 \dots f_r} := \left\{ \left(1 + \sum_{i=1}^r t_i f_i \right) g \mid t_1, \dots, t_r \in \mathbb{R} \right\} \cap R(M),$$

where $r \in \mathbb{N} \setminus \{0\}$ and $f_1, \dots, f_r \in C^\infty(M, \mathbb{R})$.

Our first aim is to construct a map, which associates to a Riemannian metric $h \in [g]$ an eigenspinor of $D^{g,h}$ in a continuous way.

Lemma 6.2.9. *Let $k, m \in \mathbb{N} \setminus \{0\}$. Let $g \in S_m(M)$ and equip $[g]$ with the C^k -topology. Let $\lambda \in \{\lambda_1^\pm(g), \dots, \lambda_m^\pm(g)\}$ and let ψ be an eigenspinor of D^g corresponding to λ . Then there exists an open neighborhood $V \subset [g]$ of g and a map $F_\psi: V \rightarrow C^\infty(\Sigma^g M)$ such that for every $h \in V$ the spinor $F_\psi(h)$ is an eigenspinor of $D^{g,h}$ and such that the map $F_\psi^{ev}: V \times M \rightarrow \Sigma^g M$ defined by $F_\psi^{ev}(h, x) := F_\psi(h)(x)$ is continuous and such that for all functions $f_1, \dots, f_r \in C^\infty(M, \mathbb{R})$ the restriction $F_\psi^{ev}|_{V_{f_1 \dots f_r} \times M}$ is differentiable.*

Proof. Let $V \subset [g]$ be an open convex neighborhood of g , which is contained in $S_m(M) \cap [g]$. Let $h \in V$ and let $I \subset \mathbb{R}$ be an open interval containing $[0, 1]$ such that for every $t \in I$ the tensor field $g_t := g + t(h - g)$ is a Riemannian metric on M . Let $\lambda \in \{\lambda_1^\pm(g), \dots, \lambda_m^\pm(g)\}$. Then there exists exactly one real-analytic family $t \mapsto \lambda_t$ of eigenvalues of D^{g_t} such that $\lambda_0 = \lambda$. Furthermore there exists a real-analytic family ψ_t of spinors, such that $\{\psi_t\}$ respectively $\{\psi_t, J\psi_t\}$ forms an L^2 -orthonormal basis of $\ker(D^{g_t} - \lambda_t)$ for every t . After possibly shrinking V we may assume that for all metrics $h \in V$ and for all $\lambda \in \{\lambda_1^\pm(g), \dots, \lambda_m^\pm(g)\}$ the families λ_t and ψ_t are defined for all $t \in [0, 1]$.

Since we are free to replace $\psi_t, J\psi_t$ by linear combinations $a\psi_t + bJ\psi_t$ with $a, b \in \mathbb{C}$, $|a|^2 + |b|^2 > 0$, we may assume that $\psi_0 = \psi$. We define $F_\psi(h) := \psi_1$.

Let $h \in V$ and let $k = fg$ for some $f \in C^\infty(M, \mathbb{R})$. With $h_t := h + tk$ consider $F_\psi(h_t)$ for small $|t|$ such that $h_t \in V$. Since $a_{g,h}$ and a_{h,h_t} commute, we have $b_{h,h_t} \circ b_{g,h} = b_{g,h_t}$ by Lemma 2.3.1. and thus $\bar{\beta}_{h,h_t} \circ \bar{\beta}_{g,h} = \bar{\beta}_{g,h_t}$. It follows that

$$D^{g,h_t} = \bar{\beta}_{h,g} \circ D^{h,h_t} \circ \bar{\beta}_{g,h}.$$

There exists a real-analytic family $t \mapsto \chi(t) \in C^\infty(\Sigma^h M)$ of eigenspinors of D^{h,h_t} with $\chi(0) = \bar{\beta}_{g,h}(F_\psi(h))$. It follows that

$$F_\psi(h_t) = \bar{\beta}_{h,g}(\chi(t)).$$

By taking $k := h - h'$ for $h, h' \in V$ we find that F_ψ^{ev} is continuous. Furthermore for all functions $f_1, \dots, f_r \in C^\infty(M, \mathbb{R})$ we can use this equation to see that the restriction $F_\psi^{ev}|_{V_{f_1 \dots f_r} \times M}$ is differentiable. \square

The strategy for the proof of Theorem 6.2.6 is based on the following remark.

Remark 6.2.10. *Let $A \subset \Sigma^g M$ be the zero section. Since the dimension of the total space $\Sigma^g M$ of the spinor bundle is*

$$\dim \Sigma^g M = n + 2^{1+[n/2]} > 2n = \dim M + \dim A,$$

a map $f: M \rightarrow \Sigma^g M$ is transverse to A if and only if $f^{-1}(A) = \emptyset$.

If for every $\lambda_i \in \{\lambda_1^\pm, \dots, \lambda_m^\pm\}$ we choose an eigenspinor ψ_i and define the map

$$F_{\psi_i} : V_{f_1 \dots f_r} \rightarrow C^\infty(\Sigma^g M)$$

as in Lemma 6.2.9, then by this remark $V_{f_1 \dots f_r} \cap N_m(M)$ is the set of all Riemannian metrics $h \in V_{f_1 \dots f_r}$ such that all the $F_{\psi_i}(h)$ are transverse to the zero section. Therefore in order to prove that $N_m(M) \cap [g]$ is dense in $[g]$ we would like to apply Theorem 6.1.6. Our aim is then to show that a suitable restriction of the map F_ψ^{ev} defined as in Lemma 6.2.9 is transverse to the zero section.

Let $p \in M$ with $\psi(p) = 0$. We have a canonical decomposition of the tangent space

$$T_{\psi(p)}\Sigma^g M \cong \Sigma_p^g M \oplus T_p M$$

and thus

$$dF_\psi^{ev}|_{(g,p)} : T_g V_{f_1 \dots f_r} \oplus T_p M \rightarrow \Sigma_p^g M \oplus T_p M.$$

For a given $h \in V_{f_1 \dots f_r}$ we will write $g_t = g + t(h - g)$ and

$$\psi_t := F_\psi(g_t), \quad \frac{d\psi_t(x)}{dt} \Big|_{t=0} := \pi_1(dF_\psi^{ev}|_{(g,x)}(h - g, 0)).$$

Then it follows that

$$0 = \left(\frac{d}{dt} D^{g, g_t} \Big|_{t=0} - \frac{d\lambda_t}{dt} \Big|_{t=0} \right) \psi + (D^g - \lambda) \frac{d\psi_t}{dt} \Big|_{t=0}. \quad (6.1)$$

Remark 6.2.11. Let $\varphi \in \Sigma_p^g M$ and $X, Y \in T_p M$. If one polarizes the identity

$$\langle X \cdot \varphi, X \cdot \varphi \rangle = g(X, X) \langle \varphi, \varphi \rangle,$$

then one obtains

$$\operatorname{Re} \langle X \cdot \varphi, Y \cdot \varphi \rangle = g(X, Y) \langle \varphi, \varphi \rangle. \quad (6.2)$$

Since Clifford multiplication with vectors is antisymmetric, it follows that $\operatorname{Re} \langle X \cdot \varphi, \varphi \rangle = 0$. Let $\varphi \neq 0$ and let $(e_i)_{i=1}^n$ be an orthonormal basis of $T_p M$. It follows that for $n = 2$ the spinors

$$\varphi, e_1 \cdot \varphi, e_2 \cdot \varphi, e_1 \cdot e_2 \cdot \varphi$$

form an orthogonal basis of $\Sigma_p^g M$ with respect to the real scalar product $\operatorname{Re} \langle \cdot, \cdot \rangle$. Similarly for $n = 3$ the spinors

$$\varphi, e_1 \cdot \varphi, e_2 \cdot \varphi, e_3 \cdot \varphi$$

form an orthogonal basis of $\Sigma_p^g M$ with respect to $\operatorname{Re} \langle \cdot, \cdot \rangle$.

The following rather long lemma is the most important step in showing that a suitable restriction of F_ψ^{ev} is transverse to the zero section.

Lemma 6.2.12. Let $n \in \{2, 3\}$ and $m \in \mathbb{N} \setminus \{0\}$. Let $g \in S_m(M)$ and let $\lambda \in \{\lambda_1^\pm(g), \dots, \lambda_m^\pm(g)\}$. Let ψ be an eigenspinor of D^g corresponding to λ and let $p \in M$ with $\psi(p) = 0$. Then there exist $f_1, \dots, f_4 \in C^\infty(M, \mathbb{R})$ such that the map $F_\psi^{ev}: V_{f_1 \dots f_4} \times M \rightarrow \Sigma^g M$ satisfies

$$\pi_1(dF_\psi^{ev}|_{(g,p)}(T_g V_{f_1 \dots f_4} \oplus \{0\})) = \Sigma_p^g M.$$

Proof. Assume that the claim is wrong. Then there exists $\varphi \in \Sigma_p^g M \setminus \{0\}$ such that for all $f \in C^\infty(M, \mathbb{R})$ we have

$$0 = \operatorname{Re} \langle \pi_1(dF_\psi^{ev}|_{(g,p)}(fg, 0)), \varphi \rangle = \operatorname{Re} \left\langle \frac{d\psi_t}{dt} \Big|_{t=0}(p), \varphi \right\rangle.$$

From the formula (5.7) for Green's function it follows that

$$0 = \operatorname{Re} \int_{M \setminus \{p\}} \langle (D^g - \lambda) \frac{d\psi_t}{dt} \Big|_{t=0}, G_\lambda^g(\cdot, p) \varphi \rangle dv^g + \operatorname{Re} \langle P(\frac{d\psi_t}{dt} \Big|_{t=0})(p), \varphi \rangle.$$

Since λ is a simple eigenvalue, all spinors in $\ker(D^g - \lambda)$ vanish at p . Thus the last term vanishes. By (5.6) and (6.1) we have

$$\begin{aligned} 0 &= -\operatorname{Re} \int_{M \setminus \{p\}} \langle (\frac{d}{dt} D^{g, g_t} \Big|_{t=0} - \frac{d\lambda_t}{dt} \Big|_{t=0}) \psi, G_\lambda^g(\cdot, p) \varphi \rangle dv^g \\ &= -\operatorname{Re} \int_{M \setminus \{p\}} \langle \frac{d}{dt} D^{g, g_t} \Big|_{t=0} \psi, G_\lambda^g(\cdot, p) \varphi \rangle dv^g \end{aligned}$$

for all $f \in C^\infty(M, \mathbb{R})$. If we use the formula (2.13) for the derivative of the Dirac operator and

$$\operatorname{grad}^g(f) \cdot \psi = (D^g - \lambda)(f\psi)$$

it follows that

$$\begin{aligned} 0 &= \frac{1}{2} \operatorname{Re} \int_{M \setminus \{p\}} \lambda f \langle \psi, G_\lambda^g(\cdot, p) \varphi \rangle dv^g \\ &\quad + \frac{1}{4} \operatorname{Re} \int_{M \setminus \{p\}} \langle (D^g - \lambda)(f\psi), G_\lambda^g(\cdot, p) \varphi \rangle dv^g. \end{aligned}$$

Using the definition of Green's function and that all spinors in $\ker(D^g - \lambda)$ vanish at p , we find that

$$\begin{aligned} 0 &= \frac{1}{2} \operatorname{Re} \int_{M \setminus \{p\}} \lambda f \langle \psi, G_\lambda^g(\cdot, p) \varphi \rangle dv^g + \frac{1}{4} \operatorname{Re} \langle (f\psi)(p) - P(f\psi)(p), \varphi \rangle \\ &= \frac{1}{2} \operatorname{Re} \int_{M \setminus \{p\}} \lambda f \langle \psi, G_\lambda^g(\cdot, p) \varphi \rangle dv^g \end{aligned}$$

for all $f \in C^\infty(M, \mathbb{R})$. Since we have $\lambda \neq 0$ it follows that $\operatorname{Re} \langle \psi, G_\lambda^g(\cdot, p) \varphi \rangle$ vanishes identically on $M \setminus \{p\}$.

Our aim is now to conclude that all the derivatives of ψ at the point p vanish. Then by Theorem 6.2.8 it follows that ψ is identically zero, which is a contradiction. In order to show this we choose a local parametrization $\rho: V \rightarrow U$ of M by Riemannian normal coordinates, where $U \subset M$ is an open neighborhood of p , $V \subset \mathbb{R}^n$ is an open neighborhood of 0 and $\rho(0) = p$. Furthermore let

$$\beta: \Sigma \mathbb{R}^n|_V \rightarrow \Sigma^g M|_U, \quad A: C^\infty(\Sigma^g M|_U) \rightarrow C^\infty(\Sigma \mathbb{R}^n|_V)$$

denote the maps which send a spinor to its corresponding spinor in the Bourguignon-Gauduchon trivialization defined in Section 5.1. We show by induction that $\nabla^r A\psi(0) = 0$ for all $r \in \mathbb{N}$, where ∇ denotes the covariant derivative on $\Sigma\mathbb{R}^n$. The case $r = 0$ is clear.

Let $r \geq 1$ and assume that we have $\nabla^s A\psi(0) = 0$ for all $s \leq r - 1$. Let $(E_i)_{i=1}^n$ be the standard basis of \mathbb{R}^n . First consider the case $n = 2$. In the Bourguignon-Gauduchon trivialization we have

$$A\psi(x) = \frac{1}{r!} \sum_{j_1, \dots, j_r=1}^2 x_{j_1} \dots x_{j_r} \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) + O(|x|^{r+1})$$

by Taylor's formula and

$$A(G_\lambda^g(\cdot, p)\varphi)(x) = -\frac{1}{2\pi|x|^2} x \cdot \gamma - \frac{\lambda}{2\pi} \ln|x|\gamma + O(|x|^0)$$

by Theorem 5.3.10, where $\gamma := \beta^{-1}\varphi \in \Sigma_n$ is the constant spinor on \mathbb{R}^n corresponding to φ . It follows that

$$\begin{aligned} 0 &= -2\pi r! |x|^2 \operatorname{Re} \langle A(G_\lambda^g(\cdot, p)\varphi)(x), A\psi(x) \rangle \\ &= \sum_{i, j_1, \dots, j_r=1}^2 x_i x_{j_1} \dots x_{j_r} \operatorname{Re} \langle E_i \cdot \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle \\ &\quad + \lambda \sum_{j_1, \dots, j_r=1}^2 x_{j_1} \dots x_{j_r} |x|^2 \ln|x| \operatorname{Re} \langle \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle + O(|x|^{r+2}) \end{aligned}$$

and therefore

$$\begin{aligned} 0 &= \operatorname{Re} \langle E_i \cdot \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle \\ &\quad + \sum_{s=1}^r \operatorname{Re} \langle E_{j_s} \cdot \gamma, \nabla_{E_i} \nabla_{E_{j_1}} \dots \widehat{\nabla_{E_{j_s}}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle \end{aligned} \quad (6.3)$$

$$0 = \operatorname{Re} \langle \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle \quad (6.4)$$

for all $j_1, \dots, j_r, i \in \{1, 2\}$, where the hat means that the operator is left out. By (6.4) and Remark 6.2.11 there exist $a_{j_1, \dots, j_r, k}, b_{j_1, \dots, j_r} \in \mathbb{R}$ such that

$$\nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) = \sum_{k=1}^2 a_{j_1, \dots, j_r, k} E_k \cdot \gamma + b_{j_1, \dots, j_r} E_1 \cdot E_2 \cdot \gamma.$$

Observe that the coefficients $a_{j_1, \dots, j_r, k}$ are symmetric in the first r indices. We insert this into (6.3) and we obtain

$$0 = a_{j_1, \dots, j_r, i} + \sum_{k=1}^r a_{i, j_1, \dots, \widehat{j_k}, \dots, j_r, j_k} \quad (6.5)$$

for all $j_1, \dots, j_r, i \in \{1, 2\}$. On the other hand since $\psi \in \ker(D^g - \lambda)$ we find using the induction hypothesis

$$\begin{aligned}
0 &= \lambda \nabla_{E_{j_1}} \dots \nabla_{E_{j_{r-1}}} A\psi(0) \\
&= \nabla_{E_{j_1}} \dots \nabla_{E_{j_{r-1}}} \sum_{i=1}^2 E_i \cdot \nabla_{E_i} A\psi(0) \\
&= \sum_{i,k=1}^2 a_{j_1, \dots, j_{r-1}, i, k} E_i \cdot E_k \cdot \gamma + \sum_{i=1}^2 b_{j_1, \dots, j_{r-1}, i} E_i \cdot E_1 \cdot E_2 \cdot \gamma \\
&= -(a_{j_1, \dots, j_{r-1}, 1, 1} + a_{j_1, \dots, j_{r-1}, 2, 2})\gamma \\
&\quad + (a_{j_1, \dots, j_{r-1}, 1, 2} - a_{j_1, \dots, j_{r-1}, 2, 1})E_1 \cdot E_2 \cdot \gamma \\
&\quad + b_{j_1, \dots, j_{r-1}, 2}E_1 \cdot \gamma - b_{j_1, \dots, j_{r-1}, 1}E_2 \cdot \gamma
\end{aligned} \tag{6.6}$$

for all $j_1, \dots, j_{r-1} \in \{1, 2\}$. We conclude $b_{j_1, \dots, j_r} = 0$ for all $j_1, \dots, j_r \in \{1, 2\}$. Next consider $a_{j_1, \dots, j_r, i}$ with fixed $j_1, \dots, j_r, i \in \{1, 2\}$. If we have $j_k = i$ for all $k \in \{1, \dots, r\}$, then by (6.5) we know that $a_{j_1, \dots, j_r, i} = 0$. If there exists k such that $j_k \neq i$ it follows from the coefficient of $E_1 \cdot E_2 \cdot \gamma$ in (6.6) that

$$a_{i, j_1, \dots, \widehat{j_k}, \dots, j_r, j_k} = a_{j_1, \dots, j_r, i}.$$

Again (6.5) yields $a_{j_1, \dots, j_r, i} = 0$. We conclude that all $a_{j_1, \dots, j_r, i}$ vanish and that $\nabla^r A\psi(0) = 0$. This proves the assertion in the case $n = 2$.

Next consider $n = 3$. In the Bourguignon-Gauduchon trivialization we have

$$\begin{aligned}
A\psi(x) &= \frac{1}{r!} \sum_{j_1, \dots, j_r=1}^3 x_{j_1} \dots x_{j_r} \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \\
&\quad + \frac{1}{(r+1)!} \sum_{j_1, \dots, j_r, i=1}^3 x_{j_1} \dots x_{j_r} x_i \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} \nabla_{E_i} A\psi(0) + o(|x|^{r+1})
\end{aligned}$$

by Taylor's formula and

$$A(G_\lambda^g(\cdot, p)\varphi)(x) = -\frac{1}{4\pi|x|^3}x \cdot \gamma + \frac{\lambda}{4\pi|x|}\gamma + o(|x|^{-s})$$

for every $s > 0$ by Theorem 5.3.10, where γ is as above. It follows that

$$\begin{aligned}
0 &= -4\pi r!|x|^3 \operatorname{Re}\langle A(G_\lambda^g(\cdot, p)\varphi)(x), A\psi(x) \rangle \\
&= \sum_{i, j_1, \dots, j_r=1}^3 x_{j_1} \dots x_{j_r} x_i \operatorname{Re}\langle E_i \cdot \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{r+1} \sum_{i,j_1,\dots,j_r,m=1}^3 x_{j_1}\dots x_{j_r} x_i x_m \operatorname{Re}\langle E_i \cdot \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} \nabla_{E_m} A\psi(0) \rangle \\
& - \lambda \sum_{j_1,\dots,j_r=1}^3 x_{j_1}\dots x_{j_r} |x|^2 \operatorname{Re}\langle \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle + o(|x|^{r+2}). \quad (6.7)
\end{aligned}$$

From the first term on the right hand side we obtain

$$\begin{aligned}
0 & = \operatorname{Re}\langle E_i \cdot \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle \\
& + \sum_{s=1}^r \operatorname{Re}\langle E_{j_s} \cdot \gamma, \nabla_{E_i} \nabla_{E_{j_1}} \dots \widehat{\nabla_{E_{j_s}}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle. \quad (6.8)
\end{aligned}$$

for all $j_1, \dots, j_r, i \in \{1, 2, 3\}$, where the hat means that the operator is left out. Our next aim is to obtain an analogue of (6.4) from the second and third term on the right hand side of (6.7). It is more difficult than in the case $n = 2$, since derivatives of both orders r and $r + 1$ appear. The equation (5.2) reads

$$\begin{aligned}
\lambda A\psi & = D^{g_{\text{eucl}}} A\psi + \sum_{i,j=1}^3 (B_i^j - \delta_i^j) E_i \cdot \nabla_{E_j} A\psi \\
& + \frac{1}{4} \sum_{i,j,k=1}^3 \tilde{\Gamma}_{ij}^k E_i \cdot E_j \cdot E_k \cdot A\psi.
\end{aligned}$$

Using (5.3), (5.4) and that $|A\psi(x)|_{g_{\text{eucl}}} = O(|x|^r)$ as $x \rightarrow 0$ we find

$$\lambda A\psi = D^{g_{\text{eucl}}} A\psi + O(|x|^{r+1})$$

and therefore

$$\nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} D^{g_{\text{eucl}}} A\psi(0) = \lambda \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \quad (6.9)$$

for all $j_1, \dots, j_r \in \{1, 2, 3\}$. Using the equation (5.1) we find

$$A\nabla_{e_i}^g \psi = \nabla_{E_i} A\psi + O(|x|^{r+1}), \quad A\nabla_{e_i}^g \nabla_{e_j}^g \psi = \nabla_{E_i} \nabla_{E_j} A\psi + O(|x|^r)$$

for all $i, j \in \{1, 2, 3\}$. Since by definition $d\rho|_x^{-1}(e_i) = E_i + O(|x|^2)$ the second term in the local formula (2.5) for $\nabla^* \nabla$ vanishes at p and therefore we find

$$A\nabla^* \nabla \psi = - \sum_{i=1}^3 \nabla_{E_i} \nabla_{E_i} A\psi + O(|x|^r).$$

From the Schrödinger-Lichnerowicz formula (2.6) it follows that

$$\lambda^2 A\psi - \frac{\text{scal}}{4} A\psi = - \sum_{i=1}^3 \nabla_{E_i} \nabla_{E_i} A\psi + O(|x|^r)$$

and thus

$$\nabla_{E_{j_1}} \dots \nabla_{E_{j_{r-1}}} \sum_{i=1}^3 \nabla_{E_i} \nabla_{E_i} A\psi(0) = 0 \quad (6.10)$$

for all $j_1, \dots, j_{r-1} \in \{1, 2, 3\}$. Now recall the second and third term on the right hand side of (6.7)

$$\begin{aligned} 0 &= \frac{1}{r+1} \sum_{j_1, \dots, j_r, i, m=1}^3 x_{j_1} \dots x_{j_r} x_i x_m \text{Re} \langle E_i \cdot \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} \nabla_{E_m} A\psi(0) \rangle \\ &\quad - \lambda \sum_{j_1, \dots, j_r=1}^3 x_{j_1} \dots x_{j_r} |x|^2 \text{Re} \langle \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle \end{aligned}$$

and let $k_1, k_2, k_3 \in \mathbb{N}$ such that $k_1 + k_2 + k_3 = r$. Then from the coefficient of $x_1^{k_1+2} x_2^{k_2} x_3^{k_3}$ we find

$$0 = \text{Re} \langle E_1 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_1} A\psi(0) \rangle \frac{r!}{(k_1+1)!k_2!k_3!} \quad (I)$$

$$+ \text{Re} \langle E_2 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2-1} \nabla_{E_3}^{k_3} \nabla_{E_1}^2 A\psi(0) \rangle \frac{r!k_2}{(k_1+2)!k_2!k_3!} \quad (III)$$

$$+ \text{Re} \langle E_3 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3-1} \nabla_{E_1}^2 A\psi(0) \rangle \frac{r!k_3}{(k_1+2)!k_2!k_3!} \quad (IV)$$

$$- \lambda \text{Re} \langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle \frac{r!}{k_1!k_2!k_3!} \quad (V)$$

$$- \lambda \text{Re} \langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2-2} \nabla_{E_3}^{k_3} \nabla_{E_1}^2 A\psi(0) \rangle \frac{r!k_2(k_2-1)}{(k_1+2)!k_2!k_3!} \quad (VII)$$

$$- \lambda \text{Re} \langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3-2} \nabla_{E_1}^2 A\psi(0) \rangle \frac{r!k_3(k_3-1)}{(k_1+2)!k_2!k_3!} \quad (VIII).$$

From the coefficient of $x_1^{k_1} x_2^{k_2+2} x_3^{k_3}$ we find

$$0 = \text{Re} \langle E_1 \cdot \gamma, \nabla_{E_1}^{k_1-1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_2}^2 A\psi(0) \rangle \frac{r!k_1}{k_1!(k_2+2)!k_3!} \quad (II)$$

$$+ \text{Re} \langle E_2 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_2} A\psi(0) \rangle \frac{r!}{k_1!(k_2+1)!k_3!} \quad (I)$$

$$+ \operatorname{Re}\langle E_3 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3-1} \nabla_{E_2}^2 A\psi(0) \rangle \frac{r!k_3}{k_1!(k_2+2)!k_3!} \quad (IV)$$

$$- \lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1-2} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_2}^2 A\psi(0) \rangle \frac{r!k_1(k_1-1)}{k_1!(k_2+2)!k_3!} \quad (VI)$$

$$- \lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle \frac{r!}{k_1!k_2!k_3!} \quad (V)$$

$$- \lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3-2} \nabla_{E_2}^2 A\psi(0) \rangle \frac{r!k_3(k_3-1)}{k_1!(k_2+2)!k_3!} \quad (VIII).$$

From the coefficient of $x_1^{k_1} x_2^{k_2} x_3^{k_3+2}$ we find

$$0 = \operatorname{Re}\langle E_1 \cdot \gamma, \nabla_{E_1}^{k_1-1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_3}^2 A\psi(0) \rangle \frac{r!k_1}{k_1!k_2!(k_3+2)!} \quad (II)$$

$$+ \operatorname{Re}\langle E_2 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2-1} \nabla_{E_3}^{k_3} \nabla_{E_3}^2 A\psi(0) \rangle \frac{r!k_2}{k_1!k_2!(k_3+2)!} \quad (III)$$

$$+ \operatorname{Re}\langle E_3 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_3} A\psi(0) \rangle \frac{r!}{k_1!k_2!(k_3+1)!} \quad (I)$$

$$- \lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1-2} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_3}^2 A\psi(0) \rangle \frac{r!k_1(k_1-1)}{k_1!k_2!(k_3+2)!} \quad (VI)$$

$$- \lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2-2} \nabla_{E_3}^{k_3} \nabla_{E_3}^2 A\psi(0) \rangle \frac{r!k_2(k_2-1)}{k_1!k_2!(k_3+2)!} \quad (VII)$$

$$- \lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle \frac{r!}{k_1!k_2!k_3!} \quad (V).$$

We multiply the first equation with $\frac{(k_1+2)!k_2!k_3!}{r!}$, the second equation with $\frac{k_1!(k_2+2)!k_3!}{r!}$ and the third equation with $\frac{k_1!k_2!(k_3+2)!}{r!}$ and then add the multiplied equations. If we consider the lines with the same Roman numbers separately and use (6.9), (6.10), then we find

$$\begin{aligned} 0 &= -2\lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle \\ &+ \operatorname{Re}\langle E_1 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_1} A\psi(0) \rangle k_1 \\ &+ \operatorname{Re}\langle E_2 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_2} A\psi(0) \rangle k_2 \\ &+ \operatorname{Re}\langle E_3 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_3} A\psi(0) \rangle k_3 \end{aligned} \quad (I)$$

$$- \operatorname{Re}\langle E_1 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_1} A\psi(0) \rangle k_1 \quad (II)$$

$$- \operatorname{Re}\langle E_2 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_2} A\psi(0) \rangle k_2 \quad (III)$$

$$- \operatorname{Re}\langle E_3 \cdot \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} \nabla_{E_3} A\psi(0) \rangle k_3 \quad (IV)$$

$$- \lambda \operatorname{Re}\langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle \sum_{i=1}^3 (k_i+2)(k_i+1) \quad (V)$$

$$+ \lambda \operatorname{Re} \langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle k_1(k_1 - 1) \quad (VI)$$

$$+ \lambda \operatorname{Re} \langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle k_2(k_2 - 1) \quad (VII)$$

$$+ \lambda \operatorname{Re} \langle \gamma, \nabla_{E_1}^{k_1} \nabla_{E_2}^{k_2} \nabla_{E_3}^{k_3} A\psi(0) \rangle k_3(k_3 - 1) \quad (VIII).$$

Therefore we obtain the analogue of (6.4) namely

$$\operatorname{Re} \langle \gamma, \nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) \rangle = 0$$

for all $j_1, \dots, j_r \in \{1, 2, 3\}$. Thus there exist $a_{j_1, \dots, j_r, k} \in \mathbb{R}$ such that

$$\nabla_{E_{j_1}} \dots \nabla_{E_{j_r}} A\psi(0) = \sum_{k=1}^3 a_{j_1, \dots, j_r, k} E_k \cdot \gamma.$$

Observe that the coefficients $a_{j_1, \dots, j_r, k}$ are symmetric in the first r indices. We insert this into (6.8) and we obtain

$$0 = a_{j_1, \dots, j_r, i} + \sum_{k=1}^r a_{i, j_1, \dots, \widehat{j}_k, \dots, j_r, j_k} \quad (6.11)$$

for all $j_1, \dots, j_r, i \in \{1, 2, 3\}$. On the other hand since $\psi \in \ker(D^g - \lambda)$ we find using the induction hypothesis

$$\begin{aligned} 0 &= \lambda \nabla_{E_{j_1}} \dots \nabla_{E_{j_{r-1}}} A\psi(0) \\ &= \nabla_{E_{j_1}} \dots \nabla_{E_{j_{r-1}}} \sum_{i=1}^3 E_i \cdot \nabla_{E_i} A\psi(0) \\ &= \sum_{i, k=1}^3 a_{j_1, \dots, j_{r-1}, i, k} E_i \cdot E_k \cdot \gamma \\ &= - \sum_{i=1}^3 a_{j_1, \dots, j_{r-1}, i, i} \gamma \\ &\quad + \sum_{\substack{i, k=1 \\ i < k}}^3 (a_{j_1, \dots, j_{r-1}, i, k} - a_{j_1, \dots, j_{r-1}, k, i}) E_i \cdot E_k \cdot \gamma \end{aligned} \quad (6.12)$$

for all $j_1, \dots, j_{r-1} \in \{1, 2, 3\}$. Consider $a_{j_1, \dots, j_r, i}$ with $j_1, \dots, j_r, i \in \{1, 2, 3\}$. If $j_k = i$ for all $k \in \{1, \dots, r\}$ then by (6.11) we know that $a_{j_1, \dots, j_r, i} = 0$. If there exists k such that $j_k \neq i$ it follows from the coefficient of $E_{j_k} \cdot E_i \cdot \gamma$ in (6.12) that

$$a_{i, j_1, \dots, \widehat{j}_k, \dots, j_r, j_k} = a_{j_1, \dots, j_r, i}.$$

Again (6.11) yields $a_{j_1, \dots, j_r, i} = 0$. We conclude that all $a_{j_1, \dots, j_r, i}$ vanish and that $\nabla^r A\psi(0) = 0$. This proves the assertion in the case $n = 3$. \square

Remark 6.2.13. *It is not clear how to prove this lemma for $n \geq 4$. Namely the condition $\operatorname{Re}\langle \gamma, \nabla_{E_i} A\psi(0) \rangle = 0$ for all i leads to*

$$\nabla_{E_i} A\psi(0) = \sum_{k=1}^n a_{ik} E_k \cdot \gamma + \sum_{k=1}^n b_{ik} \cdot \gamma$$

with $a_{ik} \in \mathbb{R}$ and $b_{ik} \in \operatorname{Cl}(n)$. As above it follows that $a_{ik} = -a_{ki}$ for all i, k and furthermore

$$\begin{aligned} 0 &= \lambda A\psi(0) \\ &= \sum_{i=1}^n E_i \cdot \nabla_{E_i} A\psi(0) \\ &= 2 \sum_{\substack{i,k=1 \\ i < k}}^n a_{ik} E_i \cdot E_k \cdot \gamma + \sum_{i,k=1}^n E_i \cdot b_{ik} \cdot \gamma. \end{aligned}$$

But for $n \geq 4$ the spinors $E_1 \cdot E_2 \cdot \gamma$ and $E_3 \cdot E_4 \cdot \gamma$ are not linearly independent in general. Thus we cannot conclude immediately that all the a_{ik} vanish.

Proof of Theorem 6.2.6. Let $k, m \in \mathbb{N} \setminus \{0\}$, let $g \in R(M)$ and equip $[g]$ with the C^k -topology. Let $U \subset [g]$ be open. In order to show that $N_m(M) \cap [g]$ is dense in $[g]$ we have to show that $U \cap N_m(M)$ is not empty. Since $S_m(M) \cap [g]$ is dense in $[g]$, there exists a metric in $U \cap S_m(M)$, which we denote again by g . Let λ be one of the eigenvalues $\{\lambda_1^\pm, \dots, \lambda_m^\pm\}$ of D^g and let ψ be an eigenspinor corresponding to λ . Choose an open neighborhood $V \subset [g]$ of g which is contained in $U \cap S_m(M)$ and define

$$F_\psi : V \rightarrow C^\infty(\Sigma^g M)$$

as in Lemma 6.2.9

Next we show that a suitable restriction of F_ψ^{ev} is transverse to the zero section of $\Sigma^g M$. Let $p \in M$ with $\psi(p) = 0$. By Lemma 6.2.12 there exists an open neighborhood $U_p \subset M$ of p and $f_{p,1}, \dots, f_{p,4} \in C^\infty(M, \mathbb{R})$ and an open neighborhood $V_p \subset V_{f_{p,1}, \dots, f_{p,4}}$ of g such that for all $(h, q) \in V_p \times U_p$ we have

$$\pi_1(dF_\psi^{ev}|_{(h,q)}(T_h V_{f_{p,1}, \dots, f_{p,4}} \oplus \{0\})) = \Sigma_q^g M. \quad (6.13)$$

Since the zero set of ψ is compact, there exist points $p_1, \dots, p_r \in M$ and open neighborhoods $U_{p_i} \subset M$ of p_i and $f_{p_i,1}, \dots, f_{p_i,4} \in C^\infty(M, \mathbb{R})$ and open neighborhoods $V_{p_i} \subset V_{f_{p_i,1}, \dots, f_{p_i,4}}$ of g , $1 \leq i \leq r$, such that the open neighborhoods U_{p_1}, \dots, U_{p_r} cover the zero set of ψ and such that for every i the equation (6.13)

holds for all $(h, q) \in V_{p_i} \times U_{p_i}$. We label the functions $f_{p_i, j}$ by f_1, \dots, f_{4r} and define

$$V_{f_1 \dots f_{4r}} := \left\{ \left(1 + \sum_{i=1}^{4r} t_i f_i \right) g \mid t_i \in \mathbb{R} \right\} \cap V.$$

and $F_\psi: V_{f_1 \dots f_{4r}} \rightarrow C^\infty(\Sigma^g M)$ as in Lemma 6.2.9. Since M is compact there exists $C > 0$ such that $|\psi|_g \geq C$ on the complement of the union of the sets U_{p_i} . Thus we can find an open neighborhood $V_\psi \subset V_{f_1 \dots f_{4r}}$ of g such that the equation (6.13) holds for all $(h, q) \in V_\psi \times M$. It follows that the restriction of F_ψ^{ev} to $V_\psi \times M$ is transverse to the zero section of $\Sigma^g M$.

Define W_ψ as the subset of all $h \in V_\psi$ such that $F_\psi(h)$ is nowhere zero on M . By Remark 6.2.10 this condition is equivalent to the condition that $F_\psi(h)$ is transverse to the zero section of $\Sigma^g M$. By Theorem 6.1.6 the set W_ψ is dense in V_ψ . Since the zero section is closed in $\Sigma^g M$ and F_ψ^{ev} is continuous, the set W_ψ is also open in V_ψ . If $h \in W_\psi$, then the eigenspinor $F_\psi(h)$ of $D^{g, h}$ is nowhere zero on M and it corresponds to a simple eigenvalue of $D^{g, h}$. Thus all the eigenspinors of $D^{g, h}$ corresponding to this eigenvalue are nowhere zero on M .

For every one of the finitely many eigenvalues $\lambda_1^\pm, \dots, \lambda_m^\pm$ of D^g we choose an eigenspinor ψ and obtain an open subset $W_\psi \subset V$ as above. Let W be the intersection of these open subsets W_ψ . It is not empty, since the W_ψ are dense in a neighborhood of g . If $h \in W$, then all the eigenspinors of D^h corresponding to the eigenvalues $\lambda_1^\pm, \dots, \lambda_m^\pm$ of D^h are nowhere zero on M . Since $W \subset U$ by construction, we have $U \cap N_m(M) \neq \emptyset$. Thus $N_m(M) \cap [g]$ is dense in $[g]$. We have already seen that $N_m(M) \cap [g]$ is open in $[g]$. \square

6.3 Mass endomorphism in dimension 3

Let M be a closed spin manifold of dimension n with a fixed spin structure and let $R(M)$ be the set of all smooth Riemannian metrics on M equipped with the C^1 -topology. We define

$$R^*(M) := \{g \in R(M) \mid \ker(D^g) = 0\}.$$

It follows from the Atiyah-Singer index theorem that there exist spin manifolds M and spin structures Θ on M such that for every Riemannian metric g on M there exist harmonic spinors on (M, g, Θ) (see e. g. [LM], Section 3 in [BD]). This shows that $R^*(M)$ can be empty. Let $p \in M$ and define

$$R_p(M) := \{g \in R^*(M) \mid g \text{ is flat on an open neighborhood of } p\}.$$

We consider the case $n = 3$. Then by Theorem 1.2 in [Ma] for every spin structure the subset $R^*(M)$ of $R(M)$ is generic. Let g be a Riemannian metric on M , which is flat on an open neighborhood of p . It is shown in [ADH2] that by changing the metric g on an arbitrarily small open subset of M one obtains a metric in $R^*(M)$. Thus for every closed spin manifold M of dimension 3 with a fixed spin structure the set $R_p(M)$ is not empty for every $p \in M$. For every $g \in R_p(M)$ we define the mass endomorphism $m_p^g \in \text{End}(\Sigma_p^g M)$ as in Section 5.4. Our aim is now to find Riemannian metrics on M , for which the mass endomorphism does not vanish. Thus we define

$$S_p(M) := \{g \in R_p(M) \mid m_p^g \neq 0\}.$$

In this section we prove the following result which has been published in [He].

Theorem 6.3.1. *Let M be a closed spin manifold of dimension 3 with a fixed spin structure and let $p \in M$. Then $S_p(M)$ is dense in $R_p(M)$.*

Remark 6.3.2. *If M is a closed spin manifold of dimension 2 and if the mass endomorphism in $p \in M$ can be defined, then by [AHM] it always vanishes. On the other hand it is conjectured that Theorem 6.3.1 holds for every closed spin manifold M of dimension $n \geq 3$ if $R^*(M)$ is not empty. A proof has been proposed in [ADHH]. The idea is to use surgery methods to construct a Riemannian metric g on M which is flat near some point p and has nonzero mass endomorphism. Then from perturbation theory it follows that $S_p(M)$ is open and dense in $R_p(M)$.*

We will need the following properties of the energy momentum tensor which was defined in Section 2.3.

Lemma 6.3.3. *Let g and $h = e^{2u}g$ be conformally related Riemannian metrics, $u \in C^\infty(M, \mathbb{R})$. Let $\psi \in C^\infty(\Sigma^g M)$ and let $\beta := \beta_{g,h}$ as in Section 2.3. Let $f \in C^\infty(M, \mathbb{R})$. Then we have*

$$Q_{f\psi} = f^2 Q_\psi, \quad Q_{\beta\psi} = e^u Q_\psi.$$

Proof. The first equation follows, since $\text{Re}\langle Y(f)X \cdot \psi, \psi \rangle = 0$ for all $X, Y \in TM$. Recall that we have $b_{g,h}X = e^{-u}X$ for all $X \in TM$ and therefore $\beta(X \cdot \psi) = e^{-u}X \cdot \beta\psi$. It follows that

$$\begin{aligned} \langle X \cdot \nabla_Y^h \beta\psi, \beta\psi \rangle_h &= \langle X \cdot \beta(\nabla_Y^g \psi - \frac{1}{2}Y \cdot \text{grad}^g(u) \cdot \psi - \frac{1}{2}Y(u)\psi), \beta\psi \rangle_h \\ &= e^u \langle X \cdot \nabla_Y^g \psi, \psi \rangle_g - \frac{e^u}{2} \langle X \cdot Y \cdot \text{grad}^g(u) \cdot \psi, \psi \rangle_g \\ &\quad - \frac{e^u}{2} Y(u) \langle X \cdot \psi, \psi \rangle_g. \end{aligned}$$

If we add the corresponding equation for $\langle Y \cdot \nabla_X^h \beta\psi, \beta\psi \rangle_h$ to this equation and use that $X \cdot Y + Y \cdot X = -2g(X, Y)$, we see that the sum of the second terms on the right hand side is purely imaginary. Since the third terms on the right hand side are each purely imaginary, the assertion follows. \square

Let $g \in R_p(M)$ and let $k \in C^\infty(\text{Sym}^2(T^*M))$ such that $k = 0$ on an open neighborhood U of p . Let $I \subset \mathbb{R}$ be an open interval around 0 such that for all $t \in I$ the tensor field $g_t := g + tk$ is a Riemannian metric on M and is in $R^*(M)$. Then we have $g_t \in R_p(M)$ for all $t \in I$. For every t we obtain an isomorphism of spinor bundles

$$\bar{\beta}_{g, g_t} : \Sigma^g M \rightarrow \Sigma^{g_t} M$$

as in Lemma 2.3.5. Let $\varepsilon > 0$ such that $B_{2\varepsilon}(p) \subset U$. Recall from Section 5.4 that Green's function G^{g_t} for D^{g_t} on $M \setminus \{p\}$ can be written as

$$G^{g_t}(x, p) \bar{\beta}_{g, g_t} \varphi = (A^{g_t})^{-1} (\eta G^{g_{\text{eucl}}}(\cdot, 0) \gamma_t)(x) + w^{g_t}(x, p) \bar{\beta}_{g, g_t} \varphi, \quad (6.14)$$

where $\eta: \mathbb{R}^n \rightarrow [0, 1]$ is a smooth function such that $\eta \equiv 1$ on $B_\varepsilon(0)$ and $\text{supp}(\eta) \subset B_{2\varepsilon}(0)$ and where $\varphi \in \Sigma_p^g M$ and $\gamma_t := (\beta^{g_t})^{-1} \bar{\beta}_{g, g_t} \varphi$. Recall that the mass endomorphism for the metric g_t at p is by definition $m_p^{g_t} = w^{g_t}(p, p)$. We define a spinor $w_\varphi^{g_t} \in C^\infty(\Sigma^g M)$ by

$$w_\varphi^{g_t}(x) := \bar{\beta}_{g_t, g} w^{g_t}(x, p) \bar{\beta}_{g, g_t} \varphi$$

and we define $m^{g, g_t} \varphi := w_\varphi^{g_t}(p) = \bar{\beta}_{g_t, g} m_p^{g_t} \bar{\beta}_{g, g_t} \varphi$ for all t .

Lemma 6.3.4. *We have*

$$\frac{d}{dt} \langle m^{g, g_t} \varphi, \varphi \rangle|_{t=0} = \frac{1}{2} \int_{M \setminus \{p\}} (k, Q_{G^g(\cdot, p)\varphi}) d\mathbf{v}^g,$$

where (\cdot, \cdot) denotes the standard pointwise inner product of $(2, 0)$ tensor fields.

Proof. Since $g_t = g$ on $\text{supp}(\eta)$ for all t , the first term on the right hand side of (6.14) is independent of t . Therefore we find

$$\bar{\beta}_{g_t, g} G^{g_t}(x, p) \bar{\beta}_{g, g_t} \varphi = (A^g)^{-1} (\eta G^{g_{\text{eucl}}}(\cdot, 0) \gamma)(x) + w_\varphi^{g_t}(x)$$

for all $x \in M \setminus \{p\}$. We apply D^{g, g_t} defined as in Section 2.3 and since $g_t = g$ on $\text{supp}(\eta)$ we find

$$\delta_p = D^g (A^g)^{-1} (\eta G^{g_{\text{eucl}}}(\cdot, 0) \gamma) + D^{g, g_t} w_\varphi^{g_t}.$$

We take the derivative with respect to t at $t = 0$ and we obtain

$$0 = \left(\frac{d}{dt}D^{g,g_t}\Big|_{t=0}\right)w_\varphi^g + D^g\left(\frac{d}{dt}w_\varphi^{g_t}\Big|_{t=0}\right).$$

By the formula for Green's function (5.7) and since $\ker(D^g) = 0$, we find that

$$\begin{aligned} \frac{d}{dt}\langle m^{g,g_t}\varphi, \varphi\rangle\Big|_{t=0} &= \left\langle \frac{d}{dt}w_\varphi^{g_t}(p)\Big|_{t=0}, \varphi\right\rangle\Big|_{t=0} \\ &= \int_{M\setminus\{p\}} \langle D^g\left(\frac{d}{dt}w_\varphi^{g_t}\Big|_{t=0}\right), G^g(\cdot, p)\varphi\rangle dv^g \\ &= - \int_{M\setminus\{p\}} \left\langle \frac{d}{dt}D^{g,g_t}\Big|_{t=0}w_\varphi^g, G^g(\cdot, p)\varphi\right\rangle dv^g \end{aligned}$$

Recall that if $(e_i)_{i=1}^n$ is a local orthonormal frame, then by (2.8) we have locally

$$\frac{d}{dt}D^{g,g_t}\Big|_{t=0}w_\varphi^g = -\frac{1}{2}\sum_{i=1}^n e_i \cdot \nabla_{a_{g,k}(e_i)}^g w_\varphi^g - \frac{1}{4}\sum_{i=1}^n \operatorname{div}^g(k)(e_i)e_i \cdot w_\varphi^g.$$

Since $k = 0$ on $B_{2\varepsilon}(p)$ this spinor vanishes on $B_{2\varepsilon}(p)$. On the other hand the spinors w_φ^g and $G^g(\cdot, p)\varphi$ coincide on $M \setminus B_{2\varepsilon}(p)$. Thus we find

$$\frac{d}{dt}\langle m^{g,g_t}\varphi, \varphi\rangle\Big|_{t=0} = - \int_{M\setminus B_{2\varepsilon}(p)} \left\langle \frac{d}{dt}D^{g,g_t}\Big|_{t=0}G^g(\cdot, p)\varphi, G^g(\cdot, p)\varphi\right\rangle dv^g.$$

The assertion is now obtained in the way that (2.10) was obtained from (2.9). \square

Proof of Theorem 6.3.1. Assume that the claim is wrong. Then there is an open subset $Q \subset R_p(M)$ such that for all $g \in Q$ we have $m_p^g = 0$. Let $g \in Q$ be flat on an open neighborhood $U \subset M$ of p . By a small deformation of g on $M \setminus U$ we obtain a Riemannian metric, which is not conformally flat on $M \setminus U$ and is in Q . We denote the deformed Riemannian metric again by g . Thus there exists an open subset $V \subset M \setminus U$, such that g is nowhere conformally flat on V .

Let $k \in C^\infty(\operatorname{Sym}^2(T^*M))$ such that $k = 0$ on an open neighborhood of p and let $I \subset \mathbb{R}$ be an open interval around 0 such that for all $t \in I$ the tensor field $g_t := g + tk$ is a Riemannian metric on M and is in Q . By our assumption we have $m_p^{g_t} = 0$ for all $t \in I$. For every $\varphi \in \Sigma_p^g M$ it follows that $\int_{M\setminus\{p\}} (k, Q_{G^g(\cdot, p)\varphi}) dv^g = 0$ by Lemma 6.3.4. Since this holds for every $k \in C^\infty(\operatorname{Sym}^2(T^*M))$ which is zero on an open neighborhood of p , it holds

for $k = fQ_{G^g(\cdot, p)\varphi}$, where f is an arbitrary smooth function with $p \notin \text{supp}(f)$. We conclude that $Q_{G^g(\cdot, p)\varphi} = 0$ on $M \setminus \{p\}$. Let

$$W := \{x \in M \setminus \{p\} \mid G^g(x, p)\varphi \neq 0\}.$$

Then W is an open subset of M . The function $u: W \rightarrow \mathbb{R}$, defined by

$$u(x) := \ln |G^g(x, p)\varphi|_g$$

is smooth on W , and thus $h := e^{2u}g$ defines a Riemannian metric on W . We obtain an isomorphism of spinor bundles

$$\beta_{g,h}: \Sigma^g W \rightarrow \Sigma^h W,$$

which is a fibrewise isometry as in Lemma 2.3.4. The spinor

$$\psi := e^{-u}\beta_{g,h}G^g(\cdot, p)\varphi \in C^\infty(\Sigma^h W)$$

satisfies $D^h\psi = 0$ by (2.12). By definition of u we have $|\psi|_h \equiv 1$ on W . Furthermore by Lemma 6.3.3 we find $Q_\psi = 0$ on W .

Let $(e_i)_{i=1}^3$ be a local orthonormal frame of TW defined on an open subset $S \subset W$. Then for every $x \in S$ the system

$$\{\psi(x), e_1 \cdot \psi(x), e_2 \cdot \psi(x), e_3 \cdot \psi(x)\}$$

is a real basis of $\Sigma_x^h W$, which is orthonormal with respect to the real scalar product $\text{Re}\langle \cdot, \cdot \rangle$ (see Remark 6.2.11). Since $|\psi|_h$ is constant on W , we have

$$\text{Re}\langle \nabla_X^h \psi, \psi \rangle = 0$$

for all $X \in TM$. It follows that there exists an endomorphism A of TW such that for all $X \in TW$ we have $\nabla_X^h \psi = A(X) \cdot \psi$.

We will now show that in each fibre the endomorphism $A: T_x W \rightarrow T_x W$ is symmetric with respect to h using an observation from [AMM]. Since $\omega_{\mathbb{C}} = \text{Id}$ we have $e_1 \cdot e_2 \cdot e_3 = -\text{Id}$. In the following we abbreviate $\psi := \psi(x)$. Using (6.2) we calculate

$$\begin{aligned} h(Ae_2, e_1) &= \text{Re}\langle Ae_2 \cdot \psi, e_1 \cdot \psi \rangle \\ &= \text{Re}\langle \nabla_{e_2}^h \psi, e_1 \cdot \psi \rangle \\ &= -\text{Re}\langle e_2 \cdot \nabla_{e_2}^h \psi, e_3 \cdot \psi \rangle \\ &= \text{Re}\langle e_1 \cdot \nabla_{e_1}^h \psi, e_3 \cdot \psi \rangle + \text{Re}\langle e_3 \cdot \nabla_{e_3}^h \psi, e_3 \cdot \psi \rangle \\ &= -\text{Re}\langle e_2 \cdot \nabla_{e_1}^h \psi, \psi \rangle \\ &= \text{Re}\langle Ae_1 \cdot \psi, e_2 \cdot \psi \rangle \\ &= h(e_2, Ae_1). \end{aligned}$$

Similarly we obtain $h(Ae_1, e_3) = h(e_1, Ae_3)$ and $h(Ae_2, e_3) = h(e_2, Ae_3)$, i. e. A is symmetric with respect to h . Therefore, we can choose the basis vectors e_1, e_2, e_3 as eigenvectors of A , such that $Ae_j = \lambda_j e_j$, where $\lambda_j \in \mathbb{R}$. It follows that

$$0 = \operatorname{Re}\langle \psi, e_j \cdot \nabla_{e_j}^h \psi \rangle = -\lambda_j.$$

Hence, ψ is a parallel spinor on W . By [F4] the Riemannian manifold (W, h) is Ricci flat. Since $\dim W = 3$ it follows that (W, h) is flat. However, by definition of the metric g there exists an open subset $V \subset M \setminus U$, such that g is nowhere conformally flat on V . By Theorem 6.2.8 the spinor $G^g(\cdot, p)\varphi$ is not identically zero on V . Thus on $V \cap W \neq \emptyset$ the metric h cannot be flat. This is a contradiction. \square

Appendix A

Analytic perturbation theory

Results in perturbation theory show that if a deformation of the Riemannian metric depends real-analytically on a parameter, then for a given eigenspinor ψ of D^g one obtains a real analytic family of eigenspinors for the perturbed Dirac operators which extends ψ . There is a short note concerning this fact in [BG], but we will give a more precise explanation here following the book [K].

In this section $(X, \|\cdot\|_X)$, $(Y, \|\cdot\|_Y)$ will be complex Banach spaces. The space of bounded linear operators $X \rightarrow Y$ is denoted by $B(X, Y)$. It is equipped with the usual operator norm $\|\cdot\|_{B(X, Y)}$. The space of closed linear operators $X \rightarrow Y$ is denoted by $C(X, Y)$. We will write $C(X) := C(X, X)$. The dual space of X is denoted by X^* . Furthermore $\Omega \subset \mathbb{C}$ will always denote an open and connected subset of the complex plane and $I \subset \mathbb{R}$ will denote an open interval.

Definition A.0.5. *A family of vectors $u(z) \in X$ which depend on $z \in \Omega$ is called holomorphic on Ω , if for each $z_0 \in \Omega$ there exists $u' \in X$ such that*

$$\lim_{h \rightarrow 0} \left\| \frac{u(z_0 + h) - u(z_0)}{h} - u' \right\|_X = 0.$$

The following theorem ([K], p. 139) gives a criterion for holomorphy of vectors.

Theorem A.0.6. *Let $u(z) \in X$ be a family of vectors which depend on $z \in \Omega$. If for all $f \in X^*$ the function $f(u(z))$ is holomorphic on Ω , then $u(z)$ is holomorphic on Ω .*

Proof. Let Γ be a positively oriented circle in Ω . The Cauchy integral formula gives

$$f(u(z)) = \frac{1}{2\pi i} \oint_{\Gamma} \frac{f(u(\zeta))}{\zeta - z} d\zeta$$

for all z in the interior of Γ . Let $h \neq 0$ such that $z + h$ is in the interior of Γ . Then we find

$$\frac{f(u(z+h)) - f(u(z))}{h} - \frac{d(f \circ u)(z)}{dz} = \frac{h}{2\pi i} \oint_{\Gamma} \frac{f(u(\zeta))}{(\zeta - z - h)(\zeta - z)^2} d\zeta.$$

But the function $f(u(z))$ is bounded on Γ since it is continuous. Thus there exists a constant $C > 0$ such that

$$\left| \frac{f(u(z+h)) - f(u(z))}{h} - \frac{d(f \circ u)(z)}{dz} \right| \leq C|h| \|f\|_{X^*}.$$

One can show that C can be chosen independently of f and that this estimate implies the convergence of $\frac{u(z+h)-u(z)}{h}$ as $h \rightarrow 0$. \square

Definition A.0.7. A family of operators $D(z) \in B(X, Y)$ which depend on $z \in \Omega$ is called holomorphic on Ω , if for each $z_0 \in \Omega$ there exists a bounded linear operator $D' \in B(X, Y)$ such that

$$\lim_{h \rightarrow 0} \left\| \frac{D(z_0 + h) - D(z_0)}{h} - D' \right\|_{B(X, Y)} = 0.$$

The next theorem ([K], p. 152) gives a criterion for holomorphy of bounded linear operators.

Theorem A.0.8. Let $D(z) \in B(X, Y)$ be a family of bounded linear operators which depend on $z \in \Omega$. If for all $u \in X$ and for all $g \in Y^*$ the function $g(D(z)u)$ is holomorphic on Ω , then $D(z)$ is holomorphic on Ω .

Proof. The proof is analogous to the proof of Theorem A.0.6. \square

We generalize the notion of a holomorphic family of operators to families of closed linear operators which are not necessarily bounded (see [K], p. 366).

Definition A.0.9. A family of operators $D(z) \in C(X, Y)$ which depend on $z \in \Omega$ is called holomorphic at $z_0 \in \Omega$, if there exists a third complex Banach space Z and two families $E(z) \in B(Z, X)$, $F(z) \in B(Z, Y)$, which are holomorphic at z_0 such that for all $z \in \Omega$ the operator $E(z)$ maps Z onto the domain of $D(z)$ one to one and such that for all $z \in \Omega$ we have $D(z)E(z) = F(z)$. The family $D(z)$ is called holomorphic on Ω , if it is holomorphic at every $z_0 \in \Omega$.

If X, Y are Hilbert spaces and $D \in C(X, Y)$, we denote by D^* the adjoint operator of D . The following definition is from [K], p. 385.

Definition A.0.10. Let H be a Hilbert space and let Ω be an open and connected subset of \mathbb{C} which is symmetric with respect to the real axis. A holomorphic family of operators $D(z) \in C(H)$ is called a self-adjoint holomorphic family, if for all $z \in \Omega$ the operator $D(z)$ is densely defined and if for all $z \in \Omega$ we have $D(z)^* = D(\bar{z})$.

If H is a Hilbert space, $D(z) \in C(H)$ a self-adjoint holomorphic family on Ω and if λ is an eigenvalue of $D(z_0)$ of finite multiplicity r for some $z_0 \in \mathbb{R}$, then one obtains families of eigenvalues $\lambda_i(z)$ of $D(z)$, $1 \leq i \leq r$, which are real-analytic in z on an open interval $I \subset \Omega \cap \mathbb{R}$ around z_0 and such that $\lambda_i(z_0) = \lambda$ for all i ([K], p. 386). In general I depends on λ . However for the so called holomorphic families of type (A), one obtains an open interval around z_0 , such that all the eigenvalues and eigenvectors are real-analytic on this interval.

Definition A.0.11. A family of operators $D(z) \in C(X, Y)$ which depend on $z \in \Omega$ is called holomorphic of type (A), if the domain $\text{dom}(D(z))$ is independent of z , and if for all $z \in \Omega$ and for all $u \in \text{dom}(D(z))$ the family of vectors $D(z)u \in Y$ is holomorphic on Ω .

Remark A.0.12. If the family $D(z) \in C(X, Y)$ is holomorphic of type (A) on Ω , then it is holomorphic on Ω in the sense of the definition above.

Proof (see [K], p. 375). Choose $z_0 \in \Omega$ and set $D := D(z_0)$. The space $Z := \text{dom}(D)$ with the norm $\|u\|_Z := \|u\|_X + \|Du\|_Y$ is then a Banach space. Define $E: Z \rightarrow X$ by $E(u) := u$. Since $\|u\|_X \leq \|u\|_Z$ we find that $E \in B(Z, X)$. For all $z \in \Omega$ define $F(z): Z \rightarrow Y$ by $F(z)u := D(z)u$. Using that $D(z)$ is closed and that $\|u\|_X \leq \|u\|_Z$ we find that $F(z) \in C(Z, Y)$. Furthermore since $\text{dom}(F(z)) = Z$ the closed graph theorem ([K], p. 166) implies that $F(z) \in B(Z, Y)$. By hypothesis $F(z)u = D(z)u$ is holomorphic for all $u \in Z$. By Theorem A.0.8 it follows that $F(z) \in B(Z, Y)$ is holomorphic. Since $E: Z \rightarrow \text{dom}(D)$ is one to one and $D(z)E = F(z)$ for all $z \in \Omega$ the assertion follows. \square

We have the following result due to Rellich.

Theorem A.0.13. Let $D(z) \in C(H)$ be a self-adjoint holomorphic family of type (A) defined on an open neighborhood $\Omega \subset \mathbb{C}$ of an open interval $I \subset \mathbb{R}$. Furthermore let $D(z)$ have compact resolvent for some $z \in \Omega$. Then there is a sequence of scalar-valued functions $(\lambda_n)_{n \in \mathbb{N}}$ and a sequence of vector-valued functions $(u_n)_{n \in \mathbb{N}}$ all real-analytic on I , such that for $t \in I$ the $\lambda_n(t)$ represent all the repeated eigenvalues of $D(t)$ and the $u_n(t)$ form a complete orthonormal family of the associated eigenvectors of $D(t)$.

Proof. see [K], Thm. VII, 3.9, p. 392. \square

The following Lemma is a criterion for a family of operators to be holomorphic of type (A).

Lemma A.0.14. *Let $D: X \rightarrow Y$ be a closable operator with domain $\text{dom}(D)$ and let $D_j, j \in \mathbb{N} \setminus \{0\}$, be operators $X \rightarrow Y$ with domain containing $\text{dom}(D)$. If there exist real constants $a, b, c \geq 0$ such that $\|D_j u\| \leq c^{j-1}(a\|u\| + b\|Du\|)$ for all $u \in \text{dom}(D)$ and all $j \in \mathbb{N} \setminus \{0\}$, then the series $D(z) = D + \sum_{j=1}^{\infty} D_j z^j$ for $z \in \mathbb{C}, |z| < (b+c)^{-1}$ defines a family of closable operators with domain $\text{dom}(D)$. Furthermore the closures of these operators form a holomorphic family of type (A).*

Proof. see [K], Thm. VII, 2.6, p. 377. \square

Now let M be a compact Riemannian spin manifold, let $I \subset \mathbb{R}$ be an open interval and let $(g_t)_{t \in I}$ be a family of smooth Riemannian metrics on M .

Definition A.0.15. *The family $(g_t)_{t \in I}$ is called real-analytic, if for every $t_0 \in I$ and for every $j \in \mathbb{N}$ there exists a section $h_j \in C^\infty(\text{Sym}^2(T^*M))$, such that for all charts (U, φ) of M and for all compact subsets $L \subset U$ there exists $\varepsilon_L > 0$, such that for all t with $|t - t_0| < \varepsilon_L$, for all coordinate vector fields $\partial_a, \partial_b, 1 \leq a, b \leq n$, and for all $r \in \mathbb{N}$ we have*

$$\|g_t(\partial_a, \partial_b) - \sum_{j=0}^N (t - t_0)^j h_j(\partial_a, \partial_b)\|_{C^r(L)} \rightarrow 0 \text{ as } N \rightarrow \infty.$$

We apply the criterion above to show that for a real-analytic family $(g_t)_{t \in I}$ of Riemannian metrics on a compact spin manifold M the family of operators D^{g, g_t} constructed in Section 2.3 can be extended to a holomorphic family of type (A). Here we consider the Dirac operator as a closed operator on the Sobolev space $H^m(\Sigma^g M)$ with domain $\text{dom}(D^g) = H^{m+1}(\Sigma^g M)$ for some $m \in \mathbb{N}$.

Lemma A.0.16. *Let (M, g, Θ) be a compact Riemannian spin manifold and let $D^g \in C(H^m(\Sigma^g M))$ be the Dirac operator. Let $(g_t)_{t \in I}$ be a real-analytic family of Riemannian metrics on M with $g_0 = g$. Then there exists an open neighborhood $\Omega \subset \mathbb{C}$ of 0 and a self-adjoint holomorphic family of type (A) of operators $D(z) \in C(H^m(\Sigma^g M))$ on Ω , such that for all $t \in \Omega \cap I$ we have $D(t) = D^{g, g_t}$.*

Proof. Since $(g_t)_{t \in I}$ is a real-analytic family of Riemannian metrics, there exist $h_j \in C^\infty(\text{Sym}^2(T^*M))$, $j \in \mathbb{N}$, such that for all charts (U, φ) of M and for all compact subsets $L \subset U$ there exists $\varepsilon_L > 0$ such that for all t with $|t| < \varepsilon_L$ and for all coordinate vector fields ∂_a, ∂_b , $1 \leq a, b \leq n$ and for all $r \in \mathbb{N}$ we have

$$\|g_t(\partial_a, \partial_b) - \sum_{j=0}^N t^j h_j(\partial_a, \partial_b)\|_{C^r(L)} \rightarrow 0, \quad N \rightarrow \infty.$$

As in Section 2.3 we define endomorphisms a_{g, g_t} and a_{g, h_j} , $j \in \mathbb{N}$, of TM such that for all v, w in TM we have

$$g(a_{g, g_t} v, w) = g_t(v, w), \quad g(a_{g, h_j} v, w) = h_j(v, w).$$

Let (U, φ) be a chart of M and let $L \subset U$ be compact. For $v \in TM$ we abbreviate $|v| := g(v, v)^{1/2}$. Let $(e_i)_{i=1}^n$ be a local g -orthonormal frame on L . Since $(g_t)_{t \in I}$ is real-analytic, there exists $\varepsilon_L > 0$ such that for all t with $|t| < \varepsilon_L$ we have

$$\begin{aligned} & \sup_{v \in TM|_L, |v| \leq 1} \left| a_{g, g_t} v - \sum_{j=0}^N t^j a_{g, h_j} v \right| \\ &= \sup_{v \in TM|_L, |v| \leq 1} \left| \sum_{i=1}^n g(a_{g, g_t} v, e_i) e_i - \sum_{i=1}^n \sum_{j=0}^N t^j g(a_{g, h_j} v, e_i) e_i \right| \\ &= \sup_{v \in TM|_L, |v| \leq 1} \left| \sum_{i=1}^n g_t(v, e_i) e_i - \sum_{i=1}^n \sum_{j=0}^N t^j h_j(v, e_i) e_i \right| \\ &\leq \sum_{i, j=1}^n \|g_t(e_j, e_i) - \sum_{m=0}^N t^m h_m(e_j, e_i)\|_{C^0(L)} \rightarrow 0, \quad N \rightarrow \infty. \end{aligned}$$

Therefore we find $\delta_L > 0$ such that for all t with $|t| < \delta_L$ and for all $v \in TM|_L$ the vector $b_{g, g_t} v$ is given by a power series, which converges uniformly in v , $|v| \leq 1$. Since M is compact there exists $\delta > 0$ and for all $j \in \mathbb{N}$ there exists $b_j \in \text{End}(TM)$ such that for all $t \in I$ with $|t| < \delta$ we have

$$\sup_{v \in TM, |v| \leq 1} \left| b_{g, g_t} v - \sum_{j=0}^N t^j b_j v \right| \rightarrow 0, \quad N \rightarrow \infty.$$

By similar arguments one finds that also for any fixed t the covariant derivatives of b_{g, g_t} of any order are given by convergent power series in this sense.

We insert the power series for b_{g,g_t} , g_t and f_{g,g_t} into (2.7) and define the operator D_j as the coefficient of t^j for $j \in \mathbb{N}$. Clearly $D_j \in C(H^m(\Sigma^g M))$ and for $\psi \in \text{dom}(D_k) = H^{m+1}(\Sigma^g M)$ the spinor $D_j\psi$ locally has the form

$$D_j\psi(x) = \sum_{i=1}^n e_i \cdot \nabla_{b_j(e_i)}^g \psi(x) + c_j(x) \cdot \psi(x),$$

where c_j are functions with values in the Clifford algebra. Since the series $\sum_{j=0}^{\infty} b_j t^j$ and $\sum_{j=0}^{\infty} c_j t^j$ converge uniformly in $x \in M$ for $|t| < \delta$, there exists $C > 0$ such that for all $j \geq 1$ we have

$$\sup\{|b_j v| \mid v \in TM, |v| \leq 1\}^{1/j} \leq \frac{C}{\delta}, \quad \sup\{|c_j(x)| \mid x \in M\}^{1/j} \leq \frac{C}{\delta}.$$

Again similar estimates hold for the covariant derivatives of b_j and c_j . We find that for all $j \geq 1$ and for all $\psi \in \text{dom}(D_j)$

$$\|D_j\psi\|_{H^m} \leq \frac{C^j}{\delta^j} (n\|D^g\psi\|_{H^m} + \|\psi\|_{H^m}).$$

By the criterion above the family $D(z) := \sum_{j=0}^{\infty} z^j D_j$ defines a holomorphic family of type (A) on $\Omega := \{z \in \mathbb{C} \mid |z| < \delta\}$ such that for all $t \in \Omega \cap I$ we have $D(t) = D^{g,g_t}$. Since D^{g,g_t} is self-adjoint by the construction in Section 2.3, all the operators D_j are self-adjoint. Therefore the family $D(z)$ is self-adjoint. \square

Bibliography

- [Ab] R. Abraham, *Transversality in manifolds of mappings*. Bull. Amer. Math. Soc. **69**, 470-474 (1963).
- [Am1] B. Ammann, *Spin-Strukturen und das Spektrum des Dirac-Operators*. Dissertation zur Erlangung des Doktorgrades, Freiburg im Breisgau (1998).
- [Am3] B. Ammann, *A spin-conformal lower bound of the first positive Dirac eigenvalue*. Differ. Geom. Appl. **18**, 1, 21-32 (2003).
- [Am2] B. Ammann, *A variational problem in conformal spin geometry*. Habilitationsschrift, Universität Hamburg (2003).
- [Am4] B. Ammann, *The smallest Dirac eigenvalue in a spin-conformal class and cmc-immersions*. Comm. Anal. Geom. **17**, 429-479 (2009).
- [AH] B. Ammann, E. Humbert, *Positive mass theorem for the Yamabe problem on spin manifolds*. GAFA. **15**, 567-576 (2005).
- [AHM] B. Ammann, E. Humbert, B. Morel, *Mass endomorphism and spinorial Yamabe type problems on conformally flat manifolds*. Comm. Anal. Geom. **14**, 1, 163-182 (2006).
- [AGHM] B. Ammann, J.-F. Grosjean, E. Humbert, B. Morel, *A spinorial analogue of Aubin's inequality*. Math. Z. **260**, 127-151 (2008).
- [ADH1] B. Ammann, M. Dahl, E. Humbert, *Surgery and harmonic spinors*. Advances in Math. **220**, 2, 523-539 (2009).
- [ADH2] B. Ammann, M. Dahl, E. Humbert, *Harmonic spinors and local deformations of the metric*. Preprint, arXiv:0903.4544 (2011).
- [ADHH] B. Ammann, M. Dahl, A. Hermann, E. Humbert, *Mass endomorphism, surgery and perturbations*. Preprint, arXiv:1009.5618 (2010).

- [AMM] B. Ammann, A. Moroianu, S. Moroianu, *The Cauchy problem for metrics with parallel spinors*. Preprint, arXiv:1106.2066 (2011).
- [Ar] N. Aronszajn, *A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order*. J. Math. Pures Appl. **36**, 235-249 (1957).
- [AS] M. Abramowitz, I. Stegun (eds.), *Handbook of mathematical functions*. Dover Publications, New York (1964).
- [Bä1] C. Bär, *Lower eigenvalue estimates for Dirac operators*. Math. Ann. **293**, 39-46 (1992).
- [Bä2] C. Bär, *The Dirac operator on space forms of positive curvature*. J. Math. Soc. Japan **48**, 69-83 (1996).
- [Bä3] C. Bär, *Metrics with harmonic spinors*. Geom. Func. Anal. **6**, 899-942 (1996).
- [Bä4] C. Bär, *On nodal sets for Dirac and Laplace operators*. Comm. Math. Phys. **188**, 3, 709-721 (1997).
- [BD] C. Bär, M. Dahl, *Surgery and the spectrum of the Dirac operator*. J. Reine Angew. Math. **552**, 53-76 (2002).
- [Be] A. Besse, *Einstein manifolds*. Reprint of the 1987 edition, Springer Verlag, Berlin, Heidelberg (2008).
- [BG] J.-P. Bourguignon, P. Gauduchon, *Spineurs, opérateurs de Dirac et variations de métriques*. Comm. Math. Phys. **144**, 3, 581-599 (1992).
- [BGM] M. Berger, P. Gauduchon, E. Mazet, *Le spectre d'une variété riemannienne*. Springer Lecture Notes, **194** (1971).
- [BGM05] C. Bär, P. Gauduchon, A. Moroianu, *Generalized cylinders in semi-Riemannian and Spin Geometry*. Math. Z. **249**, 3, 545-580 (2005).
- [BN] H. Bray, A. Neves, *Classification of prime 3-manifolds with Yamabe invariant greater than $\mathbb{R}P^3$* . Ann. of Math. (2) **159**, 1, 407-424 (2004).
- [BU] S. Bando, H. Urakawa, *Generic properties of the eigenvalue of the Laplacian for compact Riemannian manifolds*. Tohoku Math. Journ. **35**, 155-172 (1983).
- [Da] M. Dahl, *Dirac eigenvalues for generic metrics on three-manifolds*. Ann. Glob. Anal. Geom. **24**, 1, 95-100 (2003).

- [DM] A. Dimakis, F. Müller-Hoissen, *On a gauge condition for orthonormal three-frames*. Phys. Lett. A **142**, 73-74 (1989).
- [EP] A. Enciso, D. Peralta-Salas, *Nondegeneracy of the eigenvalues of the Hodge Laplacian for generic metrics on three-manifolds*. Preprint, arXiv:1003.4206 (2010).
- [FNS] J. Frauendiener, J. M. Nester, L. B. Szabados, *Witten spinors on maximal, conformally flat hypersurfaces*. Class. Quantum Grav. **28**, 185004 (2011).
- [F1] T. Friedrich, *Der erste Eigenwert des Dirac-Operators einer kompakten Riemannschen Mannigfaltigkeit nichtnegativer Skalarkrümmung*. Math. Nachr. **97**, 117-146 (1980).
- [F2] T. Friedrich, *Zur Abhängigkeit des Dirac-Operators von der Spin-Struktur*. Colloq. Math. **48**, 1, 57-62 (1984).
- [F3] T. Friedrich, *On the spinor representation of surfaces in Euclidean 3-space*. J. Geom. Phys. **28**, 1-2, 143-157 (1998).
- [F4] T. Friedrich, *Dirac Operators in Riemannian geometry*. American Mathematical Society, Providence (2000).
- [G] N. Ginoux, *The Dirac spectrum*. Springer Verlag, Berlin, Heidelberg (2009).
- [He] A. Hermann, *Generic metrics and the mass endomorphism on spin three-manifolds*. Ann. Glob. Anal. Geom. **37**, 2, 163-171 (2010).
- [Hij1] O. Hijazi, *A conformal lower bound for the smallest eigenvalue of the Dirac operator and Killing spinors*. Comm. Math. Phys. **104**, 1, 151-162 (1986).
- [Hij2] O. Hijazi, *Première valeur propre de l'opérateur de Dirac et nombre de Yamabe*. C. R. Acad. Sci. Paris **313**, 12, 865-868 (1991).
- [Hij3] O. Hijazi, *Spectral properties of the Dirac operator and geometrical structures*. Geometric methods for quantum field theory (Villa de Leyva, 1999) World Sci. Publ., River Edge, NJ, 116-169 (2001).
- [Hir] M. Hirsch, *Differential topology*. Springer Verlag, New York (1976).
- [Hit] N. Hitchin, *Harmonic spinors*. Advances in Math. **14**, 1-55 (1974).

- [K] T. Kato, *Perturbation theory of linear operators*. Reprint of the 1980 edition, Springer Verlag, Berlin, Heidelberg (1995).
- [Ki1] K.-D. Kirchberg, *An estimation for the first eigenvalue of the Dirac operator on closed Kähler manifolds of positive scalar curvature*. Ann. Glob. Anal. Geom. **4**, 3, 291-325 (1986).
- [Ki2] K.-D. Kirchberg, *Compact six-dimensional Kähler spin manifolds of positive scalar curvature with the smallest possible first eigenvalue of the Dirac operator*. Math. Ann. **282**, 1, 157-176 (1988).
- [KS] R. Kusner, N. Schmitt, *The spinor representation of surfaces in space*. Preprint, arXiv:dg-ga/9610005 (1996).
- [KSW1] W. Kramer, U. Semmelmann, G. Weingart, *The first eigenvalue of the Dirac operator on quaternionic Kähler manifolds*, Comm. Math. Phys. **199**, 2, 327-349 (1998).
- [KSW2] W. Kramer, U. Semmelmann, G. Weingart, *Eigenvalue estimates for the Dirac operator on quaternionic Kähler manifolds*, Math. Z. **230**, 4, 727-751 (1999).
- [La] S. Lang, *Fundamentals of Differential Geometry*. Springer Verlag, New York (1999).
- [Li] A. Lichnerowicz, *Spineurs harmoniques*. C. R. Acad. Sci. Paris **257**, 7-9 (1963).
- [LM] H. B. Lawson, M.-L. Michelsohn, *Spin geometry*. Princeton University Press, Princeton (1989).
- [Lo] J. Lott, *Eigenvalue bounds for the Dirac operator*. Pac. J. Math. **125**, 1, 117-126 (1986).
- [LP] J. M. Lee, T. H. Parker, *The Yamabe problem*. Bull. Amer. Math. Soc., New Ser. **17**, 1, 37-91 (1987).
- [Ma] S. Maier, *Generic metrics and connections on Spin- and Spin^c-manifolds*. Comm. Math. Phys. **188**, 2, 407-437 (1997).
- [Mi] J. Milnor, *Topology from the differentiable viewpoint*. Virginia University Press (1965).
- [N] J. M. Nester, *A positive gravitational energy proof*. Phys. Lett. A **139**, 112-114 (1989).

- [PT] T. Parker, C. H. Taubes, *On Witten's proof of the positive energy theorem*. Comm. Math. Phys. **84**, 2, 223-238 (1982).
- [Ru] W. Rudin, *Functional analysis*. 2nd edition, McGraw Hill, New York (1991).
- [Sm] S. Smale, *An infinite dimensional version of Sard's theorem*. Amer. J. of Math. **7**, 861-866 (1965).
- [St] E. Stiefel, *Richtungsfelder und Fernparallelismus in n -dimensionalen Mannigfaltigkeiten*. Comment. Math. Helv. **8**, 1, 305-353 (1935).
- [TT] D. Tausk, R. Tumulka, *Can we make a Bohmian electron reach the speed of light, at least for one instant?*. J. Math. Phys. 51: 122306 (2010).
- [U] K. Uhlenbeck, *Generic properties of eigenfunctions*. Amer. J. Math. **98**, 4, 1059-1078 (1976).
- [Wh] J. H. C. Whitehead, *The immersion of an open 3-manifold in euclidean 3-space*. Proc. London Math. Soc. (3) **11**, 81-90 (1961).
- [Wi] E. Witten, *A new proof of the positive energy theorem*. Comm. Math. Phys. **80**, 3, 381-402 (1981).

Index

conformal class, 31

divergence, 26

energy momentum tensor, 27, 79

generic, 65

harmonic spinor, 17

Killing spinor, 17

parallel spinor, 17

residual, 64

structure

 quaternionic, 18

 real, 18

trace, 26

transverse, 63