



Mendelian breeding units versus standard sampling strategies: Mitochondrial DNA variation in southwest Sardinia

Daria Sanna¹, Maria Pala², Piero Cossu¹, Gian Luca Dedola¹, Sonia Melis³, Giovanni Fresu¹, Laura Morelli¹,
Domenica Obinu¹, Giancarlo Tonolo^{4,5}, Giannina Secchi⁶, Riccardo Triunfo⁷, Joseph G. Lorenz⁸, Laura
Scheinfeldt⁹, Antonio Torroni², Renato Robledo¹⁰, Paolo Francalacci¹

¹*Dipartimento di Zoologia e Genetica Evoluzionistica, Università di Sassari, Sassari, Italy.*

²*Dipartimento di Genetica e Microbiologia, Università di Pavia, Pavia, Italy.*

³*Department of Neuroscience and Center for Neurovirology, Temple University School of Medicine, Philadelphia, Pennsylvania, USA.*

⁴*Servizio di Diabetologia, Azienda ASL 2 Olbia, Olbia, Italy.*

⁵*Department of Clinical Sciences, Lund University, Malmö, Sweden.*

⁶*Servizio di Diabetologia, Università di Sassari, Sassari, Italy.*

⁷*CRS4, Centro di Ricerca, Sviluppo e Studi Superiori in Sardegna, Cagliari, Italy.*

⁸*Department of Anthropology, Central Washington University, Ellensburg, Washington, USA.*

⁹*Department of Genome Sciences, University of Washington, Seattle, Washington, USA.*

¹⁰*Dipartimento di Scienze e Tecnologie Biomediche, Università di Cagliari, Monserrato, Cagliari, Italy.*

Abstract

We report a sampling strategy based on Mendelian Breeding Units (MBUs), representing an interbreeding group of individuals sharing a common gene pool. The identification of MBUs is crucial for case-control experimental design in association studies. The aim of this work was to evaluate the possible existence of bias in terms of genetic variability and haplogroup frequencies in the MBU sample, due to severe sample selection. In order to reach this goal, the MBU sampling strategy was compared to a standard selection of individuals according to their surname and place of birth. We analysed mitochondrial DNA variation (first hypervariable segment and coding region) in unrelated healthy subjects from two different areas of Sardinia: the area around the town of Cabras and the western Campidano area. No statistically significant differences were observed when the two sampling methods were compared, indicating that the stringent sample selection needed to establish a MBU does not alter original genetic variability and haplogroup distribution. Therefore, the MBU sampling strategy can be considered a useful tool in association studies of complex traits.

Key words: breeding units strategy, mtDNA haplogroup distribution, association studies.

Received: July 8, 2010; Accepted: February 8, 2011.

Introduction

Population definition, sample selection and choice of markers are crucial points in human population genetics studies, and the sampling strategy depends principally on the questions being asked. In addition to biological aspects, such studies should also take into account important socio-cultural parameters, such as language and religion, along with social and self-identity affiliation. If a human population is clearly ethnically-identified and recent admixture is negligible, sampling strategies based only on surname (whenever distinctive) and place of birth are preferred,

since they allow exclusion of recent immigrants, not yet blended into the gene pool, from the analysis. Moreover, surname and place of birth criteria can be extended from the DNA donors to their ancestors, provided that genealogical information is available.

A more stringent sampling strategy is required in studies based on genome-wide association scans, which look for different allele distributions between individuals with (cases) or without (controls) a phenotype of interest. The case-control experimental design is expected to be appropriate in surveys on homogeneous populations, whereas both false-positive and false-negative results may occur in heterogeneous or substructured populations, if cases and controls are not carefully sampled according

to their origin. This scenario is likely to occur in an island like Sardinia, where the majority of the present population is distributed among 363 isolated villages (Siniscalco *et al.*, 1999) which, while sharing common ancestry, might have diversified during many centuries of isolation. Therefore, it is important to identify true Mendelian Breeding Units (MBUs), *i.e.* interbreeding groups of individuals sharing a common ancestral gene pool. In Sardinia, the most practical way to define a MBU is to derive a direct estimate of the percentage of endogamous mating occurring in the last 200 years. This information was obtained anonymously from municipal and ecclesiastical marriage registers (Siniscalco *et al.*, 1999). However, rigorous sample selection for reconstructing MBUs led to a conspicuous reduction in sample size, which might have significantly skewed haplotypic or allelic frequencies. In a previous paper (Siniscalco *et al.*, 1999), we reported a pilot study on 55 unrelated controls belonging to the MBU of Carloforte, who were genotyped at six markers. We showed there that the allele frequencies, and therefore the genomic profile, remained constant even when only a subset of 20 individuals was analysed.

The main goal of this work was to evaluate the reliability of the MBU approach in describing genetic variation in human populations, particularly regarding its application to association studies of complex traits.

We compared genetic variability in two sets of samples which included different individuals recruited from the same areas, using two diverse sampling strategies. With the Standard (STD) Method, individuals unrelated for at least two generations were selected on the basis of the surname and place of birth of their grandparents, depicting present-day genetic variation, with the sole exclusion of the most recent immigrants. Using the MBU Method, the selected DNA donors were proven to be descendants of individuals present in the 17th century archives, with no common ancestors for up to at least five generations. This was ascertained by means of a complete genealogical history checking, based on the official records made available to us by the City Halls. Samples collected using the latter method, being representative of population settlements before the migratory events of the last few centuries, allow an extension of the temporal resolution of genetic variability. Therefore, comparison of the two sampling methods might also reveal possible occurrences of diachronic genetic variation in the analysed areas, due to micro-evolutionary dynamics such as drift or gene flow from neighbouring populations.

The analysed samples belong to two different socio-cultural areas, Cabras and western Campidano, whose cultural traits differentiated around the second half of the 19th century: the former, and its neighbouring area, became a flourishing fishing centre, while the latter consists of rural villages whose economy is based on farming and sheep raising.

We studied mitochondrial DNA (mtDNA), since it has been extensively used as a molecular marker during the past 20 years, is maternally inherited, does not recombine and is in a haploid state; thus it is more sensitive than nuclear DNA to the effects of genetic drift and gene flow, and any discrepancy between the two sampling methods is expected to be enhanced.

Materials and Methods

Sample selection

Using the MBU strategy, we analysed 85 unrelated healthy subjects from two areas located in southwestern Sardinia: 35 individuals from Cabras and 50 individuals from western Campidano (Figure 1). Using the STD strategy, we analysed 71 unrelated individuals coming from the same areas. Comparison was performed between 48 samples from Cabras and its neighbouring area (up to 50 km) and 23 samples from the western Campidano area.




Figure 1 - Map showing the distribution of the two areas analysed in southwest Sardinia. Cas: Cabras. W Camp: western Campidano.

mtDNA analysis

Whole genomic DNA was extracted using standard procedures. For each individual, mitochondrial haplogroup affiliation was determined by both sequencing of the first hypervariable segment (HVS-I) of the control region from position 15997 to 16399 bp (Anderson *et al.*, 1981) and RFLP (Restriction Fragment Length Polymorphism) analysis of the coding region for the presence/absence of haplogroup diagnostic markers (see Table 1 for details).

Data analysis

BioEdit software 7.0.5.2 (Hall, 1999) was used to align the sequences obtained. To characterise genetic variation among sampling sites, estimates of the number of polymorphic sites (S), the number of haplotypes (h), the nucleotide diversity (Pi), and the haplotype diversity (Hd) were obtained using the DnaSP 4.10 software (Rozas and Rozas, 1999). Pearson chi-square (χ^2) values (Pearson, 1900) were calculated in order to assess whether there was any difference between the haplotype frequency distributions obtained for the same areas by means of different sampling strategies (MBU and STD). Principal Coordinate Analysis (PCoA) was carried out on the matrix of DNA pairwise differences, using the Genalex 6.3 software (Peakall and Smouse, 2006). The method based on the covariance matrix with data standardisation was applied. In order to assess the occurrence of significant genetic structuring among samples, analysis of molecular variance (AMOVA) was performed on the matrix of pairwise DNA distances among haplotypes, using the Arlequin 3.1 computer package (Excoffier *et al.*, 2005). Furthermore, genetic differentiation between pairs of samples was estimated by pairwise Φ_{ST} values, computed from the matrix of haplotype DNA pairwise differences. The significance of variance components and F -statistic was assessed by a random permutation test (10,000 replicates).

A Median-Joining network was drawn for each sampling strategy using Network 4.2.0.1 software (<http://www.fluxus-engineering.com>).

Results

Nucleotide sequence analysis of HVS-I (GenBank accession numbers: HM584611-HM584695 for MBU samples, and HM594952-HM595022 for STD samples) combined with RFLP analysis allowed the clustering of samples from both MBU and STD strategies into nine main haplogroups. They increased to eleven when sub-haplogroups K and U5b3 were also considered (Table 2). Haplogroup H, which includes the Cambridge Reference Sequence (CRS) (Anderson *et al.*, 1981), proved to be the most common. Haplogroup U5b3, reported as Sardinian-specific (Fraumene *et al.*, 2006; Pala *et al.*, 2009), was found in Cabras MBU, western Campidano MBU and Cabras STD, missing in western Campidano STD only. The values of genetic diversity, calculated for the dataset of HVS-I, were similar for all regions and sampling strategies considered, showing a high level of variability (Table 3). Furthermore, we found a total of 82 different haplotypes. Those whose occurrence was detected by both sampling methods (MBU and STD) showed comparable relative frequency distributions, with no significant Pearson chi-square values (Table 4).

Table 2 - MtDNA haplogroup distribution obtained using the two sampling methods (values are expressed as relative distribution frequencies). Cas: Cabras; W Camp: western Campidano; B: MBU method; S: STD method.

Haplogroup	Cas-B	Cas-S	W Camp-B	W Camp-S
V	2.9	6.3	10.0	4.4
H tot	37.1	29	50.0	56.5
T tot	17.1	14.6	6.0	21.7
J tot	5.7	16.7	14.0	8.7
U ¹ (x U5b3, K)	17.1	16.7	8.0	8.7
U5b3	5.7	8.3	10.0	-
K	2.9	-	-	-
I	11.4	-	-	-
W	-	4.2	-	-
X	-	4.2	-	-
M tot	-	-	2.0	-

¹This haplogroup does not include the sub-haplogroups U5b3 and K.

Table 1 - Oligonucleotide pairs used and polymorphic sites investigated to classify mitochondrial coding regions into haplogroups H, V and U/K.

Haplogroup	Primer sequences	Polymorphic site	Enzyme
H	L: aagcaatatgaaatgatctgc	-7025	<i>AhaI</i>
	H: gcgtaggtttggtctag		
V	L: gagcttaaaccccttattt	-4577	<i>NlaIII</i>
	H: gtattgattgtagtattggttattggtca		
U/K	L: ctcaaccccacatcattacc	+12308	<i>HinfI</i>
	H: attacttttttggagttgcaccaagatt		

Table 3 - Estimates of genetic diversity among samples analysed. *S*: segregating sites; *h*: number of haplotypes; *Hd*: haplotype diversity; *Pi*: nucleotide diversity; *Uh*: frequency of unique haplotypes *per* total number of haplotypes; *Sh*: frequency of individuals with shared haplotypes; *Rh*: frequency of haplotypes observed in more than one individual.

	<i>S</i>	<i>h</i>	<i>Hd</i>	<i>Pi</i>	<i>Uh</i>	<i>Sh</i>	<i>Rh</i>
Cas-B	38	22	0.946	0.019	0.629	0.486	0.114
Cas-S	45	37	0.976	0.021	0.771	0.333	0.104
W Camp-B	55	32	0.937	0.018	0.640	0.460	0.100
W Camp-S	35	16	0.913	0.018	0.696	0.391	0.087

Cas: Cabras; W Camp: western Campidano; B: MBU method; S: STD method.

Nucleotide sequences from the control region were combined with RFLP data on the coding region to obtain a single dataset for the following analysis.

The first two coordinates of PCoA, which account for 62.39% of the total variability, identify two main groups of haplotypes. However, haplotypes were not grouped either according to the geographic area of origin (Cabras or western Campidano) or to the sampling strategy adopted (MBU versus STD) (Figure 2).

Accordingly, the analysis of molecular variance (AMOVA) did not indicate significant genetic differentiation among samples ($\Phi_{ST} = 0.0096$, $p > 0.05$). Indeed, nearly all variance was found within samples (99.04%), whereas differences among samples accounted for only 0.96% of the total variation. These results were further confirmed by the pairwise comparison of samples, which did not show any significant genetic differentiation (Table 5).

Furthermore, network analysis showed similar relationships among haplogroups without geographical structuring when the two sampling methods were compared (Figure 3).

Discussion

Estimates of genetic diversity (Table 2) obtained for the two sampling strategies were compatible with no occurrence of high levels of repeated haplotypes in the STD strategy, as could be expected. This finding supports the possible occurrence of a homogeneous population shared by both the western Campidano and Cabras areas, with a constant high level of genetic variability in the samples obtained by the two sampling methods and low levels of stochastic forces.

The similarity of genetic diversity values between areas and sampling strategies may be explained considering the lack of diachronic divergence between the present and past genetic settlement of the western Campidano and Cabras areas. Furthermore, this finding is attributable to the absence of genetic drift in the analysed areas. Indeed, this stochastic force, if present, could lead to genetic heterogeneity due to random loss of haplogroups and alteration of their frequencies. The absence of higher levels of identical haplotypes among the STD samples suggests that no significant founder effects affected the population recently. Consistently, the result of PCoA applied to the combined dataset (control region + coding region) (Figure 2) contributed to group MBU and STD samples without genetic structuring. Such similarity was also confirmed by the corresponding, not significant, P values of Φ_{ST} .

Network analysis was also consistent with the results above. The two sampling strategies displayed similar global relationships among mitochondrial haplogroups without geographical structuring, showing that mtDNA haplogroup frequencies and distribution obtained by the MBU method were not skewed by the severe sample selection of the method used.

Overall, these results suggest a lack of genetic variation in southwest Sardinia, probably due to a continuous




Figure 2 - Principal Coordinate Analysis (PCoA) plot: the first PC accounts for 37.82% of variance, while the second PC accounts for 25.07% (Cas: Cabras; W Camp: western Campidano; B: MBU method; S: STD method).

Table 4 - MtDNA haplotype distribution obtained using the two sampling methods (values are expressed as relative distribution frequencies). N: number of individuals; Cas: Cabras; W Camp: western Campidano; B: MBU method; S: STD method; χ^2 : values of Pearson chi-square (significance level with p values ≤ 0.05).

Haplotype	Haplogroup	Cas-B N = 35	Cas-S N = 48	χ^2	W Camp-B N = 50	W Camp-S N = 23	χ^2	GenBank accession n°
Hap1	J	-	4.2	1.46	2.0	-	0.46	HM594952
Hap2	X	-	2.1	0.73	-	-	-	HM594953
Hap3	V	2.9	2.1	0.05	2.0	-	0.46	HM594954
Hap4	U	-	2.1	0.73	-	-	-	HM594955
Hap5	J	-	2.1	0.73	-	-	-	HM594956
Hap6	U5b3	2.9	6.2	0.48	8.0	-	1.84	HM594957
Hap7	W	-	2.1	0.73	-	-	-	HM594958
Hap8	U	-	2.1	0.73	-	-	-	HM594960
Hap9	H	17.1	14.6	0.08	24.0	30.4	0.25	HM594961
Hap10	H	-	2.1	0.73	-	-	-	HM594962
Hap11	X	-	2.1	0.73	-	-	-	HM594963
Hap12	T	14.3	2.1	0.30	-	-	-	HM594964
Hap13	U5b3	-	2.1	0.73	-	-	-	HM594965
Hap14	T	-	2.1	0.73	-	-	-	HM594966
Hap15	H	-	2.1	0.73	-	-	-	HM594967
Hap16	U	-	4.2	1.46	-	-	-	HM594969
Hap17	H	2.9	4.2	0.10	-	4.3	2.17	HM594972
Hap18	H	-	2.1	0.73	-	-	-	HM594973
Hap19	U	-	2.1	0.73	-	-	-	HM594974
Hap20	T	-	2.1	0.73	-	-	-	HM594976
Hap21	U	-	2.1	0.73	-	4.3	2.17	HM594978
Hap22	J	2.9	2.1	0.05	6.0	-	1.38	HM594979
Hap23	T	-	2.1	0.73	-	-	-	HM594980
Hap24	J	-	2.1	0.73	2.0	4.3	0.32	HM594981
Hap25	V	-	2.1	0.73	-	-	-	HM594982
Hap26	V	-	2.1	0.73	-	-	-	HM594984
Hap27	T	2.9	2.1	0.05	-	4.3	2.17	HM594986
Hap28	W	-	2.1	0.73	-	-	-	HM594987
Hap29	T	-	2.1	0.73	-	-	-	HM594988
Hap30	U	-	2.1	0.73	-	-	-	HM594989
Hap31	J	-	2.1	0.73	2.0	-	0.46	HM594990
Hap32	U	-	2.1	0.73	-	-	-	HM594991
Hap33	T	-	2.1	0.73	2.0	4.3	0.32	HM594993
Hap34	J	-	2.1	0.73	-	-	-	HM594994
Hap35	J	-	2.1	0.73	-	-	-	HM594997
Hap36	H	-	2.1	0.73	4.0	4.3	0.005	HM594998
Hap37	H	-	2.1	0.73	-	4.3	2.17	HM594999
Hap38	H	-	-	-	-	4.3	2.17	HM595001
Hap39	U	-	-	-	-	4.3	2.17	HM595002
Hap40	T	-	-	-	-	8.7	4.35 ¹	HM595003
Hap41	V	-	-	-	-	4.3	2.17	HM595005
Hap42	J	-	-	-	-	4.3	2.17	HM595010
Hap43	H	-	-	-	-	4.3	2.17	HM595014
Hap44	H	-	-	-	-	4.3	2.17	HM595015

Table 4 (cont.)

Haplotype	Haplogroup	Cas-B N = 35	Cas-S N = 48	χ^2	W Camp-B N = 50	W Camp-S N = 23	χ^2	GenBank accession n°
Hap45	T	-	-	-	-	4.3	2.17	HM595018
Hap46	U	5.7	-	2.74	4.0	-	0.92	HM584612
Hap47	M	-	-	-	2.0	-	0.46	HM584613
Hap48	H	-	-	-	2.0	-	0.46	HM584615
Hap49	H	-	-	-	2.0	-	0.46	HM584621
Hap50	J	-	-	-	2.0	-	0.46	HM584622
Hap51	V	-	-	-	2.0	-	0.46	HM584623
Hap52	H	-	-	-	2.0	-	0.46	HM584624
Hap53	V	-	-	-	2.0	-	0.46	HM584626
Hap54	U	-	-	-	2.0	-	0.46	HM584629
Hap55	T	-	-	-	2.0	-	0.46	HM584631
Hap56	H	-	-	-	2.0	-	0.46	HM584632
Hap57	H	-	-	-	2.0	-	0.46	HM584633
Hap58	U	-	-	-	2.0	-	0.46	HM584634
Hap59	U5b3	-	-	-	2.0	-	0.46	HM584635
Hap60	J	-	-	-	2.0	-	0.46	HM584639
Hap61	V	-	-	-	2.0	-	0.46	HM584644
Hap62	H	-	-	-	2.0	-	0,46	HM584646
Hap63	H	-	-	-	2.0	-	0.46	HM584650
Hap64	V	-	-	-	2.0	-	0.46	HM584653
Hap65	H	-	-	-	2.0	-	0.46	HM584654
Hap66	T	-	-	-	2.0	-	0.46	HM584656
Hap67	H	-	-	-	2.0	-	0.46	HM584657
Hap68	H	-	-	-	2.0	-	0.46	HM584660
Hap69	H	2.9	-	1.37	-	-	-	HM584667
Hap70	I	11.4	-	5.49 ¹	-	-	-	HM584668
Hap71	U	2.9	-	1.37	-	-	-	HM584669
Hap72	U	2.9	-	1.37	-	-	-	HM584674
Hap73	U	2.9	-	1.37	-	-	-	HM584676
Hap74	H	2.9	-	1.37	-	-	-	HM584677
Hap75	U	2.9	-	1.37	-	-	-	HM584679
Hap76	H	2.9	-	1.37	-	-	-	HM584683
Hap77	J	2.9	-	1.37	-	-	-	HM584685
Hap78	H	2.9	-	1.37	-	-	-	HM584686
Hap79	K	2.9	-	1.37	-	-	-	HM584687
Hap80	H	2.9	-	1.37	-	-	-	HM584690
Hap81	U5b3	2.9	-	1.37	-	-	-	HM584692
Hap82	H	2.9	-	1.37	-	-	-	HM584694

¹Significant values of χ^2 .

gene flow between the areas, either in the past or more recently, which may have counterbalanced the development of microheterogeneity due to genetic drift.

Previous studies carried out on the paternal unilinear marker Y-chromosome pointed out a similar trend for the entire Sardinian population, suggesting an initial settlement

of a relatively large number of individuals with a common origin (Contu *et al.*, 2008) and conspicuous genetic variability.

The presence of genetic structuring is the major obstacle in association studies based on genome-wide scans searching for linkage disequilibrium (LD) between patients

Table 5 - Population pairwise Φ_{ST} values among samples obtained from MBU and STD strategies. Population codes are reported as in Table 2. Conventional Φ_{ST} values are shown below the diagonal and corresponding P values with significance level ≤ 0.05 are shown above the diagonal.

	Cas-B	Cas-S	W Camp-B	W Camp-S
Cas-B	-	0.4150	0.0940	0.2286
Cas-S	-0.0003	-	0.1552	0.1082
W Camp-B	0.0150	0.0087	-	0.2020
W Camp-S	0.0096	0.0181	0.0098	-

and controls (Risch and Botstein, 1996; Terwilliger and Weiss, 1998), even in isolated populations like Finns and Sardinians (Eaves *et al.* 2000; Taillon-Miller *et al.*, 2000). Pooling individuals belonging to different breeding units may merge alleles that might have different frequencies in different villages, as we have previously reported for some common polymorphisms in Sardinian villages (Robledo *et al.*, 2002).

As previously shown, in a well-defined breeding unit, a small sample was sufficient to describe the genomic profile of the population, which was not affected by severe re-

duction of sample size (Siniscalco *et al.*, 1999). More importantly, the repeated application of our strategy in different MBUs offers the advantage of reducing the risk of false-positive results due to population stratification, since obtaining similar artifactual results in different MBUs is not anticipated.

In conclusion, the comparison of the variability detected by means of the MBU and STD sampling methods points to a diachronic continuity of the genetic structure of southwestern Sardinia. The benefit of the MBU sampling strategy lies in the possibility of: i) selecting the original population on the basis of written documents and not by inferring surname monophyly, and ii) not excluding from the analysis unrelated individuals with polyphyletic surnames, when present, in the founder families.

Our results confirm that the MBU sampling strategy, despite the drastic reduction in sample size, does not introduce deviations in gene frequencies, even if haploid markers such as mtDNA are used. Therefore it can be considered a useful tool in association studies of complex traits, making it possible to infer the genetic settlement of the population, recovering the deepest branches of a genealogy and avoiding the recent contribution of foreign peopling.




Figure 3 - Networks obtained from combined dataset (control region and coding region) for MBU (A and C) and STD (B and D) strategies. A and B: phylogenetic relationships among mitochondrial haplogroups; C and D: geographic distribution of mitochondrial haplogroups. Cas: Cabras; W Camp: western Campidano; B: MBU method; S: STD method.

Acknowledgments

We wish to thank all the participants who made this study possible. We are also grateful to Marcello Siniscalco and Marco Casu for helpful discussions and criticism and to Mary Ann Groeneweg for revising the manuscript. This work was supported by funds from the Fondazione Golinelli made available by Marcello Siniscalco, Compagnia di San Paolo (to A.T.) and grants from the Italian Ministry of Research, MIUR (funds ex 60% to R.R. and P.F.).

References

- Anderson S, Bankier AT, Barrell BG, de Bruijn MHL, Coulson AR, Drouin J, Eperon IC, Nierlich DP, Roe BA, Sanger F, *et al.* (1981) Sequence and organization of the human mitochondrial genome. *Nature* 290:457-465.
- Contu D, Morelli L, Santoni F, Foster JW, Francalacci P and Cucca F (2008) Y-chromosome based evidence for pre-Neolithic origin of the genetically homogeneous but diverse Sardinian population: Inference for association scans. *PLoS One* 3:e1430.
- Eaves IA, Merriman TR, Barber RA, Nutland S, Tuomilehto-Wolf E, Tuomilehto J, Cucca F and Todd JA (2000) The genetically isolated populations of Finland and Sardinia may not be a panacea for linkage disequilibrium mapping of common disease genes. *Nat Genet* 25:320-323.
- Excoffier L, Laval G and Schneider S (2005) Arlequin v. 3.0: An integrated software package for population genetics data analysis. *Evol Bioinform Online* 1:47-50.
- Fraumene C, Belle EMS, Castri L, Sanna S, Mancosu G, Cosso M, Marras F, Barbujani G, Pirastu M and Angius A (2006) High resolution analysis and phylogenetic network construction using complete mtDNA sequences in Sardinian genetic isolates. *Mol Biol Evol* 23:2101-2111.
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95-98.
- Pala M, Achilli A, Olivieri A, Hooshiar Kashani B, Perego UA, Sanna D, Metspalu E, Tambets K, Tamm E, Accetturo M, *et al.* (2009) Mitochondrial haplogroup U5b3: A distant echo of the epipaleolithic in Italy and the legacy of the early Sardinians. *Am J Hum Genet* 84:814-821.
- Peakall R and Smouse PE (2006) Genalex 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes* 6:288-295.
- Pearson K (1900) On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *Phil Mag Ser* 5:157-175.
- Risch N and Botstein D (1996) A manic depressive history. *Nat Genet* 12:351-353.
- Robledo R, Orrù S, Sidoti A, Muresu R, Esposito D, Grimaldi MC, Carcassi C, Rinaldi A, Bernini L, Contu L, *et al.* (2002) A 9.1-kb gap in the genome reference map is shown to be a stable deletion/insertion polymorphism of ancestral origin. *Genomics* 80:585-592.
- Rozas J and Rozas R (1999) DnaSP v. 3: An integrated program for molecular population genetics and molecular evolution analysis. *Bioinformatics* 15:174-175.
- Siniscalco M, Robledo R, Bender P, Carcassi C, Contu L and Beck J (1999) Population genomics in Sardinia: A novel approach to hunt for genomic combinations underlying complex traits and diseases. *Cytogenet Cell Genet* 86:148-152 (and Erratum 87:296).
- Taillon-Miller P, Bauer-Sardiña I, Saccone NL, Putzel J, Laitinen T, Cao A, Kere J, Pilia G, Rice JP and Kwok PY (2000) Juxtaposed regions of extensive and minimal linkage disequilibrium in human Xq25 and Xq28. *Nat Genet* 25:324-328.
- Terwilliger JD and Weiss KM (1998) Linkage disequilibrium mapping of complex disease: Fantasy or reality? *Curr Opin Biotechnol* 9:578-594.

Associate Editor: Angela M. Vianna-Morgante

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.