

This is an author produced version of a paper published in *Nordic Hydrology*. This paper has been peer-reviewed and is proof-corrected, but does not include the journal pagination.

Citation for the published paper:

Ulén, B., Johansson, G., Simonsson, M. (2008) Changes in nutrient leaching and groundwater quality during long-term studies of an arable field on the Swedish south-west coast. *Nordic Hydrology*.

Volume: 39 Number: 1, pp 63-77.

<http://dx.doi.org/10.2166/nh.2008.030>

Access to the published version may require journal subscription.

Published with permission from: IWA Publishing



Epsilon Open Archive <http://epsilon.slu.se>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

**Changes in nutrient leaching and groundwater quality during long-term studies of an arable field on the Swedish south-west coast**

Barbro Ulén\*, Göran Johansson\* & Magnus Simonsson\*\*

\* Department of Soil Sciences, Division of Water Quality Management, Swedish University of Agricultural Sciences  
\*\* Department of Soil Sciences, Division of Soil Chemistry and Pedology, Swedish University of Agricultural Sciences

Corresponding author: B. Ulén, Department of Soil Sciences, Division of Water Quality Management, Swedish University of Agricultural Sciences, PO Box 7014, SE-750 07 Uppsala, Sweden.  
E-Mail [barbro.ulen@mv.slu.se](mailto:barbro.ulen@mv.slu.se)

1 **Abstract** The aim of this study was to evaluate the long-term (1977-2004) effects of new  
2 agricultural practices and reduced acid rain on drainwater and groundwater chemistry for an  
3 intensely cultivated arable field with sandy soil in south-west Sweden. Trends in chemical  
4 composition of the drainwater were compared with those of atmospheric deposition and  
5 groundwater. A modified crop rotation including catch crops significantly decreased the  
6 average concentration of nitrate nitrogen ( $\text{NO}_3\text{-N}$ ) in drainwater from 13.0 to 7.2  $\text{mg l}^{-1}$ . This  
7 rotation was also found to be a very effective measure against high  $\text{NO}_3\text{-N}$  concentrations in  
8 shallow groundwater ( $>1.7$  m below the soil surface). The degree of phosphorus saturation  
9 (DPS) in the subsoil, calculated to be 10% and 9% by two different laboratory methods,  
10 corresponded to an average and constant concentration of dissolved reactive phosphate (DRP)  
11 in drainwater of 0.006  $\text{mg l}^{-1}$ . Generally lower inputs of acid deposition to the soil were  
12 confirmed by a decreasing  $\text{SO}_4\text{-S}$  trend (by 3% over 24 years) in drainwater. Changes in  
13 cropping had reduced the effect of acid load to the soil, while drainwater alkalinity showed a  
14 slow but significant positive trend amounting to 0.4% over 24 years.

15 **Key words:** *Drainwater, groundwater, ionic balance, nutrients, trends, water chemistry.*  
16  
17

## 18 **Introduction**

19 Environmental aspects relating to leaching of nutrients to surrounding surface water and  
20 groundwater bodies and countermeasures have become a major issue for agriculture. Thus,  
21 since the 1970s there has been a great concern in Sweden about nitrogen (N) and phosphorus  
22 (P) losses to surface waters (Gustafson, 1987). In contrast to the situation in Denmark where  
23 reduction of groundwater  $\text{NO}_3$  by pyrite oxidation is a well-known phenomenon (Postma *et*  
24 *al.*, 1991) leaching of nitrate ( $\text{NO}_3$ ) to groundwater is also of great concern in Sweden  
25 especially in groundwater intake areas with sandy soils without a protective clay layer  
26 (Gustafson, 1983). In both countries a proposed measure to reduce  $\text{NO}_3$  leaching both to drain

1 water and groundwater is growing catch crops; fast-growing crops used between seasons of  
2 regular planting to reduce the concentration of water-soluble NO<sub>3</sub> and other mineral nitrogen  
3 in the soil. Perennial ryegrass (*Lolium perenne* L.) may be undersown in winter crops during  
4 spring. Several studies from drained experimental plots have reported that this grass has an  
5 appropriate growth pattern for capturing mineral nitrogen in the soil, since it grows fast  
6 during autumn (Torstensson, 1998; Torstensson and Aronsson, 2000), and does not compete  
7 with the main crop (Kvist, 1992; Andersen and Olsen, 1993; Ohlander *et al.*, 1996; Bergqvist  
8 *et al.*, 2002). Repeated cropping with ryegrass for several years may also preserve, or even  
9 improve, the soil organic matter content (Aronsson, 2000). In the 1990s, Sweden joined the  
10 EU and subsidies for catch crops were introduced in the southern part of the country, but take-  
11 up by farmers was low until 2001, when new and improved subsidies became available (Ulén  
12 and Fölster, 2007). Estimates of NO<sub>3</sub> leaching based on the database programme SOIL-NDB  
13 (Mårtensson and Johansson, 2006) have revealed an essential reduction in NO<sub>3</sub>-N in water  
14 leaving the root-zone when catch crops are grown compared with ordinary crops. However in  
15 agricultural streams with decreasing trends in NO<sub>3</sub>-N concentrations these improvements  
16 were not clearly correlated to the acreage of arable land with catch crops in the river basin  
17 (Ulén and Fölster, 2007).

18 Phosphorus losses from single fields in Sweden have primarily been evaluated by a P risk  
19 index (Djordjic and Bergström, 2005a) based on results from monitored fields (Djordjic and  
20 Bergström, 2005b). Phosphorus in the water may be either bound to particles and colloids, or  
21 exist in dissolved forms. Dissolved reactive phosphorus (DRP) may be fixed by soil particles  
22 either through anion adsorption or through precipitation of ferric minerals. Consequently one  
23 important factor for the P risk index is the P sorption capacity in the soil profile, which is  
24 related to the degree of P saturation (DPS). Methods proposed for this characterisation DPS  
25 include (i) a laboratory test by which the P sorption of a certain amount of soil (PSI) is

1 deduced from the P concentration remaining in solution after addition of one large dose of P  
2 (Börling *et al.*, 2001); and (ii) an extraction method based on the molar ratio of P to the sum  
3 of aluminium (Al) and iron (Fe) in an acid extract (Ulén, 2006). However, no comparison of  
4 of these methods has been done. Neither has the spatial variability of DPS been investigated  
5 on a single field in order to suggest soil sampling strategies for this soil indicator.

6 To be an efficient and environmentally-friendly fertiliser, manure should deliver soluble  
7 inorganic nutrients at a time when the crop needs them. If nutrients from manure are released  
8 too late in the growing season, or after the crop has been harvested, they can potentially leach  
9 through the unsaturated zone and cause large leaching loads via tile drains (Bergström and  
10 Kirchmann, 1999; Kirchmann *et al.*, 2002). In the 1990s, several pieces of Swedish legislation  
11 relating to nutrient losses were introduced regarding livestock density and handling of manure  
12 (Ulén *et al.*, 2004). Through this legislation heavy manure applications are avoided at farm  
13 level but for a precision manure application, DPS tests on single fields might be  
14 recommended.

15 The relationships between contamination of agricultural land by mineral fertiliser/manure  
16 and nutrient/ionic concentrations in drainwater and groundwater are complex. When  
17 groundwater resources are evaluated on the basis of nutrient loads from agriculture,  
18 simultaneous changes in acid and ionic load to the soil should be taken into account. Drained  
19 agricultural land tends to have high hydraulic permeability in the subsurface horizon, which  
20 may cause enhanced downward chemical transport from the topsoil. High concentrations of  
21 chloride (Cl<sup>-</sup>) may be linked to high intensity of cattle production (Hornish *et al.*, 2002). In  
22 addition, agricultural application of potassium (K<sup>+</sup>) as a plant nutrient may result in Cl<sup>-</sup>  
23 contamination of recharging groundwater, since K is commonly applied in the form of KCl.  
24 In clay soils K may exchange with the clay mineral, so that K/Cl ratios in agricultural  
25 recharge are generally less than 1 (Böhlke, 2001). Several cations such as ammonium (NH<sub>4</sub><sup>+</sup>),

1 calcium ( $\text{Ca}^{2+}$ ) and magnesium ( $\text{Mg}^{2+}$ ) from agricultural activities have been shown to be  
2 capable to displace other cations from exchange sites in soils (Parkhurst and Plummer, 1993).  
3 Furthermore sulphate, ( $\text{SO}_4^{2-}$ ) added with  $\text{NH}_4^+$  and  $\text{Ca}^{2+}$  may be retarded by anion sorption.

4 Groundwater contamination of  $\text{SO}_4^{2-}$  is most likely in areas with preferential flow  
5 pathways, high recharge rates and coarse-grained superficial deposits lacking sesquioxides.  
6 Recharging groundwater with high acidity or ionic strength may increase the weathering rate  
7 and solubility of minerals. Direct acid loads of the soils have been of particular concern for  
8 the Swedish south-west coast. For several decades, this region was clearly affected by acid  
9 deposition, which carried high amounts of sulphur (S) and N compounds originating from  
10 atmospheric reactions with S and N oxides emitted during industrial combustion. In general,  
11 atmospheric acid deposition has decreased considerably in recent years and there has also  
12 been a slight reduction in N deposition (Lövblad *et al.*, 2002). Presently Sweden is in the  
13 early stages of recovery from acidification (Warfvinge and Bertills, 2000). However, in  
14 agriculture intensive areas N emission of ammonia from livestock and manure spreading  
15 should still be of concern.

16 In addition to loads of base cations by fertilisers and direct acidification the netto-removal  
17 of base cations from arable land by the crop may have an indirect effect on weathering  
18 processes and ion exchange. This removal has an acidifying effect unless compensated for by  
19 liming (Bergström and Gustafson, 1985; Debreczeni and Kismányoky, 2005) and has been  
20 demonstrated to decrease pH in drainage water from a Swedish clay soil in southern Sweden  
21 (Ulén *et al.*, 2007).

22 The aims of the present study were to confirm or reject the following hypotheses: 1)  
23 phosphorus retention, proposed to be related to the P sorption ability of the soil, may be  
24 indicated by appropriate soil sampling strategy and a well-chosen laboratory method; 2)  
25 modified farming practices may reduce nitrogen load to drainwater and surface groundwater;

1 and 3) crop production and acidification may interact based on drainwater and surface  
2 groundwater chemistry.

3

#### 4 **Materials and methods**

5 An arable field (code 12N) situated seven kilometres from the shore of Laholm Bay in the  
6 county of Halland in Sweden (Figure 1) was chosen to test the hypotheses. The agriculture in  
7 this area with coarse-textured soil is intense and the livestock density high. The monitored  
8 field lies six kilometres from the long-term plot experiments at Mellby, where leaching of  
9 nutrients has been studied since 1986. The field and underlying groundwater have been  
10 monitored since 1977. It belongs to a monitoring system based on studies of individual  
11 Swedish agricultural fields, established in order to systematically evaluate the long-term  
12 environmental trends in nutrient leaching in relation to agricultural practices and to determine  
13 general quality and trends in drainwater and groundwater chemistry. This programme  
14 currently consists of 13 drained fields many of which have been monitored for more than 28  
15 years.

16

#### 17 *Measuring devices and geochemical description*

18 The field (14.5 ha) is situated 12-15 m above sea level, and has an average slope of 1.4%  
19 (Figure 1). The upper 1-2 m layer of the profile contains both fine and coarse water-sorted  
20 sand, underlain by deep marine clay. The latter is very dense and especially in the upper part  
21 of the field the groundwater is very shallow. Tile drains were installed in a herringbone  
22 pattern in 1975 leading the drainwater to a Thomson weir in an underground measuring  
23 station. Pairs of groundwater pipes were also installed close to the water monitoring station in  
24 1975. These pipes are at three different depths below the surface (1.7 m, 2.2 m and 5.5 m).  
25 The intake of the pipe pair at 1.7 m depth is situated in coarse sand just above the clay layer,

1 while the intakes for the other two pairs of pipes are situated in the clay matrix underlying the  
2 drained sandy layers. The infiltrating water forming groundwater mainly streams in the sandy  
3 overlayer to a valley south of the field. Groundwater in the sandy layer has been indicated to  
4 be young (Gustafson *et al.*, 1984) and nearly all has its origin in precipitation that infiltrates  
5 the field. The clay layer clearly prevents the shallow groundwater from percolating down into  
6 the deeper glacial deposits. Since this water does not percolate down through the clay layer,  
7 there is always much water flowing through the drainpipe system. Annual average discharge  
8 is 420 mm equal to 58% of the mean annual precipitation. The bedrock contains some pyrite  
9 in addition to the main components quartz, K-feldspar and plagioclase (Sjöström, 1993).

10

#### 11 *Soil quality parameters and field management*

12 Intensive soil sampling took place in 1993 and was extensively followed up in 2005. The  
13 topsoil, classified as loamy sand, has a texture of 13% clay, 10% silt and 77% sand. It has an  
14 organic carbon (C) content of 3.7% and a C/N ratio of 18. The soil is slightly acid; the pH is  
15 6.0 in topsoil and 6.3 in the subsoil. The soil cannot be defined as calcareous since it does not  
16 visibly react with acid and since the CaO concentration is low (0.2%). Effective base  
17 saturation is high; 94% in the topsoil and 86% in the upper subsoil, calculated from  
18 exchangeable base cations extracted in 0.1 M barium chloride and divided by effective cation  
19 exchange capacity (CEC, the sum of exchangeable base cations and exchangeable acidity). In  
20 order to evaluate P sorption capacity, 136 soil samples evenly distributed over the field and  
21 representing 4 depth layers were investigated. P sorption index (PSI) was determined  
22 according to the method of Börling *et al.* (2001), which involves addition of one single dose  
23 of P to a soil-water mixture. An acid ammonium lactate solution method according to Egnér  
24 *et al.* (1960) was used for extraction of the soil. In addition to the common chemical  
25 determination of phosphorus (P-AL) and calcium (Ca-AL) in these extracts, iron (Fe-AL) and



1 aluminium (Al-AL) were also analysed. The results were calculated to the degree of  
2 phosphorus saturation (DPS) as the P-AL ratio divided by the sum of Fe-AL and Al-AL on a  
3 molar basis according to Ulén (2006). Comparison was made between two independent sets  
4 of differently distributed soil samples. One set of soil samples was taken in a regular square  
5 grid with 77 m spacings , while the other set was taken at 50 m intervals along three 200-260  
6 m transects that covered the main slopes of the field.

7 The farmer in charge of the field provided information about fertilisation, harvesting and  
8 other agricultural practices. The field is included in regular farm operations. The farm had a  
9 pig production unit until 2002 and during the period 1976-1990, pig slurry was relatively  
10 frequently applied. When Sweden joined the EU, the field was under green fallow (1991-  
11 1992). Thereafter it has been managed as a modified crop rotation with frequent sowing of  
12 catch crops of perennial ryegrass (*Lolium perenne* L.). However, the field still receives much  
13 fertiliser since potatoes and sugarbeet are intensively cropped (Table 1). In 2002 the field  
14 changed ownership and became part of a dairy farm, with application of solid cattle manure  
15 in recent years.

16

#### 17 *Changes in sampling and analytical programme*

18 Water level over a Thomson weir has been recorded continuously with a water stage recorder  
19 since installation. The chart has been digitalised on an hourly basis and the water flow has  
20 been calculated. During recent years, the chart has been complemented with a datalogger  
21 (Thalimedes). Drainwater has been sampled twice a month throughout all years. Groundwater  
22 sampling has been monthly in the first years (1977-1982), four times per year in the period  
23 1983-1986 and six times per year from 1987. Water samples have immediately been sent to  
24 the Water Laboratory at the Division of Water Quality Management, Swedish University of  
25 Agricultural Sciences, where they have been analysed according to the European Committee

1 for Standardisation. The laboratory is accredited for analysis of nutrients, as well as many  
2 other water analyses according to SWEDAC (Swedish Board for Accreditation and  
3 Conformity Assessment). Total nitrogen was analysed until 2002 after oxidation with  
4 persulphate ( $K_2S_2O_8$ ) and thereafter together with organic carbon with a CN analyser  
5 (Schimadzu). Nitrate nitrogen and nitrite nitrogen have been analysed together and referred to  
6 as nitrate-nitrogen ( $NO_3-N$ ). Organic nitrogen (ON) is calculated as the difference between  
7 Total N and  $NO_3-N$ . Ammonium-nitrogen ( $NH_4-N$ ) was found to be a minor component in  
8 both the drainwater and groundwater according to analyses carried out in the first years.  
9 Phosphorus fractions have been analysed in the drainwater, but not in the groundwater, since  
10 the concentrations were low. Total P has been analysed as soluble molybdate-reactive P after  
11 acid oxidation with  $K_2S_2O_8$ . Dissolved reactive phosphorus (DRP) was analysed after pre-  
12 centrifugation until 2001, and subsequently after pre-filtration. For clay soils in Sweden, these  
13 different pre-treatments have been found to be of major importance for DRP determination,  
14 since the colloidal clay particles are retained more efficiently by filtration (filters from  
15 Schleicher and Schüll, Germany, with pore diameter  $0.2\ \mu m$ ) than by centrifugation (3000  
16 rpm during 20 minutes). However, in this non-clay soil area differences in pre-treatments  
17 were of minor importance and on average less than  $0.002\ mg\ P\ l^{-1}$ . Nevertheless older RP  
18 results were recalculated as if filtration had been performed before analysis using regression  
19 equation from analyses with the two pretreatments. Non-reactive phosphorus (NRP) was  
20 calculated as the difference between Total P and RP and may consist of both organically- and  
21 inorganically-bound P.

22 Total ionic composition was analysed from 1980 in both drainwater and groundwater and  
23 the alkalinity was determined by titration. Analytical procedures for calcium (Ca), magnesium  
24 (Mg), sodium (Na) and potassium (K) were changed in 2002 from AAS (atomic adsorption  
25 spectrophotometry) to ion chromatography (IC). At the same time determination of sulphur in

1 sulphate (SO<sub>4</sub>-S) changed from a colorimetric determination by autoanalyser to analysis with  
2 IC, and determination of chloride (Cl) from flow injection analysis to IC. Tests indicate that  
3 these changes in analytical procedures are of minor importance, in quantitative terms less than  
4 the general uncertainty of laboratory handling and analytical determination (10%).

#### 6 *Average water concentrations and ionic composition*

7 The amount of precipitation was measured at a local station close to the field but data on the  
8 ion components were obtained from a monitoring station for air and precipitation (Lövblad *et*  
9 *al.*, 2002) 62 kilometres north of the field (Figure 1). The average concentrations are values  
10 weighted to the amount of precipitation. Similarly all concentrations in drainwater were  
11 weighted to the amount of water, i.e. the total element transport was divided by the total  
12 drainage amount. Concentrations in groundwater were calculated as straight average values.  
13 Flow-weighted average concentrations in drainwater were compared with management factors  
14 in a simple way by testing for significance (T-test). The annual flow-weighted concentrations  
15 for periods characterised by one factor were added and compared with annual concentrations  
16 for periods characterised by another factor. Similar comparisons have been made based on  
17 results from the long-term experimental plots at Mellby (Torstensson *et al.*, 1992; Torstensson  
18 *et al.*, 2001; Torstensson and Ekre, 2003; Torstensson, 2003).

19 Data on ionic composition of the mineral fertilisers were obtained from the manufacturers  
20 and the concentrations in manure and crop were calculated based on standard composition  
21 data (Eriksson *et al.*, 1997; Steineck *et al.*, 1999). However, the sodium concentration is not  
22 reported in the present paper since this concentration varied widely. All major ions were  
23 expressed in millimoles of charge in order to check the ionic balance of the drainwater and  
24 groundwater. Concentrations from three drainwater samples taken during snowmelt were  
25 excluded based on a large ionic imbalance under these conditions. For the rest of the time, the

1 error in ionic balance was generally low, 1-3%, which indicates no analytical problems.  
2 Precipitation water was checked for negative alkalinity (-Alk) as the difference in the  
3 equivalent sum of base cations (Ca, Mg, Na, K, NH<sub>4</sub>-N) and acid anions (Cl, SO<sub>4</sub> and NO<sub>3</sub>).  
4 The neutralising capacity of calculated alkalinity was also roughly estimated in the fertilisers  
5 and in harvested products but without taking into consideration any sodium or nitrogen ions.

6

### 7 *Trend analysis*

8 All concentrations in the drainwater and the three levels of groundwater were examined for  
9 seasonality according to the Kruskal-Wallis test. All concentrations in the drainwater were  
10 flow-normalised with the non-linear method LOWESS (Locally Weighted Scatter-plot  
11 Smoothing) before trend analysis (Cleveland, 1979). The recommended factor 0.5 was used  
12 for adjustment. Flow-normalised values were the residuals from the curve adjusted to  
13 LOWESS. The significance of the trend was tested according to Hirsch and Slack (1984) using  
14 a non-parametric method (Mann Kendall) in the Visual-Basic program MiniTab<sup>®</sup>. The  
15 'Theil's slope' (Helsel and Hirsch, 1992) was used for calculating the magnitude of the trend.  
16 The method assumes no change in the relationship between concentration and discharge over  
17 time and no trends in the amount of precipitation or drainwater discharge during the period of  
18 investigation, conditions that were completely fulfilled. The concentrations in the  
19 groundwater were normalised against the level of the groundwater, which usually involved  
20 significant seasonality. Only the directions of the trends in the groundwater, if any, are  
21 reported here.

22

### 23 *Theoretical estimations of nitrate concentrations leaving the root-zone*

24 Beside actual measurement of NO<sub>3</sub> in tile drain water and groundwater, annual theoretically  
25 leaching NO<sub>3</sub> concentrations for water leaving the root-zone have also been estimated for a

1 loamy sand in the actual production area (Mårtensson and Johnsson, 2006). These  
2 concentrations have been normalised according to the climate at the actual site. In addition,  
3 these annual leaching coefficients estimated with a combined leaching and database  
4 programme (SOIL-NDB) have been generalised for normal crops and for possible crop  
5 combinations in the production area (Mårtensson and Johnsson, 2006).

6

## 7 **Results**

### 8 *Degree of P saturation in the soil*

9 The topsoil of the field had a P-AL number (9 mg 100 g soil<sup>-1</sup>) that is typical for Sweden. In  
10 contrast, both the Fe-AL number and the Al-AL number (13 and 55 respectively) were four  
11 times higher than average values based on more than 200 soils in the south of Sweden (Ulén,  
12 2006). The Fe-AL value varied in different parts of the field and the variation was high  
13 throughout the entire soil profile (Table 2). This may be a result of the tendency of Fe and  
14 iron hydroxide to occur as discrete mineral particles in the soil. DPS calculated with the PSI  
15 method was higher than that calculated from the Fe-AL and Al-AL concentrations in the  
16 topsoil layer and the layer below plough mixing depth (23-35 cm). The abundant Ca ions in  
17 the topsoil may have bound some P in the neutral solution used for the PSI test, resulting in  
18 relatively low PSI values. The two methods for calculating DPS gave very similar results in  
19 the subsoil and were also well correlated with each other, with a Pearson's correlation  
20 coefficient of 0.92 (p = 0.00). The low values in the 35-65 cm layer (5.0 and 5.5%) and the  
21 35-100 cm layer (10 and 9%) corresponded well with the low RP concentration in the  
22 drainwater (average 0.006 mg l<sup>-1</sup>).

23 Soil sampling in transects along the slopes of the field gave slightly higher DPS values in  
24 the topsoil compared to sampling an evenly distributed net of squares (+3 and +7% for the  
25 extraction method and the PSI method respectively. This was the result of higher P-AL

1 numbers at the lower end of the slope in the south-east part of the field compared with the rest  
2 of the field. The general variation in the ability of the field topsoil to sorb P was quite  
3 moderate; it varied between 22-44% for the two sampling strategies combined with the two  
4 laboratory methods.

5

#### 6 *Characterisation of precipitation, drainwater and groundwater*

7 Average Na concentration in the precipitation was high in this coastal area ( $0.10 \text{ mmol}_e \text{ l}^{-1}$ ).  
8 Precipitation water had a similar large proportion of sulphate and nitrate as the drainwater and  
9 the shallow groundwater at the monitored field (24-28%). The precipitation water can be  
10 described as a water of Na-SO<sub>4</sub>-Cl type, while the drainwater and shallow groundwater  
11 represent Ca-NO<sub>3</sub> water types. There was a weak indication of impact from pig slurry since  
12 the K:Na ratio in the drainwater was slightly higher than 0.4 (based on weight) and since the  
13 average chloride value was relatively high ( $18 \text{ mg l}^{-1}$ ). However, pig manure was applied less  
14 often during later years and the two indications became even weaker with time. Similarly a  
15 low K:Cl ratio in drainwater (0.2 based on weight) tended to become even lower during recent  
16 times.

17 Chemical data for deep groundwater (5.5 m) revealed a water of sodium (Na-Cl) type,  
18 reflecting the marine origin of the clay. This saline groundwater is most probably relict water  
19 often found at depth in this low-lying area. After the last ice age, the sea covered a large  
20 proportion of this landscape. As the land rose out of the sea, groundwater was displaced by  
21 precipitation at different rates depending on local conditions.

22

#### 23 *Nutrient trends in drainwater and groundwater*

24 A substantial reduction in N leaching with catch crops was clearly demonstrated, as was an  
25 effect of modified land use of the field (Tables 3 and 4). Comparing rotations with catch crops

1 to rotations without and to applying less pig slurry revealed a reduction in the NO<sub>3</sub>-N  
2 concentration in the drainwater. Trend analysis also revealed this reduction mainly to take  
3 place in September-November, when catch crops are most effective. Similarly, potatoes  
4 seemed to be a risk crop compared to cereals, a finding which was confirmed in the  
5 experimental plots (Table 3). In addition, organic nitrogen (ON) showed a tendency for lower  
6 leaching from the field when catch crops were grown, but this was not evident from the  
7 experimental plots. A trend for decreasing NO<sub>3</sub>-N in the drainwater equivalent to 5.3 mg l<sup>-1</sup>  
8 over the study period was also apparent, together with the more conservative parameter Na  
9 (Figure 2a). Corresponding time-series indicated an even faster NO<sub>3</sub>-N-reduction for the  
10 groundwater at 1.7 m depth (Figure 2b). The latter was also evident when mean annual  
11 concentrations were compared with estimates based on generalising leaching coefficients for  
12 the actual soil in this agricultural production area (Figure 3). Measured NO<sub>3</sub>-N concentrations  
13 in both drainwater and surface groundwater were also comparatively lower than theoretically  
14 estimated concentrations in root-zone leaching water.

15 Different cropping had no impact on the P concentrations in the drainwater, of the field  
16 which is not surprising based on the demonstrated good sorption capacity of the subsoil. In  
17 addition, Fe in groundwater transported from longer distances in the experimental plots at  
18 Mellby may bind P. Based on the results from the experimental plots, catch crops may reduce  
19 the NRP concentration in the drainwater. However, there was no such indication in the arable  
20 field studied, nor was any trend in P recorded in the drainwater during the entire period.

21

## 22 *Soil nutrient and ionic accumulation*

23 Fertilisation generated P accumulation in the soil at a rate of 0.4 kg ha<sup>-1</sup>yr<sup>-1</sup>, which is a low  
24 value for a Swedish farm with livestock. The P-AL index of the topsoil was found to have  
25 remained unchanged during the twelve-year modified cropping period 1993-2005. Potassium

1 was removed slightly faster by the crops than applied by the fertiliser during the first  
2 production specialisation (1977-1990) but not during the second (Table 5). This may be a  
3 result of more potassium-demanding leys being grown during earlier years (Table 1). The  
4 composition of the used mineral fertilisers changed. During the first period, with the initial  
5 crop rotation, addition of base cations took place in the form of Ca, while there was more net  
6 gain of Mg during later years. During both original and modified crop rotation substantial  
7 underbalances of  $\text{Ca}^{2+}$  and alkalinity were estimated based on inputs and removals from the  
8 soil (Table 5).

9

#### 10 *Ionic composition and trends in precipitation, drainwater and groundwater*

11 The ionic composition in precipitation may have differed slightly between the measured site  
12 and the observation field. The latter is situated closer to the shore but is on the other hand  
13 more protected from western winds by a large spit of land (Figure 1). The ions  $\text{Na}^+$  and  $\text{Cl}^-$  in  
14 the precipitation were indicated to be influenced by sea-salt laden storm episodes over a  
15 number of years. Such temporary and frequent storms were obvious during the earlier part of  
16 this study, namely 1984 and during the period 1989-1992. The episodes were also followed  
17 by short-term increases in  $\text{Cl}^-$  concentration of the drainwater. Lack of such episodes in recent  
18 years was probably the main reason for the apparent decrease in  $\text{Cl}^-$  concentration in  
19 drainwater (Table 4). In contrast concentrations of  $\text{Na}^+$  did not decreased significantly in the  
20 drainwater. These concentrations were positively related to concentrations of Na in the  
21 shallow groundwater were the concentrations were relatively high and even increased (Figure  
22 2b). The relationship between  $\text{NO}_3\text{-N}$  and Na concentrations in the shallow groundwater was  
23 negative and had a Pearson's correlation coefficient of -0.77 ( $p = 0.00$ ). During late summer  
24 (July-September), when there was a generally low groundwater level, this relationship was  
25 even stronger (Pearson's correlation coefficient of -0.87).



1        Decreasing trends in S and increasing trends in alkalinity in the precipitation were quite  
2 clear. The composition of the used mineral fertilisers changed so that more S was applied  
3 during the modified cropping system in later years (Table 5). Some commercial mineral  
4 fertilisers are now enriched in S as compensation for a generally smaller deposition of S from  
5 the air. However, the  $\text{SO}_4^{2-}$  concentration in the drainwater decreased in the period studied  
6 (Table 4) along with reduced load by precipitation. In contrast, the complementary studies in  
7 deeper groundwater showed that  $\text{SO}_4\text{-S}$  concentrations remained constant in these waters  
8 (Table 4).

9        Deeper groundwater had different chemistry and ionic trends than the shallow groundwater  
10 (Table 4). Deeper groundwater mainly provides information about the older long-term  
11 translocations. Moderate and decreasing concentrations of Ca indicate relatively old water  
12 without contact with Ca-rich layers.

13

#### 14 *Soil acid load*

15        The high leaching of Ca removed by drainwater (Table 5) might enhance soil acidification.  
16 However, the documented high effective base saturation of the soil, topsoil as well as subsoil,  
17 indicates a good ability to buffer acidity loads and average pH was satisfactorily high, namely  
18 6.6 in the drainwater and 6.5 in the surface groundwater. Plant uptake of base cations and  
19 their removal in harvested products might also involve substantial soil acidification but were  
20 approximately counteracted by the added fertilisers, as indicated by the rough estimates of  
21 calculated alkalinity (Table 5). Removal of excess base cations by drainwater and by  
22 harvested products was in both cases similar to an approximately ten times higher acid load  
23 than the negative alkalinity via precipitation (Table 5). The latter was calculated to be reduced  
24 by  $6.3 \text{ kmol}_c \text{ ha}^{-1}$  during the modified cropping system compared to the original, equal to a  
25 reduction of more than 50%, while acidification by plant uptake of base-cations remained

1 constantly high. A slow but significant increase in  $\text{HCO}_3^-$  ions was indicated to occur in the  
2 drainwater, as well as a faster increase in the groundwater (Figure 4). The increase in the  
3 drainwater was equal to 0.4% as calculated for the entire period 1977-2004. Meanwhile the  
4 catch crops had efficiently taken up  $\text{NO}_3^-$  ions but the soil had delivered more  $\text{HCO}_3^-$  ions  
5 instead. Intensive catch cropping during the modified period reduced the  $\text{NO}_3\text{-N}$  leaching  
6 from 44-to 32  $\text{kmol}_c \text{ ha}^{-1}$  in comparison with the original period. The reduction in  $\text{NO}_3^-$  ions  
7 (equal to 12  $\text{kmol}_c \text{ ha}^{-1}$ ) loaded the soil, while leached alkalinity increased from 35 to 42  
8  $\text{kmol}_c \text{ ha}^{-1}$ . Based on these estimates, changes in cropping were indicated to be a more  
9 important mitigation of water acidification than the reduction in acid rain. This is also  
10 indicated by the negative correlation between  $\text{HCO}_3^-$  and  $\text{NO}_3^-$  ions in the drainwater during  
11 the periods with catch crops (Pearson's correlation coefficient = -0.60,  $p = 0.00$ ).

12

### 13 **Discussion**

14 Even for this sandy soil some minor lateral long-transport of P may have taken place since  
15 higher P-AL numbers were found at the topsoil lower end of the slope in the south-east part of  
16 the field compared with the rest of the field. Since the two sampling strategies gave similar  
17 results, sampling along slopes is of less importance for this loamy and sandy soil. In contrast  
18 sorption capacity in the subsoil is more important since P probably sorbs to the soil matrix  
19 when water is moving slowly down through the profile. Subsoil values have also proposed for  
20 use in calculating the risk of P leaching in sandy soils in general (Ulén, 2006). Furthermore,  
21 since the ability to retain P in the subsoil based on chemical analysis was indicated to be  
22 similar by the two laboratory methods the most appropriate method may be chosen.

23 Similar sorption abilities are often found in this region. In contrast the degree of P  
24 saturation (DPS) of 5% in the soil extract of the subsoil was low for the condition of other  
25 counties of southern Sweden, where median values for sandy soil are more than twice as high

1 (Ulén, 2006). For clay soils significantly lower sorption ability of the topsoil were often  
2 found. In addition, for clay soils susceptible to erosion less Ca addition might influence the  
3 future capacity of the topsoil to bind P, since formation of Ca-P complexes has been found to  
4 be important for retention of colloidal P even on non-calcareous soils (Ulén and Snäll, 2007).  
5 For the present field, with documented very low losses of colloidal particles, this change  
6 should be of minor importance. Consequently this soil seems to have a very good ability to  
7 sorb P in drainwater, and agricultural practices such as changes in manure application based  
8 on in-field variations in DPS values seem to be of minor importance for P leaching via field  
9 drains at this part of the region. However, sampling along slopes should be recommended for  
10 more fine-textured soils (Heckrath *et al.*, 2005; McDowell *et al.*, 2001), and P manuring  
11 application rate should be adapted to site conditions based on the ability to retain P  
12 determined from chemical analysis of the soil.

13 The findings from the present site demonstrate the general differences between P and N  
14 retention mechanisms in agricultural water from a sandy soil. Even after storm- or  
15 snowmelting events no abrupt changes in P concentration took place, no general macropore  
16 flow might have occur but adsorption may take place of P during water percolating through  
17 the soil profile. In contrast, reduced N mass transport is an essential part of reduced  $\text{NO}_3^-$   
18 concentration from the root zone. In certain areas groundwater may constitute an important N  
19 store for receiving streams (Ruiz *et al.*, 2002).

20 The magnitude of  $\text{NO}_3\text{-N}$  reduction in the present study was faster than indicated by  
21 recent (1993-2004) trends in agricultural rivers in SW Sweden, where catch crops have been  
22 cropped on 13-17% of the arable land for five years (Ulén and Fölster, 2007). This is not  
23 surprising since the present field represent more than ten years of modified crop rotation and,  
24 in addition, more frequent cultivation of catch crop, equal to 41% of the area. Theoretically  
25 and generalised relative reduction in  $\text{NO}_3\text{-N}$  leaving the root-zone as a result of the

1 corresponding crop rotation including catch crops was also slower, only 13% during the entire  
2 period, based on estimates from model calculations. The apparent reductions in drainage and  
3 shallow groundwater were even faster (Figure 3). Generally sandy soils with good water  
4 drainage are relatively unlikely to promote fast microbial denitrification and this process has  
5 also been estimated to be of minor importance at the Mellby experimental field (Aronsson and  
6 Torstensson, 1998). Since  $\text{NO}_3$  generally follows advective forces of the water, a main reason  
7 for fast reduction in concentrations may be a generally short transit time in the present surface  
8 groundwater moving just above a clay layer. In this shallow groundwater K decreased in a  
9 similar way as in the drainwater, as did  $\text{NO}_3$  - N. These similarities in observed concentration  
10 changes supports the general finding that the transit time for the surface groundwater is short  
11 at the present site, in contrast to several other Swedish sites (Gustafson, 1983). Chemical  
12 reduction of  $\text{NO}_3$ -N by pyrite may also have taken place in the groundwater near the clay  
13 layer in the field studied here. This cannot be clearly verified since no significant increase in  
14 the pyrite oxidation product  $\text{SO}_4^{2-}$  could be estimated in the shallow groundwater. The  
15 concentration of this ion that decreased in the drainwater showed no trends in the shallow  
16 groundwater. These results were most probably the result of negative S load to the soil  
17 together with retardation by sorption.

18 The direct impact from fertilisers on ionic composition of the groundwater was small in  
19 the present study but. Furthermore indirect changes in weathering rates and ion-exchange  
20 equilibrium of most constituents in groundwater as a result of agricultural impacts on the  
21 recharge fluxes of  $\text{NO}_3^-$  and  $\text{H}^+$  were apparently small except for  $\text{HCO}_3^-$ . Based on the poor  
22 correlation no exchange of  $\text{Na}^+$  ion apparently occurs between drainwater and surface water.  
23 In contrast, the relatively high and increasing concentrations of Na in the shallow  
24 groundwater may indicate a net accumulation of this ion.

1 The clay layer in the subsoil of the agricultural field bottom mainly seems to form a  
2 hydraulic and transport barrier since more constant concentrations occurred in the deep  
3 groundwater. However the  $\text{Na}^+$  ion may be an exception and increasing concentration of Na in  
4 the shallow groundwater at the present site might have been a result of mass transfer from  
5 deeper to more shallow groundwater, the former being quite rich in Na. Alternatively ionic  
6 exchange may occur between the lower layer of shallow groundwater and the clay matrix  
7 layer. The latter suggestion may explain the fact that the apparent relationship between  $\text{NO}_3^-$   
8 and  $\text{Na}^+$  concentrations in the shallow groundwater was negative and was even stronger when  
9 groundwater level was generally low. At the deepest sampling depth (5.5 m), the Na  
10 concentration probably increases as more water is pumped out from the groundwater pipes by  
11 sampling. The very sampling may be a significant water movement in this rather stagnant  
12 water with slow lateral transport.

13 Generally groundwater reaction rates may be estimated based on tracers and not on  
14 conservative ions. Separation of the components derived from groundwater and soilwater is  
15 also desirable for a deeper understanding of the processes (Langan and Hirst, 2004), but was  
16 outside the scope of the present study.

17

## 18 **Conclusions**

19

20

- 21 • Phosphorus losses were related to the ability of the subsoil to retain P at the present  
22 site, whereas soil sampling strategy and laboratory method were of minor importance  
23 for risk estimates of P losses.
- 24 • Modification of the crop rotation can rapidly decrease surface water and shallow  
25 groundwater nitrogen loads by leaching, even under intense farming. At the present  
26 site this might have been promoted by a short transit time of the shallow groundwater.

- Modification of the crop rotation in the present study, i.g. inclusion of a catch crop, also prevented acidification of the drainwater since the alkalinity increased while the nitrate concentration decreased. Compared with this, reduced load of acid rain was indicated to be of less importance.
- Deeper knowledge of processes resulting in changed ionic composition of the groundwater should be obtained by tracer studies.

## Acknowledgements

Our sincere thanks for long-term discussions of the results to Arne Gustafson. He also established the monitoring arable field. The long-term monitoring system of arable fields is funded by the Swedish Environmental Protection Agency. Financial support for the evaluation was also provided by the Swedish Farmers' Foundation for Agricultural Research (SLF).

## References

- Andersen, A. & Olsen, C. C. 1993 Rye grass as a catch crop in spring barley. *Acta Agriculturae Scandinavica* **43**, 218-230.
- Aronsson, H. 2000 Nitrogen turnover and leaching in cropping systems with ryegrass catch crops. Dissertation. *Agraria* **214**, SLU, Uppsala, Sweden.
- Aronsson, H. & Torstensson, G. 1998 Measured and simulated availability and leaching of nitrogen associated with frequent use of catch crops. *Soil Use and Management* **14**, 6-13.
- Aronsson, H. & Torstensson, G. 2004 Beräkning av olika odlingsåtgärders inverkan på kväveutlakningen. Beskrivning av ett pedagogiskt verktyg för beräkning av kväveutlakning från enskilda fält och gårdar. *Ekohydrologi* **78**. Div. Water Quality Management, Swedish University of Agricultural Sciences, Uppsala. (In Swedish.)
- Bergqvist, G., Ohlander L. & Rydberg, T. 2002 Insådd av mellangrödor i höstsäd. Inst. Ecology and Crop production, Report 4, SLU, Uppsala, Sweden. (In Swedish.)
- Bergström, L. and Gustafson, A. 1985 Hydrogen ion budgets of four small runoff basins in Sweden. *Ambio* **14**, 346-348.
- Bergström, L. and Kirchmann, H. 1999 Leaching of total nitrogen from 15-labeled poultry manure and inorganic nitrogen fertilizer. *Journal Environmental Quality* **28**, 1283-1290.
- Böhlke, J-K. 2002 Groundwater recharge and agricultural contamination. *Hydrogeology Journal* **10**, 153-179.
- Börling, K., Otabbong, E. & Berberis, E. 2001 Phosphorus sorption in relation to soil properties in some cultivated soils. *Nutrient Cycling in Agroecosystems* **59**, 39-46.
- Cleveland, W. S. 1979 Robust locally weighted regression and smoothing scatter plots. *Journal of the American Statistical Association* **74**, 829-836.

- 1 Debreczeni, K. & Kismányoky, T. 2005 Acidification of soils in long-term field experiments.  
2 *Communications in Soil Science and Plant Analysis* **36**, 321–329.
- 3 Djodjic F. & Bergström L. 2005a Conditional phosphorus index as an educational tool for risk  
4 assessment and phosphorus management. *Ambio* **34** (4), 296–300.
- 5 Djodjic, F. & Bergström, L. 2005b Phosphorus losses from arable fields in Sweden—effects  
6 of field-specific and long-term trends. *Environmental Monitoring and Assessment* **102**, 103–  
7 117.
- 8 Egnér, H., Riehm, H. & Domingo, W. R. 1960 Untersuchungen über die chemische  
9 Bodenanalyse als Grundlage für die Beurteilung des Nährstoffzustandes der Böden. II  
10 Chemische Extraktionsmethoden zur Phosphor- und Kaliumbestimmung. *Kungliga*  
11 *Lantbrukshögskolans Annaler* **26**, 199–215.
- 12 Eriksson, J., Andersson, A., Stenberg, B. & Andersson, R. 1997 Tillståndet i svensk åkermark  
13 och spannmålsgröda. *Swedish Environmental Protection Agency Report 5062*. 106–48  
14 Stockholm, 65 pp. (In Swedish).
- 15 Gustafson, A. 1983 Leaching of nitrate from arable land into groundwater in Sweden.  
16 *Environmental Geology* **5** (2), 65–71.
- 17 Gustafson, A. 1987 Nitrate leaching from arable land in Sweden under four cropping systems.  
18 *Swedish Journal of Agriculture Research* **17**, 169–177.
- 19 Gustafson, A., Gustavsson, A. & Torstensson, G. 1984 Intensity and duration of drainage  
20 discharge from arable land. *Ekohydrologi* **16**, 75 pp. Division of Water Quality  
21 Management, Swedish University of Agricultural Sciences (In Swedish).
- 22 Helsel, D. R. & Hirsch, R. M. (1992) *Statistical Measures in Water Research*. 529 pp.  
23 Elsevier Science B. V., Amsterdam.
- 24 Hessel Tjell, K., Aronsson, H., Torstensson, G., Gustafson, A., Lindén, B., Stenberg, M. &  
25 Rydberg, T. 1999 Mineralkvävedynamik och växtnäringsutlakning i handels- och  
26 stallgödslade odlingsssystem med och utan fänggröda. Resultat från en grovmjord i södra  
27 Halland, perioden 1990–1998. *Ekohydrologi* **50**. Div. Water Quality Management, Swedish  
28 University of Agricultural Sciences, Uppsala. (In Swedish.)
- 29 Hirsch, R. M. & Slack, J. R. 1984 A nonparametric trend test for seasonal data with serial  
30 dependence. *Water Resources Research* **20**, 727–732.
- 31 Hirsch, R. M., Slack, J. R. & Smith, R. A. 1982 Techniques of trend analysis for monthly  
32 water quality data. *Water Resources Research* **18**, 107–121.
- 33 Hornish, M., Hellmeier, C. & Weiss, K. 2002 Response of surface and subsurface water  
34 quality to land use change. *Geoderma* **104**, 277–298.
- 35 Johansson, H. & Mårtensson, K. 2006 Calculation of nitrogen leaching effects from  
36 environmental subsidies 'reduced nitrogen leaching' year 2001 based on TRK calculations  
37 for year 1999. Technical report Division of Water Quality Management, Swedish  
38 University of Agricultural Sciences, Uppsala, Sweden (in Swedish).
- 39 Kirchmann, H., Johnson, A. E. J. & Bergström L. F. 2002 Possibilities for reducing nitrate  
40 leaching from agricultural land. *Ambio* **31**, 404–408.
- 41 Kvist, M. 1992 Catch crops undersown in barley—compensative effects and cropping  
42 methods. *Dissertation, Crop Production Science 15*, Department of Crop Production  
43 Sciences SLU, Uppsala.
- 44 Langan, S. L. & Hirst, D. 2004 An analysis of the long-term variation in stream water quality  
45 for three upland catchments at Loch Dee (Galloway, SW Scotland) under contrasting land  
46 management. *Hydrology and Earth System Sciences* **8** (3), 422–435.
- 47 Lövlblad, G., Henningsson, E., Sjöberg, K., Brorström-Lundén E., Lindskog, A. & Munthe, L.  
48 2002 *Trends in Swedish Background air 1980–2000*. IVL, Swedish Environmental  
49 Research Institute, PO Box 5302, SE 40014 Göteborg.

- 1 [McDowell, R., Sharpley, A. and Folmar, G. 2001. Phosphorus export from an agricultural](#)  
2 [watershed Debreczeni, K. & Kismányoky, T. 2005 Acidification of soils in long-term field](#)  
3 [experiments. \*Communications in Soil Science and Plant Analysis\* \*\*36\*\*, 321-329.](#)
- 4 [Djodjic F. & Bergström L. 2005a Conditional phosphorus index as an educational tool for risk](#)  
5 [assessment and phosphorus management. \*Ambio\* \*\*34\*\* \(4\), 296-300.](#)
- 6 [Djodjic, F. & Bergström, L. 2005b Phosphorus losses from arable fields in Sweden – effects](#)  
7 [of field-specific and long-tem trends. \*Environmental Monitoring and Assessment\* \*\*102\*\*, 103-](#)  
8 [117.](#)
- 9 [Egnér, H., Riehm, H. & Domingo, W. R. 1960 Untersuchungen über die chemische](#)  
10 [Bodenanalyse als Grundlage für die Beurteilung des Nährstoffzustandes der Böden. II](#)  
11 [Chemische Extraktions-methoden zur Phosphor och Kaliumbestimmung. \*Kungliga\*](#)  
12 [\*Lantbrukshögskolans Annaler\* \*\*26\*\*, 199-215.](#)
- 13 [Eriksson, J., Andersson, A., Stenberg, B. & Andersson, R. 1997 Tillståndet i svensk åkermark](#)  
14 [och spannmålsgröda. \*Swedish Environmental Protection Agency Report 5062\*. 106 48](#)  
15 [Stockholm , 65 pp. \(In Swedish.\).](#)
- 16 [Gustafson, A. 1983 Leaching of nitrate from arable land into groundwater in Sweden.](#)  
17 [\*Environmental Geology\* \*\*5\*\* \(2\), 65-71.](#)
- 18 [Gustafson, A. 1987 Nitrate leaching from arable land in Sweden under four cropping systems.](#)  
19 [\*Swedish Journal of Agriculture Research\* \*\*17\*\*, 169-177.](#)
- 20 [Gustafson, A., Gustavsson, A. & Torstensson, G. 1984 Intensity and duration of drainage](#)  
21 [discharge from arable land. \*Ekohydrology\* \*\*16\*\*, 75 pp. Division of Water Quality](#)  
22 [Management. Swedish University of Agricultural Sciences \(In Swedish\).](#)
- 23 [Helsel, D. R. & Hirsch, R. M. \(1992\) \*Statistical Measures in Water Research\*. 529 pp.](#)  
24 [Elsevier Science B. V., Amsterdam.](#)
- 25 [Hessel Tjell, K., Aronsson, H., Torstensson, G., Gustafson, A., Lindén, B., Stenberg, M. &](#)  
26 [Rydberg, T. 1999 Mineralkvävedynamik och växtnäringsutlakning i handels- och](#)  
27 [stallgödslade odlingssystem med och utan fånggröda. Resultat från en grovmjord i södra](#)  
28 [Halland, perioden 1990-1998. \*Ekohydrologi\* \*\*50\*\*. Div. Water Quality Management, Swedish](#)  
29 [University of Agricultural Sciences, Uppsala. \(In Swedish.\)](#)
- 30 [Hirsch, R. M. & Slack, J. R. 1984 A nonparametric trend test for seasonal data with serial](#)  
31 [dependence. \*Water Resources Research\* \*\*20\*\*, 727-732.](#)
- 32 [Hirsch, R. M., Slack, J. R. & Smith, R. A. 1982 Techniques of trend analysis for monthly](#)  
33 [water quality data. \*Water Resources Research\* \*\*18\*\*, 107-121.](#)
- 34 [Hornish, M., Hellmeier, C. & Weiss, K. 2002 Response of surface and subsurface water](#)  
35 [quality to lands use change. \*Geoderma\* \*\*104\*\*, 277-298.](#)
- 36 [Johansson, H. & Mårtensson, K. 2006 Calculation of nitrogen leaching effects from](#)  
37 [environmental subsidies 'reduced nitrogen leaching' year 2001 based on TRK calculations](#)  
38 [for year 1999. Technical report Division of Water Quality Management. Swedish](#)  
39 [University of Agricultural Sciences. Uppsala, Sweden \(in Swedish\).](#)
- 40 [Kirchmann, H., Johnson, A. E. J. & Bergström L. F. 2002 Possibilities for reducing nitrate](#)  
41 [leaching from agricultural land. \*Ambio\* \*\*31\*\*, 404-408.](#)
- 42 [Kvist, M. 1992 Catch crops undersown in barley – compensative effects and cropping](#)  
43 [methods. \*Dissertation, Crop Production Science 15\*, Department of Crop Production](#)  
44 [Sciences SLU, Uppsala.](#)
- 45 [Langan, S. L. & Hirst, D. 2004 An analysis of the long-term variation in stream water quality](#)  
46 [for three upland catchments at Loch Dee \(Galloway, SW Scotland\) under contrasting land](#)  
47 [management. \*Hydrology and Earth System Sciences\* \*\*8\*\* \(3\), 422-435.](#)
- 48 [Lövlblad, G., Henningsson, E., Sjöberg, K., Brorström-Lundén E., Lindskog, A. & Munthe, L.](#)  
49 [2002 \*Trends in Swedish Background air 1980-2000\*. IVL. Swedish Environmental](#)  
50 [Research Institute, PO Box 5302, SE 40014 Göteborg.](#)



- 1 [McDowell, R., Sharpley, A. and Folmar, G. 2001. Phosphorus export from an agricultural –](#)  
2 linking sources and transport mechanisms. *Journal of Environmental Quality* **30**, 1587-  
3 1595.
- 4 Ohlander, L., Bergkvist, G., Stendahl, F. & Kvist, M. 1996 Yield of catch crops and spring  
5 barley as affected by time of undersowing. *Acta Agriculturae Scandinavica* **456**, 161-168.
- 6 Parkhurst, D.L. & Plummer, L.N. 1993. Geochemical models in: in Alley, W.D. (ed) Regional  
7 Groundwater Quality. Van Nostrand Reinhold, New York, pp 199-225.
- 8 Postma, D., Boesen, C, Kristiansen, H. & Larsen, F. 1991 Nitrate reduction in an unconfined  
9 sandy aquifer: Water chemistry, reduction processes, and geochemical modeling. *Water*  
10 *Resource Research* **27**, 2027-2045.
- 11 Ruiz, L., Abiven, S., Martin, C., Durand, P., Beaujouan, V. & Molenat, J. 2002. Effect on  
12 nitrate concentration in stream water of agricultural practices in small catchments in  
13 Brittany (France). II Temporal variations and mixing processes. *Hydrology & Earth*  
14 *System Sciences* **6** 507-503.
- 15 Sjöström, J. 1993 Ionic composition and mineral equilibria of acid groundwater on the west  
16 coast of Sweden. *Environmental Geology* **21**, 219-226.
- 17 Steineck, S., Gustafson, G., Andersson, A., Tersmeden, M. & Bergström, J. 1999 Animal  
18 manure – contents of nutrients and trace elements. *Swedish Environmental Protection*  
19 *Agency Report 4794*, 106 48 Stockholm, 28 pp. (In Swedish.)
- 20 Torstensson, G. 1998 Nitrogen availability for crop uptake and leaching. Dissertation,  
21 *Agraria* **98**, SLU, Uppsala, Sweden.
- 22 Torstensson, G. 2003. Ekologisk odling – Utlakningsrisker och kväveomsättning i ekologiska  
23 odlingssystem med resp. utan djurhållning på sandig grovmo i södra Halland. Resultat från  
24 perioden 1991 – 2002. *Ekohydrologi* **72**, Avd. för vattenvårdslära, Sveriges  
25 Lantbruksuniversitet, Uppsala. (In Swedish.)
- 26 Torstensson, G. & Aronsson, H. 2000 Nitrogen leaching and crop availability in manured  
27 catch crop systems. *Nutrient Cycling in Agroecosystems* **56** (2) 139-152.
- 28 Torstensson, G., Gustafson, A., Lindén, B. & Skyggesson, G. 1992 Mineralkvävedynamik  
29 och växtnäingsutlakning på en grovmojord med handels- och stallgödslade odlingssystem  
30 i södra Halland. *Ekohydrologi* **28**, Div. Water Quality Management, Swedish University of  
31 Agricultural Sciences, Uppsala. (In Swedish.)
- 32 Torstensson, G. and Håkansson, M. 2001 Kväveutlakning på sandjord – motåtgärder med ny  
33 odlingsteknik. Miljöanpassad stallgödselanvändning och odling i realistiska odlingssystem.  
34 Resultat från perioden 1991-1999. *Ekohydrologi* **57**, Div. Water Quality Management,  
35 Swedish University of Agricultural Sciences, Uppsala. (In Swedish.)
- 36 Torstensson, G., and Ekre, E. 2003 Kväveutlakning på sandjord - motåtgärder med ny  
37 odlingsteknik. Miljöanpassad stallgödselanvändning och odling i realistiska odlingssystem.  
38 Resultat från en grovmojord i södra Halland, perioden 1999-2002. *Ekohydrologi* **71**, Div.  
39 Water Quality Management, Swedish University of Agricultural Sciences, Uppsala. (In  
40 Swedish.)
- 41 Torstensson, G., Gustafson, A., Lindén, B. and Skyggesson, G. (1992) Mineralkvävedynamik  
42 och växtnäingsutlakning på en grovmojord med handels- och stallgödslade odlingssystem  
43 i södra Halland. *Ekohydrologi* **28**, Div. Water Quality Management, Swedish University of  
44 Agricultural Sciences, Uppsala. (In Swedish.)
- 45 Ulén, B. 2006 A simplified risk assessment for losses of dissolved reactive phosphorus  
46 through drainage pipes from agriculture soils. *Acta Agriculturae Scandinavica B* **56**, 307-  
47 314.

- 1 Ulén, B. & Fölster, J. 2007 Recent trends in nutrient concentrations in Swedish agricultural  
2 rivers. *Science of the Total Environment* **373**, 473-487.
- 3 Ulén, B. and Snäll, S. 2007 Forms and retention of phosphorus in an illite-clay soil profile  
4 with a history of fertilisation by pig manure and mineral fertilisers. *Geoderma* **137**, 455-  
5 465.
- 6 Ulén, B., Carlsson, C. & Lidberg, B. 2004 Recent trends and pattern in agricultural streams.  
7 *Environmental Monitoring and Assessment* **101**, 103-117.
- 8 Ulén, B., Aronsson, H., Bergström, L., Gustafson, A., Larsson, M. & Torstensson, G. 2006  
9 Swedish long-term experimental sites for studying nutrient losses, nutrient turnover and  
10 model developments. *Ekohydrologi* **90**, Div. Water Quality Management, Swedish  
11 University of Agricultural Sciences, Uppsala.
- 12 Ulén, B., Johansson, G. & Simonsson, M. 2008. Leaching of nutrients and major ions from an  
13 arable field with an unfertilized fallow as infield buffer zone. *Acta Agriculturae*  
14 *Scandinavica B* **58**, 51-59..
- 15 Warfvinge, P. and Bertills, U. (2000) Recovery from acidification in the natural environment.  
16 *Swedish Environmental Protection Agency Report 5034*. Berglings skogs, Trelleborg,  
17 Sweden.  
18

1  
 2 Table 1. Acreage of crops (%) grown during the  
 3 original cropping period (1977-1990) and in the modified crop  
 4 rotation (1992-2004), including catch crops

Crop/Year	Original	Modified
	1977-1990	1991-2004
Oats	13	14
Barley	3	11
Spring wheat	7	19
Winter wheat	25	7
Spring oilseeds	14	0
<i>Catch crop</i>	0	41
Peas	10	0
Potatoes	14	21
Sugarbeet	0	13
Ley	13	0
Fallow	0	14

5  
 6  
 7

1  
2  
3  
4  
5

Table 2. Average (Av) and variance (Var) of calcium, phosphorous, iron and aluminium extracted in acid ammonium lactate (Ca-AL, P-AL, Fe-AL and Al-AL) (mg 100 g dry soil<sup>-1</sup>). Phosphorus saturation index (PSI) and calculated degree of phosphorus saturation (%) based on the extraction (DPS-EXT) and on phosphorus sorption index (DPS-PSI) at different soil depths based on 34 samples

Depth	Ca-AL		P-AL		Fe-AL		Al-AL		PSI		DPS-EXT		DPS-PSI	
	Av	Var	Av	Var	Av	Var	Av	Var	Av	Var	Av	Var	Av	Var
0-23	147	26	8.8	27	12.5	62	54.9	36	9.5	13	13.6	34	30.6	32
23-35	92	73	2.8	57	10.8	120	56.0	57	10.1	33	5.2	89	10.0	75
35-65	36	81	0.9	87	7.4	123	27.0	70	6.6	42	5.0	172	5.5	105
65-100	26	92	1.3	73	3.5	95	11.9	81	4.5	40	15.2	97	12.4	89

6  
7

1 Table 3. Average concentrations ( $\text{mg l}^{-1}$ ) of nitrate nitrogen ( $\text{NO}_3\text{-N}$ ), organic nitrogen (ON), dissolved  
 2 reactive phosphorus (RP), non-reactive phosphorus (NRP) and potassium (K) in drainwater during  
 3 1976-2005. Comparisons were made between: periods/years of original (1977-90) crop rotation and  
 4 modified (1992-2004) crop rotation; loadings of pig slurry (Pig slurry); growing cereals with catch  
 5 crops (Catch crop); growing potatoes and growing cereals/oilseeds without catch crops (Cereals).  
 6 Concentrations with non-significant differences are given within brackets. The last column shows  
 7 number of years (Y) and number of fields/plots (No) with the different agricultural managements

	$\text{NO}_3\text{-N}$	ON	RP	NRP	K	YxNo
<i>Arable field years 1977-2005</i>						
Pig slurry, original	13.0	1.2	(0.005)	(0.014)	5.4	14x1
Catch crop, dairy, modified	7.2 <sup>b</sup>	0.9 <sup>b</sup>	(0.006)	(0.015)	4.0 <sup>b</sup>	14x1
Pig slurry	11.8	1.1	(0.007)	(0.018)	5.0	8x1
Catch crop	6.2 <sup>b</sup>	0.8 <sup>b</sup>	(0.005)	(0.016)	3.9 <sup>b</sup>	7x1
Potatoes	13.9	(1.2)	(0.007)	(0.014)	(5.3)	6x1
Cereals	10.3 <sup>a</sup>	(1.1)	(0.005)	(0.014)	(5.1)	9x1
<i>Experimental plots 1984-1997</i>						
Pig slurry	19.4	(1.7)	(0.032)	0.047	15.0	4x4
Catch crop	3.6 <sup>b</sup>	(1.5)	(0.040)	0.028 <sup>b</sup>	8.0 <sup>b</sup>	7x2
Potatoes	17.6	(1.8)	(0.038)	(0.044)	(6.1)	6x3
Cereals	11.1 <sup>a</sup>	(1.8)	(0.032)	(0.053)	(9.4)	10x2

<sup>b</sup> Highly significant ( $p \leq 0.01$ )

<sup>a</sup> Significant ( $0.01 < p \leq 0.05$ )

8  
 9  
 10  
 11

1 Table 4. Average concentration (Conc.) in precipitation, drainwater (*Drainwater*), and in groundwater  
 2 at 1.7 m, 2.2 and 5.5 m depth (*Groundwater*) of calcium (Ca<sup>2+</sup>), magnesium (Mg<sup>2+</sup>), potassium (K<sup>+</sup>),  
 3 sodium (Na<sup>+</sup>), nitrogen as nitrate (N<sup>-</sup>), chloride (Cl<sup>-</sup>) sulphate as sulphur (S<sup>2-</sup>) and alkalinity (Alk) in  
 4 millimole of charge per litre (mmol<sub>c</sub> l<sup>-1</sup>) during 1981-2004. Estimated significant (z > 1.78) trends in  
 5 drainwater (non-significant trend within brackets) and the direction of significant trends (Trend direc.)  
 6 for precipitation and groundwater. Significant seasonality (ssy) according to Kruskal-Wallis test is  
 7 indicated for the parameters in the groundwater

Period	Ca <sup>2+</sup>	Mg <sup>2+</sup>	K <sup>+</sup>	Na <sup>+</sup>	N <sup>-</sup>	Cl <sup>-</sup>	S <sup>2-</sup>	Alk
<i>Precipitation*</i>								
Conc.	0.008	0.012	0.003	0.050	0.043	0.060	0.052	-0.035
Trend direc.	0	0	0	0	0	0	-	+
<i>Drainwater</i>								
Conc.	1.79	0.28	0.13	0.52	0.55	0.50	0.72	0.88
Trend direc.	-0.025	(-0.001)	-0.001	(-0.001)	-0.38	-0.005	-0.022	+0.004
<i>Groundwater (1.7 m)</i>								
Conc.	1.65	0.40	0.12	0.91	0.42	0.63	1.30	0.73
Trend direc.	-	+	-	+(ssy)	-(ssy)	(ssy)	(ssy)	0
<i>Groundwater (2.2 m)</i>								
Conc.	2.76	0.77	0.12	2.45	0.054	1.10	1.42	3.41
Trend direc.	-	+	0	+	-(ssy)	+	(ssy)	+
<i>Groundwater (5.5 m)</i>								
Conc.	1.66	1.30	0.31	11.91	0.026	5.85	1.47	7.72
Trend direc.	-(ssy)	+	-(ssy)	+(ssy)	-	+(ssy)	0	+

8 \* in addition H<sup>+</sup> 0.035 and NH<sub>4</sub><sup>+</sup> 0.046 (mmol<sub>c</sub> l<sup>-1</sup>)

9 \*\* Negative alkalinity calculated from the ionic composition

10

11

1 Table 5. Inputs by precipitation (+*Precipitation*) and by fertilisers and manure (+*Fertilisation*),  
 2 removed by crops (-*Crop*) and loss by drainage (-*Drain*) of calcium ( $\text{Ca}^{2+}$ ), magnesium ( $\text{Mg}^{2+}$ ),  
 3 potassium ( $\text{K}^+$ ), nitrogen as ammonium, ( $\text{N}^+$ ), nitrogen as nitrate ( $\text{N}^-$ ), phosphorus as hydrogen  
 4 phosphate ( $\text{P}^-$ ), chloride ( $\text{Cl}^-$ ) and sulphate as sulphur ( $\text{S}^{2-}$ ) and alkalinity (Alk) in kilomoles of charge  
 5 per hectare ( $\text{kmol}_c \text{ ha}^{-1}$ ) during two (14 year) periods of different crop rotations, original (1977-1990)  
 6 and modified (1992-2004)

Period	$\text{Ca}^{2+}$	$\text{Mg}^{2+}$	$\text{K}^+$	$\text{N}^+$	$\text{N}^-$	$\text{P}^-$	$\text{Cl}^-$	$\text{S}^{2-}$	Alk
<i>+Precipitation<sup>a</sup></i>									
Original	1.2	2.6	0.3	4.9	4.6	-	15.2	8.5	-11.7 <sup>b</sup>
Modified	1.0	2.0	0.3	3.8	4.1	-	9.8	5.2	-5.4 <sup>b</sup>
<i>+Fertilisation</i>									
Original	33.9	11.6	21.6	48.6	49.9	10.8	7.5	17.6	(31) <sup>b</sup>
Modified	25.6	23.8	31.1	67.8	44.4	11.6	6.6	27.0	(35) <sup>b</sup>
<i>-Crop</i>									
Original	8.6	10.0	23.9	-	-	6.3	2.0	4.2	(43) <sup>b</sup>
Modified	9.5	10.0	22.2	-	-	6.9	2.5	2.1	(41) <sup>b</sup>
<i>-Drain*</i>									
Original	104.3	14.0	7.4	0.1	43.5	0.1	28.6	43.4	34.5
Modified	89.6	15.1	6.0	0.1	31.5	0.1	24.4	36.8	42.1

<sup>a</sup> Estimates from the period 1980-2002 and 1980-2004 respectively

<sup>b</sup> Calculated from the ionic composition. The calculated alkalinity is negative in precipitation water

7  
8  
9  
10  
11





Figure 1. Central county of Halland in SW Sweden with the arable monitored field N12, the experimental site Mellby and precipitation station used for chemical determinations (P chem). To the left tile drainage map, soil map and soil profiles with groundwater

pipes.


- 1
- 2






1

1  
2  
3  
4  
5



6  
7  
8  
9  
10  
11  
12


Figure 2. a) Measured concentration (mmol<sub>c</sub> l<sup>-1</sup>) of nitrate nitrogen (NO<sub>3</sub>-N) and sodium (Na) in drainwater. b) Measured concentration (mmol<sub>c</sub> l<sup>-1</sup>) of nitrate nitrogen (NO<sub>3</sub>-N) and sodium (Na) in the shallow groundwater pipe with the inlet at 1.7 m depth.



13  
14  
15  
16  
17

Figure 3. Theoretically estimated yearly concentrations of nitrate nitrogen (NO<sub>3</sub>-N in mg l<sup>-1</sup>) leaving the root-zone together with actual measured concentrations in drainwater (flow-weight averaged) and shallow groundwater (right averages).

1  
2



3  
4  
5  
6  
7  
8  
9

Figure 4. Concentrations (mmol<sub>c</sub> l<sup>-1</sup>) of hydrocarbonate-ions (HCO<sub>3</sub><sup>-</sup>) in drainwater and in shallow groundwater.