

Acknowledgements

This research was sponsored by the EPSRC. T.W.F. first suggested the electrochemical deoxidation of titanium metal. G.Z.C. was the first to observe that it was possible to reduce thick layers of oxide on titanium metal using molten salt electrochemistry. D.J.F. suggested the experiment, which was carried out by G.Z.C., on the reduction of the solid titanium dioxide pellets. M. S. P. Shaffer took the original SEM image of Fig. 4a.

Correspondence and requests for materials should be addressed to D. J. F. (e-mail: djf25@hermes.cam.ac.uk).

Reduced calcification of marine plankton in response to increased atmospheric CO₂

Ulf Riebesell*, Ingrid Zondervan[†], Björn Rost[‡], Philippe D. Tortell[†], Richard E. Zeebe^{*‡} & François M. M. Morel[†]

* Alfred Wegener Institute for Polar and Marine Research, P.O. Box 120161, D-27515 Bremerhaven, Germany

[†] Department of Geosciences & Department of Ecology and Evolutionary Biology, Princeton University, Princeton, New Jersey 08544, USA

[‡] Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York 10964, USA

The formation of calcareous skeletons by marine planktonic organisms and their subsequent sinking to depth generates a continuous rain of calcium carbonate to the deep ocean and underlying sediments¹. This is important in regulating marine carbon cycling and ocean-atmosphere CO₂ exchange². The present rise in atmospheric CO₂ levels³ causes significant changes in surface ocean pH and carbonate chemistry⁴. Such changes have been shown to slow down calcification in corals and coralline macroalgae^{5,6}, but the majority of marine calcification occurs in planktonic organisms. Here we report reduced calcite production at increased CO₂ concentrations in monospecific cultures of two dominant marine calcifying phytoplankton species, the coccolithophorids *Emiliania huxleyi* and *Gephyrocapsa oceanica*. This was accompanied by an increased proportion of malformed coccoliths and incomplete coccospheres. Diminished calcification led to a reduction in the ratio of calcite precipitation to organic matter production. Similar results were obtained in incubations of natural plankton assemblages from the north Pacific ocean when exposed to experimentally elevated CO₂ levels. We suggest that the progressive increase in atmospheric CO₂ concentrations may therefore slow down the production of calcium carbonate in the surface ocean. As the process of calcification releases CO₂ to the atmosphere, the response observed here could potentially act as a negative feedback on atmospheric CO₂ levels.

By the end of the next century, the expected increase in atmospheric CO₂ (ref. 3) will give rise to an almost threefold increase in surface ocean CO₂ concentrations relative to pre-industrial values. This will cause CO₃²⁻ concentrations and surface water pH to drop by about 50% and 0.35 units, respectively⁴. Changes of this magnitude have been shown to significantly slow down calcification of temperate and tropical corals and coralline macroalgae^{5,6}. Although coral reefs are the most conspicuous life-supporting calcareous structures, the majority of biogenic carbonate precipitation (>80%) is carried out by planktonic microorganisms¹, particularly coccolithophorids⁷. These unicellular microalgae are major contributors to marine primary production and an important component of open ocean and coastal marine ecosystems⁸. Two prominent

representatives of the coccolithophorids, *Emiliania huxleyi* and *Gephyrocapsa oceanica*, are both bloom-forming and have a world-wide distribution. *G. oceanica* is the dominant coccolithophorid in neritic environments of tropical waters⁹, whereas *E. huxleyi*, one of the most prominent producers of calcium carbonate in the world ocean¹⁰, forms extensive blooms covering large areas in temperate and subpolar latitudes^{9,11}.

The response of these two species to CO₂-related changes in seawater carbonate chemistry was examined under controlled




Figure 1 Response of organic and inorganic carbon production to CO₂ concentration in laboratory-cultured coccolithophorids. **a**, Particulate organic carbon (POC) production; **b**, calcification; and **c**, the ratio of calcification to POC production (calcite/POC) of the coccolithophorids *Emiliania huxleyi* (circles) and *Gephyrocapsa oceanica* (squares) as a function of CO₂ concentration, [CO₂]. Bars denote ± 1 s.d. ($n = 3$); dotted lines represent linear regressions. Also indicated are corresponding ranges of pH, pCO₂, and [CO₃²⁻]. We note that DIC and total alkalinity differed slightly between experimental sets for the two species; values for pH, pCO₂, and [CO₃²⁻], therefore, are only approximations. Vertical lines indicate pCO₂ values of 280, 365 and 750 p.p.m.v., representing pre-industrial, present day and future concentrations.

laboratory conditions. The carbonate system of the growth medium was manipulated by adding acid or base to cover a range from pre-industrial CO₂ levels (280 p.p.m.v.) to approximately triple pre-industrial values (about 750 p.p.m.v.). Over this range, *E. huxleyi* and *G. oceanica* experience a slight increase in photosynthetic carbon fixation of 8.5% and 18.6%, respectively (Fig. 1a), and a comparatively larger decrease in the rate of calcification of 15.7% and 44.7%, respectively (Fig. 1b). The ratio of calcite to organic matter production (calcite/POC) for the two species decreased by 21.0% and 52.5%, respectively, between 280 and 750 p.p.m.v. (Fig. 1c). Since calcite production has been shown to vary with ambient light conditions¹², we have grown *E. huxleyi* under different light/dark cycles and photon flux densities. A similar decrease in the calcite/POC ratio in response to CO₂-related changes in carbonate chemistry was obtained over a fivefold range in photon flux densities (Fig. 2).

Scanning electron microscopy indicated that malformed coccoliths and incomplete coccospheres increased in relative numbers

with increasing CO₂ concentrations (Fig. 3). Coccolith undercalcification and malformation is a common phenomenon frequently observed both in natural environments and under laboratory conditions¹³. The systematic trend in the relative abundance of malformed coccoliths and coccospheres observed here, however, suggests a direct effect of seawater carbonate chemistry on the regulatory mechanisms controlling coccolith production inside the cell. Based on light microscopic analysis, no consistent trend was obtained in the number of attached or free coccoliths per coccosphere.

Our laboratory results are consistent with CO₂-related responses of natural plankton assemblages collected in the subarctic north Pacific, a region where coccolithophorids are major contributors to primary production¹⁴. After incubation of replicate samples at pCO₂ levels of about 250 p.p.m.v. and about 800 p.p.m.v. for 1.5 to 9 days, the rate of calcification was reduced by 36% to 83% in high-CO₂ relative to low-CO₂ treatments in four independent experiments (Fig. 4). A similar CO₂-dependent response was obtained under




Figure 2 Ratio of calcification to POC production (calcite/POC) of *Emiliania huxleyi* as a function of CO₂ concentration, [CO₂]. Cells were incubated at photon flux densities of 30, 80 and 150 μmol m⁻² s⁻¹ (denoted by circles, squares and triangles and corresponding

solid, dashed, dash-dotted regression lines, respectively). Bars denote ±1 s.d. (n = 3); lines represent linear regressions. Vertical lines indicate pCO₂ values of 280, 365 and 750 p.p.m.v.




Figure 3 Scanning electron microscopy (SEM) photographs of coccolithophorids under different CO₂ concentrations. **a, b, d, e**, *Emiliania huxleyi*; and **c, f**, *Gephyrocapsa oceanica* collected from cultures incubated at [CO₂] ≈ 12 μmol l⁻¹ (**a-c**) and at [CO₂] ≈ 30–33 μmol l⁻¹ (**d-f**), corresponding to pCO₂ levels of about 300 p.p.m.v. and 780–850 p.p.m.v., respectively. Scale bars represent 1 μm. Note the difference in the

coccolith structure (including distinct malformations) and in the degree of calcification of cells grown at normal and elevated CO₂ levels. Pictures are selected from a large set of SEM photographs to depict the general trend in coccolith calcification. As the culture medium was super-saturated with respect to calcium carbonate under all experimental conditions, post-formation calcite dissolution is not expected to have occurred.

reduced light intensities (10% surface irradiance, data not shown). No significant differences were obtained between short and longer-term incubations of the natural assemblages. As short-term incubations are not likely to experience large changes in species composition, the observed response most probably reflects a reduction in carbonate precipitation of the calcifying organisms in the plankton assemblage.

The observed decrease in calcification with increasing p_{CO_2} , if representative of biogenic calcification in the world's ocean, has significant implications for the marine carbon cycle. Owing to its effect on carbonate system equilibria, calcification is a source of CO_2 to the surrounding water¹⁵, whereby the increase in CO_2 concen-

tration, due to calcification is a function of the buffer capacity of sea water. Theoretically, the buffer state of pre-industrial sea water resulted in 0.63 mole CO_2 released per mole CaCO_3 precipitated¹⁶ (assuming temperature $T = 15^\circ\text{C}$, and salinity $S = 35$). Following the predictions of future atmospheric CO_2 rise, this value will increase to 0.79 in 2100 (assuming Intergovernmental Panel on Climate Change (IPCC) scenario IS92a, ref. 3). At constant global ocean calcification this results in an additional source of CO_2 to the atmosphere. In the case of reduced calcification, this positive feedback is reversed. Assuming a pre-industrial pelagic inorganic carbon production of 0.86 Gt C y^{-1} (ref. 17) and a CO_2 -related decrease in planktonic calcification as observed in our laboratory and field experiments (ranging between 16% and 83%), model calculations yield an additional storage capacity of the surface ocean for CO_2 between 6.2 Gt C and 32.3 Gt C for the period of 1950 to 2100.

Our results indicate that the ratio of calcite to organic matter production in cultured coccolithophorids and in oceanic phytoplankton assemblages is highly sensitive to the seawater p_{CO_2} . Although it is presently not clear what the physiological and ecological role of coccolith formation is¹⁸, we propose that CO_2 -dependent changes in calcification may affect cellular processes such as acquisition of inorganic carbon¹⁹ and nutrients²⁰ as well as trophic interactions, and particle sinking rate^{21,22}. These, in turn, may influence the structure and regulation of marine ecosystems in which coccolithophorids are dominant. From a geochemical viewpoint, a decrease in global ocean calcification would enhance CO_2 storage in the upper ocean^{3,15,23}, thus providing a negative feedback for changes in atmospheric p_{CO_2} . Such a feedback should be taken into account when predicting the role of the ocean in mitigating future anthropogenic CO_2 increases or in reconstructing the relation between ocean productivity and glacial–interglacial variations in p_{CO_2} . □

Methods

Laboratory

Monospecific cultures of the coccolithophorids *Emiliania huxleyi* (strain PML B92/11A) and *Gephyrocapsa oceanica* (strain PC7/1) were grown in dilute batch cultures at 15°C in filtered ($0.2 \mu\text{m}$) sea water enriched with nitrate and phosphate to concentrations of 100 and $6.25 \mu\text{mol l}^{-1}$, respectively, and with metals and vitamins according to the *fl2* culture medium (ref. 24). The carbonate system was adjusted through addition of 1 N HCl or 1 N NaOH to the medium. Cells were acclimated to the experimental conditions for 7–9 generations and allowed to grow for about 8 cell divisions during experiments. Cultures were incubated in triplicate at photon flux densities of $150 \mu\text{mol m}^{-2} \text{ s}^{-1}$, light/dark (L/D) cycle = 16/8 h (Fig. 1) and of $150, 80$ and $30 \mu\text{mol m}^{-2} \text{ s}^{-1}$, L/D = 24/0 (Fig. 2). Dissolved inorganic carbon (DIC) was measured coulometrically in duplicate (UIC model 5012)²⁵. Alkalinity was determined in duplicate through potentiometric titration²⁶. pH, CO_2 and CO_3^{2-} concentrations were calculated from alkalinity, DIC and phosphate concentrations ($T = 15^\circ\text{C}$; $S = 31$) using the dissociation constants of ref. 27. Subsamples for total particulate carbon (TPC) and particulate organic carbon (POC), which in L/D = 16/8 were taken at the end of the dark phase, were filtered onto pre-combusted (12 h, 500°C) QM-A filters (pore width is about $0.6 \mu\text{m}$) and stored at -25°C . Before analysis, POC filters were fumed for 2 h with saturated HCl solution in order to remove all inorganic carbon. TPC and POC were subsequently measured on a mass spectrometer (ANCA-SL 20-20 Europa Scientific). Particulate inorganic carbon (PIC) was calculated as the difference between TPC and POC. Cell counts obtained with a Coulter Multisizer at the beginning and the end of incubations were used to calculate specific growth rates. PIC and POC production rates were calculated from cellular inorganic and organic carbon contents and specific growth rates.

Field

Ship-board productivity and calcification experiments were conducted at three stations in the subarctic North Pacific Ocean in June of 1998 (one experiment) and September of 1999 (four experiments). Station locations are given in the legend of Fig. 4. Surface seawater (10–20 m) was collected using a trace-metal-clean *in situ* pumping system and dispensed into acid-soaked polycarbonate bottles (3–4 replicate bottles per treatment). Samples were incubated on deck at about 30% surface irradiance levels in a flow-through incubator at *in situ* temperatures ($13 \pm 1^\circ\text{C}$). CO_2 concentrations in samples were manipulated by either bubbling with commercially prepared CO_2 /air mixtures (Station P26, 1998/1999) or by the addition of high-purity (trace-metal-clean) HCl/NaOH. Low CO_2 samples contained about $10 \mu\text{M CO}_2$ (~ 250 p.p.m.v.) with a pH of about 8.20 while high CO_2 treatments contained approximately $33 \mu\text{M CO}_2$ (~ 800 p.p.m.v.) with a pH of about 7.75. Total alkalinity ($\sim 2180 \mu\text{Eq l}^{-1}$) was unaffected by CO_2 bubbling while HCl and NaOH additions changed alkalinity by -6.4% and $+3.4\%$, respectively. CO_2




Figure 4 Effects of CO_2 manipulations on POC production, calcification and the ratio of calcification to POC production (calcite/POC) in subarctic North Pacific phytoplankton assemblages. **a**, POC production; **b**, calcification; and **c**, the calcite/POC ratio. Station P26 (50°N , 145°W) 1998, 6.8-day CO_2 pre-conditioning (filled circles). Station P26 1999, 2-day CO_2 pre-conditioning (squares). Station P26 1999, 9-day CO_2 pre-conditioning (filled triangles). Station P20 ($43^\circ 30' \text{N}$, $138^\circ 40' \text{W}$) 1999, 1.5-day CO_2 pre-conditioning (open triangles). Station Z9 (55°N , 145°W) 1999, 1.5-day pre-conditioning (open circles). In all five experiments, POC production did not differ significantly between CO_2 treatments (t -test, $p \geq 0.4$). The statistical significance of calcification rate differences (t -test) is as follows: P26, all experiments and measurements ($p < 0.05$); P20, calcification ($p = 0.056$); calcite/POC ($p = 0.074$); Z9, calcification ($p = 0.135$); calcite/POC ($p = 0.11$). Error bars represent standard errors of means.

concentrations in samples were calculated from measurements of pH and alkalinity using the algorithm developed by ref. 28. After a pre-conditioning period ranging from 1.5–9 days, samples were incubated with 40 $\mu\text{Ci } ^{14}\text{C}$ (50 mCi mmol^{-1}) for 6–9 hours, harvested onto 0.4- μm polycarbonate filters, and immediately frozen in scintillation vials at -20°C . In the laboratory, filter samples were acidified with either 10% HCl or H_3PO_4 to measure acid-stable (organic) carbon. The liberated acid-labile (particulate inorganic) carbon was trapped in NaOH solution contained either in small vials suspended in the primary vials or soaked into small GF/D glass fibre filters stuck onto the caps of the primary vials. Samples were measured on a liquid scintillation counter and corrected for ^{14}C uptake in dark control bottles as well as filtered seawater blanks.

Received 2 August 1999; accepted 18 July 2000.

1. Milliman, J. D. Production and accumulation of calcium carbonate in the ocean: budget of a nonsteady state. *Glob. Biogeochem. Cycles* **7**, 927–957 (1993).
2. Holligan, P. M. & Robertson, J. E. Significance of ocean carbonate budgets for the global carbon cycle. *Glob. Change Biol.* **2**, 85–95 (1996).
3. Houghton, J. T. et al. (eds) *Climate Change 1994: Radiative Forcing of Climate Change and an Evaluation of the IS92 Emission Scenario* (Cambridge Univ. Press, Cambridge, 1995).
4. Wolf-Gladrow, D. A., Riebesell, U., Burkhardt, S. & Bijma, J. Direct effects of CO_2 concentration on growth and isotopic composition of marine plankton. *Tellus* **51B**, 461–476 (1999).
5. Gattuso, J.-P., Frankignoulle, M., Bourge, I., Romaine, S. & Buddemeier, R. W. Effect of calcium carbonate saturation of seawater on coral calcification. *Glob. Planet. Change* **18**, 37–46 (1998).
6. Langdon, C., Takahashi, T., Sweeney, C., Chipman, D., Goddard, J., Marubini, F., Aceves, H., Barnett, H. & Atkinson, M. Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef. *Global Biogeochem. Cycles* **14**, 639–654 (2000).
7. Westbroek, P., Young, J. R. & Linschooten, K. Coccolith production (biomineralisation) in the marine alga *Emiliana huxleyi*. *J. Protozool.* **36**, 368–373 (1989).
8. Westbroek, P. et al. A model system approach to biological climate forcing. The example of *Emiliana huxleyi*. *Glob. Planet. Change* **8**, 27–46 (1993).
9. Winter, A., Jordan, R. W. & Roth, P. H. in *Coccolithophores* (eds Winter, A. & Siesser, W. G.) 161–179 (Cambridge Univ. Press, Cambridge, 1994).
10. Westbroek, P. et al. Strategies for the study of climate forcing by calcification. *Bull. Inst. Oceanogr. Monaco* (Spec. Issue) **13**, 37–60 (1994).
11. Holligan, P. M. et al. A biogeochemical study of the coccolithophore, *Emiliana huxleyi*, in the North Atlantic. *Glob. Biogeochem. Cycles* **7**, 879–900 (1993).
12. Nielsen, M. V. Growth, dark respiration and photosynthetic parameters of the coccolithophorid *Emiliana huxleyi* (Prymnesiophyceae) acclimated to different day length-irradiance combinations. *J. Phycol.* **33**, 818–822 (1997).
13. Young, J. R. Variation in *Emiliana huxleyi* coccolith morphology in samples from the Norwegian EHUX Experiment, 1992. *Sarsia* **79**, 417–425 (1994).
14. Booth, B. C., Lewin, J. & Postel, J. R. Temporal variation in the structure of autotrophic and heterotrophic communities in the subarctic Pacific. *Prog. Oceanogr.* **32**, 57–99 (1993).
15. Purdie, D. A. & Finch, M. S. Impact of a coccolithophorid bloom on dissolved carbon dioxide in sea water enclosures in a Norwegian fjord. *Sarsia* **79**, 379–387 (1994).
16. Frankignoulle, M., Canon, C. & Gattuso, J.-P. Marine calcification as a source of carbon dioxide: Positive feedback of increasing atmospheric CO_2 . *Limnol. Oceanogr.* **39**, 458–462 (1994).
17. Morse, J. W. & Mackenzie, F. T. in *Developments in Sedimentology 48: Geochemistry of Sedimentary Carbonates* (Elsevier, Amsterdam, 1990).
18. Young, J. R. in *Coccolithophores*. (eds Winter, A. & Siesser, W. G.) 63–82 (Cambridge Univ. Press, Cambridge, 1994).
19. McConnaughey, T. A. Calcification, photosynthesis, and global cycles. *Bull. Inst. Oceanogr. Monaco* (Spec. Issue) **13**, 137–161 (1994).
20. McConnaughey, T. A. & Whelan, J. F. Calcification generates protons for nutrient and bicarbonate uptake. *Earth Sci. Rev.* **42**, 95–117 (1997).
21. Harris, R. P. Zooplankton grazing on the coccolithophorid *Emiliana huxleyi* and its role in inorganic carbon flux. *Mar. Biol.* **119**, 431–439 (1994).
22. Fritz, J. & Balch, W. M. A light-limited continuous culture study of *Emiliana huxleyi*: determination of coccolith detachment and its relevance to cell sinking. *J. Exp. Mar. Biol. Ecol.* **207**, 127–147 (1996).
23. Ksheshgi, H. S., Flannery, B. P. & Hoffert, M. I. Marine biota effects on the compositional structure of the world oceans. *J. Geophys. Res.* **96**, 4957–4969 (1991).
24. Guillard, R. R. L. & Rytner, J. H. Studies of marine planktonic diatoms. I. *Cyclotella nana* (Hustedt) and *Detonula confervacea* (Cleve). *Can. J. Microbiol.* **8**, 229–239 (1962).
25. Johnson, K. M., Wills, K. D., Butler, D. B., Johnson, W. K. & Wong, C. S. Coulometric total carbon dioxide analysis for marine studies: maximizing the performance of an automated gas extraction system and coulometric detector. *Mar. Chem.* **44**, 167–187 (1993).
26. Bradshaw, A. L., Brewer, P. G., Shafer, D. K. & Williams, R. T. Measurements of total carbon dioxide and alkalinity by potentiometric titration in the GEOSECS program. *Earth Planet. Sci. Lett.* **55**, 99–115 (1981).
27. Goyet, C. & Poisson, A. New determination of carbonic acid dissociation constants in seawater as a function of temperature and salinity. *Deep-Sea Res.* **36**, 1635–1654 (1989).
28. Lewis, E. & Wallace, D. W. R. Program Developed for CO_2 System Calculations. ORNL/CDIAC-105. (Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 1998).

Acknowledgements

We thank A. Dauelsberg, B. Höhnisch, A. Terbrüggen, and K.-U. Richter for laboratory assistance, F. Hinz for REM analyses, D. Crawford, M. Lipsen, F. Whitney and C. Mayfield for invaluable help at sea and C. S. Wong for providing space on RV J.P. Tully. The *E. huxleyi* strain PML B92/11A was generously supplied by J. Green, Plymouth Marine Laboratory, and the *G. oceanica* strain PC 7/1 by the CODENET Algae collection in Caen. This work was supported by the Netherlands-Bremen Cooperation in Oceanography (NEBROC).

Correspondence and requests for materials should be addressed to U. R. (e-mail: uriebesell@awi-bremerhaven.de).

.....
Increased dissolved oxygen in Pacific intermediate waters due to lower rates of carbon oxidation in sediments

Lowell D. Stott, William Berelson, Robert Douglas & Donn Gorsline

Department of Earth Sciences, University of Southern California, Los Angeles, California 90089, USA

.....
Concentrations of dissolved oxygen in the ocean seem to correlate well with climate instabilities over the past 100,000 years. For example, the concentration of dissolved oxygen in Pacific intermediate waters was considerably higher during Pleistocene glacial periods than it is today^{1–4}. This has been inferred from the presence of bioturbated sediments, implying that oxygen levels were sufficient for burrowing organisms to live. Today, basins in the northeastern Pacific Ocean are floored by laminated sediments implying lower oxygen levels, which may be explained by reduced ventilation^{2–4}. Here we report a recent return to bioturbated sediments in the northeastern Pacific Ocean since the late 1970s. From the carbon isotope composition of benthic foraminifers living in the sediment, we infer a twofold decrease in the carbon oxidation rate occurring within sediments, equivalent to an increase in dissolved oxygen concentration of 15–20 micromoles per litre. These changes, at the edges of the Santa Barbara, Santa Monica and Alfonso basins, are coincident with a change in North Pacific climate which has reduced upwelling by 20–30% and increased sea surface temperatures by 1.5–3 °C. This suggests that climate effects on surface productivity, reducing the supply organic matter to sediments, may have had a greater effect on benthic oxygen levels than changes in ocean circulation patterns.

Laminated sediments have been accumulating in the Santa Barbara and Santa Monica basins, and basins within the Gulf of California, for the past few centuries. However, the spatial extent of laminated sediments within these basins has been changing⁵. In Santa Monica basin, a study of Holocene sediment patterns revealed that between AD 1600 and the 1970s, laminated sediments spread systematically outwards and upwards into shallower waters from the centre of the basin, so as to encompass the entire basin floor by the 1970s (ref. 5). During this same period, laminated sediments were also forming in Santa Barbara basin and in the basins along the margins of the Gulf of California^{6–11}. The temporal record of expanding laminated sediments in Santa Monica basin suggests that there are factors operating on centennial timescales that affect bottom-water oxygen levels, which probably control the distribution of bioturbating organisms. It is also now clear that there are shorter-timescale environmental changes that also affect the North Pacific marine environment, and that these are superimposed on the longer-scale patterns of variability. In particular, the mean climate state of the North Pacific, as measured by a number of different variables, has varied between a warm and a cold phase with a quasi-regular decadal oscillation^{12–19}. This decadal pattern has been referred to as the Pacific Decadal Oscillation¹⁹ (PDO). In the ocean, the PDO is characterized by a shift in the bifurcation point of the Sub-arctic Current as it approaches North America. This point of bifurcation strongly influences oceanographic, as well as weather patterns along the coast of the western USA^{15–18}. The last clear PDO shift was coincident with a 1976–77 El Niño event. Since that time, the average sea surface temperatures within the southern California Current during the months of upwelling (spring/summer) have increased by 1.5–3 °C (Fig. 1). Upwelling along the coast at the latitude of Santa Monica basin during the spring has also decreased (Fig. 1). The result of these changes has been a systematic decline in marine fisheries in the southern California Current,