

Macro-epibenthic communities and diversity of Arctic Kongsfjorden, Svalbard, in relation to depth and substrate

Ricardo Sahade^{1*}, Alexia Stellfeldt¹, Marcos Tatián¹ and Jürgen Laudien²

¹ Ecología Marina Universidad de Córdoba, Av. Vélez Sársfield 299. CP 5000. Córdoba, Argentina.

² Alfred Wegener Institute for Polar and Marine Research, 27515 Bremerhaven, Germany.

*E-mail: rsahade@efn.uncor.edu

Introduction

Low and relatively constant temperatures, seasonal pulses of primary production, changes in ice cover, siltation and salinity are the main characteristics of polar marine ecosystems. Shallow coastal systems are especially affected by ice as a disruptive factor, which has a huge effect in structuring the established benthic communities, both in the Antarctic (Dayton 1990, Clarke 1996a, Grebmeier & Barry 1991, Sahade *et al.* 1998, Gutt 2001, Gutt & Piepenburg 2003) and the Arctic (Laudien *et al.*, this issue). Despite these similarities the two polar ecosystems differ considerably in their evolutionary histories. While the Antarctic has been evolving isolated the last 20 million years, the Arctic Holocene history only reaches back some 10 thousand years (Clarke & Crame 1992, Dunton 1992, Dayton *et al.* 1994). These differences are reflected in the recent biota with the Arctic being impoverished in respect to the Antarctic in terms of species richness, diversity and abundance of organisms (Poore & Wilson 1993, Rex *et al.* 1993, Clarke 1996a,b, Arntz *et al.* 1997, Gray 2002). However, at small-scale these differences are not evident and several Arctic benthic communities exhibiting high abundance and diversity have been described (e.g., Grebmeier *et al.* 1989, Kendall 1996, Sejr *et al.* 2000). The major difficulties in comparing both polar ecosystems are due to the use of different methods, depth ranges, scales, and analytical procedures.

Photo-transects allow the quantitative analysis of communities, providing information about the habitat, abundances, percentage cover, species associations of epibenthic assemblages and a fast data acquisition in the field. Although this method underestimates abundances of small, cryptic and highly mobile individuals (Barthel *et al.* 1991, Roberts *et al.* 1994, Jørgensen & Gulliksen 2001), it has been successfully used to define polar benthic assemblages (Dayton *et al.* 1974, Barthel *et al.* 1991, Barthel & Gutt 1992, Jørgensen & Gulliksen 2001, Teixido *et al.* 2002).

The present study is a preliminary analysis of epi-macro-benthic community structures in relation to substrate types and depth in the Arctic Kongsfjorden. The application of the same methods (photo-transects), environmental variables (depth and substrate) and analytical procedures as in Antarctic Potter Cove (Sahade 1999) will allow to further establish valid comparisons between both polar benthic systems.

Material and methods

Study area

The study area was the Arctic glacial Kongsfjorden located on the western coast of Spitsbergen (79°N, 12°E). The fjord is 20km long and 4 to 10km wide and has a maximum depth close to 350m. It is directly connected to the North Atlantic Ocean via the Kongsfjord-Renna trough (Bluhm *et al.* 2001, Jørgensen & Gulliksen 2001, Svendsen *et al.* 2002). For a detailed review of environmental characteristics and a description of the marine ecosystem see Svendsen *et al.* (2002) and Hop *et al.* (2002). Three stations with different substrate types were selected, Prins Heinrichøya (S1), mainly composed of a sand-clay mixture with occasional ice-rafted stones, Hansneset (S2), rocky bottoms interspersed with sediment pools and Kongsfjordneset (S3) pure bedrock (Fig. 1).

Sampling

Sampling was carried out during the boreal summer 2001 by means of photo-transects taken by SCUBA-divers with a Nikonos V camera, a 15-mm lens and a Nikonos SB-104 strobe, mounted on an aluminium frame (50 x 50cm) at particular depth profiles 15, 20, 25 and 30m (Kühne 1992, Sahade *et al.* 1998). Percentage cover of the main taxonomic groups was obtained from the slides taken at each of the three stations. Thereafter data were analysed to assess diversity and community structure. In order to compare diversity patterns for both the different stations and the depth gradient, K-dominance curves were plotted. Multivariate analysis as classification (clustering) and ordination MDS (non-metric multidimensional analysis) followed to evaluate community structures using *PRIMER* v5 (Clarke & Gorley 2001).




Fig. 1: Maps of the study area, sampling stations S1 (mainly soft bottom), S2 (mixed substrates) and S3 (bedrock) are indicated.

Results

A total of 47 species or morphospecies of 18 higher taxa (order to class) were identified. Table 1 list the species and their occurrences at the three stations.

Table 1: List of taxa identified, * indicates the presence of taxa at station S1 (mainly soft bottom), S2 (mixed substrates) and S3 (bedrock).

	Taxa	S1	S2	S3		Taxa	S1	S2	S3	
Phaeophyta	<i>Laminaria</i> sp.	*		*	Polyplacophora	<i>Tonicella</i> sp.	*	*		
	<i>Desmarestia aculeata</i>	*		*		Cirripedia	<i>Balanus balanus</i>	*	*	*
	<i>Desmarestia viridis</i>	*	*	*	Decapoda		<i>Lebbeus polaris</i>	*	*	
	<i>Acrosiphonia</i> aff. <i>penicilliformis</i>	*		*		<i>Eupagurus</i> sp.	*	*	*	
Rhodophyta	<i>Polysiphonia arctica</i>	*		*		Unidentified sp. 1		*		
	<i>Phyllophora</i> sp.	*		*		Unidentified sp. 2	*	*		
Porifera	Unidentified sp. 1	*		*	Pycnogonida	Unidentified sp. 1			*	
	<i>Haliclona</i> sp.	*	*			Bryozoa	<i>Reteporella beaniana</i> ¹	*	*	*
	Unidentified sp. 1		*		<i>Crisia denticulata</i> ¹		*	*	*	
Unidentified sp. 2		*		<i>Porella</i> cf. <i>compressa</i>			*			
Anthozoa	<i>Urticina equees</i>	*	*	*	Unidentified sp. 1				*	
	<i>Hormathia nodosa</i>	*	*		Unidentified sp. 2		*			
	Unidentified sp. 1		*		Asteroidea	<i>Crossaster papposus</i>		*	*	
Hydrozoa	<i>Eudendrium</i> sp.		*			<i>Henricia</i> sp. ¹		*		
	Polychaeta	<i>Thelepus</i> cf. <i>cincinnatus</i>		*			<i>Pteraster</i> sp. ¹		*	
Serpulidae			*		Echinoidea	<i>Strongylocentrotus</i> sp.	*	*	*	
Sabellidae			*			Ophiuroidea	<i>Ophiopholis aculeata</i>	*	*	*
Unidentified sp. 1			*		Ascidiacea		<i>Boltenia echinata</i>		*	
Unidentified sp. 2		*				<i>Styela rustica</i>		*	*	
Gastropoda	<i>Buccinum</i> sp.	*				<i>Halocynthia pyriformis</i>		*	*	
	<i>Neptunea</i> sp.	*				Unidentified sp. 1		*	*	
Bivalvia	<i>Mya truncata</i>	*	*	*	Pisces	Unidentified sp. 1			*	
	<i>Chlamys islandica</i>	*	*	*		Total		47	25	34
	<i>Hiatella arctica</i>	*	*	*						
	Unidentified sp. 1	*		*						

K-dominance curves indicated that diversity was lower at the station mainly composed of soft bottom (S1) than at stations dominated by hard substrate elements (S2 and S3) (Fig. 2). Diversity increased with depths at all three stations. This pattern was more evident at station S1, which showed a constant increase. At station S2 and S3 the differences were clear between 15m and 30m, but species composition was more similar at 20m and 25m (Fig. 2).

Multivariate analysis of classification and ordination (MDS) clearly separated the samples in first term by location, substrate type and later by depth (Figs. 3 and 4). Three communities were separated according to the sampling stations. The two sites with significant hard bottom occurrences (S2 and S3) were further grouped and separated from S1. Not one single sample was included among the samples of another station. Additionally, there was a distinction following the bathymetric gradient at each station.

All stations showed a change from macroalgal dominance in shallow waters to faunal communities in deeper waters. This pattern was especially clear at stations S1 and S2, while at S3 macroalgae were less abundant (even at 15m).




Fig. 2: K-dominance curves showing cumulative ranked abundances plotted against log species rank (following Lambshead *et al.* 1983). a: The most elevated curve (mainly soft bottom S₁) shows a lower diversity than the two lower ones (S₂ and S₃). b-d: At all three stations cline of increasing diversity with depth was observed.

Discussion

The epibenthic community structures analysed in Kongsfjorden are coincident with previous studies of the area (Jørgensen & Gulliksen 2001, Lippert *et al.* 2001, Hop *et al.* 2002). However, communities showed marked differences in relation to substrate type and depth. Higher diversities of organisms were found in habitats mainly composed of hard bottoms (S₂ and S₃), while epifauna and -flora of habitats dominated by soft bottoms were less diverse. This observation reflects that most of the organisms inhabiting hard substrates are epifaunal, while the epifauna of soft bottoms in Kongsfjorden is a rather small proportion of the total soft bottom inhabitants, which are mainly infaunal species. Therefore, the macro-epibenthic communities of soft habitats may be less diverse than the communities of hard substrates, as only a special group is using the surface of the sediment. Correspondingly, Jørgensen and Gulliksen (2001) state that the infauna was underestimated by their photographic method. However, this non-destructive sampling provides reliable information about differences of larger epibenthic taxa between locations and along the bathymetric gradient.

The comparison of the present results with those achieved from a soft-bottom study of macrobenthos, which was located at Brandal, approximately 5km down-fjord (north-west) of our study location (Laudien *et al.*, this issue), revealed significant differences in species composition. The present species list includes rare, large species like the anthozoan *Hormathia nodosa*, the gastropods




Fig. 3: Dendrogram resulting from classification analysis (Bray-Curtis similarities) of macrobenthos from Kongsfjorden based on the UPGMA method. All samples are separated according to the stations (substrate type); the two communities inhabiting habitats dominated by hard bottom (S2 and S3) are more closely related and separated from the community inhabiting soft-bottom dominated habitats (S1).



Fig. 4: MDS plot, circles are the superimposed results of the cluster analysis, ●: station with mainly soft bottom (S1), ■: station with mixed substrate (S2) and ▲: station with bedrock (S3). The shading of the sample symbols indicates depth, following the legend given in Figure 3.

Buccinum sp. and *Neptunea* sp., as well as highly mobile species like the decapods *Lebbeus polaris* and *Eupagurus* sp., which were not reported in the Brandal study. The latter sampled five replicates of a 20 × 20cm corer (surface: 0.2m²) by means of a suction method, while the present study covered a much wider area of 59 photo quadrates of 50 × 50cm (surface: 14.75m²). However, according to species-accumulation curves, the previous study already detected the majority of the total soft bottom fauna. Therefore both studies are contributing to a detailed description of the macrofauna of Kongsfjorden. Moreover, four species found in this study had been reported for Svalbard, but not for Kongsfjorden before (Gulliksen *et al.* 1999, Jørgensen & Gulliksen 2001) (Table 1).

Differences in species diversity might not only be substrate related, but also be correlated with environmental parameters (e.g., salinity, turbidity, temperature) following a gradient along the mayor axis of the fjord (for review: Svendsen *et al.* 2002). This explanation agrees well with the general trend that deposit feeding infauna (not quantitatively detectable with the applied photographic method) increases in dominance and filter feeders decrease in abundance – due to unfavourable conditions – with a decreasing distance from the glacier front (Farrow *et al.* 1983, Syvitsky *et al.* 1989, Holte *et al.* 1996, Wlodarska *et al.* 1996). Low species richness near the glacier fronts has also been related to the scarcity of food available to substrate detritivorous species as a consequence of low levels of primary production and the dilution of organic matter in the substrate by high sedimentation (Görlich *et al.* 1987). The same trend can be seen by comparing the two stations dominated by hard bottom substrates, where the station located down-fjords (S3, pure bedrock) showed a higher diversity than the intermittent station (S2). Under constant environmental conditions the opposite pattern would have been expected as the habitat heterogeneity of station S2 should result in a higher diversity (Gray 2001). Therefore, the environmental parameters could also have a strong effect in determining these communities. Both, this hypothesis and the explanation of substrates playing a major role in determining the benthic community structure will be analyzed in future studies.

Besides substrate and/or the location, depth appears important in structuring epibenthic communities. We detected increasing diversity along the bathymetric gradient at each of the three stations. This trend is most obvious in the station mainly composed of soft bottom showing a constant increase of diversity from 15m to 30m. The communities were characterized by a dominance of macroalgae at 15m and a shift to faunal dominance at 20m, diversity was highest at 30m. As boulders colonised by a typical rocky community (barnacles, actinians, ascidians and sponges) were more frequent at 30m, again the substrate could explain increasing diversity with depths. The previous study conducted at Brandal (Laudien *et al.*, this issue) revealed that species richness was lowest at 5m but highest at an intermittent depth of 10-15m; thereafter species richness steadily decreased (20, 25, 30m). This observation was explained according to the ‘intermediate disturbance hypothesis’ (Connell 1978): locations with minimal

disturbance show reduced species richness because of competitive exclusion between species. With increasing intensity or frequency of iceberg scouring (more icebergs ground in shallower areas) competition is relaxed, reflected in increasing species richness; while at higher or more frequent levels of disturbance species start to be eliminated by stress. These findings appear to be true for homogeneous substrates as sampled in the Brandal study. However, the results of the present study indicate that such a trend may be masked by substrate heterogeneity, especially when the density of hard bottom elements increase with depths.

The up-fjord bedrock station (S3) showed considerably lower abundances of macroalgae than the other stations. This observation could be explained with high densities of the green sea urchin *Strongylocentrotus droebachiensis* foraging on macroalgae (H. Wessels, Univ. Bremen, unpubl. data). In the year previous to this study abundances of 80 ind. m⁻² were observed at the same location (F. Beuchel and B. Gulliksen, Univ. Tromsø pers. comm. in Hop *et al.* 2002). Grazed areas are commonly populated by sessile organisms such as actinians, barnacles and bryozoans. At deeper waters from 20m to 30m the increase in diversity is less marked than at the other stations and the community is dominated by suspension feeders (barnacles, actinians, bryozoans and ascidians). These organisms are favoured by rich Atlantic waters entering into the fjord. However, in the central and inner part of Kongsfjorden filter feeders might be under permanent stress, as the high inorganic load (800 g m⁻² d⁻¹ at the glacier front, Svendsen *et al.* 2002) is likely to clog their filter organs (e.g., Moore 1977).

The comparison with Potter Cove (King George Island, Antarctica), although still preliminary, appears to be coincident with the general trend of Antarctic communities being more diverse than the Arctic ones. There are striking differences: In the Potter Cove the highest diversities were found on soft bottoms (Sahade 1999) with pennatulaceans, bivalves and ascidians as dominant organisms, while in Kongsfjorden hard bottom habitats were more diverse. Besides that, soft bottom communities at Potter Cove were dominated by suspension feeders, especially ascidians and sponges, which are characteristic for many Antarctic faunal communities, but rarely seen in other areas. Diversity patterns along the major axis of both fjords are also different. While diversity is lower at the outer station of Potter Cove (hard bottom) compared with inner stations, the opposite is the case for Kongsfjorden. Future work will focus on further comparisons of both polar systems.

Conclusions

Photo-transects (15, 20, 25 and 30m) of macro-epibenthic invertebrate communities and algae from Arctic glacial Kongsfjorden (Spitsbergen) revealed a species list comprising 47 taxa. Mean species richness (17-20 species) was similar at all three stations. K-dominance curves showed higher diversity in stations

dominated by hard substrates than in habitats mainly composed of soft sediment. A bathymetric pattern was evident showing increasing diversity with depth, which is apparently related to increasing densities in hard bottom elements. The comparison with studies conducted in the Antarctic fjordic system Potter Cove, revealed major differences between the two polar systems.

Acknowledgements We are especially grateful to our dive companions, Hendrik Wessels, Jens Meyer, Sabine Bauer, Heike Lippert and to our fellows at the Koldewey-Station. Our thanks are due to Hendrik Wessels and Prof. Dr. Christian Wiencke who helped with the identification of macro-algae. Dr. Eike Rachor and Prof. Dr. Christian Wiencke gave valuable comments on an earlier draft of the manuscript. AWI provided working facilities at the Koldewey-Station. Financial support was supplied by AWI, DAAD and SECyT-IB.

References

- Arntz WE, Gutt J, Klages M (1997) Antarctic marine biodiversity: an overview. In: B Battaglia, J Valencia, D Walton (eds) Antarctic communities. Proc 6th SCAR Biology Symposium, Venice 1994 Cambridge Univ Pr, Cambridge pp 3-14
- Barthel D, Gutt J (1992) Sponge associations in the eastern Weddell Sea. *Antarct Sci* 4:137-150
- Barthel D, Gutt J, Tendal OS (1991) New information on the biology of Antarctic deep-water sponges derived from underwater photography. *Mar Ecol Prog Ser* 69:303-307
- Bluhm B, Iken K, Laudien J, Lippert H (2001) German activity in cold water scientific diving, S.C. Jewett (ed.). *Cold Water Diving for Science. Proceedings of the 21st Annual Scientific Diving Symposium, American Academy of Underwater Sciences. University of Alaska Sea Grant, AK-SG-01-06, Fairbanks* pp 1-4
- Clarke A (1996a) Marine benthic populations in Antarctica: Patterns and Processes. In: *Foundations for Ecological Research West of the Antarctic Peninsula*. Antarctic Research Series Ross R. M., Hofmann E. E., Quetin L. B. (eds). 70:373-388
- Clarke A (1996b) The distribution of Antarctic marine communities. In: *Foundations for Ecological Research West of the Antarctic Peninsula*. Antarctic Research Series Ross R. M., Hofmann E. E., Quetin L. B. (eds.) 70:219-230
- Clarke A, Crame JA (1992) The Southern Ocean benthic fauna and climate change: a historical perspective. *Phil. Trans. R. Soc. Lond. B* 338:299-309
- Clarke KR, Gorley RN (2001) *PRIMER v5: User Manual/Tutorial*, PRIMER-E: Plymouth.
- Connell JH (1978) Diversity in tropical rain forests and coral reefs. *Science* 199:1303-1310
- Dayton PK (1990) Polar benthos. In: WE Smith (ed) *Polar Oceanography, part B: Chemistry, Biology and Geology*. Academic Press, London pp 631-685
- Dayton PK, Robilliard GA, Paine RT, Dayton LB (1974) Biological accommodation in the benthic community at McMurdo Sound, Antarctica. *Ecol Monogr* 44:105-128
- Dayton PK, Mordida BJ, Bacon F (1994) Polar marine communities. *Amer zool* 34:90-99
- Dunton K (1992) Arctic biogeography: The paradox of the marine benthic fauna and flora. *Trends Ecol Evol* 7(6):183-189
- Farrow GE, Syvitsky JPM, Tunnicliffe V (1983) Suspended particulate loading on the macrobenthos in a highly turbid fjord: Knight Inlet British Columbia. *Can J Fish Aquat Sci* 40 (Suppl 1):273-288
- Görlich K, Weslawski JM, Zajaczkowski M (1987) Suspension settling effect on macrobenthos biomass distribution in the Hornsund fjord Spitsbergen. *Polar Res* 5:175-192
- Gray JS (2001) Marine diversity: the paradigms in patterns of species richness examined. *Sci Mar* 65(2):41-56
- Gray JS (2002) Species richness of marine soft sediments. *Mar Ecol Prog Ser* 244:285-297

- Grebmeier JM, Barry JP (1991) The influence of oceanographic processes on pelagic-benthic coupling in polar regions: A benthic perspective. *J Mar Syst* 2(3-4):495-518
- Grebmeier JM, Feder HM, McRoy CP (1989) Pelagic-benthic coupling on the shelf of the northern Bering and Chukchi Seas. 2. Benthic community structure. *Mar Ecol Prog Ser* 51(3):253-268
- Gulliksen B, Palerud R, Brattegard T, Sneli J-A (eds) (1999) Distribution of marine benthic macroorganisms at Svalbard (including Bear Island) and Jan Mayen. Research Report for DN 1999-4. Directorate for Nature Management, Trondheim pp 1-148
- Gutt J (2001) On the direct impact of ice on marine benthic communities, a review. *Polar Biology* 24:553-564
- Holte B, Dahle S, Gulliksen B, Naes K (1996) Some macrofaunal effects of local pollution and glacier-induced sedimentation with indicative chemical analyses in the sediments of two Arctic fjords. *Polar Biol* 16:549-557
- Hop H, Pearson T, Hegseth EN, Kovacs KM, Wiencke C, Kwasniewski S, Eiane K, Mehlum F, Gulliksen B, Wlodarska-Kowalczyk M, Lydersen C, Weslawski JM, Cochrane S, Gabrielsen GW, Leakey RJG, Lønne OJ, Zajaczkowski M, Falk-Petersen S, Kendall M, Wängberg S-A, Bischof K, Voronkov AY, Kovaltchouk NA, Wiktor J, Poltermann M, di Prisco G, Papucci C, Gerland S (2002) The marine ecosystem of Kongsfjorden, Svalbard. *Polar Res* 21(1):167-208
- Jørgensen LL, Gulliksen B (2001) Rocky bottom fauna in arctic Kongsfjord (Svalbard) studied by means of suction sampling and photography. *Polar Biol* 24:113-121
- Kendall MA (1996) Are Arctic soft-sediment macrobenthic communities impoverished? *Polar Biol* 16(6):393-399
- Kühne S (1992) Die Fauna des Steingrundes in der Deutschen Bucht. Diploma Thesis, University of Bonn pp 1-128
- Lambshhead PJD, Platt HM, Shaw KM (1983) The detection of differences among assemblages of marine benthic species based on an assessment of dominance and diversity. *J nat Hist* 17:859-874
- Lippert H, Iken K, Rachor E, Wiencke C (2001) Macrofauna associated with macroalgae in the Kongsfjord (Spitsbergen), *Polar Biol* 24:512-522
- Moore PG (1977) Inorganic particulate suspensions in the sea and their effects on marine animals. *Oceanogr Mar Biol Annu Rev* 15:225-363
- Poore GCB, Wilson GDF (1993) Marine species richness. *Nature* 361:597-598
- Rex MA, Stuart CT, Hessler RR, Allen JA, Sanders HL, Wilson GDF (1993) Global-scale latitudinal patterns of species diversity in the deep-sea benthos. *Nature* 365:636-639
- Roberts DE, Fitzhenry SR, Kennelly SJ (1994) Quantifying subtidal macrobenthic assemblages on hard substrata using a jump camera method *J Exp Mar Biol Ecol* 177:157-170
- Sahade R, Tatián M, Kowalke J, Kühne S, Esnal GB (1998) Benthic faunal associations on soft substrates at Potter Cove, King George Island, Antarctica. *Polar Biol* 19:85-91
- Sahade R (1999) Patterns and Processes in an Antarctic Epi-Benthic Community: the Case of Potter Cove. PhD Thesis, Universidad Nacional de Córdoba
- Sejr MK, Jensen KT, Rysgaard S (2000) Macrozoobenthos in a Northeast Greenland Fjord: Structure and Diversity. *Polar Biol* 23:792-801
- Svendsen H, Beszczynska-Møller A, Hagen JO, Lefauconnier B, Tverberg V, Gerland S, Ørbæk JB, Bischof K, Papucci C, Zajaczkowski M, Azzolini R, Bruland O, Wiencke C, Winther J-G, Dallmann W (2002) The physical environment of Kongsfjorden-Krossfjorden, an Arctic fjord system in Svalbard. *Polar Res* 21(1):133-166
- Syvitski JPM, Farrow GE, Atkinson RJA, Moore PG, Andrews JT (1989) Baffin Island fjord macrobenthos: bottom communities and environmental significance. *Arctic* 42:232-247
- Teixido N, Garrabou J, Arntz WE (2002) Spatial pattern quantification of Antarctic benthic communities using landscape indices. *Mar Ecol Prog Ser* 242:1-14
- Wlodarska M, Weslawski JM, Gromisz S (1996) A comparison of the macrofaunal community structure and diversity in two arctic glacial bays – a 'cold' one off Franz Josef Land and a 'warm' one off Spitsbergen. *Oceanologia* 38:251-283