


Our reference: JEMBE 49563

P-authorquery-v8

**AUTHOR QUERY FORM**

 ELSEVIER	<b>Journal: JEMBE</b>  <b>Article Number: 49563</b>	<b>Please e-mail or fax your responses and any corrections to:</b> <b>E-mail: <a href="mailto:corrections.esil@elsevier.spitech.com">corrections.esil@elsevier.spitech.com</a></b> <b>Fax: +1 619 699 6721</b>
---	---	--

Dear Author,

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.


<b>Location in article</b>	<b>Query / Remark: <a href="#">click on the Q link to go</a> Please insert your reply or correction at the corresponding line in the proof</b>
Q1	Please confirm that given names and surnames have been identified correctly.
Q2	Highlights should consist of only 85 characters per bullet point, including spaces. However, the highlights provided for this item exceed the maximum requirement; thus, they were not captured. Kindly provide the necessary corrections. For more information, please see <a href="#">Guide for Authors</a> .
Q3	Please confirm the changes made in the spelling of " <i>Laniche conchilega</i> ".
Q4	Uncited references: This section comprises references that occur in the reference list but not in the body of the text. Please position each reference in the text or, alternatively, delete it. Any reference not dealt with will be retained in this section. Thank you.

Thank you for your assistance.



Contents lists available at SciVerse ScienceDirect

## Journal of Experimental Marine Biology and Ecology

journal homepage: [www.elsevier.com/locate/jembe](http://www.elsevier.com/locate/jembe)

# Oxygen and capacity limited thermal tolerance of the lugworm *Arenicola marina*: A seasonal comparison

Mareike Schröer, Julia Saphörster, Christian Bock\*, Hans-O. Pörtner

Integrative Ecophysiology, Alfred Wegener Institute for Polar and Marine Research, Am Handelshafen 12, D-27570 Bremerhaven, Germany

## ARTICLE INFO

## Article history:

Received 31 March 2011

Received in revised form 5 August 2011

Accepted 18 September 2011

Available online xxxx

## Keywords:

<sup>13</sup>C NMR spectroscopy*Arenicola marina*

Exercise performance

Growth rate

Protein synthesis

Seasonal acclimatisation

## ABSTRACT

Lugworms *Arenicola marina* were collected from Arcachon Bay in two summers and winters of consecutive years. The worms were acclimated to different temperatures (5 and 10 °C for winter animals and 15 °C for summer animals). Each group was investigated over an experimental temperature range concerning its optimum in exercise performance, acute growth rate as well as respiration and ventilation activities to reveal seasonal acclimatisation effects, potential inter-annual differences and the influence of laboratory acclimation temperatures on the parameters of interest. The groups investigated at the two consecutive summers yielded nearly identical results for ventilation and respiration activities. A clear seasonal difference developed in exercise performance, with an optimum at lower temperatures in winter than in summer, irrespective of acclimation temperature. Respiration and ventilation activities showed no significant differences between winter specimens acclimated to 10 °C and summer specimens acclimated to 15 °C. However, an acclimation temperature of 5 °C for winter animals caused noticeable differences to those acclimated at 10 °C. Acute growth rates differed seasonally as well as between acclimation temperatures with the highest rates found around 10 °C in summer and around 15 °C in winter. The lowest rates were recorded in winter worms acclimated to 5 °C. These acute patterns may reflect high thermal limits in warm acclimated winter worms and temperature dependent shifts in energy demand in summer animals.

© 2011 Published by Elsevier B.V.

## 1. Introduction

*Arenicola marina* is one of the most eminent secondary producers in the intertidal habitat. It is important for other epibenthic and infaunal animals aerating the habitat by sediment dwelling. The fresh weight of individuals may rise up to more than 30 g. Field studies in northern France showed that up to 346.6 kg lugworm faeces were produced per m<sup>2</sup> and year, which corresponds to a sediment layer of 21.5 cm height (Pollack, 1979). *A. marina* feeds on ciliates, microalgae and bacteria in the sediment and the overlying water. Bacterial biomass and chlorophyll a concentration in the sediment show a clear seasonal variability with a maximum occurring between April and October. In contrast, coastal surface waters show the highest bacterial biomass and concentrations of chlorophyll a in March to July (Hubas et al., 2007). Ciliates and other mesosammon-organisms also show the highest density in summer (Pollack, 1979). The main food uptake by *A. marina* occurs during high tide, when the sediment is covered by surface water. *A. marina* not only exploits the nutrients enclosed in the upper sand layer by consuming the sand caving in from the surface like a funnel, but also extracts those from the surface water (Pollack, 1979) by generating a headward directed water current through its

burrow (Wells, 1945). Suspended substances and planctonic organisms are trapped in the sand of the burrow headshaft, which acts as a filter (Krüger, 1957). The worm emits mucus to fill the interstices so that small colloidal particles are also retained. This way, the food region is enriched of organic material, which the lugworm ingests together with the sand.

*A. marina* is very abundant in Arcachon Bay. It is found at any beach (Boisseau, 1962) in densities up to 20 or sometimes 30 to 40 individuals per m<sup>2</sup>, mostly in the intertidal zone and around the island Ile aux Oiseaux even in direct neighbourhood of oyster and eelgrass beds (Amoureux, 1966). The studied population is located at La Hume, a sheltered beach at the southern coast of Arcachon Bay. The surface inhabited by lugworms is a band of approximately 300 m width, bordered by a *Zostera nana* bed at the lower margin and a salt marsh vegetation at the upper margin (Cazaux, 1966). The population consists of at least three generations. The youngest generation is found from the end of March onward in the highest zone of the intertidal, close to the sandy beach. The older individuals with a body weight of up to 6 g inhabit the lower intertidal (Cazaux, 1966). At La Hume lugworms start the production of gametes around the end of April, spawning occurs between the end of August and the beginning of October. After reproduction, the oldest generation disappears and the younger ones resume growing until spawning of the next year (Cazaux, 1966).

\* Corresponding author. Tel.: +49 471 4831 1288; fax: +49 471 4831 1149.  
E-mail address: [Christian.Bock@awi.de](mailto:Christian.Bock@awi.de) (C. Bock).

This strong seasonal pattern mirrors the temperature changes between seasons which are accompanied by a shift in the widths and positioning of thermal tolerance windows on a temperature scale. Seasonal changes in *A. marina* thermal tolerance have already been investigated in a North Sea population (Sommer, 2001; Sommer and Pörtner, 2004; Sommer et al., 1997; Wittmann, 2005; Wittmann et al., 2008). According to the concept of oxygen and capacity limited thermal tolerance, critical temperatures represent the threshold beyond which anaerobic metabolism is necessary for survival because the oxygen demand can no longer be met by aerobic processes (Frederich and Pörtner, 2000; Pörtner, 2001). A more or less parallel shift of both low and high critical temperature values was found with seasonal acclimatisation, characterised by anaerobic end product accumulation (Sommer et al., 1997). Investigations at the mitochondria level revealed that seasonal cold acclimatisation in North Sea lugworms involved a drop in mitochondrial density below summer values, combined with an increasing efficiency of aerobic energy production of each individual mitochondrion (Sommer, 2001; Sommer and Pörtner, 2004). Wittmann et al. (2008) compared thermal tolerance windows of North Sea lugworms during winter to those during summer and demonstrated a widening of the window, accompanied by a shift of critical temperatures towards higher values.

The present study was designed to assess the physiological responses accompanying seasonal acclimatisation and the biochemical changes reported above. Effects of temperature acclimation and inter-annual variation were investigated in a Southern population of the lugworm. Methods were chosen to investigate exercise performance capacity, metabolic energy demand and supply as well as somatic growth. Recordings of digging periods have already been established as a measure for muscular performance and were successfully applied to show differences in performance levels of lugworm populations from various latitudes (Schröer et al., 2009). The water volume which the lugworm pumps through its burrow gives some information about oxygen demand, as oxygen is extracted by gills and body surface from the bypassing water. Pumping frequency is changed to enhance or reduce the water volume flow through the burrow in order to adjust oxygen availability to the respective demand. The experimental setup simultaneously recorded water volume flow and oxygen content and also provided data for pumping frequency and oxygen extraction efficiency determination. Pumping frequency has already been used as a measure for performance depending on salinity (Shumway and Davenport, 1977) and temperature (Schröer et al., 2009; Wittmann, 2005; Wittmann et al., 2008). The mechanical aspects of the lugworm pump (Riisgård et al., 1996) and the biogeochemical consequences for the vented burrow fluids (Davey et al., 1990) have also been of interest. For the determination of acute growth optima, the incorporation of  $^{13}\text{C}$  labelled phenylalanine was tracked by  $^{13}\text{C}$  NMR spectroscopy as introduced by Wittmann et al. (2008).

## 2. Materials and methods

### 2.1. Animals

Specimens of the polychaete *A. marina* (L.) were collected in the intertidal zone at the sampling site in La Hume (44.65° N, 1.17° W) near Arcachon at the French Atlantic coast. Lugworms collected in August 2005 were used for investigations of respiration, ventilation and protein syntheses. For the same experiments during winter, animals were collected in January/February 2006. For yearly comparison the collection of animals was repeated in August 2006 and reinvestigated for potential annual changes. Digging performance was studied on animals from August 2006 and February 2007, for summer and winter, respectively. All worms were maintained in basins filled with natural sediment in a natural seawater flow-through

aquarium system at the Alfred Wegener Institute until experimental use. The specimens collected in winter were divided between incubation temperatures of 5 °C and 10 °C, those collected in summer were kept at 15 °C. All animals were exposed to a salinity of 32‰ and a 12 h/12 h light/dark cycle in the aquaria and fed with ground and soaked Tetramin® flakes every other week.

### 2.2. Field measurements

In parallel to animal collection, biotic and abiotic field parameters were recorded at the sampling site as described by Schröer et al. (2009). In particular, temperatures in air, tidal puddles and sediment were recorded with a thermometer (Testo 925, Testo, Lenzkirch, Germany) using a special temperature-receiving element (6 mm diameter, 500 mm length, TC Direct, Mönchengladbach, Germany). Salinity was measured in tidal puddles by use of a multiple parameter pocket measurement device (Multi 340i, WTW, Weilheim, Germany). The length of the tail shaft of the worm's burrow was taken as a measure of burrow depth by using a scaled metal stick inserted into the opening of the burrow. For abundance recordings, a 1 m × 1 m wooden frame was placed onto the intertidal sediment at haphazard and the number of faecal piles therein was counted. Bodyweight was measured using a scale (EMB 220-I, Kern, Balingen-Frommern, Deutschland).

### 2.3. Digging performance

Experimental temperatures were chosen at 4 °C steps, between –1 and 19 °C for the winter worms. Specimens sampled in summer were measured at 7 to 27 °C. For initial short-term acclimation animals were transferred to a plastic container placed into a temperature controlled aerated seawater bath. Temperature was changed at 2 °C h<sup>-1</sup> starting from maintenance conditions and kept constant for at least 12 h at the new experimental temperature. After acclimation for at least 12 h, animals were transferred into the experimental setup 1 h prior to measurements.

The experimental setup and procedure were the same as described by Schröer et al. (2009). Briefly, the animals were positioned on the sediment surface and the duration of each digging period was recorded using a stopwatch. For an analysis of burrowing capacity during a limited time window this routine was repeated for 90 min and the number of digging periods was recorded. In total, five specimens of each group were examined at each temperature (n=5). Summer (15 °C acclimated) data were published in our previous work recently (Schröer et al., 2009).

### 2.4. Respiration and ventilation experiment

Analyses of respiration and ventilatory activity were carried out as described previously (Wittmann et al., 2008). Briefly, measurements were performed in the dark using artificial burrows consisting of straight Perspex tubes with a rough inner surface. As in their natural burrows, animals generated a water current to provide themselves with oxygen. Air saturation of incurrent and excurrent water was monitored continuously with oxygen micro-optodes (PreSens, Regensburg, Germany). The volume flow produced by the worms was measured using an electromagnetic flowmeter (inner diameter of probe head: 3 mm, RT-500, Hugo Sachs Elektronik, March-Hugstetten, Germany).

Experiments started at a temperature of 4.5 °C for winter animals kept at 5 °C and at 10.7 °C for those kept at 10 °C. Summer animals were tested beginning at 15 °C. Temperature was changed at a rate of 1 °C h<sup>-1</sup> (first lowered) by steps of 3 °C and kept constant for 6 h. Lugworms collected in winter were exposed to a temperature range from –0.0 to 22.8 °C, summer worms experienced a range from 2.8 to 26.1 °C. Mean oxygen partial pressure of incurrent and excurrent water ( $P_{\text{IO}_2}$  and  $P_{\text{EO}_2}$ , kPa) and weight specific volume flow ( $V_w$ , ml h<sup>-1</sup> g<sup>-1</sup>) were calculated for the last 3 h of each incubation

209 period. From these data oxygen consumption ( $M_{O_2}$ ,  $\mu\text{mol O}_2 \text{ h}^{-1} \text{ g}^{-1}$ )  
 210 and the extraction coefficient (%) were determined. The analysis  
 211 also included the mean volume flow during the phases of active ven-  
 212 tilation (active volume flow,  $\text{ml min}^{-1}$ ) and the number of contrac-  
 213 tion waves of the body wall (pumping frequency,  $\text{min}^{-1}$ ). The water  
 214 volume transported per peristaltic wave of the body wall musculature  
 215 (wave volume, ml, see Wittmann et al., 2008) was calculated from  
 216 volume flow recordings during active ventilation periods. Results  
 217 from summer 2006 animals were published before (Schröer et al.,  
 218 2009).

## 219 2.5. Protein biosynthesis

220 Analyses of protein biosynthesis were carried out as described  
 221 previously (Wittmann et al., 2008) and following the principles out-  
 222 lined by Langenbuch et al. (2006). Briefly, uniformly labelled  $^{13}\text{C}$ -L-  
 223 phenylalanine, dissolved in filtered seawater ( $75 \text{ mmol l}^{-1}$ ), was  
 224 injected at  $40 \mu\text{l g}^{-1}$  body weight into the coelomic cavity of the  
 225 lugworms. The animals were then inserted into their artificial bur-  
 226 rows and incubated at temperatures of  $-0.9$ ,  $3.5$ ,  $6.9$ ,  $11$ ,  $15.2$  and  
 227  $19.2$  °C for winter animals and  $7.4$ ,  $10.2$ ,  $14.3$ ,  $17.8$  and  $22.8$  °C for  
 228 summer animals while they were actively ventilating their bur-  
 229 rows. For animals collected in summer the experiment was carried  
 230 out with 8 animals at each temperature. Animals were exposed to  
 231 incubation temperatures for 24 h, then injected and incubated for  
 232 another 30, 60, 90, 120, 180, 240, 300 and 360 min, respectively.  
 233 For winter animals, 12 specimens were used at each temperature  
 234 and 2 of them pooled after 30, 60, 120, 180, 240 and 360 min,  
 235 respectively.

236 After the respective incubation time, the cuticulo-muscular tube  
 237 was frozen in liquid nitrogen. The frozen tissue was ground under  
 238 liquid nitrogen and extracted with TCA (trichloroacetic acid). The  
 239 homogenate was centrifuged and supernatant and pellet were treated  
 240 differently. The supernatant representing the cytosolic fraction  
 241 and containing low molecular weight constituents was neutralised,  
 242 dried and dissolved in  $\text{D}_2\text{O}$  (deuterium oxide).

243 The pellet was washed, suspended in distilled water and neutral-  
 244 ised. After centrifugation, the supernatant containing water-soluble  
 245 proteins was removed and stored while the pellet was boiled in a  
 246 water bath with 1 M NaOH. The water-insoluble proteins were then  
 247 added to the water-soluble protein fraction and both were dried.  
 248 The total protein was dissolved in  $\text{D}_2\text{O}$ .

249 Both cytosolic and protein extracts were measured in a NMR Spec-  
 250 trometer (9.4 T Avance, Bruker, Rheinstetten, Germany) at frequen-  
 251 cies of 100.6 MHz for  $^{13}\text{C}$  spectra. The same parameters of the NMR  
 252 recordings were used as described in Wittmann et al. (2008).

253 The protein contents of the protein extracts were quantified  
 254 according to Bradford (1976). The amount of incorporated  $^{13}\text{C}$ -L-  
 255 phenylalanine was obtained as described by Wittmann et al. (2008).  
 256 The calculated amount of incorporated  $^{13}\text{C}$ -L-phenylalanine was related  
 257 to the respective protein content of the sample and plotted against  
 258 incubation temperature for incubation time. Due to the insensitiv-  
 259 ity of the NMR technique and the fact that only one sample per indivi-  
 260 duum could be analysed we could not track the time-dependent  
 261 increase of incorporated  $^{13}\text{C}$ -L-phenylalanine in sufficiently high resolu-  
 262 tion. Inter-individual differences exceeded those determined over  
 263 time, which is why data points were pooled into two time frames:  
 264 short incubation period (30 to 120 min) and long incubation period  
 265 (180 to 360 min) (see also Wittmann et al., 2008).

## 266 2.6. Statistics

267 Statistical analysis was performed using GraphPad Prism ver-  
 268 sion 4.0c for Macintosh (GraphPad Software, San Diego, California,  
 269 USA). Nonlinear regression curves were fitted to temperature depen-  
 270 dent burrowing activities, using the equation  $EP(T) = F_1(T) + F_2(T) =$

$(A_1 e^{B_1 T} + C_1) + (A_2 e^{B_2 T} + C_2)$  according to Pörtner and Knust (2007).  
 271 For weight specific volume flow data the equation  $V_w(T) = Ae^{BT} + C$   
 272 was used as described in Wittmann et al. (2008). Significant devia-  
 273 tions of  $V_w$  values from the hypothetical value extrapolated from  
 274 the exponential relationship were identified by one-sample Student's  
 275 t-test. One-way ANOVA was performed to detect significant changes  
 276 in pumping frequency, extraction coefficient and protein synthesis  
 277 values over the temperature range within each group (F-test). Two-  
 278 way ANOVA combined with a Bonferroni posthoc test was applied  
 279 to detect significant differences in pumping frequency between the  
 280 groups at each temperature.  $Q_{10}$  values were calculated for specific  
 281 temperature ranges using standard procedures as in Sommer and  
 282 Pörtner (2002). One-way ANOVA and Tukey's posthoc test were  
 283 used to identify differences in field data between the groups. Statis-  
 284 tical significance was identified at the  $p \leq 0.05$  level. All data are  
 285 given as means  $\pm$  SE if not stated otherwise.  
 286

## 287 3. Results

### 288 3.1. Field measurements

289 Biotic and abiotic field parameters (Table 1) were recorded to  
 290 characterise the natural habitat conditions of the population at the  
 291 time of collection during the respective season. Salinity was signifi-  
 292 cantly ( $p < 0.001$ ) lower in winter than in summer, whereas vari-  
 293 ability between the two consecutive years was small. Temperatures  
 294 in air, tidal puddles and sediment (20 cm depth) differed significantly  
 295 ( $p < 0.001$ ) between winter and summer. Differences between the two  
 296 winters or the two summers, respectively, were much smaller, although  
 297 some significant deviations could still be found (air temperature in  
 298 winter 2006 vs. winter 2007:  $p < 0.001$ ; temperature in tidal puddles  
 299 summer 2005 vs. summer 2006:  $p < 0.001$ ; sediment temperature in  
 300 20 cm depth:  $p < 0.001$  between all groups), as data were only avail-  
 301 able for the week of animal collection. Burrow depth (length of the  
 302 tail shaft) did not show any seasonal differences. Abundance also  
 303 did not vary between winter and summer. A seasonal variation in  
 304 animal weight could be detected; specimens collected in summer  
 305 2006 were significantly heavier than animals collected in winter  
 306 (vs. winter 2007:  $p < 0.001$ ; vs. winter 2006:  $p < 0.05$ ). The value for  
 307 summer 2005 is not representative, as juveniles were comprised in  
 308 the measurement.

### 309 3.2. Digging performance


310 Fig. 1 shows the temperature dependent number of digging periods  
 311 performed during 90 minutes experimental intervals. Lugworms col-  
 312 lected in winter and acclimated to 5 °C showed up to  $8.60 \pm 0.24$  dig-  
 313 ging periods  $90 \text{ min}^{-1}$  at 15.4 °C. Digging performance decreased  
 314 rapidly towards higher temperatures and more slowly towards lower  
 315 temperatures. At  $-0.8$ ,  $3.0$  and  $7.0$  °C, digging activity stayed on the  
 316 same level of about 4 periods  $90 \text{ min}^{-1}$ . The regression analysis of  
 317 performance exhibited a maximum of 7.35 digging periods  $90 \text{ min}^{-1}$   
 318 at 14.2 °C.

319 Winter animals acclimated to 10 °C showed similar digging per-  
 320 formances. They displayed the highest number of digging periods  
 321 ( $7.80 \pm 0.37$   $90 \text{ min}^{-1}$ ) at 15.3 °C. As seen in 5 °C acclimated worms,  
 322 digging performance decreased rapidly towards higher temperatures  
 323 and more slowly towards lower temperatures. In contrast to the lat-  
 324 ter, values at  $-1.1$ ,  $3.0$  and  $6.9$  °C did not stay at the same level but  
 325 increased steadily. The performance model indicated a maximum of  
 326 7.48 periods  $90 \text{ min}^{-1}$  at 13.3 °C.

327 Lugworms collected in summer and acclimated to 15 °C per-  
 328 formed the highest number of digging periods ( $5.80 \pm 0.80$   $90 \text{ min}^{-1}$ )  
 329 at 23.2 °C, while the model predicted a maximum of 5.26 periods  
 330  $90 \text{ min}^{-1}$  at 19.6 °C. As observed in winter worms, performance ca-  
 331 pacity decreased more rapidly towards higher than towards lower

**Table 1**  
Abiotic and biotic field parameters at the time of collection of *Arenicola marina* specimens at the sampling site in La Hume near Arcachon (Atlantic, France), mean values  $\pm$  SD.

Sampling month	Salinity (‰)	T (°C) air	T (°C) tidal puddles	T (°C) in 20 cm depth	Length of tail shaft (cm)	Abundance ( $m^{-2}$ )	Weight (g)
August 2005	35.8 $\pm$ 1.2 n = 17	20.2 $\pm$ 2.4 n = 13	20.7 $\pm$ 2.4 n = 13	20.3 $\pm$ 0.6 n = 13	8.0 $\pm$ 2.4 n = 6	22.7 $\pm$ 5.0 n = 6	4.1 $\pm$ 1.4 n = 29
January 2006	26.7 $\pm$ 0.5 n = 25	7.6 $\pm$ 2.4 n = 23	9.5 $\pm$ 2.3 n = 23	7.1 $\pm$ 0.8 n = 23	13.0 $\pm$ 2.4 n = 10	23.0 $\pm$ 9.1 n = 20	4.7 $\pm$ 1.6 n = 72
August 2006	34.1 $\pm$ 0.6 n = 20	22.1 $\pm$ 2.4 n = 20	25.8 $\pm$ 4.1 n = 20	22.2 $\pm$ 0.8 n = 20	12.9 $\pm$ 2.9 n = 20	28.4 $\pm$ 11.2 n = 20	5.7 $\pm$ 1.8 n = 30
February 2007	25.1 $\pm$ 4.2 n = 10	12.5 $\pm$ 0.4 n = 10	12.1 $\pm$ 0.8 n = 10	9.0 $\pm$ 0.3 n = 10	12.8 $\pm$ 2.0 n = 10	8.2 $\pm$ 3.2 n = 13	4.1 $\pm$ 1.3 n = 30



**Fig. 1.** Temperature dependent burrowing capacity in *Arenicola marina*. Mean values  $\pm$  SE for animals collected in summer 2006 and in winter 2007 at the French Atlantic coast,  $n=5$ . Winter animals were acclimated to 5 and 10 °C and tested between  $-1.1$  and  $18.9$  °C. Summer animals were acclimated to 15 °C and investigated between 6.8 and 26.8 °C (depicted from Schröder et al., 2009). Data were fitted to the equation  $EP(T) = F_1(T) + F_2(T) = (A_1e^{B_1T} + C_1) + (A_2e^{B_2T} + C_2)$  with  $EP(T)$  = temperature dependent muscle exercise performance capacity. The first term,  $F_1(T) = A_1e^{B_1T} + C_1$ , represents the temperature dependence of aerobic processes supporting exercise performance. The second term,  $F_2(T) = A_2e^{B_2T} + C_2$ , represents the parallel exponential rise in processes limiting aerobic scope and thus exercise performance capacity. For the winter data (5 °C acclimated):  $A_1 = -12.02$ ,  $B_1 = 0.1385$ ,  $C_1 = 1.800$ ,  $A_2 = 15.23$ ,  $B_2 = 0.1276$ ,  $C_2 = -1.807$ ,  $r = 0.6472$ . For the winter data (10 °C acclimated):  $A_1 = 7.403$ ,  $B_1 = 0.1173$ ,  $C_1 = 0.9182$ ,  $A_2 = -3.791$ ,  $B_2 = 0.1494$ ,  $C_2 = -0.9866$ ,  $r = 0.6810$ . For the summer data:  $A_1 = 4.083$ ,  $B_1 = 0.06221$ ,  $C_1 = -0.4552$ ,  $A_2 = -1.081$ ,  $B_2 = 0.1038$ ,  $C_2 = 0.1652$ ,  $r = 0.3723$ . Shaded area: naturally experienced habitat temperatures in the respective season.

temperatures. Temperature dependence was less pronounced and lower maximum values were reached in summer animals, all values stayed between 3.6 and 5.8 digging periods  $90 \text{ min}^{-1}$ .

### 3.3. Respiration and ventilation experiment

Fig. 2 depicts temperature dependent weight specific volume flow ( $\text{ml g}^{-1} \text{h}^{-1}$ ) in winter animals acclimated to 5 and 10 °C and in summer worms acclimated to 15 °C from two years. Specimens collected in winter and acclimated to 5 °C showed an exponential increase in weight specific volume flow over the experimental temperature range with a  $Q_{10}$  value of  $5.57 \pm 0.63$  and remarkably larger error bars at temperatures  $\geq 17.2$  °C.

Winter animals acclimated to 10 °C also exhibited an exponential increase over the experimental temperature range, but the  $Q_{10}$  of  $2.38 \pm 0.38$  was significantly lower than in 5 °C acclimated specimens.


Animals collected in summer 2005 and acclimated to 15 °C displayed a  $Q_{10}$  of  $2.36 \pm 0.12$  within the temperature range of exponential increase, similar to 10 °C acclimated winter specimens. At 2.8 °C only one value was available, as the other worms stopped ventilation activity during cooling. No statistics could be applied here. The value recorded at 2.8 °C was lower than all values at 5.4 °C, therefore, the one at 2.8 °C was excluded from the exponential range.

Summer lugworms from 2006 (also acclimated to 15 °C) displayed the same exponential range as those from the preceding year, the value at 2.9 °C differed significantly from the hypothetical value predicted by the regression curve. The  $Q_{10}$  of  $2.24 \pm 0.23$  was comparable to the one evaluated in 2005.


In general, weight specific volume flow curves were similar in winter and summer animals acclimated to 10 °C and 15 °C, respectively. Only the data collected in winter animals acclimated to 5 °C differed remarkably with around two times higher values in the temperature range above 15 °C (Fig. 2).

Fig. 3 shows temperature dependent pumping frequencies ( $\text{min}^{-1}$ ) during the phases of active ventilation. Winter specimens that were acclimated to 5 °C showed a significant increase (F-test) over the experimental temperature range. Values at 19.7 and 22.6 °C displayed large variability with means significantly higher than in all other groups. Animals acclimated to 10 °C in winter and to 15 °C in summer both displayed a significant increase (F-test) with warming. Pumping frequency in lugworms collected in summer 2005 did differ significantly neither from the frequency recorded in winter animals acclimated to 10 °C nor from data collected in summer 2006. The only significant differences observed were the values at 10.9 and 23.2 °C from summer 2006 animals in comparison to both winter groups.


Fig. 4 depicts the temperature dependent extraction coefficient (%), i.e. the proportion of oxygen that the animals were able to withdraw from the inflowing water. Lugworms sampled in winter and acclimated to 5 °C showed a more or less stable extraction coefficient in the temperature range between 0 and 5.1 °C while a slight decrease was observed upon warming. In contrast, animals acclimated to 10 °C displayed a significant rise in the extraction coefficient from 0 to 17.7 °C followed by a slight decline. Summer specimens



**Fig. 2.** Temperature ( $^{\circ}\text{C}$ ) dependent weight specific volume flow ( $\text{ml g}^{-1} \text{h}^{-1}$ , mean values  $\pm$  SE) in *Arenicola marina* dwelling in an artificial burrow. Animals were collected in summer 2005, in winter 2006 and in summer 2006 at the French Atlantic coast.  $n=5$  except for winter ( $5^{\circ}\text{C}$  acclimated) at  $22.6^{\circ}\text{C}$ , winter ( $10^{\circ}\text{C}$  acclimated) at  $10.7^{\circ}\text{C}$  as well as summer 2005 at  $15.4$  and  $5.4^{\circ}\text{C}$  with  $n=4$  and summer 2005 at  $2.8^{\circ}\text{C}$  with  $n=1$ . Winter worms were acclimated to  $5$  and  $10^{\circ}\text{C}$  one half each, summer animals to  $15^{\circ}\text{C}$ . The winter specimens were investigated from  $-0.0$  to  $22.8^{\circ}\text{C}$ , both summer groups were exposed to a temperature range from  $2.8$  to  $26.1^{\circ}\text{C}$ . (Summer 2006 data depicted from Schröer et al., 2009). Data were fitted to  $V_w(T) = Ae^{BT} + C$  (Wittmann et al., 2008). For the winter  $5^{\circ}\text{C}$  acclimated group:  $A = 0.8083$ ,  $B = 0.1952$ ,  $C = 4.497$ ,  $r = 0.8118$ . For the winter  $10^{\circ}\text{C}$  acclimated group:  $A = 19.41$ ,  $B = 0.02816$ ,  $C = -16.36$ ,  $r = 0.6654$ . For the summer 2005 group:  $A = 0.2620$ ,  $B = 0.1694$ ,  $C = 5.714$ ,  $r = 0.6989$ . For the summer 2006 group:  $A = 0.08428$ ,  $B = 0.2273$ ,  $C = 7.832$ ,  $r = 0.7495$ . Asterisks (\*) designate data points, which are significantly different from the exponential regression curve ( $p = 0.0234$  for summer 2006 at  $2.9^{\circ}\text{C}$ ). Shaded area: naturally experienced habitat temperatures in the respective season.



**Fig. 3.** Ventilatory performance of *Arenicola marina* dwelling in an artificial burrow. For each group (winter  $5^{\circ}\text{C}$  acclimated, winter  $10^{\circ}\text{C}$  acclimated, summer 2005 and summer 2006 (after Schröer et al., 2009)) pumping frequency ( $\text{min}^{-1}$ ) is plotted against incubation temperature ( $^{\circ}\text{C}$ ). Mean values  $\pm$  SE;  $n=5$  except for winter ( $5^{\circ}\text{C}$  acclimated) at  $19.7$  and  $22.6^{\circ}\text{C}$ , winter ( $10^{\circ}\text{C}$  acclimated) at  $10.7^{\circ}\text{C}$  as well as summer 2005 at  $15.4$  and  $5.4^{\circ}\text{C}$  with  $n=4$  and summer 2005 at  $2.8^{\circ}\text{C}$  with  $n=1$ . Sampling time and acclimation see legend Fig. 2. Incubation temperatures ranged from  $2.8$  to  $26.1^{\circ}\text{C}$  for summer worms and from  $-0.0$  to  $22.8^{\circ}\text{C}$  for winter animals. Shaded area: naturally experienced habitat temperatures in the respective season.



**Fig. 4.** Oxygen extraction by *Arenicola marina* dwelling in an artificial burrow. The oxygen extraction coefficient (%) versus incubation temperature (°C) is shown for the four groups (winter 5 °C acclimated, winter 10 °C acclimated, summer 2005, summer 2006 (from Schröder et al., 2009), mean values  $\pm$  SE, for further information see Fig. 2. \* = significantly higher than at  $-0.0$  °C.

384 from both years showed a slight increase between 2.8 and 14.7 °C. With  
385 further warming, values stayed more or less constant. Again, the  
386 extraction coefficient in warm acclimated winter lugworms and  
387 summer specimens shows similar trends and differed from data  
388 obtained in winter animals acclimated at 5 °C.

### 389 3.4. Protein biosynthesis

390 Fig. 5 shows the temperature dependent incorporation of  $^{13}\text{C}$ -  
391 phenylalanine into proteins of the body wall after short and long  
392 term incubation times. A comparison of the three graphs reveals an  
393 overall increase of the amount of incorporated  $^{13}\text{C}$ -phenylalanine  
394 with acclimation temperature. In particular, winter animals accli-  
395 mated to 5 °C displayed no increase in  $^{13}\text{C}$ -phenylalanine content  
396 over time at  $-0.9$  °C. Data points for short and long incubation pe-  
397 riods resulted similarly at around  $4.4$  nmol  $\text{mg}^{-1}$  protein. At all  
398 other incubation temperatures, at least a slight increase between  
399 short and long incubation times was detectable with a significant dif-  
400 ference at the highest incubation temperature (19.2 °C). In winter  
401 worms acclimated to 10 °C, the rise in the amount of incorporated  
402 amino acids over time was seen at all incubation temperatures. The  
403 maximum of  $24.26 \pm 1.55$  nmol  $\text{mg}^{-1}$  protein was reached at 15.2 °C  
404 after long incubation times. Summer animals also showed an increasing  
405 incorporation of labelled phenylalanine over experimental time at all  
406 incubation temperatures. The largest amount of  $^{13}\text{C}$ -phenylalanine  
407 was incorporated at 10.2 °C, reaching  $32.48 \pm 1.24$  nmol  $\text{mg}^{-1}$  protein,  
408 a value significantly higher than at all other temperatures after short as  
409 well as long incubation periods.


## 410 4. Discussion

411 The aim of this study was to investigate potential seasonal accli-  
412 matisation effects in physiological performances of the lugworm  
413 *A. marina* from a southern population and to distinguish possible

414 effects of acclimation temperature and inter-annual variability. Recent-  
415 ly, we could show that animals from the southern-most population dis-  
416 played a lower performance level than their counterparts from higher  
417 latitudes (Schröder et al., 2009).

418 Our results show clear seasonal differences in temperature tol-  
419 erance and performance of *A. marina*. These are most likely due to  
420 seasonal temperature changes, however, other biotic and abiotic  
421 parameters also differ seasonally with potential influences on organ-  
422 ismic performance. We observed a higher animal weight in summer  
423 2006 compared to both winter samplings (Table 1), which might be  
424 explained by a higher food availability in summer and more impor-  
425 tantly, the onset of gamete production. In summer 2005, juveniles  
426 were included in the samples and resulted in a lower average weight  
427 than in 2006 and not higher than in winter. Under field conditions  
428 growth in *A. marina* was only observed during spring and summer  
429 until October (Beukema and de Vlas, 1979; Newell, 1948; Smidt, 1951;  
430 Wolff and de Wolf, 1979) whereas during autumn and winter growth  
431 was absent. Moreover, the weight of adults tends to decrease in winter  
432 (Beukema and de Vlas, 1979; Newell, 1948). Pollack (1979) argued  
433 that the lower mean animal weight in winter is not necessarily due  
434 to individual weight loss but may also result from the immigration  
435 of smaller worms to the sampling area and/or from a higher mortal-  
436 ity among older and thereby larger worms. Any effects of lower  
437 food availability in winter are exacerbated by a higher precipitation  
438 in this season, easily visible in the changing salinity of the surface  
439 water (Table 1). The lugworm shelters from the influence of fresh-  
440 water by closing its burrow and reducing its pumping and feeding  
441 activities during rainfall (Pollack, 1979). The maximally tolerated  
442 daily salinity variation amounts to 4–6‰ (Amoureux, 1966).

443 Maximum food availability from April to October coincides perfectly  
444 with the period of reproductive growth from the end of April until the  
445 beginning of October (Cazaux, 1966). Although water temperature  
446 has been well correlated with reproductive cycles and has often been  
447 considered to be a major factor in their control, studies have also



**Fig. 5.** Amount of incorporated <sup>13</sup>C-phenylalanine (nmol mg<sup>-1</sup> protein, means ± SE) into protein of the cuticulo-muscular tube of lugworms, dependent on incubation temperature. Specimens were collected in August 2005 and January/February 2006. Winter worms were acclimated to 5 and 10 °C one half each and summer animals to 15 °C. Incubation temperatures ranged from -0.9 to 19.2 °C for the winter groups and from 7.4 to 22.8 °C for summer animals. Open diamonds: amount of incorporated <sup>13</sup>C-phenylalanine after 30 to 120 min of incubation; closed diamonds: amount after 180 to 360 min; n = 4 for summer data except for short incubation times at 22.8 °C with n = 3, n = 3 for winter data. \*1 = significantly higher than the values after short incubation times at -0.9, 3.5 and 19.2 °C as well as at -0.9 °C after long incubation times. \*2 = significantly higher than the values after short incubation times at -0.9 and 6.9 °C. \*3 = significantly higher than all other values.

emphasised the importance of local food conditions (MacDonald and Thompson, 1986), as demonstrated by Newell et al. (1982) in *Mytilus edulis* populations, which experienced nearly identical temperature cycles but different regimes of food availability.

Our measurements of acute somatic growth rates by tracking the incorporation of phenylalanine into the body wall revealed an optimum temperature for growth without substrate limitation at 15.2 °C in winter animals and at 10.2 °C in summer animals (Fig. 5). Interestingly, neither 15.2 °C is experienced in winter nor 10.2 °C in summer in the natural habitat. A look at the sea surface temperature (Ifremer, 2007), which correlates well with the sediment temperature in 20 cm depth (Nießing, 2006), showed that temperatures between 10 and 16 °C are experienced in March, April and May as well as in October, November and December. Our winter animals which were acclimated

to 10 °C showed maximum somatic growth at 15.2 °C and thereby, the capacity to tolerate higher temperatures. This might indicate an early shift to sustain spring conditions. Somatic growth is expected to take place in spring (March to May) when temperature is at its optimum and food is already available, as was the case in our aquarium system. Phytoplanktonic and therewith phytobenthic growth takes place a bit earlier in the year at Arcachon (Ifremer, 2007) compared to the more northern beaches, for which the food availability data are mentioned in the Introduction. The fact that *A. marina* juveniles are found already at the end of March (Cazaux, 1966) also argues for food availability already in March.

Our summer animals which were acclimated to 15 °C showed maximum acute somatic growth at an exposure temperature of 10.2 °C. Following the same rationale as before, this might be seen as a preparation for autumn conditions. This might also involve constraints on energy budget in the warmth where cold exposure slows motor activity and supports a shift of resources to growth. After spawning in autumn, the worms are expected to replenish their glycogen reserves for winter survival (see below). So growth is controlled by exposure rather than acclimation temperature. Similarly complex interactions were previously shown for juveniles of the oyster *Ostrea edulis*, which displayed a maximum scope for growth at an acclimation temperature of approximately 17 °C and an exposure temperature of approximately 25 °C, a common condition found in shallow waters during the summer months (Buxton et al., 1981). In *A. marina*, all energy is invested into reproductive growth during summer, while somatic growth takes place in spring and autumn, suggesting a close relationship between energy available for growth, the reproductive cycle and thermal limits. In fact, Füllner (2009) observed a higher sensitivity to temperature changes in North Sea summer worms compared to North Sea spring animals, which might be due to the increased energy expenditure for reproductive growth.

Bayne and Newell (1983) suggested that scope for growth and growth efficiency are more dependent on food availability than on temperature. Both, a trade-off between growth and reproduction and a strong dependence on food availability, were found in the giant scallop *Placopecten magellanicus*: scope for growth was low or negative during winter, rapid gamete maturation was observed during the spring bloom and somatic weight declined during gamete development while it increased after spawning and during periods of low gametogenic activity (MacDonald and Thompson, 1986). Similarly, low or negative growth during the winter and generally higher values in the spring and/or summer are also described for *Chlamys islandica* (Vahl, 1980), *M. edulis* (Bayne and Widdows, 1978; Thompson, 1984) and *Mya arenaria* (Gilfillan et al., 1976). Whether somatic tissue weight declines during gamete development in *A. marina* has not been investigated so far and may depend on local temperature and food conditions. For example, *Chlamys varia* shows both, simultaneous reproductive and somatic growth as well as gamete development fuelled by somatic reserves, depending on environmental conditions (Shafee, 1980). In *Macoma balthica*, somatic growth becomes negative during gametogenesis when temperatures are high, but continues despite gamete development at low temperatures (DeWilde, 1975). High food availability combined with low temperatures and thereby a reduced metabolic demand results in high scope for growth. These conditions would allow for simultaneous somatic and reproductive growth and might be found at Arcachon in April and May. In these months, gamete development already begins (Cazaux, 1966) and temperature ranges around the optimum for somatic growth (see above).

Laboratory experiments on somatic growth in *A. marina* were already carried out by de Wilde and Berghuis (1979). Their study was performed on animals from the Netherlands with constant food supply and excluding the influence of reproduction by using juvenile worms. In that study, maximum growth rates in length and weight were observed at 20 °C with a small influence between 5 and 20 °C and a strong influence between 20 and 25 °C, as at 25 °C



the growth rate was considerably lower than at 5 °C (de Wilde and Berghuis, 1979). It should be noted that these were acclimated, not acute growth rates. In natural populations of juvenile lugworms growth rates were usually found much lower than in the experiments. It may be assumed that the main limiting factor for growth in *Arenicola* is food limitation, caused either by food competition with other small mud flat inhabiting organisms or by a poor quantity and/or quality of the organic matter available (Boon and Haverkamp, 1979; de Wilde and Berghuis, 1979). Interestingly, the increase in biomass, i.e. the product of numerical density and mean weight, was highest at 10 °C in laboratory experiments, indicating a higher mortality with rising temperature, probably caused by permanently low oxygen concentrations induced by the combination of high amounts of organic matter and high temperatures (de Wilde and Berghuis, 1979). In field studies, the highest mortality rates were found from January to March, decreasing during spring and resulting in considerably lower values in summer than in winter (Pollack, 1979). This observation suggests once more that food availability is a limiting factor for growth in the field.

The only remarkable seasonal difference was found in exercise performance (Fig. 1). Corresponding to the concept of oxygen and capacity limited thermal tolerance (Pörtner and Knust, 2007; Pörtner et al., 2004), digging activity displays an asymmetric bell shaped curve, as already shown in a previous study (Schröder et al., 2009). This study demonstrated clear seasonal acclimatisation with a more pronounced optimum at lower temperatures in winter and a wider and less distinct performance curve with an optimum at higher temperatures in summer. The data suggest a trend for curve width to increase from winter to summer at the expense of a decrease in performance amplitude. Comparing cold- and warm-adapted lugworms from different latitudes (Schröder et al., 2009) a trade-off between the width and the amplitude of the performance curve also became apparent (cf. Angilletta et al., 2002; Huey and Hertz, 1984; Pörtner, 2006). For ectothermic vertebrates, it has already been shown that muscle twitch tension decreases with increasing temperature in fast-twitch muscles (Bennett, 1984). In addition, viscoelastic properties of cell membranes change with temperature, as seen for example in human red blood cells (Hochmuth et al., 1980). So higher temperatures in summer might result in a higher elasticity of cell membranes and change the worm's whole bodywall. For this reason, only a lower internal resting pressure (turgor) might be achieved in lugworms, causing lower amplitudes of body wall contraction pressure resulting in reduced digging performance. Seymour (1971) observed that a higher resting pressure resulted in a higher peak pressure in burrowing lugworms. Our observations of the internal pressure in burrowing lugworms by use of a catheter (data not shown) in fact exhibited a tendency of higher pressures at lower temperatures, but rising contraction frequencies with rising temperature. Therefore, the internal pressure might be the limiting factor at higher temperatures, while contraction frequency may be the limiting factor at lower temperatures.

Temperature fluctuation is higher in summer (16 to 29 °C) than in winter (4 to 14 °C), which might be one reason for a broader performance curve in summer. The lower performance amplitude in summer argues for a trade-off between reproductive growth and exercise performance. Performance curves of 5 and 10 °C acclimated specimens in winter were nearly identical leading to the conclusion that acclimation temperature alone has no effect on performance capacity. Consequently, seasonal acclimatisation does not only depend on temperature, but other seasonal changes (e.g. photoperiod, precipitation and food availability), as well as competing physiological processes like reproductive growth can also play an important role for the observed seasonal differences in muscular performance (i.e. digging activity).

Thermal acclimatisation of muscle performance has already been shown before in other marine invertebrates as in the European queen scallop, *Aequipecten opercularis* (Bailey and Johnston, 2005) and the

giant scallop, *P. magellanicus* (Guderley et al., 2008). In these studies, higher performance amplitudes also result in cold acclimated animals. Cold acclimated *A. opercularis* attained higher swimming velocities and accelerated faster at winter temperatures than warm acclimated animals at summer temperatures (Bailey and Johnston, 2005). A study by Guderley et al. (2008) reported that even handling stress had less impact on cold than on warm acclimated animals of the species *P. magellanicus*. Guderley (2004) suggested that locomotor performance and reproduction are closely coupled, as muscle metabolic capacities fall in parallel with glycogen mobilisation for gametogenesis, and reproductive fitness will be favoured more than maintenance of performance. This interpretation might explain our observations on lugworms as well. In our study, winter acclimated worms displayed the highest performance optima, as reproductive growth takes place in summer (from end of April to the beginning of October, Cazaux, 1966) and spawning occurs during autumn (October to November). So in summer animals the gonads take up most of the body mass, whereas winter animals do not invest any energy into reproduction. Also *A. opercularis* and *P. magellanicus* both have their spawning period until October and achieve performance maxima in winter after spawning. Füßner (2009) observed a much narrower thermal tolerance window in *A. marina* from the North Sea during the time of spawning than in specimens during the early stage of gamete production.

Consistent with previous findings (Schröder et al., 2009), performance optima were found within the naturally experienced temperature range during the respective season in this study. In winter, modelled performance maxima of 14.2 and 13.3 °C, respectively, were found close to the upper limit of naturally experienced habitat temperatures of around 14 °C. In contrast, the modelled performance maximum of 19.6 °C in summer was found close to the lower limit of habitat temperatures during the respective season (around 17 °C). This leads to the conclusion that performance, despite its seasonal acclimatisation, is well adapted to the yearly mean habitat temperature of approximately 16 °C ensuring constant performance during the whole year. By extrapolating the model curve, it also becomes obvious that the upper critical temperature in summer, which is predicted by the model at 31.7 °C, nearly falls into the naturally experienced range of habitat temperatures of up to 29 °C in summer. This suggests that the population is more sensitive to warming effects in summer than in winter.

Effects of temperature acclimation without the influence of reproduction have been investigated in tadpoles of *Limnodystes peronii* as well as tadpoles and adults of *Xenopus laevis* (Wilson and Franklin, 1999; Wilson et al., 2000). Cold acclimated organisms reached a higher swimming velocity at low temperatures than warm acclimated specimens, while at high temperatures warm acclimated animals showed a higher swimming performance than cold acclimated ones. This simple relationship is overlaid with the effects of reproductive growth in our study when comparing digging activity in winter and summer acclimated lugworms.

Despite a nearly identical performance curve, the group of winter animals acclimated to 5 °C showed a much higher volume flow than those acclimated to 10 °C. Differences became obvious especially at temperatures above 15 °C, resulting in a high  $Q_{10}$  value of  $5.57 \pm 0.63$  compared to values between 2.24 and 2.38 in the other winter and summer groups. 5 °C acclimated winter specimens showed their highest extraction coefficients at temperatures from 0.0 to 5.1 °C, whereas 10 °C acclimated winter worms and both groups of summer worms exhibited their lowest extraction coefficient values in this temperature range. Cold compensation at the cellular level likely occurred at 5 °C. Sommer and Pörtner (2004) found changes in mitochondrial functions taking place in *A. marina* from the North Sea, which were acclimated to 0 °C in comparison to those acclimated to 5 and 11 °C. These findings suggest that cold acclimatisation occurs below a threshold temperature rather than progressively

with falling environmental temperatures. The other way round, returning to warm acclimatisation also seems to occur stepwise, as Wittmann (2005) found different thermal tolerance windows in worms which experienced a warming pulse of 3 °C compared to those which were investigated before. Somatic growth (Fig. 5) was minimal in our 5 °C acclimated compared to 10 °C acclimated winter lugworms. These findings could be an indication for a temperature induced dormant condition as it is found in the common shallow-water and littoral bivalve *Cardium* (= *Cerastoderma*) *edule* L. in winter (Newell and Bayne, 1980). Dormancy has also been found in other marine polychaetes like *Lanice conchilega* (Herses, 1997). Dormant individual relies on carbohydrate reserves for maintenance energy requirements (Newell and Bayne, 1980). Lugworms also show a decrease in the glycogen content of the body wall between November and February (Nießing, 2006), which coincides with the time of low food availability (see above). Accordingly, winter specimens in our study displayed a significantly lower fresh weight than summer animals (Table 1). Field observations revealed a lower faeces production and hence food uptake from October to January compared to the spring and summer months (Pollack, 1979), possibly initiated by a drop in temperature and/or a reduced food availability. It was shown in laboratory experiments that faeces production and hence feeding activity is dependent on the food content of the sediment. Considerable differences between actual numbers of lugworms and number of faecal casts occurred, indicating part of the animals to be inactive at less favourable conditions in poor sediments (de Wilde and Berghuis, 1979).

Also the organic matter content of the water was shown to influence the pumping activity (Krüger, 1964). Our 5 °C cold-acclimated winter lugworms pursued the strategy of a “conserver”, which lives under conditions of short food, feeds at a low rate because a higher rate might increase the costs of foraging more than the gains, and has a low growth rate in favour of longevity (Branch et al., 1988). In contrast to volume flow, pumping frequency, extraction coefficient and growth, the modelled exercise performance curves and maxima (Fig. 1) were nearly identical for both winter groups. This emphasises that maintenance of muscle exercise capacity seemed to be important for winter lugworms. An explanation might be the observation of active winter migrations in *A. marina*, initiated by temperatures close to critical values (Werner, 1956). When the limits of acclimatory adjustment are reached, the animals leave their habitat and rebury in a lower zone of the intertidal. De Wilde and Berghuis (1979) made a similar observation in a laboratory experiment, when lugworms fed and maintained at 5 °C left the sediment for migration.

## 5. Conclusions

Seasonal differences become obvious in exercise performance curves of *A. marina*. Winter animals exhibit an optimum at around 14 °C and a high performance amplitude. In summer, the optimum is shifted towards 19 °C, accompanied by a widening of the performance window and lower performance amplitudes. A trade-off between exercise capacity and reproductive growth seems to take place in summer. In addition to temperature food availability is likely an important factor controlling seasonal acclimatisation processes. Somatic growth may occur mostly in spring and autumn, when food is available, and outside the reproductive period. The experimentally determined optimum combination of acclimation and exposure temperatures for growth matches well with the temperature conditions found in spring and autumn in the worms' natural habitat. Altogether, the present results show that temperature determines the metabolic state of lugworms despite of the season. Furthermore, our data confirm that the investigated lugworm population from Arcachon lives at the upper level of its thermal limit, making it most susceptible for warming waters in the near future.

## 6. Uncited references

Augee et al., 1984  
 Brockington and Clarke, 2001  
 Clarke, 1993  
 De Vooy, 1975  
 Duncan, 1960  
 Ehrenhauss et al., 2004  
 Foster-Smith, 1975  
 Kristensen, 1988  
 Longbottom, 1970  
 Urban and Mercuri, 1998

## Acknowledgements

This work was supported by grants from the Deutsche Forschungsgemeinschaft (Aquashift Po 278/11). The experiments described here comply with the current laws of the country in which they were performed. We would like to thank A. Wittmann for her work in the respiration and ventilation activity experiments. We are very grateful for the generous support by the Biological Station at Arcachon (Atlantic, France). Special thanks go to the T. Hirse, B. Klein and R. Wittig for technical support and G. Lannig for assistance in revising the manuscript. [SS]

## References

- Amoureux, L., 1966. Etude bionomique et écologique de quelques Annélides Polychètes des sables intertidaux des côtes ouest de la France, Faculté des Sciences. Université de Paris, Paris, pp. Ed. C.N.R.S., 218 pp.  
 Angilletta Jr., M.J., Niewiarowski, P.H., Navas, C.A., 2002. The evolution of thermal physiology in ectotherms. *J. Therm. Biol.* 27, 249–268.  
 Augee, M.L., Pehowich, D.J., Raison, J.K., Wang, L.C.H., 1984. Seasonal and temperature-related changes in mitochondrial membranes associated with torpor in the mammalian hibernator *Spermophilus richardsonii*. *Biochim. Biophys. Acta* 776, 27–36.  
 Bailey, D.M., Johnston, I.A., 2005. Temperature acclimatisation of swimming performance in the European Queen Scallop. *J. Therm. Biol.* 30, 119–124.  
 Bayne, B.L., Newell, R.C., 1983. Physiological energetics of marine molluscs. *The Mollusca*. Academic Press, pp. 407–515.  
 Bayne, B.L., Widdows, J., 1978. The physiological ecology of two populations of *Mytilus edulis* L. *Oecologia* 37, 137–162.  
 Bennett, A.F., 1984. Thermal dependence of muscle function. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 247/2, R217–R229.  
 Beukema, J.J., de Vlas, J., 1979. Population parameters of the lugworm, *Arenicola marina*, living in the Dutch Wadden Sea. *Neth. J. Sea Res.* 13, 331–353.  
 Boisseau, J., 1962. Contribution à la faune du Bassin d'Arcachon. P.V. Soc. linn. Bordeaux 99, 113–126.  
 Boon, J.J., Haverkamp, J., 1979. Pyrolysis mass spectrometry of a benthic marine ecosystem – the influence of *Arenicola marina* on the organic matter cycle. *Neth. J. Sea Res.* 12, 58–77.  
 Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.  
 Branch, G.M., Borchers, P., Brown, C.R., Donnelly, D., 1988. Temperature and food as factor influencing oxygen consumption of intertidal organisms, particularly limpets. *Am. Zool.* 28, 137–146.  
 Brockington, S., Clarke, A., 2001. The relative influence of temperature and food on the metabolism of a marine invertebrate. *J. Exp. Mar. Biol. Ecol.* 258, 87–99.  
 Buxton, C.D., Newell, R.C., Field, J.G., 1981. Response-surface analysis of the combined effects of exposure and acclimation temperatures on filtration, oxygen consumption and scope for growth in the oyster *Ostrea edulis*. *Mar. Ecol. Prog. Ser.* 6, 73–82.  
 Cáceres, C.E., 1997. Dormancy in invertebrates. *Invert. Biol.* 116, 371–383.  
 Cazaux, C., 1966. Evolution d'une population d'*Arenicola marina* (L.) à Arcachon. Actes de la Société Linnéenne de Bordeaux, 103, pp. 3–18.  
 Clarke, A., 1993. Seasonal acclimation and latitudinal compensation: do they exist? *Funct. Ecol.* 7, 139–149.  
 Davey, J.T., Watson, P.G., Bruce, R.H., Frickers, P.E., 1990. An instrument for the monitoring and collection of the vented burrow fluids of benthic infauna in sediment microcosms and its application to the polychaetes *Hediste diversicolor* and *Arenicola marina*. *J. Exp. Mar. Biol. Ecol.* 139, 135–149.  
 De Vooy, C.G.N., 1975. Glycogen and total lipids in the lugworm (*Arenicola marina*) in relation to reproduction. *Neth. J. Sea Res.* 9, 311–319.  
 De Wilde, P.A.W.J., Berghuis, E.M., 1979. Laboratory experiments on growth of juvenile lugworms, *Arenicola marina*. *Neth. J. Sea Res.* 13, 487–502.

- 793 DeWilde, P.A.W.J., 1975. Influence of temperature on behaviour, energy metabolism  
794 and growth of *Macoma balthica* (L.). In: Barnes, H. (Ed.), 9th Eur. Mar. Biol. Symp.  
795 Aberdeen University Press, Aberdeen, pp. 239–256.
- 796 Duncan, A., 1960. The spawning of *Arenicola marina* (L.) in the British Isles. Proc. zool.  
797 Soc. Lond. 134, 137–156.
- 798 Ehrenhauss, S., Witte, U., Bühring, S., Huettel, M., 2004. Effect of advective pore water  
799 transport on distribution and degradation of diatoms in permeable North Sea  
800 sediments. Mar. Ecol. Prog. Ser. 271, 99–111.
- 801 Foster-Smith, R.L., 1975. The effect of concentration of suspension on the filtration  
802 rates and pseudofaecal production for *Mytilus edulis* (L.), *Cerastoderma edule* (L.)  
803 and *Venerupis pullastra* (Montagu). J. Exp. Mar. Biol. Ecol. 17, 1–22.
- 804 Frederich, M., Pörtner, H.O., 2000. Oxygen limitation of thermal tolerance defined by  
805 cardiac and ventilatory performance in spider crab, *Maja Squinado*. Am. J. Physiol.  
806 279, R1531–R1538.
- 807 Fülner, V., 2009. Online Bestimmung der kritischen Temperatur beim Wattwurm  
808 *Arenicola marina* (L.). Universität Bremen, Bremen.
- 809 Gilfillan, E.S., Mayo, D., Hanson, S., Donovan, D., Jiang, L.C., 1976. Reduction in carbon  
810 flux in *Mya arenaria* caused by a spill of no. 6 fuel oil. Mar. Biol. 37, 115–123.
- 811 Guderley, H., 2004. Locomotor performance and muscle metabolic capacities: impact of  
812 temperature and energetic status. Comp. Biochem. Physiol. B 139, 371–382.
- 813 Guderley, H., Janssoone, X., Nadeau, M., Bourgeois, M., Cortés, H.P., 2008. Force record-  
814 ings during escape responses by *Placopecten magellanicus* (Gmelin): seasonal  
815 changes in the impact of handling stress. J. Exp. Mar. Biol. Ecol. 355, 85–94.
- 816 Hochmuth, R.M., Buxbaum, K.L., Evans, E.A., 1980. Temperature dependence of the  
817 viscoelastic recovery of red cell membrane. Biophys. J. 29, 177–182.
- 818 Hubas, C., Lamy, D., Artigas, L.F., Davoult, D., 2007. Seasonal variability of intertidal bac-  
819 terial metabolism and growth efficiency in an exposed sandy beach during low  
820 tide. Mar. Biol. 151, 41–52.
- 821 Huey, R.B., Hertz, P.E., 1984. Is a jack-of-all-temperatures a master of none? Evolution  
822 38, 441–444.
- 823 Ifremer, 2007. Résultats de la surveillance de la qualité du milieu marin littoral. In:  
824 Ifremer/RST.LERAR/07.002 (Ed.), Bulletin de la surveillance. Laboratoire Environ-  
825 nement Ressources d'Arcachon, pp. 1–81.
- 826 Kristensen, E., 1988. Benthic fauna and geochemical processes in marine sediments:  
827 microbial activities and fluxes. In: Blackburn, T.H., Sorensen, J. (Eds.), Nitrogen Cycling  
828 in Coastal Marine Environments. John Wiley & Sons, pp. 275–299.
- 829 Krüger, F., 1957. Der Wohnbau von *Arenicola* als Filtereinrichtung. Naturwissenschaften  
830 44, 597.
- 831 Krüger, F., 1964. Versuche über die Abhängigkeit der Atmung von *Arenicola marina*  
832 (Annelida, Polychaeta) von Größe und Temperatur. Helg. Wiss. Meeresuntersuch.  
833 10, 38–63.
- 834 Langenbuch, M., Bock, C., Leibfritz, D., Pörtner, H.-O., 2006. Effects of environmental  
835 hypercapnia on animal physiology: a <sup>13</sup>C NMR study of protein synthesis rates  
836 in the marine invertebrate *Sipunculus nudus*. Comp. Biochem. Physiol. 144,  
837 479–484.
- 838 Longbottom, M.R., 1970. The distribution of *Arenicola marina* (L.) with particular  
839 reference to the effects of particle size and organic matter of the sediment. J.  
840 Exp. Mar. Biol. Ecol. 5, 138–157.
- 841 MacDonald, B.A., Thompson, R.J., 1986. Influence of temperature and food availability on  
842 the ecological energetics of the giant scallop *Placopecten magellanicus*. III. Physiological  
843 ecology, the gametogenic cycle and scope for growth. Mar. Biol. 93, 37–48.
- 844 Newell, G.E., 1948. A contribution to our knowledge of the life history of *Arenicola*  
845 *marina*. J. Mar. Biol. Assoc. U. K. 27, 554–580.
- 846 Newell, R.I.E., Bayne, B.L., 1980. Seasonal changes in the physiology, reproductive  
847 condition and carbohydrate content of the cockle *Cardium* (= *Cerastoderma*)  
848 *edule* (Bivalvia: Cardiidae). Mar. Biol. 56, 11–19.
- 849 Newell, R.I.E., Hilbish, T.J., Koehn, R.K., Newell, C.J., 1982. Temporal variation in the  
850 reproductive cycle of *Mytilus edulis* (L.) (Bivalvia: Mytilidae) from localities on  
851 the east coast of the United States. Biol. Bull. Mar. Biol. Lab., Woods Hole 162,  
852 299–310.
- 853 Nießing, V., 2006. Biochemische, physiologische und ökologische Aspekte der  
854 Thermotoleranz von *Arenicola marina*. Institut für Zoophysologie, Westfälische  
855 Wilhelms-Universität, Münster, pp. 1–70.
- Pollack, H., 1979. Populationsdynamik, Produktivität und Energiehaushalt des  
856 Wattwurms *Arenicola marina* (Annelida, Polychaeta). Helgol. Mar. Res. 32, 313–358.  
857
- Pörtner, H.-O., 2001. Climate change and temperature-dependent biogeography:  
858 oxygen limitation of thermal tolerance in animals. Naturwissenschaften 88,  
859 137–146.
- Pörtner, H.-O., 2006. Climate-dependent evolution of Antarctic ectotherms: an integra-  
861 tive analysis. Deep Sea Res. Part II 53, 1071–1104.
- Pörtner, H.-O., Knust, R., 2007. Climate change affects marine fishes through the oxygen  
863 limitation of thermal tolerance. Science 315, 95–97.
- Pörtner, H.O., Mark, F.C., Bock, C., 2004. Oxygen limited thermal tolerance in fish?  
865 Answers obtained by nuclear magnetic resonance techniques. Respir. Physiol.  
866 Neurobiol. 141, 243–260.
- Riisgård, H.U., Berntsen, I., Tarp, B., 1996. The lugworm (*Arenicola marina*) pump:  
868 characteristics, modelling and energy cost. Mar. Ecol. Prog. Ser. 138, 149–156.  
869
- Schröder, M., Wittmann, A.C., Grüner, N., Steeger, H.-U., Bock, C., Paul, R.J., Pörtner, H.O.,  
870 2009. Oxygen limited thermal tolerance and performance in the lugworm *Arenicola*  
871 *marina*: a latitudinal comparison. J. Exp. Mar. Biol. Ecol. 372, 22–30.
- Seymour, M.K., 1971. Burrowing behaviour in the European lugworm *Arenicola marina*  
873 (Polychaeta: Arenicolidae). J. Zool. 164, 93–132.
- Shafee, M.S., 1980. Ecophysiological Studies on a Temperate Bivalve *Chlamys varia* (L.)  
875 from Lanevoc (Bay of Brest). Université de Bretagne Occidentale, France.  
876
- Shumway, S.E., Davenport, J., 1977. Some aspects of the physiology of *Arenicola marina*  
877 (Polychaeta) exposed to fluctuating salinities. J. Mar. Biol. Assoc. U. K. 57, 907–924.  
878
- Smidt, E.L.B., 1951. Animal production in the Danish Wadden Sea. Meddr Kommn  
879 Danm. Fisk-og Havunders. Serie Fiskeeri, 11, pp. 1–151.
- Sommer, A., 2001. Russian/German cooperation at the White Sea: physiological adap-  
881 tations of marine invertebrates to a life at different climatic zones. Mitt. Pollichia  
882 88, 95–100.
- Sommer, A.M., Pörtner, H.-O., 2002. Metabolic cold adaptation in the lugworm  
884 *Arenicola marina*: comparison of a North Sea and a White Sea population.  
885 Mar. Ecol. Prog. Ser. 240, 171–182.
- Sommer, A.M., Pörtner, H.O., 2004. Mitochondrial function in seasonal acclimatization  
887 versus latitudinal adaptation to cold in the lugworm *Arenicola marina* (L.). Physiol.  
888 Biochem. Zool. 77, 174–186.
- Sommer, A., Klein, B., Pörtner, H.O., 1997. Temperature induced anaerobiosis in two  
890 populations of the polychaete worm *Arenicola marina* (L.). J. Comp. Physiol. B 167,  
891 25–35.
- Thompson, R.J., 1984. The reproductive cycle and physiological ecology of the mussel  
893 *Mytilus edulis* in a subarctic, non-estuarine environment. Mar. Biol. 79, 277–288.  
894
- Urban, H.J., Mercuri, C., 1998. Population dynamics of the bivalve *Laternula elliptica*  
895 from Potter Cove, King George Island, South Shetland Islands. Antart. Sci. 10,  
896 153–160.
- Vahl, O., 1980. Seasonal variations in seston and in the growth rate of the iceland  
898 scallop *Chlamys islandica* (O.F. Müller) from Balsfjord, 70°N. J. exp. mar. Biol.  
899 Ecol. 48, 195–204.
- Wells, G.P., 1945. The mode of live of *Arenicola marina*. J. Mar. Biol. Assoc. U. K. 26,  
901 170–207.
- Werner, B., 1956. Über die Winterwanderung von *Arenicola marina* L. (Polychaeta  
903 sedentaria). Helgoländer Wiss. Meeresunters 5, 353–378.
- Wilson, R.S., Franklin, C.E., 1999. Thermal acclimation of locomotor performance in  
905 tadpoles of the frog *Limodynastes peronii*. J. Comp. Physiol. B 169, 445–451.
- Wilson, R.S., James, R.S., Johnston, I.A., 2000. Thermal acclimation of locomotor per-  
907 formance in tadpoles and adults of the aquatic frog *Xenopus laevis*. J. Comp. Physiol. B  
908 170, 117–124.
- Wittmann, A.C., 2005. Bestimmung des sauerstofflimitierten Temperaturtoleranzfensters  
910 beim Wattwurm *Arenicola marina* (L.). Universität Bremen, Bremen.
- Wittmann, A.C., Schröder, M., Bock, C., Steeger, H.-U., Paul, R.J., Pörtner, H.-O., 2008. 912  
913 Indicators of oxygen- and capacity-limited thermal tolerance in the lugworm  
*Arenicola marina*. Climate Res. 37, 227–240.
- Wolff, W.J., de Wolf, L., 1979. Biomass and production of zoobenthos in the Grevelingen  
915 estuary, the Netherlands. Estuar. Coast. Mar. Sci. 5, 1–24.  
916