

Methods for ^{137}Cs and $^{239,240}\text{Pu}$ analysis

Patricia Cámara Mor - Polarstern ARK-XXII/2, (GEOTRACES GIPY11)

Sampling

Artificial radionuclides to be measured (Plutonium isotopes and Cesium-137) are present at ultra trace levels, which makes difficult their measurement and thus forces to collect large volume samples. 100 L water samples were transferred into a plastic barrel and acidified with 32% HCl. ^{242}Pu and stable Be spikes as chemical yield tracers, and an iron carrier solution (FeCl_3) were added under constant stirring. After waiting 12 hours in order to get a chemical equilibrium, the pH was then made 9 with NaOH in order to precipitate the iron as $\text{Fe}(\text{OH})_3$. This scavenges beryllium and plutonium onto the precipitate while cesium stays in solution. After that the supernatant was then transferred into a second barrel while the precipitate was stored in small plastic bottles for further processing at laboratory. By addition of 65% HNO_3 , the pH was lowered again and subsequently stable Cs was added as a chemical yield tracer. Then, a pre-weighed sample of ammonium molybdophosphate was added while stirring thoroughly. This produced a yellow precipitate that scavenges Cs. The precipitate was left to settle and then transferred into smaller bottles.

Cs-137

A pre-weighed sample of ammonium molybdophosphate (AMP) was added while stirring thoroughly. This produced a yellow precipitate that scavenges Cs. The precipitate was left to settle and then transferred into vials for gamma spectrometry.

Pu-239,240

The method used to quantify Pu isotopes in sediment samples by ICP-MS at LDEO was modified after Maxwell [1997] and Kenna [2002]. About 2 g of dry sediment were spiked with ^{242}Pu as a yield monitor. Sediments were completely digested using a mixture of HNO_3 (65%) and HF (40%) under heating at 180°C in a hot plate. After matrix dissolution, HClO_4 (1 mL) was added in order to remove HF by fuming. This step was repeated three times and the solution was evaporated to dryness. Sample was redissolved with 1 mL HCl (%) and 5 mL d.i. H_2O . After that, Pu was co-precipitated with $\text{Fe}(\text{OH})_3$, adding concentrated NH_4OH . After cooling, the resulting precipitate was centrifuged and the supernate removed. The precipitate was rinsed and re-centrifuged three additional times with deionized water. The final iron precipitate was dissolved in concentrated 1 mL HNO_3 (65%) and taken to dryness. This step was done 3 times. After that the iron precipitate was dissolved in 40 mL 3M HNO_3 . Prior to ion exchange purification, the solution was prepared to contain Pu in the IV oxidation state by adding 1.3 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ and 2.2 mL 4.2 M NaNO_2 . Ion exchange separation was carried out with TEVA- resins and TEVA-pre-filter resin placed on a vacuum system. The 30 mL 3M HNO_3 solution containing the digested sample is loaded to the TEVA resin after its preconditioning with 25 mL of 3M HNO_3 . The Pu fraction was isolated from the other actinides by washing the column with 9 M HCl (2.5 mL) to remove Th and 8 M HNO_3 (15 mL) to remove U. Before Pu was eluted, a pre-filter resin was placed between column and solution

collector to minimize the contribution of other components and then Pu was eluted with a mixture of 0.13 M HF and 0.02 M HNO₃ (25 mL). The eluate was evaporated to dryness and redissolved with 7 mL HNO₃ (65%) and 2 mL 3M HNO₃. Then, it was prepared for a second purification adding 0.65 g of FeSO₄·7H₂O and 2 ml 4.2M NaNO₂ to ensure that Pu was in IV oxidation stage. The radiochemical separation with TEVA-resins was identical to the first step. After collecting the final Pu fraction, three drops of concentrated HClO₄ was added to remove the rest of resin, and then the solution was evaporated to almost dryness. The drop was dissolved in 1mL HNO₃ (65%) and transferred to small Teflon vial and heated to dryness at 200°C. Finally, 2 mL of an acid mixture 1% HNO₃/0.1% HF was added to adjust and transfer the solution to the ICP-MS vial and solution was heated few seconds.

Kenna T. (2002). Determination of plutonium isotopes and neptunium-237 in environmental samples by inductively coupled plasma mass spectrometry with total sample dissolution. *Journal of Analytical Atomic Spectrometry*, 17(11), doi:[10.1039/b205001c](https://doi.org/10.1039/b205001c)

Maxwell III S.L. (1997), Rapid actinide separation methods. *Radioactivity and Radiochemistry*, 8, 36-44, <http://sti.srs.gov/fulltext/ms2005386/ms2005386.pdf>