

# A STUDY OF GEOPOTENTIAL GEOID IN THE PENINSULAR OF MALAYSIA

by

**Shahrin Ses**  
Jabatan Geodesi & Astronomi  
Fakulti Ukur  
Universiti Teknologi Malaysia

## Abstract

Geoidal heights can be computed for a single point value or a grid of values. A program to compute geoidal heights from a set of high degree potential coefficients was developed. Backward recurrence formulae have been used to evaluate the value of normalized Legendre functions. The regional and global geopotential geoid evaluated from the available sets of potential coefficients were shown in the form of contoured maps. The computed geoidal heights derived from different sets of potential coefficients were compared with Doppler derived values at six points in Peninsular Malaysia. The results indicate that, of the models tested OSU86 gives the best solution to the geopotential geoid in the region.

## 1.0 Introduction

The geoid has been loosely defined as the equipotential surface of the earth's gravity field which would coincide with the mean sea level if the latter were undisturbed and affected only by the earth's gravity field. It is an important surface to which many geodetic observations are related. While the geodetic coordinates of the point are referred to the ellipsoid, orthometric heights are referred to the geoid. The relationship between the terrain, geoid, and ellipsoid is shown in figure 1. The geoid is of increasing importance in modern development of geodesy, as it needs to be known in order to convert ellipsoidal heights to orthometric heights.

The above equation can be used to calculate the geoidal heights  $N(\theta, \lambda, r)$  by using the Brun's formula  $N=T/\gamma$ , where  $\gamma$  is normal value of gravity at the given

and

GM geocentric gravitational constant  
 $r, \theta, \lambda$  geocentric coordinates  
 $C_{nm} \cdot S_{nm}$  fully normalized potential coefficients  
 $P_{nm}$  fully normalized Legendre function of degree  $n$   
 $a$  equatorial radius of a reference ellipsoid  
 $m$  order

where

$$T(\theta, \lambda, r) = \frac{r}{GM} \sum_{n=2}^{\infty} \left[ \frac{r}{a} \right]^n \sum_{m=0}^n [C_{nm} \cos m\lambda + S_{nm} \sin m\lambda] P_{nm}(\cos \theta) \dots (1)$$


The earth's disturbing potential (T) is given by a set of fully normalized potential coefficients:

### 2.0 Approach used to compute geoidal heights

Geoidal heights may be computed if a global estimate of the gravity anomaly field is used in the Stokes' equation. However, if we were given a set of potential coefficients describing the gravitational potential of the earth, the geoidal heights may also be computed. These coefficients are determined from a combination of satellite orbit perturbations, satellite altimeter data, and mean terrestrial gravity data. Once determined, they are valid everywhere on the earth's surface, i.e. they can be used to compute geoidal height for any given latitude and longitude. Examples of such coefficients are the GEM10B model (Lerch et al [1981]), the OSU81 and OSU86 models (Rapp [1981] and Rapp et al [1986]) and the GPM2 model (Wanzel [1985]).

If the terrain point P is to have all three defining parameters referred to the ellipsoid, a knowledge of the geoidal height is required.

Figure 1 : Topographic surface, geoid and ellipsoid



Geoidal heights can be computed for two different cases. These include a single point value or a grid of values. The data required is depending upon the case selected by the user.

A. Choice of case

The program is written to be interactive and prompts the user throughout. The program offers the following options:

**4.0 Description of the Program**

t      cos θ  
θ      colatitude of computation point

where

$$P_{nm}(t) = [2(2n+1) 2n(4n)^{-1} 2n^{-1} (4n-4)^{-1} \dots n+1(4)^{-1}]^{1/2} \sin^n \theta$$

$$P_{n-1}(t) = [2n]^{1/2} P_{nm}(t) \cot \theta$$

$$P_{nm}(t) = 2^{(m+1)} \cot \theta [(n-m)^{-1} (n+m+1)^{-1}]^{1/2} P_{nm+1}(t)$$

$$[(n-m-1) (n+m+2) (n-m)^{-1} (n+m+1)^{-1}]^{1/2} P_{nm+2}(t) \dots (4)$$

The normalized Legendre functions are required for the geoidal height computation. These normalized values may be computed by using either a direct or recursive method. The following backward recurrence formulas derived from a combination of the two methods will be used to compute the normalized Legendre functions:

**3.0 Method to compute the Legendre functions**

$\Delta C_{2n}$       fully normalized correction term  
e      first eccentricity

where


$$\Delta C_{2n} = (4n+1)^{-1/2} (-1)^n 3e^{2n} (2n+1)^{-1} (2n+3)^{-1} [1 - n + (5n C_2/e^2)] \dots (3)$$

The lower even degree zonal coefficients i.e.  $C_2, C_4$  and  $C_6$  have to be corrected to remove the effect of the normal gravity field. The correction can be computed using the series expansion of the normal gravity field (Heiskanen and Moritz [1967]):

$$N(\theta, \lambda, r) = \frac{r}{GM} \sum_{n=2}^{\infty} \left[ \frac{r}{a} \right]^n \sum_{m=0}^n [ \bar{C}_{nm} \cos m\lambda + \bar{S}_{nm} \sin m\lambda ] \bar{P}_{nm}(\cos \theta) \dots (2)$$

point:

Figure 2: C.P.U. time for the normalized Legendre functions calculation



Next, the subroutine was timed for the calculation of the normalized values for varying degree and order. The following graph in Figure 2 shows the C.P.U. times required to compute the normalized values for different N max.

In order to check the stability of computing the normalized Legendre functions using equations (4), the related subroutine was tested for different latitudes and varying degree and order. The normalized values obtained from different values of N max were compared. For each case the normalized Legendre functions agreed. This shows the stability of the method being used to compute these values.

5.0 Discussion on the method of computation

The parameters defining the geocentric reference system used in this program (a, f, C2 and GM) could be changed if desired. By changing f or C2, the coefficients of the normal gravity field (equation (3)) are altered.

C. Normal gravity field

Although equation (2) indicates a sum to infinity, in practice the sum is up to a finite degree such as 36, 180, 360, etc. The user may specify any value for N max, up to a maximum of 360, for the computation of geoidal height.

B. Maximum degree (N max) required

The results shown in Table 1 indicate that the best solution considering the root mean square of the difference is a given by OSU86 with expansion complete to degree and order 360 (rms = 0.69).


Geoidal heights at six Doppler points in Peninsular Malaysia were computed from the different sets of potential coefficients. The results were then compared with the geoidal heights derived from satellite Doppler derived positions. Prior to this, the Doppler derived cartesian coordinates which are given in WGS72 have been transformed to GRS 80 using the parameters given by Cross [1987].

### 6.0 Discussion of the results

The accuracy of the computed geoidal heights is mainly dependent upon two factors. First, the accuracy of the potential coefficients being used and second, the degree and order at which the infinite series in equation (2) is truncated.

The main program was also timed for a single geoidal height calculation and the second graph in Figure 3 shows the results for varying degree and order. These times include the computation of the normalized Legendre functions.

**Figure 3:** C.P.U. time for geoidal height calculation.



The geoid determined here can only represent the long wavelength features of the geoid in the regions. Since the current holding of gravity data is lacking, the remaining features of the local geoid in the Peninsular of Malaysia cannot be evaluated. In future, the long wavelength geoidal heights information could be combined with the remaining features evaluated gravimetrically to determine a local gravimetric geoid in the region.

This paper has discussed a computer program that can be used for the calculation of geoidal heights from a set of potential coefficients. Several computations were carried out to determine the geoidal heights using three sets of potential coefficients with the expansions up to degree and order 360. The comparison with the Doppler derived geoidal heights indicates that OSU86 gives the best solution to the geopotential geoid in Peninsular Malaysia.

**7.0 Conclusions**

Finally the program was used to compute the global geopotential geoid from the GEM10B lower degree field which is complete to degree and order 36. A contour of the geoid above the reference ellipsoid used in GEM10B ( $a=6378138$  m,  $f=1/298.257$ ) is shown in Figure 5 together with its block diagram. The contoured map was then compared with the one prepared by Lerch et al [1981] and showing a good agreement.

The program was also used to evaluate the geopotential geoid in the Malaysian region ( $0^\circ < \phi < 8^\circ$ ,  $96^\circ < \lambda < 120^\circ$ ) using the OSU86 potential coefficients set which is complete to degree and order 360. A map showing the geoid above the GRS80 ellipsoid which has been constructed from points of  $0.5^\circ \times 0.5^\circ$  grid intersection is shown in Figure 4. We can see that the Malaysian region has a steep geoid in the east-west direction.

**Table 1 :** Comparison of Doppler derived geoidal heights in Peninsular Malaysia with values from different sets of potential coefficients

Doppler points		Lat	Long	1 Dopp	2 GEM10B	3 OSU81	4 OSU86	RMS	2.89	1.00	0.69
Geoidal heights (m)											
3.4638	102.6217	0.80	3.31	-0.12	0.25	-2.51	0.92	0.41	0.98	-0.73	0.04
3.0247	101.4456	-3.22	0.31	-3.17	-2.25	-3.53	-0.05	-0.98	-0.98	-0.73	0.04
6.0387	102.3205	-6.12	-3.09	-6.19	-5.39	-3.03	0.07	-0.98	-0.98	-0.73	0.04
1.3765	103.6080	6.50	8.21	7.99	6.46	-1.71	-1.49	0.04	0.04	-0.73	0.04
6.1397	100.3849	-12.28	-8.70	-12.82	-11.26	-3.58	0.54	-1.02	-0.43	-0.43	0.04
1.4689	103.2564	4.57	7.13	6.22	5.00	-2.56	-1.65	-0.43	-0.43	-0.43	0.04

**ACKNOWLEDGEMENTS**

I wish to acknowledge the guidance given by Prof. Dr. Paul A. Cross at the Department of Surveying, University of Newcastle upon Tyne.  
The Doppler data used in this study is provided by The Department of Surveying and Mapping, Malaysia.

**REFERENCES**

- Cross, P.A., private communication, Dept. of Surveying, University of Newcastle upon Tyne, 1987.
- Heiskanen, W.A. and H. Moritz, Physical Geodesy. W.H. Freeman and Company, San Francisco and London, 1967.
- Lerch, F.J., F. Putney, S. Klosko and C. Wagner, Earth Models for Oceanographic Applications (GEM10B and 10C), Marine Geodesy, Vol. 5, 145-187, 1981.
- Rapp, R.H., The Earth Gravity Field to Degree and Order 180 Using SEASAT Altimeter Data, Terrestrial Gravity Data, and Other Data, Report No. 322, Dept. of Geodetic Science and Surveying, The Ohio State University, Columbus, 1981.
- Rapp, R.H., and Cruz, J.Y., Spherical Harmonic Expansions of the earth's Gravitational Potential to Degree 360 Using 30' Mean Anomalies, Report No. 376, Dept. of Geodetic Science and Surveying, The Ohio State University, Columbus, 1986.




Figure 4 : Malaysian Geopotential Geoid Computed From The OSU86 Model  
(C.I. = 1 metre, Ref. Ellipsoid = GRS80)






Figure 5 : Global Geopotential Geoid Computed From The GEM10B Model  
(C.I. = 5m Ref. Ellipsoid  $a = 6378138m$  and  $f = 1/298.257$ )