

NONLINEAR POLYMERIC
ARCHITECTURES VIA OLEFIN
METATHESIS

Thesis by

Irina A. Gorodetskaya

In Partial Fulfillment of the Requirements for the

degree of

Doctor of Philosophy

CALIFORNIA INSTITUTE OF TECHNOLOGY

Pasadena, California

2009

(Defended November 25, 2008)

© 2009

Irina A. Gorodetskaya

All Rights Reserved

To my grandmother Yadviga

ACKNOWLEDGEMENTS

First, I would like to thank my research advisor, Professor Grubbs, for accepting me into his group. Being a part of the Grubbs groups was everything I sought in a graduate school experience. I am very grateful to Bob for his seemingly infinite patience, which allowed me the time and resources to pursue the projects that were interesting to me at my own pace. Bob's guidance, combined with the consistently diverse and stimulating environment of the group, provided an invaluable learning experience on both scientific and personal levels, and it has made me a smarter, better, stronger person.

I am grateful to Dr. Tae-Lim Choi for believing in me and for being so persistently "annoying" with his suggestions on the potential of olefin metathesis chemistry for hyperbranched polymerization. This thesis would not be quite the same without the research that evolved from the TLC's nagging—thank you so much!

Next, I thank my most immediate collaborators over the years at Caltech. Lucia Fernandez-Ballister from the Kornfield group at Caltech and Jian Wang from the McKenna group at Texas Tech taught me everything I know about the rheology of polymer melts. They greatly contributed to the cyclic polymers research presented in Chapter 5 of this thesis (while making me secretly happy I did not choose chemical engineering as my major). Alon Gorodetsky provided his invaluable expertise in the study of pyrene-functionalized polymers described in Chapter 3. Finally, Katya Vinogradova worked very long hours on the hyperbranched project over the summer of 2007 as a summer undergraduate research fellow. Thank you all! I greatly benefited from your skill, knowledge, and time, and it has been a pleasure to know you and work with you.

I would also like to thank Alon again, Dr. Rose Conrad, and Dr. Ian Stewart for their time and help in proof-reading this thesis. I am sorry, but despite all of the language coaching you provided, I don't have enough fancy English words in my vocabulary to express my gratitude to you for reading it all (including the experimentals!) and providing numerous helpful suggestions. I also enjoyed sharing the lab with Rose and Ian, and our non-thesis- and even non-science-related(!) conversations.

I am very grateful to Professor Tobias Ritter, the most brilliant synthetic chemist I have ever met, for a number of key research suggestions, but also, and more importantly, for teaching me one of the most valuable skills I gained while in graduate school—driving. Having a car not only greatly enriched my life in Southern California, but also, finally, made me feel truly American. On this note, I would also like to thank Dr. Donde Anderson for risking her life as my diving buddy and navigator in my early driving days.

I am at loss of words, again, to describe the depth of my gratitude to my family, whether relatively close (within the US borders) or far abroad (Russia and Ukraine). Your unconditional love and support (moral and financial) made all the difference and got me through the grad school: mom, dad, grandma Lyuba, Natasha, tyetushka, Bobochka—I love you all. Special thanks go to cousin Pitrovich, who is currently the farthest away relative, but used to be the closest in San Francisco, where he was always happy to see me, even if for a few minutes of a practically unannounced, 3 am on a week-night visit of a camping gear emergency. Thank you for all your help, advice, and genuine interest in my well-being; and also for introducing me to Mitya, Olya, Fedor, Sasha, who became my adoptive Pasadena family.

I want to thank my thesis committee: Professor Tirrell, Professor Heath, and Professor Barton.

Finally, I wish to thank all of the friends and acquaintances, who are not singled out on this undeservingly short list, but whose friendships, love, care and kindness have touched my life over the past few years, making my not-so-short stay in Pasadena special and memorable.

ABSTRACT

The research presented in this thesis focuses on application of different forms of olefin metathesis, in conjunction with judicious choices of catalysts and monomers, to the construction of hyperbranched and cyclic macromolecules. This multifaceted reaction is briefly reviewed in Chapter 1, along with ruthenium-based metathesis catalysts and applications of olefin metathesis in polymer synthesis.

Hyperbranched polymers are curious materials which feature multiple end groups and possess a host of desirable physical properties; potential applications stemming from the unique properties of these macromolecules include their use as viscosity lowering additives and analyte carriers. In general, the major drawbacks faced by the classical, AB_n monomer-based hyperbranched polymers are the limited availability of specially designed monomers, harsh synthetic conditions, and poor control of the required step-growth polymerization methods. A very mild, simple, and modular, olefin metathesis-based hyperbranched polymerization route, which addresses some of these challenges, is presented in Chapter 2. This method utilizes the cross metathesis selectivity of the functional group tolerant N-heterocyclic carbene ruthenium catalyst towards different types of alkenes, and it can be applied to the polymerization of many easily prepared AB_n monomers. Moreover, the same method can be used to post-synthetically functionalize such polymers for realization of their substrate carrying potential. Chapter 3 describes one functionalization example—a pyrene analyte is attached to a metathesis derived hyperbranched polymer. This modification of the polymer provides insight into its solution structure relative to a linear analog. In addition, molecular weight control of the metathesis hyperbranched polymerization is discussed in detail in Chapter 4. The careful choice of the catalysts loading and the use of a multifunctional core are found to be important parameters in preparation of polymers which span a range of molecular weights.

Even well-established materials, such as polyethylene, can benefit from olefin metathesis and the unusual polymeric architectures it can efficiently create. For example, a cyclic polymer which lacks end groups, as opposed to having many end groups like a

hyperbranched polymer, is expected to possess unique physical properties. The preparation of cyclic and linear polyethylenes and the study of their relative rheological properties are described in Chapter 5. The polymerization methodology outlined in this Chapter takes advantage of ring-expansion metathesis polymerization—a facile method for the synthesis of cyclic macromolecules. Some efforts directed at molecular weight control of this cyclic polymerization are also discussed.

Taken together, the findings presented in this thesis emphasize the utility of olefin metathesis for the preparation of nonlinear polymers. The unusual polymeric architectures available through this chemical transformation may lead to a host of new materials with unique properties.

TABLE OF CONTENTS

CHAPTER 1: <i>Introduction to Olefin Metathesis and Its Applications in</i>	
<i>Polymer Synthesis</i>	1
Olefin Metathesis.....	2
Polymer Synthesis Applications of Olefin Metathesis.....	6
<i>ADMET</i>	6
<i>ROMP</i>	7
Summary and Thesis Research	8
References.....	11
CHAPTER 2: <i>Hyperbranched Polymers via Acyclic Diene Metathesis</i>	
<i>Polymerization</i>	14
Abstract.....	15
Introduction.....	16
Results and Discussion.....	17
Conclusion	23
Experimental Procedures.....	24
Representative Polymerization Procedure.....	31
References.....	33

CHAPTER 3: <i>An Olefin Metathesis Route to the Preparation of Functionalized Hyperbranched Polymers</i>	35
Abstract	36
Introduction.....	37
Results and Discussion	38
Functionalization of the Hyperbranched Polymer.....	38
Preparation of the Pyrene Modified Linear Analog	43
Fluorescence Properties of Pyrene-Functionalized Hyperbranched and Linear Polymers.....	45
Conclusion	46
Experimental Procedures.....	48
References.....	52
CHAPTER 4: <i>Towards Molecular Weight Control of the Hyperbranched ADMET Polymerization</i>	54
Abstract	55
Introduction.....	56
Results and Discussion	58
Catalyst Loading.....	58
Reaction Time	62
Addition of a Multifunctional Core	64
Conclusion	66
Experimental Procedures.....	67

References.....	68	
CHAPTER 5: <i>The Importance of Molecular Weight Control for Cyclic</i>		
<i>Polymers Prepared via Ring-Expansion Metathesis Polymerization</i>		69
Abstract.....	70	
Introduction.....	71	
Results and Discussion.....	72	
Part 1: Studies of shear-induced crystallization processes in polyethylene and the need for large cyclic polymers.....	72	
Part 2: Studies of the viscoelastic properties of cyclic polyethylene and the need for small cyclic polymers	77	
Conclusion	82	
Polymerization Experimental Procedures.....	84	
References.....	86	
Appendix A: <i>Towards AB_n-Based Hyperbranched Polyethylene</i>		88
Experimental Procedures.....	91	
Notes and References	93	

LIST OF FIGURES, SCHEMES AND TABLES

CHAPTER 1

Figures

Figure 1.1. Ruthenium-based olefin metathesis catalysts	5
---	---

Schemes

Scheme 1.1. General mechanism of olefin metathesis.....	2
Scheme 1.2. Types of olefin metathesis reactions.....	3
Scheme 1.3. Selectivity of 1 st and 2 nd generation ruthenium catalysts.	6
Scheme 1.4. Olefin metathesis routes to polyethylene of different architectures.....	9

Tables

Table 1.1. Functional group tolerance of olefin metathesis catalysts.	4
--	---

CHAPTER 2

Figures

Figure 2.1. Acyclic diene metathesis polymerization catalyst.....	18
Figure 2.2. Monomers for hyperbranched ADMET polymerization.....	18
Figure 2.3. ¹ H NMR spectra of monomer 4 and hyperbranched polymer 4a	19
Figure 2.4. Mark-Houwink-Sakurada plots for polymers 2a–7a	21
Figure 2.5. Representative MALS-SEC traces for polymerization of 4 with consecutive batch catalyst addition.	32

Schemes

Scheme 2.1. Synthesis of dendritic polymers.....	16
Scheme 2.2. Synthesis of AB ₂ monomer 4 and its hyperbranched ADMET polymerization.	18

Scheme 2.3. AB ₂ monomers which were found to be challenging for ADMET.....	22
Scheme 2.4. Synthesis of AB ₄ monomer for hyperbranched ADMET polymerization.....	23
Scheme 2.5. Synthesis of monomer 2	24
Scheme 2.6. Synthesis of monomers 3 and 5	25
Scheme 2.7. Synthesis of monomer 6	27
Scheme 2.8. Synthesis of monomer 7	28
Scheme 2.9. Synthesis of monomer 8	30
Scheme 2.10. Synthesis of monomer 9	30

Tables

Table 2.1. Results for polymerization of 2–7	20
---	----

CHAPTER 3

Figures

Figure 3.1. Imidazolynylidene-based ruthenium olefin metathesis catalyst 1	38
Figure 3.2. SEC traces for 3 before and after functionalization with 0.5 equivalents of 10-bromo-1-decene.....	40
Figure 3.3. ¹ H NMR spectra with integration values for 2 , 3 , and 5	42
Figure 3.4. SEC traces for crude 3 , crude and purified 5	43
Figure 3.5. ¹ H NMR spectra with integration values for 6 , 7 , and 8	44
Figure 3.6. (A) UV-visible absorbance and fluorescence emission spectra for 4 , 5 , and 8 . (B) A plot of the monomer to excimer intensity emission ratio at various concentrations.....	46

Schemes

Scheme 3.1. Hyperbranched ADMET polymerization	
--	--

and subsequent end group functionalization.	39
Scheme 3.2. Hyperbranched polymer 3 functionalization with pyrene....	41
Scheme 3.3. Synthesis of the pyrene-functionalized linear polymeric analog.....	44
Scheme 3.4. Synthesis of the monomers for linear ROMP.....	49

CHAPTER 4

Figures

Figure 4.1. ADMET catalyst 1	57
Figure 4.2. SEC traces for 3 made with different amounts of 1	58
Figure 4.3. SEC traces for 5 made with different amounts of 1	60
Figure 4.4. Molecular weight and PDI timeline of 3 at 0.5 mol % of 1 . ..	62
Figure 4.5. Proposed polymerization “error” correction mechanism.	63
Figure 4.6. The dependence of the observed M_w at 0.5 mol % of 1 on the addition time of 8-bromo-1-octene.....	64
Figure 4.7. SEC traces for 3 made with a fixed amount of 1 but different amounts of 4	65

Schemes

Scheme 4.1. Hyperbranched ADMET polymerization.....	57
Scheme 4.2. Linear ADMET polymerization.....	59
Scheme 4.3. The “chain-walking” mechanism for hyperbranched ADMET polymerization at low loading of 1	61

CHAPTER 5

Figures

Figure 5.1. The effect of shear stress on polymer chain orientation.	73
Figure 5.2. Catalysts required for the preparation of linear and cyclic polyalkenamers.....	74
Figure 5.3. Polarized optical microscopy of selected samples.....	77

Figure 5.4. The results of the viscoelastic properties investigation of cyclic PE prepared by REMP.....	79
Figure 5.5. Proposed 2 nd generation REMP catalysts.....	80
Figure 5.6. First accomplished 2 nd generation REMP catalyst.....	83

Schemes

Scheme 5.1. Ring-expansion polymerization.....	72
Scheme 5.2. Synthesis of long and short linear PE chains.....	75
Scheme 5.3. Synthesis of large PE rings.....	76
Scheme 5.4. Synthetic route towards 2 nd generation REMP catalyst 4	81
Scheme 5.5. Synthetic route towards 2 nd generation REMP catalyst 5	82

Tables

Table 5.1. Results for REMP of cyclic alkenes with 3	76
---	----

Appendix A

Figures

Figure A1. Hyperbranched ADMET catalysts	90
Figure A2. H NMR evidence for polymerization of 3 to 4	90

Schemes

Scheme A1. Synthetic route towards the hyperbranched polyethylene via ADMET.....	90
---	----

