



UNIVERSITY
OF TRENTO

DEPARTMENT OF INFORMATION AND COMMUNICATION TECHNOLOGY

38050 Povo – Trento (Italy), Via Sommarive 14
<http://www.dit.unitn.it>

INDECOMPOSABLE r -GRAPHS AND SOME
OTHER COUNTEREXAMPLES

Romeo Rizzi

September 1999

Technical Report # DIT-02-0052

Also: in *Journal of Graph Theory* 32 (1) (1999) 1--15.

Indecomposable r -graphs and some other counterexamples

Romeo Rizzi*

September 13, 1999

**Dipartimento di Matematica Pura ed Applicata, Università di Padova*
Via Belzoni 7, 35131 Padova, Italy
romeo@euler.math.unipd.it

Abstract

An r -graph is any graph that can be obtained as a conic combination of its own 1-factors. An r -graph $G(V, E)$ is said *indecomposable* when its edge set E cannot be partitioned as $E = E_1 \cup E_2$ so that $G_i(V, E_i)$ is an r_i -graph for $i = 1, 2$ and for some r_1, r_2 . We give an indecomposable r -graph for every integer $r \geq 4$. This answers a question raised in [11, 12] and has interesting consequences for the Schrijver System of the T -cut polyhedron to be given in [9]. A graph in which every two 1-factors intersect is said to be *poorly matchable*. Every poorly matchable r -graph is indecomposable. We show that for every $r \geq 4$ "being indecomposable" does not imply "being poorly matchable". Next we give a poorly matchable r -graph for every $r \geq 4$. The paper provides counterexamples to some conjectures of Seymour [11, 12].

Key words: r -graph; indecomposable; Petersen graph; Fulkerson Coloring.

1 Introduction

In this article graphs may have parallel edges but contain no loop. The set of edges with precisely one endpoint in S is denoted by $\partial(S)$. To specify the graph, say G , we write $\partial_G(S)$. Let r be a positive integer. The notion of r -graph is due to Seymour [12]: an r -graph is a regular graph of valency r such that $|\partial(S)| \geq r$ for every set of nodes S with $|S|$ odd.

We rely on standard notation $d(S) = |\partial(S)|$ and $d_G(S) = |\partial_G(S)|$. Moreover $\partial(v) = \partial(\{v\})$ and $d(v) = d(\{v\})$. If G is an r -graph then G has an even number of nodes, since $d(V(G)) = 0$.

Given a graph G , a *1-factor* of G is a spanning subgraph of G which is a 1-graph.

The celebrated Edmonds' matching polytope theorem [1] states that for every graph G the vertices of the following polytope are integral:

$$\begin{cases} x_e \geq 0 & \forall e \in E(G) \\ x(\partial(v)) = 1 & \forall v \in V(G) \\ x(\partial(S)) \geq 1 & \forall S \subseteq V(G) \text{ with } |S| \text{ odd} \end{cases} \quad (1)$$

Seymour [12] observed that Edmonds' theorem is equivalent to the following statement: a graph G is an r -graph if and only if G can be obtained as a conic combination of its own

1-factors. As a consequence, for every r -graph G and for every edge e of G , there exists a 1-factor of G containing e (see [12, 6]).

Let $G_1(V, E_1), \dots, G_k(V, E_k)$ be graphs on a common node set V but with disjoint edge sets E_1, \dots, E_k . We denote by $G_1 + \dots + G_k$ the graph $G(V, E_1 \cup \dots \cup E_k)$ and say that G is the *sum* of G_1, \dots, G_k . Note that if $G_i(V, E_i)$ is an r_i -graph for $i = 1, \dots, k$ then $G_1 + \dots + G_k$ is an $(r_1 + \dots + r_k)$ -graph. For $k \in \mathbb{N}$, we denote by kG the graph obtained by summing up k copies of G (that is, replacing every edge of G by k parallel edges).

An *unslicable* r -graph is an r -graph, which cannot be expressed as the sum of an $(r - 1)$ -graph and a 1-factor. An r -graph G , which can be expressed as the sum of an r_1 -graph and an r_2 -graph, is said to be *decomposable*. When no such decomposition exists G is called *indecomposable*. Finally, a graph in which every two 1-factors intersect is called *poorly matchable*.

In [11, 12], Seymour raised the following question:

Question 1 *Does there exist a constant \bar{r} such that every unslicable r -graph has $r < \bar{r}$?*

In the same articles, he proposed the following.

Conjecture 1.1 *The answer to Question 1 is positive and in fact we can take $\bar{r} = 4$.*

Conjecture 1.1 implies Conjecture 1.2 and makes Conjecture 1.3 imply Conjecture 1.4.

Conjecture 1.2 (Seymour [12]) *Every r -graph is $r + 1$ edge colorable.*

A *Fulkerson coloring* of an r -graph G is a decomposition of $2G$ into 1-factors.

Conjecture 1.3 (Berge-Fulkerson) *Every 3-graph has a Fulkerson coloring.*

Conjecture 1.4 (Seymour [12]) *Every r -graph has a Fulkerson coloring.*

The author, while working on a bound for the size of the coefficients in the Schrijver System for the T -cut polyhedron (see [9]), became interested in the following question.

Question 2 *Does there exist a constant \bar{r} such that every r -graph with $r \geq \bar{r}$ is decomposable?*

This article presents a counterexample to Conjecture 1.1. In fact, we settle Question 2 (and, hence, Question 1) in the negative by constructing for every r an indecomposable r -graph. More surprisingly, we exhibit for every r a poorly matchable r -graph.

2 Preliminary Observations

The Petersen graph is the 3-graph \mathcal{P} shown in Fig. 1. The six 1-factors of \mathcal{P} are all equivalent under isomorphisms of \mathcal{P} . Let M be a 1-factor of \mathcal{P} . The essentially unique r -graph $\mathcal{P}(r) = \mathcal{P} + (r - 3)M$, shown for $r = 4$ in Fig. 1 and dating back to Meredith (see [7]), acts as a fundamental component in three of the constructions presented in this article.

Every edge of \mathcal{P} belongs to precisely two distinct 1-factors of \mathcal{P} . Conversely, every two of the six 1-factors of \mathcal{P} have precisely one edge in common. This is expressed more formally by the following proposition.




Figure 1: The Petersen Graph \mathcal{P} and the Meredith Graph $\mathcal{P}(4)$.

Proposition 2.1 *Associate to every pair of 1-factors of \mathcal{P} the edge they have in common. This is a one to one correspondence between edges and pairs of distinct 1-factors.*

Proof: The Petersen graph can be defined as follows (see [3]): the nodes of \mathcal{P} are the pairs of elements in $\mathbb{N}_5 = \{1, 2, 3, 4, 5\}$, where two nodes $\{i, j\}$ and $\{h, k\}$ of \mathcal{P} are adjacent if and only if they are disjoint. Thus \mathcal{P} is a regular graph of valency $\binom{5-2}{2} = 3$ with $\binom{5}{2} = 10$ nodes and $\frac{10 \cdot 3}{2} = 15$ edges.

Let $e = \{i, j\}\{h, k\}$ be any edge of \mathcal{P} and let x be the single element in $\mathbb{N}_5 \setminus \{i, j, h, k\}$. A 1-factor containing e can not contain any other edge with an endpoint in $\{i, j\}$ or $\{h, k\}$. Moreover, no 1-factor containing e can contain $\{i, h\}\{j, k\}$ or $\{i, k\}\{j, h\}$. Indeed, assume on the contrary and without loss of generality to have a 1-factor containing both e and $\{i, h\}\{j, k\}$. Then nodes $\{j, x\}$ and $\{h, x\}$ are both matched with node $\{i, k\}$, a contradiction.

We conclude that any 1-factor containing e has precisely 4 edges among the remaining 8 edges. These 8 edges form a circuit, whose 8 nodes appear in the following order:

$$\{j, k\}, \{i, x\}, \{j, h\}, \{k, x\}, \{i, h\}, \{j, x\}, \{i, k\}, \{h, x\}$$

Therefore, the 1-factors of \mathcal{P} containing e are precisely two and have no edge in common other than e .

The number of distinct 1-factors of \mathcal{P} is, therefore, $\frac{2|E(\mathcal{P})|}{5} = 6$. The function which associates to each edge e of \mathcal{P} the pair of 1-factors containing e is injective, for we said that the two 1-factors have no edge in common other than e . Since $|E(\mathcal{P})| = 15 = \binom{6}{2}$ is the number of pairs of 1-factors, the function is also surjective and bijective. \square

An immediate consequence of Property 2.1 is the following.

Lemma 2.2 *Let M_1, M_2 be two edge-disjoint 1-factors of $\mathcal{P}(r) = \mathcal{P} + (r-3)M$. Then either $M_1 = M$ or $M_2 = M$.*

The following lemma is involved in a first construction of indecomposable r -graphs.

Lemma 2.3 *Assume $\mathcal{P}(r) = G_1 + G_2$, where, for $i = 1, 2$, G_i is an r_i -graph. Then there exist k_1, k_2 such that $G_1 = \mathcal{P} + k_1M$ and $G_2 = k_2M$ or vice versa.*

Proof: It suffices to show that $\mathcal{P} \setminus M$ is contained in either G_1 or G_2 . Assume the contrary and let e_i ($i = 1, 2$) be an edge of G_i contained in $\mathcal{P} \setminus M$. Let M_i be a 1-factor of G_i containing

e_i . Thus M_1 and M_2 are two edge-disjoint 1-factors of $\mathcal{P}(r)$ contradicting Lemma 2.2. \square

A *tight cut* in an r -graph is an edge set of the form $\partial(S)$ where S is a set of nodes of odd cardinality and $d(S) = r$. The following proposition plays a central role in proving that the graphs to be constructed in the next section are indecomposable.

Proposition 2.4 *Let $\partial(S)$ be a tight cut in an r -graph G . Then the graph G^* obtained from G by identifying all nodes in S is an r -graph.*

Assume $G = G_1 + \dots + G_h$ where G_i is an r_i -graph ($i = 1, \dots, h$). Note that $d_{G_i}(S) = r_i$ ($i = 1, \dots, h$). Let G_1^, \dots, G_h^* be the graphs obtained from G_1, \dots, G_h by identifying all nodes in S . As above G_i^* is an r_i -graph ($i = 1, \dots, h$). Moreover $G^* = G_1^* + \dots + G_h^*$.*

Lemma 2.5 *Let G be a graph and $S \subseteq V(G)$ with $|S|$ odd. Let G_S and $G_{\overline{S}}$ be the graphs obtained from G by identifying all nodes in S and in $\overline{S} = V(G) \setminus S$ respectively. If G_S and $G_{\overline{S}}$ are both r -graphs then G is an r -graph.*

Proof: Obviously G is r -regular. Assume $d_G(X) < r$ and $|X|$ odd. Exchanging S and \overline{S} , if necessary, we can assume that $|X \cap S|$ and $|X \cup S|$ are odd. By submodularity of d_G , $d_G(X \cap S) + d_G(X \cup S) \leq d_G(X) + d_G(S) < r + r$. So, either $d_{G_{\overline{S}}}(X \cap S) = d_G(X \cap S) < r$ or $d_{G_S}(X \cup S) = d_G(X \cup S) < r$ contrary to the assumption that both G_S and $G_{\overline{S}}$ are r -graphs. \square

3 An infinite family of counterexamples

Let r be any integer with $r \geq 4$. In this section we construct an unslicable r -graph $U(r)$.

By Lemma 2.3, the only way to decompose the r -graph $\mathcal{P}(r) = \mathcal{P} + (r - 3)M$ into an $(r - 1)$ -graph and a 1-factor is $\mathcal{P}(r) = \mathcal{P}(r - 1) + M$. Let $e = uv$ be any edge of \mathcal{P} which is not in M (all such edges are equivalent by symmetry). Take r distinct copies C_1, \dots, C_r of $\mathcal{P}(r) \setminus e$. For $i = 1, \dots, r$, copy C_i contains two nodes of degree $r - 1$, namely u_i and v_i . Let x and y be two nodes not belonging to $V(C_i)$ for any i . The r -graph $U(r)$ is obtained from the components C_1, \dots, C_r and the nodes x, y by adding all edges xu_i and yv_i for $i = 1, \dots, r$.

When $G = G_1 + G_2$ we say that G_2 is the *complement* of G_1 in G .

Claim 3.1 *The r -graph $U(r)$ is unslicable.*

Proof: Any 1-factor F of $U(r)$ contains an edge incident with x . Assume without loss of generality that $xu_1 \in F$. For parity reasons, $yv_1 \in F$. Therefore, $F \cap E(C_1) + u_1v_1$ is a 1-factor of $C_1 + u_1v_1$. Moreover, the complement of $F \cap E(C_1) + u_1v_1$ in $C_1 + u_1v_1$ is an $(r - 1)$ -graph. Apply Lemma 2.3. \square

Evidently, "being indecomposable" is a stronger property than "being unslicable". Since every 2-graph is decomposable, the two properties are equivalent for $r < 6$. To prove them to be distinct for every $r \geq 6$, we show that the unslicable r -graph $U(r)$ is decomposable whenever $r \geq 6$. Indeed, $U(r) = G_1(r) + G_2(r)$, where $G_1(r)$ is the 3-graph collecting a copy

of \mathcal{P} from components C_1, C_2 and C_3 and a $3M$ from every other component. Also $G_2(r)$, which results as the complement of $G_1(r)$ in $U(r)$, is an $(r-3)$ -graph.

4 The indecomposable r -graph $G(r)$

Let r be any integer with $r \geq 4$. In this section, we construct an indecomposable r -graph $G(r)$.

Let z be any node of $\mathcal{P}(r) = \mathcal{P} + (r-3)M$ (all nodes are equivalent under isomorphism). Let x, a, b be the neighbors of z , where $xz \in M$. We indicate with $\langle a, x, b \rangle^{(r)}$ the graph obtained from $\mathcal{P}(r)$ by removing node z . Symbolic representation of $\langle a, x, b \rangle^{(r)} = \mathcal{P}(r) \setminus z$ is indicated in Fig. 2.




Figure 2: $\mathcal{P}(r) \setminus z = \langle a, x, b \rangle^{(r)}$.

Let $\langle a_1, x_1, b_1 \rangle_1^{(r)}, \dots, \langle a_r, x_r, b_r \rangle_r^{(r)}$ be distinct copies of $\langle a, x, b \rangle^{(r)}$. Define C to be the set of edges $\{b_i a_{i+1} : i = 1, \dots, r-1\} \cup \{b_r a_1\}$. Let $\{v_1, \dots, v_{r-2}\}$ be a set of nodes disjoint from all the $V(\langle a_i, x_i, b_i \rangle_i^{(r)})$. For $i = 1, \dots, r-2$ define E_i as the set of edges $\{v_i x_j : j = 1, \dots, r\}$. The graph $G(r)$ is obtained from the components $\langle a_1, x_1, b_1 \rangle_1^{(r)}, \dots, \langle a_r, x_r, b_r \rangle_r^{(r)}, v_1, \dots, v_{r-2}$ by adding all the edges in $C \cup E_1 \cup E_2 \cup \dots \cup E_{r-2}$.

For example $G(4)$ and $G(5)$ are shown in Fig. 3.

By Lemma 2.5, $G(r)$ is an r -graph.

Claim 4.1 *For every integer $r \geq 4$, $G(r)$ is indecomposable.*

Proof: Assume $G(r) = G_1 + G_2$ with G_1 r_1 -graph and G_2 r_2 -graph. By Lemma 2.3 and Proposition 2.4, either $C \subseteq G_1$ or $C \subseteq G_2$. Assume without loss of generality that $C \subseteq G_1$. Proposition 2.4 implies:

$$\begin{aligned} |E(G_2) \cap (E_1 \cup \dots \cup E_{r-2})| &= \sum_{i=1}^r d_{G_2 \setminus C}(V(\langle a_i, x_i, b_i \rangle_i^{(r)})) = \\ &= \sum_{i=1}^r d_{G_2}(V(\langle a_i, x_i, b_i \rangle_i^{(r)})) = rr_2 \end{aligned} \quad (2)$$

However, $|E(G_2) \cap E_i| = d_{G_2}(v_i) = r_2$ ($i = 1, \dots, r-2$) implies $|E(G_2) \cap (E_1 \cup \dots \cup E_{r-2})| = (r-2)r_2$, in contradiction with (2). We conclude that $G(r)$ is indecomposable. \square




Figure 3: Graphs $G(4)$ and $G(5)$.

5 More indecomposable r -graphs

Let G_1, G_2 be two node-disjoint r -graphs. Choose $v_1 \in V(G_1)$ and $v_2 \in V(G_2)$. Let G be any r -regular graph obtained from G_1 and G_2 by first removing nodes v_1, v_2 , and then adding some new edges with one endpoint in $V(G_1) \setminus \{v_1\}$ and the other in $V(G_2) \setminus \{v_2\}$. We say that G has been obtained by *splicing* G_1 and G_2 (at v_1, v_2). Proposition 2.4 and Lemma 2.5 imply the following.

Lemma 5.1 *If G has been obtained by splicing two r -graphs G_1 and G_2 , then G is an r -graph. Moreover, if G_1 is indecomposable, then G is indecomposable.*

Hence, we have an infinite number of indecomposable r -graphs for any given integer $r \geq 4$. Let K_n be the complete graph on n nodes. When r is odd, then $S_r = K_{r+1}$ is a *simple* (no parallel edges) r -graph. When r is even, then let M be any matching of K_{r+1} with $|M| = \frac{r}{2}$. Let S_r be the graph obtained from K_{r+1} by first subdividing every edge in M into two edges, and next identifying all nodes of degree two so introduced. Again S_r is a simple r -graph. To obtain a simple indecomposable r -graph, start from any indecomposable r -graph and, while some parallel edges are incident with a node x , splice at x with some simple r -graph like S_r .

The smallest indecomposable r -graphs (for $r = 4, 5, 6$), we were able to construct, are given in Fig. 4. The first graph in Fig. 4 is, in fact, the smallest possible counterexample to Conjecture 1.1 (see [8]).

6 Poorly matchable r -graphs: a recursive construction

An r -graph G is said to be *poorly matchable* if G does not contain two disjoint 1-factors. Since every r -graph has a 1-factor, every poorly matchable r -graph is indecomposable. Thus, "being poorly matchable" is a stronger property than "being indecomposable". For $r = 3$, the two properties are equivalent, because the presence of two disjoint 1-factors implies 3




Figure 4: Small indecomposable r -graphs.

edge colorability. However, the two properties are distinct for every $r \geq 4$. This is proven in Subsection 6.1 by showing that, for all $r \geq 4$, the indecomposable r -graph $G(r)$ from Section 4 has two edge-disjoint 1-factors.

Subsection 6.2 gives a poorly matchable r -graph G^r for every integer $r \geq 4$. The construction we propose is, however, recursive, and the size of G^r is probably exponential in r . (Whereas, for $G(r)$, we have $|V(G(r))| = (10 - 1)r + (r - 2) = 10r - 2$, which is linear in r).

The following three statements are equivalent for an r -graph G : (i) G is poorly matchable; (ii) G does not contain a spanning 2-graph; (iii) G does not contain two disjoint spanning r -graphs.

Therefore, the existence of a poorly matchable r -graph for every integer $r \geq 3$ has the following consequence.

Proposition 6.1 *There exists no constant K such that every r -graph with $r > K$ can be expressed as the sum of a K -regular graph and an $(r - K)$ -graph.*

Proof: For any given $K \in \mathbb{N}$, consider a poorly matchable $(K + 2)$ -graph. □

We propose the following conjecture.

Conjecture 6.2 *Every $3r$ -graph is the sum of r 3-regular graphs.*

6.1 Two edge-disjoint 1-factors in $G(r)$

This subsection gives two edge-disjoint 1-factors in $G(r)$: $M_1(r)$ and $M_2(r)$. To specify $M_1(r)$ and $M_2(r)$, we rely on the description of $G(r)$ in Section 4.

- $M_1(r) \cap C = \{b_{r-1}a_r\}$, whereas $M_2(r) \cap C = \{b_1a_2\}$.
- For $1 \leq i \leq r - 2$ $\partial_{M_1(r)}(v_i) = v_i x_i$ whereas $\partial_{M_2(r)}(v_i) = v_i x_{i+2}$.
- For $3 \leq i \leq r - 2$ and $j = 1, 2$ $M_j(r) \cap E(\langle a_i, x_i, b_i \rangle_i^{(r)})$ is a copy of M with z removed. (Remember $\langle a, x, b \rangle^{(r)}$ is $\mathcal{P}(r) = \mathcal{P} + (r - 3)M$ with a node z removed).

It remains to determine $M_1(r)$ and $M_2(r)$ on $E(\langle a_r, x_r, b_r \rangle_r^{(r)}) \cup E(\langle a_{r-1}, x_{r-1}, b_{r-1} \rangle_{r-1}^{(r)})$ and $E(\langle a_1, x_1, b_1 \rangle_1^{(r)}) \cup E(\langle a_2, x_2, b_2 \rangle_2^{(r)})$: both of them are described by Fig. 5.




Figure 5: $M_1(r)$ and $M_2(r)$ on the first and last two $\langle a, x, b \rangle^{(r)}$ components.

As an example, Fig. 6 shows $M_1(r)$ and $M_2(r)$ in $G(r)$ for $r = 4, 5$.




Figure 6: M_1 and M_2 in $G(4)$ and $G(5)$.

6.2 Constructing the poorly matchable r -graph G^r

The Petersen graph \mathcal{P} is an indecomposable 3-graph and, hence, a poorly matchable 3-graph. Thus, let $G^3 = \mathcal{P}$. This subsection shows how to construct a poorly matchable r -graph G^r from a poorly matchable $(r - 1)$ -graph G^{r-1} whenever $r \geq 4$.

Let G be an r -graph, and let $e, f = uv$ be two parallel edges of G . Take a copy of $\mathcal{P}(r)$ node-disjoint from G and choose a node z in $\mathcal{P}(r)$. Let x be the node of $\mathcal{P}(r)$ joined to z by $r - 2$ edges, and let a, b be the other two nodes of $\mathcal{P}(r)$ adjacent to z . Remove node v from G and node z from $\mathcal{P}(r)$. Next, add edges ua and ub . Finally, add a set of edges with one endpoint in $V(\mathcal{P}(r)) \setminus \{z\}$ and the other in $V(G) \setminus \{v\}$ to obtain an r -regular graph G^* . We say that G^* is obtained from G by \mathcal{P} -splicing at v distinguishing e and f .

Note that \mathcal{P} -splicing is a particular instance of the splice operation defined in Section 5. Hence, by Lemma 5.1, G^* is an r -graph. Moreover, we have the following.

Lemma 6.3 *If G^* has two edge-disjoint 1-factors, then G has two edge-disjoint 1-factors M_1 and M_2 such that $\{e, f\} \not\subseteq M_1 \cup M_2$.*

Proof: Let M_1^*, M_2^* be two edge-disjoint 1-factors of G^* . Let $K = \partial_{G^*}(V(\mathcal{P}(r)) \setminus \{z\})$ denote the set of edges which have been added by \mathcal{P} -splicing. Since $|V(\mathcal{P}(r)) \setminus \{z\}| = 9$ is odd, then $|M_1^* \cap K|$ and $|M_2^* \cap K|$ are both odd. Hence, $|M_1^* \cap K|, |M_2^* \cap K| \geq 1$. In fact, $|M_1^* \cap K|, |M_2^* \cap K| = 1$, since all edges in K are incident either with x or with u . Therefore, after identifying all nodes in $V(\mathcal{P}(r)) \setminus \{z\}$, M_1^* and M_2^* become two edge-disjoint 1-factors M_1 and M_2 of G .

If $\{e, f\} \subseteq M_1 \cup M_2$, then $\{e, f\} \subseteq M_1^* \cup M_2^*$ and, after identifying in G^* all nodes in $V(G) \setminus \{v\}$, M_1^* and M_2^* become two edge-disjoint 1-factors of $\mathcal{P}(r)$ contradicting Lemma 2.2. \square

We are now ready for the recursive construction: Let G^{r-1} be a poorly matchable $(r-1)$ -graph. Let M be a 1-factor of G^{r-1} . Then $H^r = G^{r-1} + M$ is an r -graph. Let \overline{M} be the set of those edges of G^{r-1} that have multiplicity 1 in G^{r-1} and 2 in H^r . (M will stand for the edges in $H^r \setminus G^{r-1}$). Let $V_{\overline{M}}$ be a node cover for \overline{M} with $|V_{\overline{M}}| = |\overline{M}|$. Obtain G^r from H^r by \mathcal{P} -splicing at every node $\overline{v} \in V_{\overline{M}}$ distinguishing the unique edge in $\partial_M(\overline{v})$ and the unique edge in $\partial_{\overline{M}}(\overline{v})$. If G^r has two edge-disjoint 1-factors, then, by Lemma 6.3, H^r has two edge-disjoint 1-factors M_1 and M_2 with $(M_1 \cup M_2) \cap (M \cup \overline{M})$ having no parallel edges. But then, by eventually substituting the edges in M with those in \overline{M} having the same endpoints, M_1 and M_2 are two edge-disjoint 1-factors of G^{r-1} . We conclude that G^r is a poorly matchable r -graph, as in Fig. 7.




Figure 7: The poorly matchable 4-graph G^4 .

The construction proposed is not deterministic. Non-isomorphic r -graphs can, in fact, be obtained starting from a same $(r-1)$ -graph. From \mathcal{P} , however, a sole graph G^4 can be derived. Graph G^4 has 50 nodes. We have found no poorly matchable 4-graph on less than

50 nodes.

7 Avoiding tight cuts

All the unslicable, indecomposable, or poorly matchable r -graphs seen until now contain some tight cuts. This section gives a poorly matchable 4-graph without tight cuts as a counterexample to the following conjectures.

Conjecture 7.1 *Every unslicable r -graph with $r \geq 4$ has a tight cut.*

Conjecture 7.2 *Every indecomposable r -graph with $r \geq 4$ has a tight cut.*

Conjecture 7.1 is still strong enough to imply Conjecture 1.2. Conjecture 7.2 is still strong enough to make Conjecture 1.3 imply Conjecture 1.4.




Figure 8: The node gadget.

We employ a technique with some similarities to *superposition*. Superposition is a method for constructing snarks introduced in [4, 5] as a practical and effective means for capturing and exploiting “global type conditions” as suggested in [2].

The idea is to take a poorly matchable 3-graph, like \mathcal{P} , as skeleton. Next, every node in the skeleton is replaced by a distinct copy of the “node gadget” shown in Fig. 8 and every edge in the skeleton is replaced by a distinct copy of the “edge gadget” shown in Fig. 9.




Figure 9: The edge gadget.

The skeleton acts like a map, telling how edge and node gadgets are mutually connected. The resulting graph G_4 is shown in Fig. 10.




Figure 10: A poorly matchable 4-graph without tight cuts.

One can check that G_4 is a 4-graph without tight cuts. Let M_1 and M_2 be two edge-disjoint 1-factors of G_4 . Let $\phi : E(G_4) \mapsto \{(0, 0), (0, 1), (1, 0)\}$ be defined as follows:

$$\phi(e) = \begin{cases} (1, 0) & \text{if } e \in M_1 \\ (0, 1) & \text{if } e \in M_2 \\ (0, 0) & \text{if } e \notin M_1 \cup M_2 \end{cases}$$

When $F \subseteq E(G_4)$ we define $\phi(F) = \sum_{e \in F} \phi(e)$, where the sum is componentwise and modulo 2. Then ϕ satisfies the following conditions:

EVEN SET: Let S be an even set of nodes. Then $\phi(\partial(S)) = (0, 0)$.

ODD SET: Let S be an odd set of nodes. Then $\phi(\partial(S)) = (1, 1)$.

EDGE GADGET: $\phi(\{e_1, e_2\}) = \phi(\{f_1, f_2\}) \neq (1, 1)$.

Proof: Edge gadgets contain an even number of nodes. Hence $\phi(\{e_1, e_2\}) = \phi(\{f_1, f_2\})$ by

the Even Set Condition. Moreover, $\phi(\{e_1, e_2\}) = \phi(\{f_1, f_2\}) \neq (1, 1)$ by Lemma 2.2. \square

NODE GADGET: $\{\phi(\{x_1, x_2\}), \phi(\{y_1, y_2\}), \phi(\{z_1, z_2\})\} = \{(0, 0), (0, 1), (1, 0)\}$.

Proof: By the Odd Set Condition, $\phi(\{x_1, x_2\}) + \phi(\{y_1, y_2\}) + \phi(\{z_1, z_2\}) = (1, 1)$. By the Edge Gadget Condition, $\phi(\{x_1, x_2\}), \phi(\{y_1, y_2\}), \phi(\{z_1, z_2\}) \neq (1, 1)$. Thus, $(1, 0), (0, 1) \in \{\phi(\{x_1, x_2\}), \phi(\{y_1, y_2\}), \phi(\{z_1, z_2\})\}$.

But then $\{\phi(\{x_1, x_2\}), \phi(\{y_1, y_2\}), \phi(\{z_1, z_2\})\} = \{(0, 0), (0, 1), (1, 0)\}$. \square

Let e be any edge of \mathcal{P} . Let e_1, e_2 be two edges of G_4 entering the edge gadget relative to e on a same side. Define $\phi'(e) = \phi(\{e_1, e_2\})$. By the Edge Gadget Condition, ϕ' is well defined. By the Edge Gadget and Node Gadget Conditions, ϕ' is a coloring of the edges of \mathcal{P} by colors $(0, 0)$, $(0, 1)$, and $(1, 0)$. Since \mathcal{P} is not 3 edge colorable, G_4 is poorly matchable.

We recall that a *Fulkerson coloring* of an r -graph G is a decomposition of $2G$ into 1-factors.

Observation 7.3 *Let H be an indecomposable 3-graph and let H_4 be the poorly matchable 4-graph obtained from H as skeleton graph through the above described construction with Node and Edge Gadgets as in Figs. 8 and 9. From a Fulkerson coloring for H , one can derive a Fulkerson coloring for H_4 as shown in Fig. 11.*




Figure 11: Deriving a Fulkerson coloring for H_4 from one for H .

8 Poorly joinable r -graphs: a positive result

Given a graph G , a *join* of G is a set of edges $J \subseteq E(G)$ such that an odd number of edges in J is incident with each node in $V(G)$. An r -graph G is *poorly joinable*, if every two joins of G intersect. By definition, "poorly joinable" \Rightarrow "poorly matchable". For a 3-graph, the two properties are equivalent. In an early attempt of extending the construction proposed in the previous section and obtain poorly matchable r -graphs without tight cuts for $r > 4$ we found ourselves looking for poorly joinable r -graphs with $r > 3$. This approach ended in the following proposition.

Proposition 8.1 *There exists no poorly joinable r -graph for $r > 3$.*

Proof: Let G be an r -graph with $r > 3$. If r is even, then let M be any 1-factor of G , and observe that M and $G \setminus M$ are two disjoint joins of G . So r is odd, and we want to prove that G is the sum of three disjoint joins of G . We can assume that G is 4-edge connected, because 2-edge cuts give rise to an easy decomposition of the problem. By [14], G contains

two disjoint spanning trees. So G contains two disjoint joins. \square

In our opinion, the next item which makes sense to attempt to pack into r -graphs are joins. To stress this belief, we pose the following question.

Question 3 *Which functions $f(r)$ are there such that every r -graph with $r \geq \bar{r}$ admits $f(\bar{r})$ disjoint joins? Could $f(r) = \lfloor \frac{r}{2} \rfloor$ possibly work? What about $f(r) = r - 2$?*

The above question becomes even more relevant in view of its extension to grafts by arguments as given in [8].

9 Open Problems

The Petersen graph seems quite unavoidable in all our counterexamples. This suggests generalizing Tutte's conjecture as follows.

Conjecture 9.1 *Every indecomposable r -graph has a Petersen minor.*

Sebő pointed out that Conjecture 9.1 is equivalent to the following conjecture of Lovász.

Conjecture 9.2 *The 1-factors of a graph with no Petersen minor form a Hilbert basis.*

The following questions are left open.

Question 4 *Does there exist a constant \bar{r} such that every unslicable r -graph with $r \geq \bar{r}$ contains some tight cuts?*

Question 5 *Does there exist a constant \bar{r} such that every indecomposable r -graph with $r \geq \bar{r}$ contains some tight cuts?*

We propose the following.

Conjecture 9.3 *The answer to Question 4 is positive and in fact we can take $\bar{r} = 5$.*

In [12], Seymour mentioned to have proven Conjecture 1.2 for $r \leq 6$. In [13], Seymour gave a second proof that Conjecture 1.2 holds for $r \leq 6$. In fact, as a consequence of the approximation algorithm to edge color multigraphs described in [10], Conjecture 1.2 holds for $r \leq 12$. Therefore, a positive answer to Question 5 with $\bar{r} \leq 13$ would imply Conjecture 1.2.

As far as we know the following is still open.

Conjecture 9.4 *Every planar r -graph is decomposable (and hence is r edge colorable).*

Conjecture 9.5 *The 1-factors of a planar graph form a Hilbert basis.*

Finally, we insist on a conjecture introduced in Section 7.

Conjecture 9.6 *Every r -graph contains $r - 2$ disjoint joins.*

10 Acknowledgments

We thank Michele Conforti for suggesting the problem and Ajai Kapoor for his assistance in simplifying the proposed family of indecomposable r -graphs. Bill Jackson suggested how to prove that there exists no poorly joinable r -graph for $r > 3$. Paul Seymour, besides posing the problem in [12], contributed with a second original and unpublished proof in [13].

References

- [1] J. Edmonds, Maximum matching and a polyhedron with 0, 1-vertices. *Journal of Research of the National Bureau of Standards — B.*, Mathematics and Mathematical Physics Vol. 69B, 125–130, 1965
- [2] M.A. Fiol, A Boolean algebra approach to the construction of snarks. *Graph Theory, Combinatorics, and Applications*, (Y. Alavi, G. Chartrand, O. R. Oellermann, A. J. Shwenk, Eds.) 493–524, Wiley, New York, 1991
- [3] M.A. Isaacs, Infinite families of snarks. *J.*, (Y. Alavi, G. Chartrand, O.R. Oellermann, A.J. Shwenk, Eds.) 493–524, Wiley, New York, 1991
- [4] M. Kochol, A cyclically 6-edge-connected snark of order 118. *Discrete Mathematics*, Vol. 161 297–300, (1996)
- [5] M. Kochol, Snarks without Small Cycles. *Journal of Combinatorial Theory Ser. B*, Vol. 67 34–47, (1996)
- [6] L. Lovász, Matching structure and the matching lattice. *Journal of Combinatorial Theory Ser. B*, Vol. 43 187–222, (1987)
- [7] G.H.J. Meredith, Regular n -Valent n -Connected NonHamiltonian Non- n -Edge-Colorable Graphs. *Journal of Combinatorial Theory Ser. B*, Vol. 14 55–60, (1973)
- [8] R. Rizzi, On packing T -joins. *manuscript 1997*
- [9] R. Rizzi, T -couplings and T -connectors. *manuscript 1997*
- [10] A. Caprara and R. Rizzi, Improving a Family of Approximation Algorithms to Edge Color Multigraphs. *Inform. Process. Lett.*, Vol. 68 11–15, (1998)
- [11] P.D. Seymour, Some unsolved problems on one-factorizations of graphs. *Graph Theory and Related Topics*, (J.A. Bondy and U.S.R. Murty, Eds.), 367–368, Academic Press, New York, 1979
- [12] P.D. Seymour, On multi-colourings of cubic graphs, and conjectures of Fulkerson and Tutte. *Proceedings of the London Mathematical Society*, Vol. 38 423–460, (1979)
- [13] P.D. Seymour, *private communication*
- [14] Y. Shiloach, Edge-disjoint branchings in directed multigraphs. *Inform. Process. Lett.*, Vol. 8 24–27, (1979)