

ON THE (UN)DECIDABILITY OF A NEAR-UNANIMITY TERM

MIKLÓS MARÓTI

ABSTRACT. We investigate the near-unanimity problem: given a finite algebra, decide if it has a near-unanimity term of finite arity. We prove that it is undecidable of a finite algebra if it has a partial near-unanimity term on its underlying set excluding two fixed elements. On the other hand, based on Rosenberg's characterization of maximal clones, we present partial results towards proving the decidability of the general problem.

1. INTRODUCTION

We call a term t of an algebra \mathbf{A} a *near-unanimity term* if it satisfies the following identities:

$$t(y, x, \dots, x) \approx t(x, y, x, \dots, x) \approx \dots \approx t(x, \dots, x, y) \approx x.$$

For brevity, we sometimes write NU-term instead of near-unanimity term. We investigate the *near-unanimity problem*: given a finite algebra, decide if it has a near-unanimity term of finite arity. Clearly, if the arity of the near-unanimity term of \mathbf{A} is known, then finding the near-unanimity term is easy by simply calculating the free algebra generated by the appropriate number of elements in the variety generated by \mathbf{A} . The difficulty lies in the fact that we do not have an upper bound for the arity of a possible near-unanimity term.

The near-unanimity problem was posed in [3] over ten years ago, and motivated by the *natural duality problem*: given a finite algebra, decide if it generates admits a natural duality (see [2] for details). B. A. Davey and H. Werner proved in [4] that in the presence of a near-unanimity term of \mathbf{A} , the quasi-variety \mathcal{Q} generated by \mathbf{A} admits a natural duality. The converse was proved by B. A. Davey, L. Heindorf and R. McKenzie in [3] under the assumption that \mathcal{Q} is congruence join-semi-distributive: if \mathcal{Q} admits a natural duality and is congruence join-semi-distributive then \mathbf{A} has a (finitary) near-unanimity term. This theorem, known as the near-unanimity obstacle theorem, implies that if the near-unanimity problem were undecidable, then the natural duality problem would also be undecidable, but not conversely.

Clearly, an algebra \mathbf{A} has a near-unanimity term operation t if and only if the equations

$$t(y, x, \dots, x) = t(x, y, x, \dots, x) = \dots = t(x, \dots, x, y) = x$$

hold for the generator elements x, y of the two-generated free algebra in the variety generated by \mathbf{A} . Probably this observation motivated R. McKenzie's unpublished result [7] where he proves that it is undecidable of a finite algebra \mathbf{A} and a pair $x, y \in A$ of fixed elements whether \mathbf{A} has a term t that behaves as a near-unanimity

While working on this paper, the author was partially supported by the Hungarian National Foundation for Scientific Research (OTKA) grant nos. T 37877 and T 60148.

term on $\{x, y\}$. His result does not imply the undecidability of the near-unanimity problem because the algebras used in his construction are not freely generated by the elements x, y in the variety they generate.

The key result presented in this paper is the improvement of R. McKenzie’s result to a fixed $|A| - 2$ element subset, and the simplification of his construction. The basic idea, however, is intact: the use of Minsky machines—which are equivalent to Turing machines—and the encoding of their computations in “slim” terms of \mathbf{A} . The method used in the proof relies on an absorbing element as the indicator of defects. The existence of this absorbing element prevents the further improvement of this approach to prove the undecidability of the near-unanimity problem. However, an improvement to $|A| - 1$ elements might be possible, which could be formulated, analogously to the results in [5], as the undecidability of the near-unanimity problem for partial algebras:

Problem. *Given a finite partial algebra, decide whether it has a term that is defined on all near-unanimous assignments and satisfies the near-unanimity identities.*

It is natural to attack the near-unanimity problem from the other perspective, as well: try to prove that it is decidable. We have tried the divide-and-conquer approach using I. Rosenberg’s characterization of maximal clones. It turns out that in three of the six classes of maximal clones the problem is decidable. If we restrict ourselves to idempotent algebras, then we can further eliminate one of the three remaining classes. In the idempotent case the best result is obtained using Á. Szendrei’s characterization of idempotent strictly simple term minimal algebras [14].

The two parts of the paper, partial results on the undecidability and decidability of the near-unanimity problem, are not dependent on each other. We assume only basic knowledge of universal algebra and direct the reader to either [1] or [8] for reference.

The author is indebted to R. McKenzie, Á. Szendrei and J. Ježek for their invaluable comments and directions.

2. UNDECIDABILITY OF A PARTIAL NU-TERM

Definition 2.1. Let \mathbf{A} be a fixed finite algebra, $t(x_1, \dots, x_n)$ be a term of \mathbf{A} , and S be a subset of A . We say that t is a *partial near-unanimity term on S* if $t(y, \dots, y, x_i, y, \dots, y) = y$ for all $1 \leq i \leq n$ and $x_i, y \in S$.

Note that a term t of \mathbf{A} is a NU-term iff it is a partial NU-term on A . Now one can ask the decidability of a partial NU-term on some subset. It was proved in [7] that the existence of a partial NU-term on a fixed two-element subset is undecidable. We will extend this result to a subset excluding two fixed elements, which is our main result in this section.

Theorem 2.2. *There exists no algorithm that can decide of a finite algebra \mathbf{A} and two fixed elements $r, w \in A$ if \mathbf{A} has a near-unanimity term on the set $A \setminus \{r, w\}$.*

Following the proof of R. McKenzie, our work is based on the undecidability of the halting problem for Minsky machines. The Minsky machine was invented by M. Minsky in 1961 (see [9, 10]), but he writes that the concept was inspired by some ideas of M. O. Rabin and D. Scott [12]. The “hardware” of a Minsky machine \mathcal{M} consists of two *registers* A and B , which can contain arbitrary natural numbers.

The “software” is a finite set S of *states* together with a list of *commands*. There are two special states: the *initial state* $q_1 \in S$, and the *halting state* $q_0 \in S$. The machine starts in the initial state, stops at the halting state, and at any given time it is in one of the states. For each state $i \in S \setminus \{q_0\}$ there is a single command which describes the state-transition from state i together with the change of the registers’ contents. There are two types of commands:

- in state i increase register X by one and go to state j , and
- in state i if register X contains zero, then go to state j , otherwise decrease X by one and go to state k .

Now we give the formal definition.

Definition 2.3. A *Minsky machine* $\mathcal{M} = \langle S, q_0, q_1, M \rangle$ is a finite set S of *states* with two distinguished elements $q_0, q_1 \in S$ together with a mapping

$$M : S \setminus \{q_0\} \rightarrow \{ \langle X, j \rangle, \langle X, j, k \rangle \mid X \in \{A, B\} \text{ and } j, k \in S \}.$$

We call q_0 the *halting state*, and q_1 the *initial state*. The symbols A and B represent the registers.

The mapping M describes the commands of \mathcal{M} in the following way. For any given state $i \in S \setminus \{q_0\}$ the tuple $M(i)$ is either of the form $\langle X, j \rangle$ or $\langle X, j, k \rangle$, which correspond to the two types of commands described earlier.

Definition 2.4. A *configuration* $\langle i, a, b \rangle$ of \mathcal{M} is an element of $S \times \mathbb{N} \times \mathbb{N}$, which specifies the current state and the values of the registers. We call $\langle i, a, b \rangle$ an *initial configuration* (*halting configuration*) if $i = q_1$ (or $i = q_0$, respectively).

For any configuration the Minsky machine \mathcal{M} uniquely determines (computes) the next configuration. By iteration, starting from the initial configuration with zero valued registers, we obtain a sequence of configurations, which will be called the computation of \mathcal{M} .

Definition 2.5. The *processor* for \mathcal{M} is a partial mapping of the set of configurations into itself denoted by $\bar{\mathcal{M}}$ and defined as

$$\bar{\mathcal{M}}(\langle i, a, b \rangle) = \begin{cases} \text{undefined} & \text{if } i = q_0, \\ \langle j, a + 1, b \rangle & \text{if } M(i) = \langle A, j \rangle, \\ \langle j, 0, b \rangle & \text{if } M(i) = \langle A, j, k \rangle \text{ and } a = 0, \\ \langle k, a - 1, b \rangle & \text{if } M(i) = \langle A, j, k \rangle \text{ and } a > 0, \\ \langle j, a, b + 1 \rangle & \text{if } M(i) = \langle B, j \rangle, \\ \langle j, a, 0 \rangle & \text{if } M(i) = \langle B, j, k \rangle \text{ and } b = 0, \\ \langle k, a, b - 1 \rangle & \text{if } M(i) = \langle B, j, k \rangle \text{ and } b > 0. \end{cases}$$

We will use iterative applications of the processor $\bar{\mathcal{M}}$ and adopt the power notation defined as $\bar{\mathcal{M}}^0(\langle i, a, b \rangle) = \langle i, a, b \rangle$ and $\bar{\mathcal{M}}^{n+1}(\langle i, a, b \rangle) = \bar{\mathcal{M}}(\bar{\mathcal{M}}^n(\langle i, a, b \rangle))$. Note that $\bar{\mathcal{M}}^n(\langle i, a, b \rangle)$ is undefined if and only if $\bar{\mathcal{M}}^m(\langle i, a, b \rangle)$ is a halting configuration for some $m < n$.

Definition 2.6. We say that \mathcal{M} *halts* if it halts on the $\langle 0, 0 \rangle$ input, that is, if $\bar{\mathcal{M}}^n(\langle q_1, 0, 0 \rangle)$ is a halting configuration for some $n > 0$.

It is proved in [9] that Minsky machines are equivalent to Turing machines in the following sense. Given a Minsky machine \mathcal{M} (or Turing machine \mathcal{T}), we can algorithmically construct a Turing machine $\mathcal{T}(\mathcal{M})$ (or Minsky machine $\mathcal{M}(\mathcal{T})$) which halts if and only if the original machine halts. This means that the *halting problem* for Minsky machines is as difficult as for Turing machines; that is, undecidable. Thus a new path opens for proving the undecidability of algebraic problems by interpreting Minsky machines. For example this route was taken in [6] to prove the undecidability of various kinds of word problems.

In the rest of this section we are going to prove Theorem 2.2 in the following way. For any Minsky machine \mathcal{M} we define an algebra $\mathbf{A}(\mathcal{M})$ with two special elements $r, w \in A(\mathcal{M})$ such that $\mathbf{A}(\mathcal{M})$ will have a partial near-unanimity term on $A(\mathcal{M}) \setminus \{r, w\}$ if and only if \mathcal{M} halts. This is clearly enough since the halting problem for Minsky machines is undecidable.

Let S be the set of states of \mathcal{M} with two special states: the initial state $q_1 \in S$ and the halting state $q_0 \in S$. Let the symbols A and B denote the registers of \mathcal{M} . For each $i \in S \setminus \{q_0\}$ there is a unique command which is either of the form

- $i : \text{inc } R, j$ (increase register $R \in \{A, B\}$ by one and go to state $j \in S$), or
- $i : \text{dec } R, j, k$ (if register $R \in \{A, B\}$ contains zero, then go to state $j \in S$, otherwise decrease register R by one and go to state $k \in S$).

By $\text{maj}(x, y, z)$ we denote the majority element of $\{x, y, z\}$, i.e., the element that appears at least twice among x, y and z if such element exists, otherwise $\text{maj}(x, y, z)$ is undefined. Formally,

$$\text{maj}(x, y, z) = \begin{cases} x & \text{if } x = y \text{ or } x = z, \\ y & \text{if } y = x \text{ or } y = z, \\ z & \text{if } z = x \text{ or } z = y, \\ \text{undef.} & \text{otherwise.} \end{cases}$$

Now we define the algebra $\mathbf{A}(\mathcal{M})$ in full detail.

Definition 2.7. Let $C = \{0, A, B, 1\}$. We define the algebra $\mathbf{A}(\mathcal{M})$ on the set $A(\mathcal{M}) = S \times C \cup \{p, r, w\}$ with the following operations

$$I(x) = \begin{cases} w & \text{if } x \in \{r, w\}, \\ \langle q_1, 0 \rangle & \text{if } x = p, \\ r & \text{if } x \in S \times C; \end{cases}$$

$$M(x, y, z, u) = \begin{cases} \text{maj}(y, z, u) & \text{if } \{y, z, u\} \cap \{w, r\} = \emptyset, \text{maj}(y, z, u) \text{ exists} \\ & \text{and } \text{maj}(y, z, u) \neq p, \\ p & \text{if } \{y, z, u\} \cap \{w, r\} = \emptyset, \text{maj}(y, z, u) \text{ exists,} \\ & \text{maj}(y, z, u) = p \text{ and } x \in \{q_0\} \times C \cup \{r\}, \\ w & \text{otherwise;} \end{cases}$$

for each state $i \in S \setminus \{q_0\}$ for which $M(i) = \langle R, j \rangle$ where $R \in \{A, B\}$ and $j \in S$, that is, for each command of \mathcal{M} of the form $i : \text{inc } R, j$, the operation

$$F_i(x, y) = \begin{cases} \langle j, c \rangle & \text{if } x = \langle i, c \rangle \text{ and } y = p, \\ \langle j, R \rangle & \text{if } x = \langle i, 0 \rangle \text{ and } y \in S \times C, \\ r & \text{if } x = r \text{ and } y = p, \\ w & \text{otherwise;} \end{cases}$$

for each state $i \in S \setminus \{q_0\}$ for which $M(i) = \langle R, j, k \rangle$ where $R \in \{A, B\}$ and $j, k \in S$, that is, for each command of \mathcal{M} of the form $i : \text{dec } R, j, k$, the operation

$$G_i(x, y) = \begin{cases} \langle k, c \rangle & \text{if } x = \langle i, c \rangle \text{ and } y = p, \\ \langle k, 1 \rangle & \text{if } x = \langle i, R \rangle \text{ and } y \in S \times C, \\ r & \text{if } x = r \text{ and } y = p, \\ w & \text{otherwise;} \end{cases}$$

$$H_i(x) = \begin{cases} \langle j, c \rangle & \text{if } x = \langle i, c \rangle \text{ and } c \neq R, \\ r & \text{if } x = r, \\ w & \text{otherwise.} \end{cases}$$

We will investigate this algebra in detail. The first important property of $\mathbf{A}(\mathcal{M})$ is that it almost has an absorbing element.

Definition 2.8. Let A be a set, and $f : A^n \rightarrow A$. An element $w \in A$ is *absorbing* for f if $f(\bar{a}) = w$ whenever $\bar{a} \in A^n$ and $w \in \{a_1, \dots, a_n\}$.

Lemma 2.9. *The element w of $\mathbf{A}(\mathcal{M})$ is absorbing for the operations I , F_i , G_i and H_i .*

Proof. One only has to check the definition of $\mathbf{A}(\mathcal{M})$. In the definition of I this is stated explicitly. In the definition of F_i , G_i and H_i only the ‘otherwise’ case can be applied. \square

Note that w is not an absorbing element for the operation M , but almost, except in the first variable. Combining this with the previous lemma one can see that $\mathbf{A}(\mathcal{M})$ cannot have a partial NU-term on a nontrivial subset that includes w . For example if the rightmost variable of a term is evaluated with w , then the term always yields w . We will use the element w to indicate some irregularity of a term.

Definition 2.10. Let $\bar{x} = (x_1, x_2, \dots)$ be a fixed countably infinite list of variables, and \bar{p} be the constant p assignment for these variables. For each element $e \in A(\mathcal{M})$ let $\bar{p}|_{x_n=e}$ be the assignment $x_n = e$ and $x_m = p$ if $m \neq n$. We say that a term $t(\bar{x})$ is *regular* if $t(\bar{p}) \neq w$ and $t(\bar{p}|_{x_n=e}) \neq w$ for each $n \in \mathbb{N}$ and $e \in S \times C$.

Definition 2.11. We define *slim* terms inductively. The term $I(x_n)$ is slim for every variable x_n . If t is slim, then so are $F_i(t, y)$, $G_i(t, y)$ and $H_i(t)$ for any state $i \in S$ and variable $y \in \{x_1, x_2, \dots\}$.

Lemma 2.12. *Every regular term t in which the operation symbol M does not appear is either slim or a variable. Moreover, if t is regular and slim, then there exists an assignment $\bar{p}|_{x_n=e}$ for some x_n and $e \in S \times C$, such that $t(\bar{p}|_{x_n=e}) = r$.*

Proof. Assume that t is regular and that the operation symbol M does not appear in t . We use induction on the complexity of t . Note that t cannot be a variable because variables are not slim by definition, therefore the leftmost symbol of t (in prefix notation) is either I , F_i , G_i or H_i .

Suppose that $t(\bar{x}) = I(t_1(\bar{x}))$. Because of Lemma 2.9 we know that t_1 must be regular, as well. If t_1 is not a variable, then according to our assumption we have an assignment $\bar{p}|_{x_n=e}$ such that $t_1(\bar{p}|_{x_n=e}) = r$. This shows that $t(\bar{p}|_{x_n=e}) = I(r) = w$, which is a contradiction. Thus t_1 must be a variable, that is, $t = I(x_n)$ for some integer n . In this case, t is clearly slim. Moreover, for any element $e \in S \times C$, $t(\bar{p}|_{x_n=e}) = I(e) = r$ by definition.

Now suppose that $t(\bar{x}) = F_i(t_1(\bar{x}), t_2(\bar{x}))$ for some $i \in S$. Again, both t_1 and t_2 must be regular. If t_1 is a variable, then $t(\bar{p}) = F_i(p, t_2(\bar{p})) = w$. Thus t_1 cannot be a variable. So there exists an assignment $\bar{p}|_{x_n=e}$ such that $t_1(\bar{p}|_{x_n=e}) = r$, which forces $t_2(\bar{p}|_{x_n=e}) = p$. But p is not in the range of any of the operations I , F_i , G_i and H_i ; thus t_2 must be a variable. This proves that t is slim and that $t(\bar{p}|_{x_n=e}) = F_i(r, p) = r$.

The same argument works if the leftmost operation symbol of t is either G_i or H_i . \square

Regular slim terms play a very important role in the proof; they essentially encode the computation of the Minsky machine \mathcal{M} . To see how this works, we describe the construction of a partial near-unanimity term from a halting computation.

Lemma 2.13. *If \mathcal{M} halts, then there exists a partial near-unanimity term on $A(\mathcal{M}) \setminus \{r, w\}$.*

Proof. We use the processor $\bar{\mathcal{M}}^n$ from Definition 2.5. Assume that \mathcal{M} halts in n steps, that is, $\bar{\mathcal{M}}^n(\langle q_1, 0, 0 \rangle) = \langle q_0, -, - \rangle$. For each natural number $m \leq n$ we define i_m , a_m and b_m by

$$\bar{\mathcal{M}}^m(\langle q_1, 0, 0 \rangle) = \langle i_m, a_m, b_m \rangle.$$

We are going to build a slim term of depth $n+1$ by induction. Put $t_0 = I(x)$. Now suppose that t_m is already defined. At step m the machine is in state i_m . There is a unique command for each state.

If the command for state i_m is of the form $i : \text{inc } R, j$, then put $t_{m+1} = F_{i_m}(t_m, y_m)$ where y_m is a new variable. Now assume that the command for state i_m is of the form $i : \text{dec } R, j, k$ where $R = A$. If $a_m = 0$, then put $t_{m+1} = H_{i_m}(t_m)$. If $a_m \neq 0$, then let $m' < m$ be the largest natural number such that $a_{m'} < a_m$, and put $t_{m+1} = G_{i_m}(t_m, y_{m'})$. The case when $R = B$ is handled similarly using b_m and $b_{m'}$ instead of a_m and $a_{m'}$.

Finally, put $t = M(t_n, z_1, z_2, z_3)$ where z_1 , z_2 and z_3 are new variables. We claim that t_n is a regular slim term and t is a near-unanimity term on $A(\mathcal{M}) \setminus \{r, w\}$.

CLAIM 1. *The term t_n is slim.*

This follows from the construction. We have used only variables in the second coordinates of F_i and G_i .

CLAIM 2. *No variable of t has more than two occurrences. If a variable has exactly two occurrences, then it is $y_{m'}$ for some m and the two occurrences are at $t_{m'+1} = F_{i_{m'}}(t_{m'}, y_{m'})$ and $t_{m+1} = G_{i_m}(t_m, y_{m'})$. If a variable y_m has exactly one occurrence, then it is at $t_{m+1} = F_{i_m}(t_m, y_m)$.*

The variables x , z_1 , z_2 and z_3 have single occurrences. At each F_i we always introduced a new variable. Now consider the case when $t_{m+1} = G_{i_m}(t_m, y_{m'})$. From the definition we know that $a_{m'} < a_m$ and $a_m \leq a_{m'+1}, \dots, a_m$ (assuming that $R = A$). Since $a_{m'} < a_m \leq a_{m'+1}$ and the machine cannot increase a register by more than one, $a_{m'} + 1 = a_m = a_{m'+1}$. This implies that the command for state $i_{m'}$ is of the form $i : \text{inc } R, j$ and $R = A$. On the other hand, the command for state i_m is of the form $i : \text{dec } A, j, k$ and $a_m \neq 0$, therefore $a_{m+1} = a_m - 1$. To summarize, for each pair $\langle m', m \rangle$

$$\begin{aligned} a_{m'} + 1 &= a_{m'+1} = a_m = a_{m+1} + 1, \text{ and} \\ (*) \quad a_m &\leq a_{m'+1}, \dots, a_m. \end{aligned}$$

Note that this condition is symmetric. If m' is in pair with some m , then m is the least natural number such that $m' < m$ and $a_{m'+1} > a_{m+1}$. Therefore, $y_{m'}$ has at most two occurrences.

CLAIM 3. $t_m(\bar{p}) = \langle i_m, 0 \rangle$ for all $m \leq n$

We prove by induction on m . For $m = 0$ this is true by definition: $I(p) = \langle q_1, 0 \rangle$. Now we prove it for $m + 1$. By definition t_{m+1} is $F_{i_m}(t_m, y_m)$, $H_{i_m}(t_m)$ or $G_{i_m}(t_m, y_{m'})$. Therefore $t_{m+1}(\bar{p})$ is $F_{i_m}(\langle i_m, 0 \rangle, p)$, $H_{i_m}(\langle i_m, 0 \rangle)$ or $G_{i_m}(\langle i_m, 0 \rangle, p)$.

If $t_{m+1}(\bar{p}) = F_{i_m}(\langle i_m, 0 \rangle, p)$, then the command for state i_m is of the form $i : \text{inc } R, j$. Thus $F_{i_m}(\langle i_m, 0 \rangle, p) = \langle j, 0 \rangle$ by Definition 2.7, and $\mathcal{M}(\langle i_m, a_m, b_m \rangle) = \langle j, -, - \rangle$ by Definition 2.5. Therefore $j = i_{m+1}$ and consequently $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$.

If $t_{m+1}(\bar{p}) = H_{i_m}(\langle i_m, 0 \rangle)$, then the command for state i_m is of the form $i : \text{dec } R, j, k$, moreover $a_m = 0$ if $R = A$, and $b_m = 0$ if $R = B$. Now $H_{i_m}(\langle i_m, 0 \rangle, p) = \langle j, 0 \rangle$ by Definition 2.7, and $\mathcal{M}(\langle i_m, a_m, b_m \rangle) = \langle j, -, - \rangle$ by Definition 2.5. Therefore $j = i_{m+1}$ and consequently $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$.

Finally, if $t_{m+1}(\bar{p}) = G_{i_m}(\langle i_m, 0 \rangle, p)$, then the command for state i_m is of the form $i : \text{dec } R, j, k$, moreover $a_m > 0$ if $R = A$, and $b_m > 0$ if $R = B$. Now $G_{i_m}(\langle i_m, 0 \rangle, p) = \langle k, 0 \rangle$ by Definition 2.7, and $\mathcal{M}(\langle i_m, a_m, b_m \rangle) = \langle k, -, - \rangle$ by Definition 2.5. Therefore $k = i_{m+1}$ and consequently $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$.

CLAIM 4. $t_m(\bar{p}|_{x=e}) = r$ for all $m \leq n$ and $e \in S \times C$ where x is the variable used to define $t_0 = I(x)$.

We prove the claim by induction. For $m = 0$, $t_0 = I(x)$ and $t_0(\bar{p}|_{x=e}) = I(e) = r$. Now assume that $t_m(\bar{p}|_{x=e}) = r$ for some $m < n$. By the construction, t_{m+1} is either $F_{i_m}(t_m, y)$, $G_{i_m}(t_m, y)$ or $H_{i_m}(t_m)$ where y is some variable different from x . Thus $t_{m+1}(\bar{p}|_{x=e})$ equals either $F_{i_m}(r, p)$, $G_{i_m}(r, p)$ or $H_{i_m}(r)$. But each of these equals r by Definition 2.7.

CLAIM 5. Let $h < n$ and $e \in S \times C$ be fixed and assume that y_h has exactly one occurrence in t_n . Let R be the register manipulated in the command for state i_h . Then

$$t_m(\bar{p}|_{y_h=e}) = \begin{cases} \langle i_m, 0 \rangle & \text{if } 0 \leq m \leq h, \\ \langle i_m, R \rangle & \text{if } h < m \leq n. \end{cases}$$

Without loss of generality we can assume that $R = A$. By Claim 2, the single occurrence of y_h is at $t_{h+1} = F_{i_h}(t_h, y_h)$. Therefore, if $m \leq h$, then $t_m(\bar{p}|_{y_h=e}) = t_m(\bar{p}) = \langle i_m, 0 \rangle$ by Claim 3. We use induction on m to prove the other case. For the base of the induction we have $t_{h+1}(\bar{p}|_{y_h=e}) = F_{i_h}(\langle i_h, 0 \rangle, e) = \langle i_{h+1}, A \rangle$.

Now consider the induction step from m to $m + 1$. Assume that $t_{m+1} = F_{i_m}(t_m, y_m)$. Since y_h has a single occurrence, $y_h \neq y_m$, and thus $t_{m+1}(\bar{p}|_{y_h=e}) = F_{i_m}(\langle i_m, A \rangle, p) = \langle i_{m+1}, A \rangle$. The same argument works when $t_{m+1} = G_{i_m}(t_m, y_{m'})$.

Now assume that $t_{m+1} = H_{i_m}(t_m)$. From equation (*) in the proof of Claim 2 we can see that $a_h < a_{h+1}, \dots, a_n$, otherwise y_h would have at least two occurrences. Therefore, $a_m \neq 0$. By the definition of t_{m+1} we know that either a_m or b_m must be zero. Thus it is register B which is manipulated in the command for state i_m . This implies that $t_{m+1}(\bar{p}|_{y_h=e}) = H_{i_m}(\langle i_m, A \rangle) = \langle i_{m+1}, A \rangle$.

CLAIM 6. *Let $h < n$ and $e \in S \times C$ be fixed and assume that $y_{h'}$ has exactly two occurrences in t_n as described in Claim 2. Let R be the register manipulated in the commands for states $i_{h'}$ and i_h . Then*

$$t_m(\bar{p}|_{y_{h'}=e}) = \begin{cases} \langle i_m, 0 \rangle & \text{if } 0 \leq m \leq h', \\ \langle i_m, R \rangle & \text{if } h' < m \leq h, \\ \langle i_m, 1 \rangle & \text{if } h < m \leq n. \end{cases}$$

Without loss of generality we can assume that $R = A$. For the first two cases of the displayed equation above the same argument works as in the previous claim, but using h' instead of h .

We prove the third case by induction on m . For the base of the induction we have $t_{h+1} = G_{i_h}(t_h, y_{h'})$. Hence $t_{h+1}(\bar{p}|_{y_{h'}=e}) = G_{i_h}(\langle i_h, A \rangle, e) = \langle i_{h+1}, 1 \rangle$. The induction step is now easy as there are no other occurrences of $y_{h'}$ along the term t_n . Therefore, we always calculate $F_{i_m}(\langle i_m, 1 \rangle, p)$, $G_{i_m}(\langle i_m, 1 \rangle, p)$, or $H_{i_m}(\langle i_m, 1 \rangle)$, which all yield $\langle i_{m+1}, 1 \rangle$.

CLAIM 7. *The term t_n is regular. Moreover, $t_n(\bar{p}|_{u=e}) \in \{q_0\} \times C \cup \{r\}$ for all variables u and all $e \in A(\mathcal{M}) \setminus \{r, w\}$.*

Take any element $e \in S \times C$. By Claims 3 and 4 we have $t_n(\bar{p}) = \langle q_0, 0 \rangle$ and $t_n(\bar{p}|_{x=e}) = r$, respectively. Now take a variable y_h . If y_h has no occurrence in t_n , then $t_n(\bar{p}|_{y_h=e}) = t_n(\bar{p}) = \langle q_0, 0 \rangle$. Otherwise y_h has one or two occurrences by Claim 2. Then by Claims 5 and 6 we have $t_n(\bar{p}|_{y_h=e}) \in \{q_0\} \times C$.

CLAIM 8. *t is a near-unanimity term on $A(\mathcal{M}) \setminus \{r, w\}$.*

Take a near-unanimous assignment \bar{a} on $A(\mathcal{M}) \setminus \{r, w\}$. If the majority element is not p , then $t(\bar{a}) = M(t_n(\bar{a}), z_1, z_2, z_3) = \text{maj}(z_1, z_2, z_3)$. If the majority element is p , then $t_n(\bar{a}) \in \{q_0\} \times C \cup \{r\}$ by Claim 7, and hence $t(\bar{a}) = p$. Therefore, t is a near-unanimity term on $A(\mathcal{M}) \setminus \{r, w\}$. \square

We have seen how to encode the halting computation into the regular slim term t_n . Our goal now is the reverse; to show that the computation of \mathcal{M} can be recovered from a regular slim term.

Lemma 2.14. *Let t_n be a regular slim term of depth $n + 1$. Then $t_n(\bar{p}) = \langle i_n, 0 \rangle$ where i_n is the state of the machine \mathcal{M} after the first n steps.*

Proof. We want to show that the term t_n behaves the same way as the one in the proof of the previous lemma. Denote by t_m the unique subterm of t_n of depth $m + 1$. That is, $t_0 = I(-)$, and t_{m+1} is $F_i(t_m, -)$, $G_i(t_m, -)$ or $H_i(t_m)$ for some $i \in S$.

CLAIM 1. *$t_m(\bar{p}|_{x=e}) \neq w$ for all $m \leq n$, $e \in S \times C$ and all variables x of t_n .*

Suppose that $t_m(\bar{p}|_{x=e}) = w$ for some $m \leq n$, $e \in S \times C$ and variable x of t_n . The element w is absorbing for the operations of t_n by Lemma 2.9, which implies that $t_n(\bar{p}|_{x=e}) = w$. This contradicts the regularity of t_n and proves the claim.

CLAIM 2. $t_m(\bar{p}) \in S \times \{0\}$ for all $m \leq n$.

We prove the claim by induction on m . For $m = 0$ we have $t_0(\bar{p}) = I(p) = \langle q_1, 0 \rangle \in S \times \{0\}$. For the induction step, assume that $t_m(\bar{p}) = \langle s, 0 \rangle \in S \times \{0\}$ and consider t_{m+1} . If $t_{m+1} = F_i(t_m, y)$ for some $i \in S$ and variable y , then $t_{m+1}(\bar{p}) = F_i(\langle s, 0 \rangle, p)$ which equals either $\langle j, 0 \rangle$ for some $j \in S$ or w . But $t_{m+1}(\bar{p}) \neq w$ as t_{m+1} is regular, so $t_{m+1}(\bar{p}) = \langle j, 0 \rangle \in S \times \{0\}$. A similar argument works for the other two types of operations: G_i and H_i .

CLAIM 3. Let x be the variable used in t_0 . Then x has no other occurrence in t_n . Moreover, $t_m(\bar{p}|_{x=e}) = r$ for all $m \leq n$ and $e \in S \times C$.

We use induction on m . For $m = 0$ we have $t_0(\bar{p}|_{x=e}) = I(e) = r$. For the induction step from m to $m + 1$ assume that $t_m(\bar{p}|_{x=e}) = r$. Thus $t_{m+1}(\bar{p}|_{x=e})$ is $F_i(r, y)$, $G_i(r, y)$ or $H_i(r)$ for some $i \in S$ and some variable y . We know that this value is not w by Claim 1. Looking up the definition of F_i , G_i and H_i , we can see that the only choice is when the result is r (and $y = p$ for F_i and G_i). This completes the induction step and proves that $x \neq y$ (as $e \neq p$) when the operation is F_i or G_i .

CLAIM 4. Assume that a variable $y \neq x$ has exactly one occurrence in t_n . Then the occurrence is at $t_{m+1} = F_i(t_m, y)$ for some $m < n$ and $i \in S$. Moreover, there exists no $h > m$ such that $t_{h+1} = H_j(t_h)$ and the command for j manipulates the same register as the one for i .

Let m be the least natural number such that t_{m+1} has an occurrence of y . Then $t_{m+1} = F_i(t_m, y)$ or $t_{m+1} = G_i(t_m, y)$ for some $i \in S$. Take an element $e \in S \times C$, and consider $t_{m+1}(\bar{p}|_{y=e})$. By the choice of m , $t_m(\bar{p}|_{y=e}) = t_m(\bar{p})$, and then by Claim 2, $t_m(\bar{p}|_{y=e}) \in S \times \{0\}$. Checking the definition of G_i we see that $G_i(t_m(\bar{p}|_{y=e}), e) = w$, a contradiction. So $t_{m+1} = F_i(t_m, y)$. Moreover, $t_{m+1}(\bar{p}|_{y=e}) \in S \times \{R\}$ where R is the register manipulated by the command for i . Now we show that $t_h(\bar{p}|_{y=e}) \in S \times \{R\}$ for all $h > m$ by induction. For $m + 1$ we already have this. For the induction step consider $a = t_{h+1}(\bar{p}|_{y=e})$. By definition a is $F_j(\langle -, R \rangle, p)$, $G_j(\langle -, R \rangle, p)$ or $H_j(\langle -, R \rangle)$ for some $j \in S$ and $a \neq w$. In the first two cases this shows that $a \in S \times \{R\}$. On the other hand, when $a = H_j(\langle -, R \rangle) \neq w$, then the command for state j cannot manipulate the register R . This concludes the proof of this claim.

CLAIM 5. Assume that a variable $y \neq x$ has at least two occurrences in t_n . Then there exist $m' < m$ such that $t_{m'+1} = F_i(t_{m'}, y)$, $t_{m+1} = G_j(t_m, y)$ for some $i, j \in S$, the commands for i and j manipulate the same register R , and y has no other occurrences than these two. Moreover, there exists no $m' < h < m$ such that $t_{h+1} = H_k(t_h)$ and the command for k manipulates the register R .

Let m' and m be the least natural numbers such that $t_{m'+1}$ has exactly one and t_{m+1} has exactly two occurrences of y . The term t_m has exactly one occurrence of y , so we can apply the previous claim. This proves half of the claim. It remains to be shown that $t_{m+1} = G_j(t_m, y)$ for some $j \in S$, that the command for j manipulates the register R , and that there are no other occurrences of y .

Fix an element $e \in S \times C$. From the proof of the previous claim we know that $t_m(\bar{p}|_{y=e}) \in S \times \{R\}$ where R is the register manipulated by the command for state i . Consider $a = t_{m+1}(\bar{p}|_{y=e})$. This element is either $F_j(\langle -, R \rangle, e)$ or $G_j(\langle -, R \rangle, e)$ for some j . Since $a \neq w$ by Claim (1), we must have $t_{m+1} = G_j(t_m, y)$, and the command for state j must manipulate register R . Therefore, we have $t_{m+1}(\bar{p}|_{y=e}) \in S \times \{1\}$.

Finally, we show that $t_h(\bar{p}|_{y=e}) \in S \times \{1\}$ for all $h > m$ by induction. We have already the basis of the induction. To show the induction step, consider t_{h+1} . If $t_{h+1} = H_k(t_h)$ for some k , then we get $t_{h+1}(\bar{p}|_{y=e}) \in S \times \{1\}$ by the definition of H_k . Now assume that $t_{h+1} = F_k(t_h, z)$. Since $t_{h+1}(\bar{p}|_{y=e}) \neq w$ we must have $z \neq y$ and $t_{h+1}(\bar{p}|_{y=e}) \in S \times \{1\}$. The same argument works for G_k , as well.

CLAIM 6. *Let i_m , a_m and b_m be defined by $\bar{\mathcal{M}}^m(\langle q_1, 0, 0 \rangle) = \langle i_m, a_m, b_m \rangle$. Then the following hold for all $0 \leq m < n$.*

- (1) *If the command for i_m is of the form $i : \text{inc } R, j$, then $t_{m+1} = F_{i_m}(t_m, -)$.*
- (2) *If the command for i_m is of the form $i : \text{dec } R, j, k$, and if $a_m \neq 0$ for $R = A$ while $b_m \neq 0$ for $R = B$, then $t_{m+1} = G_{i_m}(t_m, -)$.*
- (3) *If the command for i_m is of the form $i : \text{dec } R, j, k$, and if $a_m = 0$ for $R = A$ while $b_m = 0$ for $R = B$, then $t_{m+1} = H_{i_m}(t_m, -)$.*

Moreover, $t_m(\bar{p}) = \langle i_m, 0 \rangle$ for all $0 \leq m \leq n$.

We prove this by induction on m . For $m = 0$ we have $t_0(\bar{p}) = I(p) = \langle q_1, 0 \rangle = \langle i_0, 0 \rangle$. For the induction step assume that (1)–(3) hold for all $m' < m$, a condition which is void if $m = 0$, and $t_m(\bar{p}) = \langle i_m, 0 \rangle$. We have to show that (1)–(3) hold for m and $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$.

Assume that $t_{m+1} = F_i(t_m, y)$ for some $i \in S$ and some variable y . We have to show that $i = i_m$ and $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$. Since the operation F_i is defined, the command for state i is $i : \text{inc } R, j$ for some $R \in \{A, B\}$ and $j \in S$. From the induction hypothesis, $t_m(\bar{p}) = \langle i_m, 0 \rangle$. Consider the element $e = t_{m+1}(\bar{p}) = F_i(\langle i_m, 0 \rangle, p)$. Since $e \neq w$ by Claim (1), we must have $i = i_m$ and $e = \langle j, 0 \rangle$. As $i_m = i$ and the command is $i : \text{inc } R, j$, we have $i_{m+1} = j$. Therefore, $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$.

Assume that $t_{m+1} = G_i(t_m, y)$ for some $i \in S$ and variable y . We have to show that $i = i_m$ and $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$. Since the operation G_i is defined, the command for state i is $i : \text{dec } R, j, k$ for some $R \in \{A, B\}$ and $j, k \in S$. Without loss of generality we can assume that $R = A$. Consider $e = t_{m+1}(\bar{p}) = G_i(\langle i_m, 0 \rangle, p)$. Since $e \neq w$, we must have $i = i_m$ and $e = \langle k, 0 \rangle$. What remains to be shown is that $i_{m+1} = k$. We know that i_{m+1} is either j or k depending on whether $a_m = 0$ or $a_m \neq 0$. We claim that $a_m \neq 0$. By the definition of the Minsky machine, we have

$$a_m = |\{h < m : \mathcal{M} \text{ has increased register } A \text{ at step } h\}| \\ - |\{h < m : \mathcal{M} \text{ has decreased register } A \text{ at step } h\}|.$$

Now using the induction hypothesis we get that

$$(S^+) \quad a_m = |\{h < m : t_{h+1} = F_{i_h}(t_h, -) \\ \text{and the command for } i_h \text{ manipulates register } A\}| \\ (S^-) \quad - |\{h < m : t_{h+1} = G_{i_h}(t_h, -) \\ \text{and the command for } i_h \text{ manipulates register } A\}|.$$

Take a number h from the second set S^- , so $t_{h+1} = G_{i_h}(t_h, z)$ for some variable z , and the command for i_h manipulates register A . By Claims 3, 4 and 5, the variable z has exactly two occurrences; the other being at $t_{h'+1} = F_{i_{h'}}(t'_h, z)$ for some $h' < h$. Moreover, the command for $i_{h'}$ manipulates the same register A . Thus h' belongs to the first set S^+ . This only shows that $a_m \geq 0$. But the same argument works for $t_{m+1} = G_i(t_m, y)$, showing that there exists an $m' < m$ which belongs to S^+ , while $m \notin S^-$. Therefore, $a_m > 0$ and $i_{m+1} = k$.

Finally, assume that $t_{m+1} = H_i(t_m)$ for some $i \in S$. We have to show that $i = i_m$ and $t_{m+1}(\bar{p}) = \langle i_{m+1}, 0 \rangle$. Since the operation H_i is defined, the command for state i is $i : \text{dec } R, j, k$ for some $R \in \{A, B\}$ and $j, k \in S$. Without loss of generality we can assume that $R = A$. Consider $e = t_{m+1}(\bar{p}) = H_i(\langle i_m, 0 \rangle)$. Since $e \neq w$, we must have $i = i_m$ and $e = \langle j, 0 \rangle$. What remains to be shown is that $i_{m+1} = j$. We know that i_{m+1} is either j or k depending on whether $a_m = 0$ or $a_m \neq 0$. To get a contradiction, suppose that $a_m \neq 0$, i.e., the set S^+ , defined in the previous subsection, has more elements than S^- . We know that each element of S^- is in pair with a unique element of S^+ . So there exists an $h < m$ such that $t_{h+1} = F_i(t_h, z)$ for some variable z , the command for i manipulates register A , and h is not in S^- . Therefore, z has exactly one occurrence in t_m . If z has two occurrences, then the other one must appear after t_{m+1} . In any case, either by Claim 4 or 5, the command for i at $t_{m+1} = H_i(t_m)$ cannot manipulate register A . But according to our assumption it does, which is a contradiction. This shows that $a_m = 0$, therefore $i_{m+1} = j$.

This finishes the proof of the last claim, which includes the statement $t_n(\bar{p}) = \langle i_n, 0 \rangle$ of the lemma. \square

The previous two lemmas give the connection between regular slim terms and halting computations. What remains to be shown is that a regular slim term can be found as a subterm of a near-unanimity term on $A(\mathcal{M}) \setminus \{r, w\}$, or at least as a subterm of a ‘‘minimal’’ near-unanimity term.

Definition 2.15. Two terms t_1 and t_2 are *p-equivalent* iff $t_1(\bar{p}) = t_2(\bar{p})$ and $t_1(\bar{p}|_{x_n=e}) = t_2(\bar{p}|_{x_n=e})$ for each $n \in \mathbb{N}$ and $e \in S \times C$. A term is *p-minimal* iff there is no *p-equivalent* term of smaller complexity.

Lemma 2.16. *Let t be a regular p-minimal term in which the operation symbol M appears. Then $\mathbf{A}(\mathcal{M})$ halts.*

Proof. We use induction on the complexity of t . If $t = F_i(t_1, t_2)$, then both t_1 and t_2 must be regular (and *p-minimal*) by Lemma 2.9. So at least in one of them the operation symbol M appears, and by induction we are done. The same argument works for the operations G_i , H_i and I , as well.

Now suppose that $t = M(t_1, t_2, t_3, t_4)$. If t_2, t_3 or t_4 is not regular, then we have some near *p-unanimous* assignment \bar{f} such that $w \in \{t_2(\bar{f}), t_3(\bar{f}), t_4(\bar{f})\}$. This forces $t(\bar{f}) = w$, which is a contradiction. So t_2, t_3 and t_4 are regular. If in one of them the operation symbol M appears, then we use induction on that subterm. So assume that M does not occur in t_2, t_3 and t_4 . By Lemma 2.12, each of them is either a slim term or a variable. If t_k is slim ($k \in \{2, 3, 4\}$), then we have an assignment $\bar{p}|_{x_n=e}$ such that $t_k(\bar{p}|_{x_n=e}) = r$. This forces a contradiction $t(\bar{p}|_{x_n=e}) = w$. Thus t_2, t_3 and t_4 must be variables. If two of them are the same variable y , then t is *p-equivalent* to y because the operation M must yield the majority element y , as it cannot return w because t is regular. This contradicts

the p -minimality, thus the terms t_2 , t_3 and t_4 are distinct variables. If t_1 is not regular, then we have an assignment $\bar{p}|_{x_n=e}$ such that $t_1(\bar{p}|_{x_n=e}) = w$. But this forces $t(\bar{p}|_{x_n=e}) = w$, a contradiction. So t_1 must be regular. If t_1 contains the operation symbol M , then we use the induction. If t_1 does not contain M , then by Lemma 2.12 it is either a slim term or a variable. It cannot be a variable because $t(\bar{p}) \neq w$. So t_1 is regular and slim term. Now by Lemma 2.14 the value $t_1(\bar{p})$ contains the last state of the correct piece of the computation. But $t(\bar{p}) \neq w$, which proves that we have reached the halting state. \square

Theorem 2.17. *Let \mathcal{M} be a Minsky machine. The algebra $\mathbf{A}(\mathcal{M})$ has a near-unanimity term on the set $A(\mathcal{M}) \setminus \{r, w\}$ iff \mathcal{M} halts.*

Proof. Suppose that t is a near-unanimity term on $A(\mathcal{M}) \setminus \{r, w\}$. Then t is regular. Let t' be a term p -equivalent to t and p -minimal. Then t' is not a variable; moreover, $t'(\bar{p}) = p$ implies that the leftmost operation symbol of t' (in prefix notation) is M . Now by Lemma 2.16, \mathcal{M} halts. The other direction is proved in Lemma 2.13. \square

This finishes the proof of Theorem 2.2, as it is undecidable of a Minsky machine if it halts.

3. TOWARDS THE DECIDABILITY OF A NU-TERM

In this section we solve the NU-problem for special classes of algebras. We start with Rosenberg's primal algebra characterization (see [13, 11]), which presents a natural framework for this. Clearly, a primal algebra has a ternary NU-term; and it is decidable if an algebra is primal. If the algebra is not primal, then its clone lies in one of the maximal clones described in Rosenberg's theorem. We solve the NU-problem in three classes of maximal clones (out of six), and present other partial results.

Rosenberg's characterization is in terms of six classes of finitary relations; a non-trivial finite algebra \mathbf{A} is preprimal (its clone is a coatom in the lattice of clones) if and only if there is a relation ϱ in one of the six classes such that the term functions of \mathbf{A} are exactly the functions preserving the relation ϱ . Now we define these classes, following Quackenbush [11].

Definition 3.1. Let A be a finite set.

A subset $\varrho \subseteq A^2$ is a *partial order* if it is *reflexive* ($\langle a, a \rangle \in \varrho$ for all $a \in A$), *antisymmetric* ($\langle a, b \rangle, \langle b, a \rangle \in \varrho$ imply that $a = b$), and *transitive* ($\langle a, b \rangle, \langle b, c \rangle \in \varrho$ imply that $\langle a, c \rangle \in \varrho$). We say that $b \in A$ is a *zero (unit)* of $\varrho \subseteq A^2$ if $\langle b, a \rangle \in \varrho$ ($\langle a, b \rangle \in \varrho$) for all $a \in A$. Note that a partial order has at most one zero and at most one unit.

A subset $\varrho \subseteq A^2$ is a *permutation* if $\varrho = \{ \langle a, \alpha(a) \rangle : a \in A \}$ where $\alpha : A \rightarrow A$ is a permutation on A . We say that the permutation ϱ is *prime* if all cycles of α have the same length that is a prime number.

A subset $\varrho \subseteq A^2$ is an *equivalence relation* if ϱ is reflexive, *symmetric* ($\langle a, b \rangle \in \varrho$ implies $\langle b, a \rangle \in \varrho$), and transitive. An equivalence relation ϱ is *non-trivial* if $\varrho \neq A^2$ and $\varrho \neq \{ \langle a, a \rangle : a \in A \}$.

A subset $\varrho \subseteq A^4$ is *affine* if we can define an abelian group operation, $+$, on A so that $\langle a, b, c, d \rangle \in \varrho$ if and only if $a + b = c + d$. An affine ϱ is *prime* if $\langle A; + \rangle$ is an elementary abelian p -group.

A subset $\varrho \subseteq A^h$ (for $h \geq 1$) is *totally symmetric* if for every permutation α on $\{1, \dots, h\}$, $\langle a_1, \dots, a_h \rangle \in \varrho$ if and only if $\langle a_{\alpha(1)}, \dots, a_{\alpha(h)} \rangle \in \varrho$. Let $A_h \subseteq A^h$ be defined by

$$(*) \quad A_h = \{ \langle a_1, \dots, a_h \rangle : a_i = a_j \text{ for some } i \neq j \}.$$

We say that ϱ is *totally reflexive* if $A_h \subseteq \varrho$. The *center* of ϱ is the set of all $a \in A$ such that for all $a_2, \dots, a_h \in A$, $\langle a, a_2, \dots, a_h \rangle \in \varrho$. We say that ϱ is *central* if it is totally symmetric, totally reflexive and has a center which is a non-empty, proper subset of A .

Let $h = \{0, 1, \dots, h-1\}$. For $1 \leq r \leq m$, let π_r^m be the r th projection of h^m onto h . Define ω_m to be the h -ary relation on h^m such that $\langle a_1, \dots, a_h \rangle \in \omega_m$ if and only if for all $1 \leq r \leq m$, $\langle \pi_r^m(a_1), \dots, \pi_r^m(a_h) \rangle \in h_h$ (where h_h is defined by (*)). A subset $\varrho \subseteq A^h$ for $h \geq 3$ is *h -regularly generated* if for some $m \geq 1$ there is a surjection $\varphi : A \rightarrow h^m$ such that $\varrho = \varphi^{-1}(\omega_m)$; i.e., $\langle a_1, \dots, a_h \rangle \in \varrho$ if and only if $\langle \varphi(a_1), \dots, \varphi(a_h) \rangle \in \omega_m$. Clearly, if ϱ is h -regularly generated, then ϱ is totally reflexive and totally symmetric.

Theorem 3.2 (Rosenberg [13]). *A finite non-trivial algebra \mathbf{A} is preprimal if and only if there exists an h -ary relation ϱ on A from the following classes*

- (1) *the set of all partial orders with a zero and unit,*
- (2) *the set of all prime permutations,*
- (3) *the set of all non-trivial equivalence relations,*
- (4) *the set of all prime affine relations,*
- (5) *the set of all central relations,*
- (6) *the set of all h -regularly generated relations,*

such that the set of term functions of \mathbf{A} is just the set of functions preserving ϱ . \square

First we show that the NU-term problem is decidable inside a maximal clone of class (1). We need the following lemma, which grew out of discussions with R. McKenzie.

Lemma 3.3. *For a finite algebra \mathbf{A} and a natural number k , it is decidable whether \mathbf{A} has a near-unanimity term in which at most k variables have repeated occurrences.*

Proof. It is enough to effectively find a number K so that if \mathbf{A} has a NU-term in which at most k variables have repeated occurrences, then it has a NU-term of depth at most K with the same property.

Suppose we do have a near-unanimity term t , and its tree has a long branch $t = t_0, t_1, \dots, t_n$. Here $t_i = g_i(t_{i+1}, -, \dots, -)$, where g_i is a basic operation with variables permuted. Let X be the tuple x_1, x_2, \dots, x_k of variables permitted to have repeated occurrences, and Y be the tuple of remaining variables.

We find a long subsequence $\{s_j\}$ of $\{t_i\}$, such that when all variables of Y are replaced by one new variable z , then $s_j(X; z) = s_l(X; z)$ for all j and l . We can also assume that $B(s_j) = B(s_l)$ for all j and l , where $B(s(X; Y))$ is the set of all term operations $b(x, z)$ of \mathbf{A} arising from the term $s(X; Y)$ by choosing some variable among Y , replacing it by z , and then replacing all other variables of Y and X by x . Also, we can assume that $s_j(x, \dots, x) = s_l(x, \dots, x)$.

Now we claim that if we create a new term t' by replacing the explicit occurrence of s_1 in t (i.e., at $t_i = g_i(s_1, -, \dots, -)$, where $s_1 = t_{i+1}$) by s_2 , then this shorter term t' is also a near-unanimity term.

Indeed, in each near-unanimous assignment in which the minority variable is from X , the terms s_1 and s_2 behave the same. If the minority variable is from Y , then it has exactly one occurrence. If this occurrence is inside of s_1 , then we use that fact that $B(s_1) = B(s_2)$. If it is outside, then we use that fact that $s_1(x, \dots, x) = s_2(x, \dots, x)$. \square

Corollary 3.4. *Given a finite algebra \mathbf{A} whose clone lies in a maximal clone of class (1). Then it is decidable if \mathbf{A} has a near-unanimity term.*

Proof. We will prove that if \mathbf{A} has a NU-term, then it has a NU-term in which no variable has multiple occurrences. By the previous lemma this is enough.

Assume that $t(x_1, \dots, x_n)$ is a NU-term of \mathbf{A} . Put

$$t'(y_{11}, \dots, y_{1m_1}, y_{21}, \dots, y_{nm_n}),$$

the term obtained from t by replacing all occurrences of each variable x_i by distinct variables y_{ij} . We claim that t' is also a NU-term. Let \leq be a compatible partial order on A with a zero element $0 \in A$ and a unit element $1 \in A$. Take elements $a, b \in A$, and consider the near-unanimous assignment $t'(a, \dots, a, b, a, \dots, a)$ where $y_{ij} = b$ for some i and j . Since \leq is compatible with t' ,

$$\begin{aligned} t'(a, \dots, a, b, a, \dots, a) &\leq t'(a, \dots, a, 1, \dots, 1, a, \dots, a) \\ &= t(a, \dots, a, 1, a, \dots, a) = a, \end{aligned}$$

where $y_{ik} = 1$ for all k , and $x_i = 1$. On the other hand, $a \leq t'(a, \dots, a, b, a, \dots, a)$ by a similar argument. Therefore

$$t'(a, \dots, a, b, a, \dots, a) = a$$

for all $a, b \in A$ and i, j . \square

Now we show that no NU-term can exist in the maximal clones of class (4) and (6), so the problem is decidable in these cases. We call an algebra \mathbf{A} *affine* if it has a compatible affine relation.

Proposition 3.5. *No finite affine algebra has a near-unanimity term. In particular, a finite algebra \mathbf{A} whose clone lies in a maximal clone of class (4), has no near-unanimity term.*

Proof. Assume the contrary, that there exists a NU-term $t(x_1, \dots, x_n)$ of \mathbf{A} . Let $0 \in A$ be the zero element of the abelian group $\langle A; + \rangle$. Fix another element $a \neq 0$ of A . For $0 \leq k \leq n$, let \bar{a}^k be the vector $\langle a, \dots, a, 0, \dots, 0 \rangle \in A^n$ with k -many a entries. We show by induction that $t(\bar{a}^k) = 0$, which is a contradiction for $k = n$. The base of the induction, $k = 0$, is true, since t is a NU-term. For the induction step

$$\begin{aligned} t(a, \dots, a, 0, 0, \dots, 0) &= 0, && \text{by the induction hypothesis,} \\ t(0, \dots, 0, a, 0, \dots, 0) &= 0, && \text{by the NU-term } t, \\ t(0, \dots, 0, 0, 0, \dots, 0) &= 0, && \text{by the NU-term } t, \text{ and} \\ t(a, \dots, a, a, 0, \dots, 0) &= b, && \text{for some } b \in A. \end{aligned}$$

On the left hand side all columns are in the relation $x + y = z + u$. Since this relation is preserved by t , $0 + 0 = 0 + b$, that is, $b = 0$. \square

Proposition 3.6. *A finite algebra \mathbf{A} whose clone lies in a maximal clone of class (6), has no near-unanimity term.*

Proof. Let h, m be natural numbers, $\varphi : A \rightarrow h^m$ be a surjection, and $\varrho \subseteq A^h$ be a relation as described in the Definition 3.1 under class (6). Assume that there exists a NU-term $t(x_1, \dots, x_n)$ of \mathbf{A} which preserves ϱ . We want to get a contradiction.

Recall that $h = \{0, 1, \dots, h-1\}$ and $h \geq 3$. Since φ is surjective, there exist $a_0, \dots, a_{h-1} \in A$ such that $\pi_1^m(\varphi(a_i)) = i$ for all $0 \leq i < h$. For $0 \leq k \leq n$ put $\bar{b}^k = \langle a_0, \dots, a_0, a_1, \dots, a_1 \rangle \in A^n$ with k many a_0 entries. We will prove by induction that $\pi_1^m(\varphi(t(\bar{b}^k))) \neq 0$ for all $0 \leq k \leq n$. For $k = 0$ this is true by definition.

For the induction step assume that the claim is true for k . Put $j = \pi_1^m(\varphi(t(\bar{b}^k)))$. By the induction hypothesis, $j \neq 0$. Consider the following tuples of A^n

$$\begin{aligned} \bar{b}^{k+1} &= \langle a_0, \dots, a_0, a_0, a_1, \dots, a_1 \rangle, \\ &\quad \langle a_1, \dots, a_1, a_0, a_1, \dots, a_1 \rangle, \\ &\quad \vdots \\ &\quad \langle a_{j-1}, \dots, a_{j-1}, a_0, a_{j-1}, \dots, a_{j-1} \rangle, \\ \bar{b}^k &= \langle a_0, \dots, a_0, a_1, a_1, \dots, a_1 \rangle, \\ &\quad \langle a_{j+1}, \dots, a_{j+1}, a_0, a_{j+1}, \dots, a_{j+1} \rangle, \\ &\quad \vdots \\ &\quad \langle a_{h-1}, \dots, a_{h-1}, a_0, a_{h-1}, \dots, a_{h-1} \rangle, \end{aligned}$$

where the i th row ($i \neq 0, j$) is the near-unanimous a_i tuple with a_0 at the $k+1$ -th coordinate. Notice that each column has a repeated entry. Indeed, for the $k+1$ -th column it is a_0 , and for all other columns it is either a_0 or a_1 from the rows \bar{b}^{k+1} and \bar{b}^k . This means that each column is in the relation ϱ . Therefore, by applying t ,

$$\langle t(\bar{b}^{k+1}), a_1, \dots, a_{j-1}, t(\bar{b}^k), a_{j+1}, \dots, a_{h-1} \rangle \in \varrho.$$

Denote this tuple by \bar{c} . By the definition of ϱ , $\varphi(\bar{c}) \in \omega_m$. Then by the definition of ω_m , $\pi_1^m(\varphi(\bar{c})) \in h_h$. But we can calculate this tuple,

$$\pi_1^m(\varphi(\bar{c})) = \langle \pi_1^m(\varphi(t(\bar{b}^{k+1}))), 1, \dots, j-1, j, j+1, \dots, h-1 \rangle.$$

By the definition of h_h , this tuple must have a repetition, thus $\pi_1^m(\varphi(t(\bar{b}^{k+1}))) \neq 0$. This completes the proof of the induction step.

We have shown that $\pi_1^m(\varphi(t(\bar{b}^n))) \neq 0$. On the other hand,

$$\pi_1^m(\varphi(t(\bar{b}^n))) = \pi_1^m(\varphi(t(a_0, \dots, a_0))) = \pi_1^m(\varphi(a_0)) = 0,$$

which is a contradiction. \square

In the rest of this section we focus on the case when the finite algebra in question is idempotent. As the first step we reduce the problem to simple algebras.

Definition 3.7. An algebra \mathbf{A} is idempotent if $f(x, \dots, x) = x$ for each basic operation f . Note that \mathbf{A} cannot have constants, by definition, if $|A| > 1$.

Lemma 3.8. *The existence of a near-unanimity term for idempotent algebras is decidable if and only if it is decidable for simple idempotent algebras.*

In order to prove this result we need the following definition and lemma which describe a way to compose NU-terms.

Definition 3.9. Let $s(x_1, \dots, x_n)$ and $t(y_1, \dots, y_m)$ be terms in n and m variables, respectively. Their *star product* $s \star t$ is a term in nm variables defined as

$$(s \star t)(z_{11}, \dots, z_{nm}) = s(t(z_{11}, \dots, z_{1m}), \dots, t(z_{n1}, \dots, z_{nm})).$$

Lemma 3.10. *Let \mathbf{A} and \mathbf{B} be similar idempotent algebras. If s and t are near-unanimity terms of \mathbf{A} and \mathbf{B} , respectively, then $s \star t$ is a near-unanimity term of both \mathbf{A} and \mathbf{B} .*

Proof. First we prove the claim for \mathbf{A} . Let $a, b \in A$, and put $c = (s \star t)(a, b, \dots, b)$. We want to show that $c = b$. Notice that this is enough, as we did not assume any ordering of the variables of s and t . By definition,

$$c = s(t(a, b, \dots, b), t(b, \dots, b), \dots, t(b, \dots, b)).$$

Since \mathbf{A} is idempotent, $t(b, \dots, b) = b$, and $c = s(t(a, b, \dots, b), b, \dots, b)$. As s is a NU-term, we conclude that $c = b$. The proof for \mathbf{B} is similar. \square

Proof of Lemma 3.8. One direction is trivial. For the other direction assume that the problem is decidable for simple idempotent algebras, and let \mathbf{A} be a finite idempotent algebra, which is not simple. The decision procedure we present is recursive; we assume that for all algebras of cardinality less than of \mathbf{A} we can decide the problem.

Let ϑ be a nontrivial congruence of \mathbf{A} , and B be a congruence block of ϑ . We claim that B is a subuniverse of \mathbf{A} . Indeed, for each basic operation f and elements $b_1, \dots, b_k \in B$, $f(b_1, \dots, b_k) \vartheta f(b_1, \dots, b_1) = b_1$. Note that, by Definition 3.7, f cannot be a constant.

Denote by \mathbf{B} the subalgebra of \mathbf{A} on the set B . If \mathbf{A} has a NU-term, then the same term is a NU-term for \mathbf{B} . Similarly, the same term is a NU-term for \mathbf{A}/ϑ . Therefore a necessary condition for \mathbf{A} to have a NU-term is that each proper subalgebra and proper homomorphic image of \mathbf{A} have a NU-term. We will show that this condition is sufficient, as well.

Let t_1, \dots, t_n be NU-terms on the nontrivial congruence blocks of ϑ , respectively, and s be a NU-term on \mathbf{A}/ϑ . By Lemma 3.10, the term $t = t_1 \star (t_2 \star (\dots (t_{n-1} \star t_n) \dots))$ is a NU-term on each congruence block of ϑ . We claim that $t \star s$ is a NU-term on \mathbf{A} . Take $a, b \in A$. Since s is idempotent on A and a NU-term of \mathbf{A}/ϑ ,

$$(t \star s)(a, \dots, a, b, a, \dots, a) = t(a, \dots, a, b', a, \dots, a) = a$$

for some element $b' = s(a, \dots, a, b, a, \dots, a) \vartheta a$. \square

We call an algebra \mathbf{A} *strictly simple* if it is simple and has no non-trivial subalgebras. By a non-trivial subalgebra we mean a proper subalgebra having at least two elements.

Theorem 3.11 (Á. Szendrei [14, 15]). *Let \mathbf{A} be a finite idempotent strictly simple algebra. Then the clone of \mathbf{A} is one of the following clones.*

- (1) $|A| = 2$ and $\text{Clo } \mathbf{A}$ is the trivial clone $[\text{id}]$.

For a vector space \mathbf{V} denote by $\text{End } \mathbf{V}$ the ring of endomorphisms of \mathbf{V} , and by ${}_{(\text{End } \mathbf{V})} \mathbf{V}$ the left module over $\text{End } \mathbf{V}$.

- (2) A finite dimensional vector space $\mathbf{V} = \langle A; +, K \rangle$ over a finite field K can be defined on A , and $\text{Clo } \mathbf{A}$ is the clone $\text{Clo}_{\text{id}}({}_{(\text{End } \mathbf{V})} \mathbf{V})$ of idempotent operations of ${}_{(\text{End } \mathbf{V})} \mathbf{V}$.

For a permutation group G on A let $\mathcal{R}_{\text{id}}(G)$ denote the clone of all idempotent operations f on A such that f admits each member of G as an automorphism.

- (3) $\text{Clo } \mathbf{A} = \mathcal{R}_{\text{id}}(G)$ for some permutation group G on A such that every non-identity member of G has at most one fixed point.

Let $0 \in A$ be some fixed element. For $k \geq 2$ put

$$\chi_k^0 = \{ (a_1, \dots, a_k) \in A^k : a_i = 0 \text{ for at least one } i, 1 \leq i \leq k \}.$$

Denote by \mathcal{F}_k^0 the clone of all operations on A preserving the relation χ_k^0 . Furthermore, put $\mathcal{F}_\omega^0 = \bigcap_{k=2}^\infty \mathcal{F}_k^0$.

- (4) $\text{Clo } \mathbf{A} = \mathcal{R}_{\text{id}}(G) \cap \mathcal{F}_k^0$ for some k ($2 \leq k \leq \omega$), some element $0 \in A$, and some permutation group G on A such that 0 is the unique fixed point of every non-identity member of G .

For a relation ϱ on A let \mathcal{P}_ϱ denote the clone of operations on A preserving ϱ .

- (5) $|A| = 2$ and $\text{Clo } \mathbf{A} = \mathcal{R}_{\text{id}}(G) \cap \mathcal{P}_\leq$ for some permutation group G on A ; or $|A| = 2$ and $\text{Clo } \mathbf{A} = \mathcal{R}_{\text{id}}(\{\text{id}\}) \cap \mathcal{P}_\leq \cap \mathcal{F}_k^0$ for some k ($2 \leq k \leq \omega$) and some element $0 \in A$.
- (6) $|A| = 2$ and $\text{Clo } \mathbf{A}$ is the clone $[\vee]$ generated by the join operation; or $|A| = 2$ and $\text{Clo } \mathbf{A}$ is the clone $[\wedge]$ generated by the meet operation. \square

Proposition 3.12. *The near-unanimity problem for idempotent, strictly simple algebras is decidable.*

Proof. Let \mathbf{A} be a idempotent, strictly simple algebra. We will use classification of Theorem 3.11 in the following decision procedure.

Assume that \mathbf{A} has a compatible partial order relation with zero and unit. Clearly, this condition is decidable. Then by Corollary 3.4 the NU-problem is decidable. This handles the cases (1), (5) and (6) of Theorem 3.11.

Recall that the algebra \mathbf{A} is called affine if it has a compatible affine relation. This is also a decidable property of \mathbf{A} . In Proposition 3.5 we have seen that if \mathbf{A} is affine, then it has no NU-term. This handles case (2) of Theorem 3.11, because in that case $\text{Clo } \mathbf{A}$ has a compatible affine relation.

If neither of the previous two conditions hold, then by Theorem 3.11 we know that $\text{Clo } \mathbf{A}$ is of type (3) or (4). In the rest of the proof we will show that the NU-problem is decidable even in these two cases.

CLAIM 1. *Assume that $\text{Clo } \mathbf{A} = \mathcal{R}_{\text{id}}(G)$ as described in case (3) of Theorem 3.11. Then \mathbf{A} has a ternary NU-term.*

Consider the function $f : A^3 \rightarrow A$, defined as

$$f(a, b, c) = \begin{cases} \text{maj}(a, b, c) & \text{if the majority exists,} \\ a & \text{otherwise.} \end{cases}$$

Clearly, f is a NU-term and admits all permutations on A .

CLAIM 2. *Assume that $\text{Clo } \mathbf{A} = \mathcal{R}_{\text{id}}(G) \cap \mathcal{F}_k^0$ as described in case (4) of Theorem 3.11, and $k < \omega$. Then \mathbf{A} has a NU-term.*

Consider the function $f : A^{k+1} \rightarrow A$, defined as

$$f(a_1, \dots, a_{k+1}) = \begin{cases} 0 & \text{if } a_i = a_j = 0 \text{ for some } i \neq j, \\ \text{maj}(a_1, \dots, a_{k+1}) & \text{else if the majority exists,} \\ a_1 & \text{otherwise.} \end{cases}$$

Clearly, f is a NU-term. By the description of case (4), the element 0 is a fixed point of every member of G . Therefore $f \in R_{\text{id}}(G)$. To show that $f \in \mathcal{F}_k^0$, take $\bar{a}^1, \dots, \bar{a}^{k+1} \in \chi_k^0$. By the Pigeon Hole Principle, there exist i, i' ($1 \leq i, i' \leq k+1$) and j ($1 \leq j \leq k$) such that $a_j^i = a_j^{i'} = 0$. This shows that $f(a_j^1, \dots, a_j^{k+1}) = 0$, therefore

$$\langle f(a_1^1, \dots, a_1^{k+1}), \dots, f(a_k^1, \dots, a_k^{k+1}) \rangle \in \chi_n^0.$$

CLAIM 3. *If $\text{Clo } \mathbf{A} \subseteq \mathcal{F}_\omega^0$ for some $0 \in A$, then \mathbf{A} has no NU-term.*

Assume the contrary, that $f \in \mathcal{F}_\omega^0$ is an n -ary NU-term. Take an element $a \in A \setminus \{0\}$, and consider the tuples $\bar{a}^i = \langle a, \dots, a, 0, a, \dots, a \rangle \in A^n$ for $1 \leq i \leq n$ where $a_i^i = 0$. Clearly, $\bar{a}^i \in \chi_n^0$, and

$$\langle f(a_1^1, \dots, a_1^n), \dots, f(a_n^1, \dots, a_n^n) \rangle = \langle a, \dots, a \rangle \notin \chi_n^0.$$

This shows that $f \notin \mathcal{F}_n^0$, which is a contradiction.

CLAIM 4. *Fix an element $0 \in A$. Then $\mathcal{F}_k^0 \supseteq \mathcal{F}_{k+1}^0$ for all $k \geq 2$.*

Take a function $f : A^n \rightarrow A$ preserving χ_{k+1}^0 . To show that it preserves χ_k^0 , as well, take $\bar{a}^1, \dots, \bar{a}^n \in \chi_k^0$. Put $\bar{b}^i = \langle \bar{a}^i, a_k^i \rangle = \langle a_1^i, \dots, a_k^i, a_k^i \rangle$ for $1 \leq i \leq n$. Clearly, $\bar{b}^i \in \chi_{k+1}^0$. Since f preserves χ_{k+1}^0 , the tuple

$$\begin{aligned} & \langle f(b_1^1, \dots, b_1^n), \dots, f(b_k^1, \dots, b_k^n), f(b_{k+1}^1, \dots, b_{k+1}^n) \rangle \\ &= \langle f(a_1^1, \dots, a_1^n), \dots, f(a_k^1, \dots, a_k^n), f(a_k^1, \dots, a_k^n) \rangle \end{aligned}$$

is in relation χ_{k+1}^0 . This means that $\langle f(a_1^1, \dots, a_1^n), \dots, f(a_k^1, \dots, a_k^n) \rangle \in \chi_k^0$, which is what we wanted to show.

CLAIM 5. *Let f be an n -ary function on A , and $0 \in A$. If $f \in \mathcal{F}_n^0$, then $f \in \mathcal{F}_k^0$ for all $n \leq k \leq \omega$.*

Fix k such that $n \leq k < \omega$, and take $\bar{a}^1, \dots, \bar{a}^n \in \chi_k^0$. By definition, there exists a “choice function” $\zeta : \{1, \dots, n\} \rightarrow \{1, \dots, k\}$ such that $a_{\zeta(i)}^i = 0$ for all $1 \leq i \leq n$. Put $\bar{b}^i = \langle a_{\zeta(1)}^i, \dots, a_{\zeta(n)}^i \rangle$ for $1 \leq i \leq n$. Since $b_i^i = 0$, $\bar{b}^i \in \chi_n^0$. By our hypothesis,

$$\langle f(b_1^1, \dots, b_1^n), \dots, f(b_n^1, \dots, b_n^n) \rangle \in \chi_n^0.$$

This means that $f(b_j^1, \dots, b_j^n) = 0$ for some $1 \leq j \leq n$, therefore $f(a_{\zeta(j)}^1, \dots, a_{\zeta(j)}^n) = 0$. Hence $f \in \mathcal{F}_k^0$. Finally, since $\mathcal{F}_2^0 \supseteq \dots \supseteq \mathcal{F}_\omega^0$ and $f \in \mathcal{F}_k^0$ for all $n \leq k < \omega$, $f \in \mathcal{F}_\omega^0$.

CLAIM 6. *Assume that $\text{Clo } \mathbf{A}$ is of type (3) or (4) as described in Theorem 3.11. Then it is decidable if \mathbf{A} has a NU-term.*

First we check if \mathbf{A} has a ternary NU-term. If it does, then we are done. Assume that \mathbf{A} has no ternary NU-term. Then by Claim 1, $\text{Clo } \mathbf{A}$ is of type (4). Moreover, by Claims 2 and 3, \mathbf{A} has no NU-term if and only if $\text{Clo } \mathbf{A} \subseteq \mathcal{F}_\omega^0$ for some $0 \in A$.

Now we show that, given $0 \in A$, it is decidable if $\text{Clo } \mathbf{A} \subseteq \mathcal{F}_\omega^0$. Take a basic n -ary operation f of \mathbf{A} . Clearly, we can decide if $f \in \mathcal{F}_n^0$. If $f \in \mathcal{F}_n^0$, then $f \in \mathcal{F}_\omega^0$, otherwise $f \notin \mathcal{F}_\omega^0$. So, $\text{Clo } \mathbf{A} \subseteq \mathcal{F}_\omega^0$ if and only if $f \in \mathcal{F}_n^0$ for all basic operations $f(x_1, \dots, x_n)$. \square

REFERENCES

- [1] S. Burris and H. P. Sankappanavar, *A course in universal algebra*. Graduate Texts in Mathematics **78**, Springer-Verlag, New York, 1981.
- [2] B. A. Davey, *Duality theory on ten dollars a day*. Algebras and Orders (Montreal, 1991), NATO Advanced Study Institute Series, Series C, **389**, 71–111.
- [3] B. A. Davey, L. Heindorf and R. McKenzie, *Near unanimity: an obstacle to general duality theory*. Algebra Universalis, **33** (1995), 428–439.
- [4] B. A. Davey and H. Werner, *Dualities and equivalences for varieties of algebras*. Contributions to lattice theory (Szeged, 1980), Colloq. Math. Soc. János Bolyai, **33** (1983), 101–275.
- [5] J. Ježek and M. Maróti, *Membership problems for finite entropic groupoids*. (manuscript).
- [6] O. G. Kharlampovich and M. V. Sapir, *Algorithmic problems in varieties*. Int. Journal of Algebra and Comp., **5** (1995), 379–602.
- [7] R. McKenzie, *Is the presence of a nu-term a decidable property of a finite algebra?* October 15, 1997 (manuscript).
- [8] R. McKenzie, G. McNulty and W. Taylor, *Algebras, Lattices, Varieties, Volume I*. Wadsworth & Brooks/Cole, Monterey, CA, 1987.
- [9] M. L. Minsky, *Recursive unsolvability of Post’s problem of “tag” and other topics in the theory of Turing Machines*. Ann. Math., **74** (1961), 437–455.
- [10] M. L. Minsky, *Computations: finite and infinite machines*. Prentice-Hall, Englewood Cliffs, N.J., 1967.
- [11] R. W. Quackenbush, *A new proof of Rosenberg’s primal algebra characterization theorem*. Finite algebra and multiple-valued logic (Szeged, 1979), Colloq. Math. Soc. János Bolyai, **28** (1981), 603–634.
- [12] M. O. Rabin and D. Scott, *Finite automata and their decision problems*. IBM Journal of Res. and Devel., **3**(2) (1969), 114–125.
- [13] I. Rosenberg, *Über die funktionale Vollständigkeit in den mehrwertigen Logiken*. Rozprawy Československe Akad. Věd., Ser. Math. Nat. Sci., **80** (1970), 3–93.
- [14] Á. Szendrei, *Idempotent algebras with restrictions on subalgebras*. Acta Sci. Math. (Szeged) **51** (1987), 251–268.
- [15] Á. Szendrei, *Term minimal algebras*. Algebra Universalis, **32** (1994), 439–477.

BOLYAI INSTITUTE, UNIVERSITY OF SZEGED, ARADI VÉRTANÚK TERE 1., 6720 SZEGED, HUNGARY

E-mail address: mmaroti@math.u-szeged.hu