

Expectancies toward the intake of alcohol in young adults of Bucaramanga, Colombia, 2005*

OSCAR F. HERRÁN, MSc EPIDEMIOLOG^{1,2}, MARÍA F. ARDILA, MSc EPIDEMIOLOG²

SUMMARY

Objective: To settle down in young adults of Bucaramanga if the statistical and conceptual performance of the test AEQ-III is similar to the one reached in young adults from Mexico, and to describe the relationship of these expectancies with some variables of the context.

Methods: Study of factorial validation and internal consistency. In Bucaramanga, Colombia, 156 young adults, answered 51 items of the alcohol expectancy questionnaire. A model of structural equation was developed to evaluate the goodness of fit of the factorial solution through the reason of verisimilitude Chi-square (X^2) and the half quadratic approach error (RMSEA). The global internal consistency and intra expectancies were carried out with the alpha of Cronbach. The scores for the expectancies were calculated with base in additives scales.

Results: The value reached for the X^2 was 6921, with 1196 grades of freedom, $p=0.000$. The RMSEA was 0.18. The global internal consistency and for the factors, it was bigger in this sample with regard to the Mexican. Equally, all the scores means was bigger in each factor and the smallest variability ($p<0.001$).

Conclusion: In young adults of Bucaramanga, the internal consistency in the eight expectancies is good. However, the statistical coherence is poor. It is necessary to adapt the questionnaire to Bucaramanga culture and in other age groups, before applying their results.

Keywords: Alcohol Expectancy Questionnaires (AEQ); Alcohol drinking; Cultural factors; Psychometrics; Colombia.

Expectativas hacia el consumo de alcohol en adultos jóvenes de Bucaramanga, Colombia, 2005

RESUMEN

Objetivos: Establecer en adultos jóvenes de Bucaramanga si el desempeño estadístico y conceptual de la prueba «Expectativas positivas hacia el consumo de alcohol -AEQ-III» es similar al alcanzado en adultos jóvenes de México, y describir la relación de estas expectativas con algunas variables del contexto.

Métodos: Estudio de validación factorial y consistencia interna. En Bucaramanga 156 adultos jóvenes contestaron 51 ítems del formulario AEQ-III. Se desarrolló una ecuación estructural para evaluar la bondad de ajuste a través de la razón de verosimilitud Chi-cuadrado (X^2) y el error de aproximación cuadrático medio (RMSEA). La consistencia interna global e intra expectativas se realizó con el alfa de Cronbach. Los puntajes para las expectativas se calcularon con base en escalas aditivas.

Resultados: El valor alcanzado para X^2 fue 6921, con 1196 grados de libertad, $p=0,000$. El RMSEA fue 0.18. La consistencia interna global y para los factores fue mayor en esta muestra con respecto a la mexicana. Igualmente, todos los puntajes medios alcanzados en las ocho expectativas fueron mayores y la variabilidad menor ($p<0.000$).

Conclusión: En adultos jóvenes de Bucaramanga, la consistencia interna en las ocho expectativas es adecuada. Sin embargo, la coherencia estadística del modelo es pobre. En este grupo de edad es necesario adaptar el formato AEQ-III antes de utilizarlo en estudios epidemiológicos.

Palabras clave: Cuestionario de Expectativas hacia el Alcohol (AEQ); Factores culturales; Consumo de bebidas alcohólicas; Psicometría; Colombia.

* This study was co-sponsored by the Epidemiologic Investigations Center (CIE-UIS) and the investigation group «Epidemiological Observatory of Cardiovascular Diseases» (COLCIENCIAS, Category A) of the Universidad Industrial de Santander (UIS), Bucaramanga, Colombia.

1. Profesor Titular, Escuela de Nutrición y Dietética Universidad Industrial de Santander, Bucaramanga, Colombia.
e-mail: herran@uis.edu.co
 2. Miembro del Grupo Observatorio Epidemiológico de Enfermedades Cardiovasculares, Centro de Investigaciones Epidemiológicas, Universidad Industrial de Santander, Bucaramanga, Colombia.
- Received for publication April 11, 2008 Accepted for publication July 1, 2009

The prevalence of alcohol intake in the Colombian population ranges between 80% and 95%¹⁻⁵. The starting age for alcohol intake (AC) as well as for the first occurrence of alcohol intoxication has descended to 13 during the last decade^{1,3,4}. Due to the cultural characteristics of the Bucaramanga population and their habits towards AC, it takes place every seven days in amounts that reach toxic levels¹. AC in toxic levels has been associated to brawls, violence in the family, problems that require the police presence, traffic accidents and unwanted situations with sexual or erotic components. All of these instances are considered as «problematic consumption» and are associated to an increase of family and social dysfunction, as well as morbidity and mortality due to trauma and violence^{4,5}.

In Colombia, problematic consumption and alcohol dependence or alcoholism are public health problems⁴. During 2003, in Bucaramanga, 15.6% of the alcohol-consuming population between the ages of 18 and 60 was classified as alcoholic³. According to National Studies on Mental Health and psychoactive substance consumption, (NSMH) carried out in 1993 and 1999, alcoholism was 7.1% and 12% respectively. According to the 1993 NSMH, for every six men there was a woman at high risk of alcoholism. In Bucaramanga, the ratio is one woman for every three men⁶.

Biological, economic and socio-demographic variables are poorly related to AC¹, for this reason AC analysis and the actions to decrease it or to delay its onset or the moment when alcohol intoxication occurs are complex. The study of problematic consumption and alcohol dependence belongs to the field of addictions⁷. A contemporary psychosocial approximation in the field of addictions is the study of positive expectancies in relation to alcohol intake. Expectancies are defined as «*the anticipation of a systematic relationship between objects or events in a future situation*»⁷. In the practice, expectancies are individual beliefs, mediated by learning, that have a causal status on the expected effects of AC^{7,8}. The development of expectancies begins at an early age with a series of diffuse and global beliefs in AC. They increase and particularize with age, exposure to AC, and the family, social and cultural contexts in which the individual learns about AC. The expectancies become relevant as they are involved in the starting and continuation of AC during adolescence and determine AC in adults differentially^{5,7-9}.

Expectancies are predictors of AC and they change with culture, AC habits and sex, and have been studied through personal information questionnaires, with the Alcohol Expectancy Questionnaire (AEQ-III) being probably the most widely used questionnaire⁷. Moderate alcohol consumers have expectancies that enhance social pleasure and the positive effects of AC, while problematic consumers and alcoholics expect that AC will lower physical and psychological stress, increase their aggressiveness and reduce their inhibitions in general. Furthermore, men reach higher scores than women in their approach to alcohol as a facilitator of group integration and the increase of sexuality⁸.

Despite the fact that research in the field of expectancies has been done for three decades in South America there are few reports relating them to AC. The study of expectancies of AC is crucial in order to understand it and carry out a rational intervention in the use and abuse of alcohol at an early age^{5,7,8}. The objectives of this study were to establish if for young adults in Bucaramanga the statistical and conceptual performance of the test AEQ-III is similar to that obtained in young adults in Mexico and to describe the relation between these expectancies and some variables of the context.

MATERIALS AND METHODS

During 2005 a study of factorial validation and internal consistency was carried out with cross section data collected in Bucaramanga, Colombia.

Selection of the subjects. A total of 156 young adults, aged 18 to 25, were selected through random sampling in different stages in order to obtain equal representation of both sexes. The size of this sample was calculated and based on the results of the study conducted in Mexico⁸. An alpha value of 0.05 and a power of 0.80 allow us to obtain score differences of 1.2 or higher between the groups for the expectancies of AC. The subjects that met the following criteria were classified as eligible:

1. To be between 18 and 25 years old.
2. To have resided in the city for two or more years.
3. Not to be attending any support group such as Alcoholics Anonymous.
4. To express their will to participate by supplying the information voluntarily.

Certain blocks of the city and some houses were randomly selected according to socioeconomic classes. For every household there was a census of eligible subjects and one subject per household was chosen at random. If the person refused to take part in the study, another person was taken.

The subjects were selected from an existing sample framework for Bucaramanga after successive random approximations: socioeconomic class, block, household and census of eligible subjects. This allows us to maintain that there were no biases in the selection process. After choosing the subjects, informing them about the objectives of the study, guaranteeing confidentiality and obtaining their written consent, they were polled. Both the project and the procedures were approved by the committee of research ethics of the Universidad Industrial de Santander.

Information sources. Pollsters who were previously trained in data collection in population studies applied three questionnaires:

1. A poll on biological variables (BV) that included the Body Mass Index (BMI), socioeconomic variables (SEV), and diet (D), including the frequency of alcoholic beverage intake and the preference for them.
2. The CAGE test, widely used in population and clinical studies⁶.
3. The AEQ-III⁷.

The AEQ-III was designed in 1980 and was perfected in 1987 by Goldman *et al.*⁷. The pollsters required no specific training, and the 120 items can be answered through a self-report or a direct 20- to 30-minute interview. The AEQ-III assesses the hope (expectancies) that determines AC and measures the degree to which an individual expects a variety of general and specific effects when consuming alcohol. There are reproducibility studies for the AEQ-III; it has a coefficient $r=0.64$ (test-retest) and an internal consistency between 0.72 and 0.92^{7,8}. Moreover, there are studies of criterion validity (concurrent and predictive) and its results are stable when they are done at a universal scale⁷⁻⁹. The response format is dichotomic. The AEQ version that was applied is the third one for adult population. It was retro-translated into English-Spanish independently by two experts based on the last original version developed in 1987⁷. The adaptation to the Bucaramanga jargon was carried out with basis on this translation and the Mexican version⁸; furthermore, it was adjusted during a pilot

study carried out with 30 adults in the target population (five subjects per socioeconomic class).

Quality of the data. The four items of CAGE were retaken from those utilized in the national consumption polls. CAGE has a study of internal validity and reproducibility in the population of Bucaramanga⁶. The polls on BV, SEV, D and CAGE were carried out sequentially. The AEQ-III was the last one to be applied. The objective of the sequential application was to reduce the likelihood of bias by the informer, as in this type of poll it is advisable to begin with the simplest polls and to finish with the most complex ones, which address aspects about the subjects' private life.

The preference shown for alcoholic beverages (AB) was the perception that every subject has about his/her intake. It corresponds to a social representation¹⁰. In order to permit further comparisons of the results, the risk of alcoholism was defined as low when a CAGE score under 3 was obtained and high when the score was 3 or 4. This cut off was proposed by the original version of CAGE⁶ and it is used in Colombia in the nationwide polls^{3,4}. The scores in the expectancies of AC were obtained in an additive scale. Neither the polled subjects nor the pollsters knew the results of CAGE or AEQ-III and the data were only gathered through codes at the moment of carrying out the statistical analysis.

Statistical analysis. To describe the categories for BV, SEV, D and CAGE, descriptive measures and their confidence intervals (CI) were calculated at 95%. In order to establish if there were statistically significant differences between BV, SEV, D and CAGE according to sex, the *t Student* and X^2 tests were performed. The factorial structure of AEQ developed in Mexico⁸ was assessed in the sample through a Structural Equation Model (SEM)¹¹ that made it possible to establish the factorial loads of the variables that constitute each of the eight expectancies or predefined factors. The goodness of fit of the factorial model was evaluated through the minimum fit function X^2 and the Root Mean Square Error of Approximation (RMSEA)¹¹. With the purpose of determining the global internal consistency and the factors (expectancies) underlying the AEQ-III, the Cronbach's alpha coefficient was calculated¹². The correlation between expectancies was determined through Pearson's correlation coefficient *r*. In order to establish if there were differences between the scores obtained in the expectancies in the Mexican factorial solution and

that of Bucaramanga, the *t Student* tests were used. Finally, with the aim of establishing if there was an association between the expectancies and the BV, SEV, D and CAGE, *t Student*, X^2 and ANOVA tests were carried out.

All of the records were typed in duplicate and debugged in the programme VALIDATE of EpiInfo 6.04d¹³. The data processing, together with the analysis, were carried out through STATA 9.1/SE¹⁴. The SEM was carried out with LISREL¹⁵.

RESULTS

All the 156 subjects answered the AEQ-III completely: between 21 and 39 per socioeconomic class. Due the estimated values were not affected by correcting for the effect of sample design is similar to those reported from the simple random sampling.

Characteristics of the target population. The mean age of the 78 polled men was not different from that of the women ($p=0.84$). The proportion of subjects with incomplete primary studies was 1.3% and that of subjects with a secondary education (11 years of schooling) was 26.3%. Only 52% of the subjects were at some level of university education and 2% had finished their higher education. The mean starting age in AC was 15.7 (CI 95%; 15.2 to 16.2), a year later for women ($p=0.01$). The age of the first occurrence of alcohol intoxication was a year after starting AC, with the difference between the sexes remaining constant ($p=0.02$). 11% of the population stated that they had not consumed any alcohol during the last year. Men have 1.7 more probabilities of being overweight and four times more probabilities of consuming alcohol every two weeks than women.

No differences were reported between the two sexes in relation to intake or non-intake of alcoholic beverages ($p=0.07$), schooling ($p=0.28$), the occurrence of any medical diagnosis ($p=1.0$), the fact of having gone on any type of diet during the last year ($p=0.81$), the preference for AC ($p=0.59$) or the high risk of alcoholism according to CAGE (0.73). Table 1 shows other characteristics of the target population.

Assessment of the factorial structure developed in Mexico⁸. Table 2 shows the variables grouped in the eight factors assessed through a SEM. The ratio of verisimilitude X^2 is the only statistically supported measure of goodness of fit in a SEM. A high value means

that the observed and estimated matrices differ considerably. The value for this statistic was 6921 with 1196 degrees of freedom, $p=0.00$. An ideal global adjustment value in a factorial model is obtained with RMSEA statistical values below 0.08. The calculated RMSEA was 0.18 (CI 95%; 0.17 to 0.18).

In order for the present study to be statistically significant, taking into account the size of the sample, the factorial load must be at least 0.41¹¹. All the factorial loads were below this value, with a minimum of 0.21 and a maximum of 0.33. However, the internal global consistency and the consistency for the factors, except the factor *Psychological Changes*, were higher in this sample than in the Mexican one⁸.

Table 3 presents the internal consistency obtained in each factor and in the AEQ-III as a global test, and the scores obtained in each of them by the young adults from Bucaramanga. All the scores in Bucaramanga were higher for each factor, and the variability was lower ($p<0.00$) than that obtained by the Mexican subjects.

Just as expected, the correlation r between the expectancies was high, with a minimum of 0.54 (between increase in the verbal expression and psychophysiological changes) and a maximum of 0.86 (between an increase in sexuality and aggressiveness and feelings of power). The matrix of correlation r between the studied factors is presented in Table 4.

Association between expectancies of AC and biological, socio-demographic and dietary variables. The score obtained in the eight evaluated expectancies does not differ according to age, intake or non-intake of alcohol, the age of the first occurrence of alcohol intoxication, the sex, the socio-economic class, or the risk of alcoholism in accordance with CAGE ($p>0.05$ for all).

The BMI and the starting age in AC were inversely associated to the score obtained in all the expectancies ($p<0.05$). The subjects who stated that they had gone on a diet during the last year obtained lower scores in the expectancies «increase in the verbal expression», «increase in sexuality», «reduction of psychological tension» and «aggressiveness and feelings of power», ($p<0.05$ for all).

The higher the level of schooling, the higher the score obtained in the expectancies «group interaction», «no inhibition», «reduction of psychological tension», and «psycho-physiological changes» ($p>0.05$ for all). The

Table 1
Characteristic of the studied population

Variable	Total	Male [n=78]	Female[n=78]
Age (years)	22.7 (22.3-23.1) ¹	22.7 (22.1-23.2)	22.8 (22.2-23.3)
Schooling (years)	12.0 (11.6-12.4)	12.3 (11.7-12.9)	11.8 (11.3-12.4)
Weight (kg)	64.8 (63.3-66.4)	70.1 (68.1-72.0)	59.6 (57.8-61.4)
BMI (kg/height ²)	23.5 (23.0-24.0)	24.2 (23.6-24.9)	22.8 (22.1-23.5)
> 25	59 {37.8} ²	37 {47.4}	22 {28.2}
Medical dx. ³ (yes)	4 {2.6}	2 {2.6}	2 {2.6}
Diet (yes)	19 {12.2}	10 {12.8}	9 {11.5}
Like for alcoholic drinks			
A little	56 {35.9}	31 {39.7}	25 {32.0}
Moderate	90 {57.7}	42 {53.9}	48 {61.6}
A lot	10 {6.4}	5 {6.4}	5 {6.4}
Frequency of intake of alcohol			
Weekly	18 {11.5}	10 {12.8}	8 {10.3}
Biweekly	35 {22.4}	28 {35.9}	7 {9.0}
Monthly	103 {66.1}	40 {51.3}	63 {80.7}
Intake of alcohol ⁴			
No	17 {10.9}	5 {6.4}	12 {15.4}
Alcoholism ⁵	9 {5.8}	4 {5.1}	5 {6.4}

1. Mean and 95% confidence interval (CI). 2. Frequency (%). 3. Illness or disorder with diagnosis carried out by a doctor. 4. In the last year. 5. High risk of alcoholism according to CAGE ≥ 3

incidence of AC every two weeks was associated with a better score in «increase in sexual activity» ($p=0.04$). A low preference for alcoholic beverages is connected to lower scores in all the expectancies ($p<0.05$).

DISCUSSION

The results were obtained from a population of young adult subjects who resided in the urban area of Bucaramanga. They were literate, consumers and non consumers of alcohol and belonged to the six different socio-economic classes. Due to the design and data collection methods, a selection or information bias is very unlikely. However, in case there is such a bias, it is not differential because there are no differences in the rates of answer or participation according to sex or socio-economic class, two of the defining variables taken into account in the selection of the subjects.

Due to the age range studied, the level of schooling was probably higher in our sample. Consequently, the

rate of incomplete primary education was 1.3%, one third of that for the general population, which is 4.7%¹⁶. The main limitation of the study is that while the Mexican factorial solution was established in university students, this level was obtained only by 54% of the studied sample. However, 26% of the population had completed their secondary education and this condition allowed us to evaluate the relationship between schooling and expectancies. The results confirmed the findings of other studies^{5,8}, which show how the population with the highest schooling (those who went to university) increased their positive expectancies towards AC; therefore, there are more probabilities of an increase in AC, problematic consumption and the beginning of dependence among university students. In Bucaramanga and Colombia there is a debate about selling alcoholic beverages in the vicinity of universities and educational centres in general because of the evidence of associated problematic consumption.

The older the subject when he/she begins AC and

Table 2
Answer in the AEQ-III (Item), grouping of variables and factors evaluated in young adults
of Bucaramanga, Colombia

Item* Interaction groupal

- 105 Alcohol enables me to have a better time at parties.
 - 86 Drinking makes get-togethers more fun.
 - 107 Drinking makes people feel more at ease in social situations.
 - 48 Having a few drinks helps me relax in a social situation.
 - 64 After a few drinks, I am usually in a better mood.
 - 99 It is easier to remember funny stories or jokes when I have been drinking.
 - 53 There is more camaraderie in a group of people who have been drinking.
 - 87 Alcohol makes it easier to forget bad feelings.
 - 58 Drinking increases female aggressiveness.
-

Item Increase in the verbal expression

- 73 A few drinks make me feel less shy.
 - 92 A couple of drinks make me feel more outgoing.
 - 32 It is easier for me to meet new people if I've been drinking.
 - 36 If I have a couple of drinks, it is easier to express my feelings.
 - 97 Opposite sex is easier for me after I have had a few drinks.
-

Item No inhibition

- 101 Alcohol makes me more talkative.
 - 120 Things seem funnier when I have been drinking, or at least I laugh more.
 - 94 Alcohol decreases my hostilities.
 - 112 Alcohol makes it easier to act impulsively or make decisions quickly.
 - 95 Alcohol makes me feel closer to people.
-

Item Increase in sexuality

- 59 I am a better lover after a few drinks.
 - 79 I enjoy having sex more if I have had some alcohol.
 - 81 I feel more masculine/feminine after a few drinks.
 - 88 After a few drinks, I am more sexually responsive.
 - 69 After I have had a couple of drinks, I feel I am more of a caring, sharing person.
 - 84 Sometimes when I drink alone or with one other person it is easy to feel cozy and romantic.
 - 35 Alcohol makes women more sensuous.
 - 66 Women can have orgasms more easily if they have been drinking.
 - 103 Men can have orgasms more easily if they have had a drink.
-

Table 2
Answer in the AEQ-III (Item), grouping of variables and factors evaluated in young adults
of Bucaramanga, Colombia (continuation)

Item Reduction of psychological tension	
74	If I am tense or anxious, having a few drinks makes me feel better.
116	A drink or two can slow me down, so I do not feel so rushed or pressured for time.
85	I feel like a more happy-go-lucky person when I drink.
111	Alcohol makes me worry less.
50	Drinking alone or with one other person makes me feel calm and serene.
15	Drinking makes me feel good.
Item Reduction of the physical tension	
109	After a drink, things like muscle aches and pains do not hurt as much.
61	Alcohol decreases muscular tension.
78	Alcohol can act as an anesthetic; that is, it can deaden pain.
57	Alcohol helps me sleep better.
Item Aggressiveness and feelings of power	
76	If I am feeling afraid, alcohol decreases my fears.
91	I become lustful when I drink.
51	After a few drinks, I feel brave and more capable of fighting.
40	After a few drinks, I feel more self-reliant than usual.
100	After a few drinks, I am less submissive to those in positions of authority.
77	A couple of drinks make me more aroused or physiologically excited.
41	After a few drinks, I don't worry as much about what other people think of me.
54	My feelings of isolation and alienation decrease when I drink.
96	I tend to be less self-critical when I have something alcoholic to drink.
52	Drinking can make me more satisfied with myself.
Item Psycho-physiological changes	
98	Drinking makes me feel flushed.
10	Drinking makes me feel physically.
89	If I am cold, having a few drinks will give me a sense of warmth.

* Order of the question or item in the AEQ-III⁸

therefore his/her first occurrence of alcohol intoxication, the lower the score in the expectancies. This finding is particularly important for the Colombian population as the starting age in AC and the first occurrence of alcohol

intoxication have been reported even in subjects under ten years of age^{3,4}. This aspect should be tackled in secondary school as top priority, if we are to anticipate one of the main aspects associated to AC.

Table 3
Internal consistency, mean and standard deviation reached for the positive expectancies toward the intake of alcohol (factors) in young adults of Bucaramanga, Colombia

Factor	Cronbach's (α)		Mean (SD) ¹	
	Méx. ²	Col.	Méx.	Col.
AEQ-III	0.93	0.97		
Interaction of group	0.79	0.84	4.03 (4.59)	6.35 (2.72)
Increase in the verbal expression	0.77	0.77	1.98 (2.88)	3.44 (2.68)
No inhibition	0.75	0.75	2.60 (2.98)	3.46 (1.63)
Increase in sexuality	0.78	0.80	2.63 (5.82)	5.99 (2.64)
Reduction of psychological tension	0.76	0.80	2.30 (3.22)	4.31 (1.91)
Reduction of the physical tension	0.58	0.69	1.67 (2.56)	2.70 (2.35)
Aggressiveness and feelings of power	0.80	0.88	2.43 (5.12)	6.87 (3.20)
Psycho-physiological changes	0.71	0.64	4.23 (1.17)	2.02 (1.07)

1. Mean of score achieved. SD standard deviation. For all the comparisons among means value of $p < 0.001$. 2. Méx. México. Col. Bucaramanga, Colombia

The fact that the subjects with a higher BMI and those who reported any type of diet during the last year obtained lower scores in the expectancies is coherent with a better perception on the need to adopt or begin habits considered as wholesome, including lower motivation to AC. If the concept of «healthy school», recommended by the Pan American Health Organization¹⁷, were adopted by the school population, it might lead to lower expectancies towards AC in adolescents.

The preference for alcohol is an expression of two types of learning: genetic and social⁷. It is possible to argue that in family and cultural contexts where AC is moderate, does not exist or is only ceremony-associated, social learning predominates over genetic aspects of preference for alcohol; in that way, it would act as a protective factor. The Mexican study did not apply CAGE nor did it investigate problematic consumption; for this reason it was not possible to compare these relationships. Contrary to what was observed in the Mexican university students, there was no evidence of association between the scores obtained in the expectancies and the sex of the participants. It is possible that as a product of modernity, education patterns and attitudes towards AC in this population are no longer differential in childhood and adolescence or that the differentiation of expectancies according to sex takes place at an older age than those studied, which is an

Table 4
Correlation matrix of Pearson (r) among the positive expectancies toward the intake of alcohol in young adults of Bucaramanga, Colombia

	F1	F2	F3	F4	F5	F6	F7	F8
F1	1.00							
F2	0.76	1.00						
F3	0.78	0.76	1.00					
F4	0.82	0.79	0.75	1.00				
F5	0.82	0.78	0.79	0.78	1.00			
F6	0.74	0.79	0.72	0.76	0.75	1.00		
F7	0.76	0.85	0.79	0.86	0.78	0.79	1.00	
F8	0.64	0.55	0.68	0.68	0.59	0.61	0.64	1.00

F1 Interaction of group
 F2 Increase in the verbal expression
 F3 No inhibition F4 Increase in sexuality
 F5 Reduction of psychological tension
 F6 Reduction of the physical tension
 F7 Aggressiveness and feelings of power
 F8 Psycho-physiological changes

unlikely occurrence.

The fact of consuming alcohol was not associated to the scores obtained in the expectancies studied. It is possible that in this age group, where AC is more a part of the adolescent's exploration to adopt or reject behaviors than a habit towards AC^{5,7,8}, it is not viable to find any differences. Furthermore, this result is coherent with the mechanism in which an expectancy is constructed. At the initial stage of consumption, expectancies are more general and diffuse; with the habit of AC they become particular and very specific^{5,7}.

The global conceptual coherence (internal consistence) and the coherence for the eight expectancies evaluated are very satisfactory, even more satisfactory than that of the reference study carried out in Mexico⁸. The values obtained in Pearson's *r* between expectancies show their interdependence and the high internal global consistency of AEQ-III. The mean scores obtained for the expectancies in the population of young adults in Bucaramanga were consistently higher and with lower variability than those in the Mexican sample, but due to the type of design, it was not possible to verify if this translates into higher AC or into the prevalence of problematic consumption or alcoholism. In this sense, the execution of multi-centric studies between countries or regions within the same country with cultural diversity is a pertinent one.

The statistical coherence of the Mexican factorial solution was poor in the studied sample. This presupposes the execution of studies of factorial validity to adapt the AEQ-III to the Bucaramanga context and other population groups as an obligatory requisite for the adoption of intervention strategies based on the results. The fact that the calculated scores of the expectancies were additive scales and not based on the factorial loads, allowed the comparison and evaluation of the solution with eight factors.

All the ideas discussed until now are valid under an underlying premise and it is that the scores obtained in all the expectancies toward intake are directly related to the amount of alcohol consumed and problematic consumption⁷. As long as there are no studies that verify, confirm or reject these assumptions in Bucaramanga population, all the previously discussed ideas must be carefully observed as a work hypothesis. In short, through the application of AEQ-III, we have explored eight expectancies towards AC and its relationships with

biological, socioeconomic and dietary variables, as well as with the risk of alcoholism according to CAGE. It is necessary to adapt the AEQ-III to our culture and to guarantee statistical, conceptual and theoretical coherence as well as the reproducibility of this document before launching intervention strategies based on it¹⁸.

Interest conflict. The authors declare that there are no interest conflicts in this manuscript.

REFERENCES

- Herrán OF, Ardila MF. Consumo de alcohol, riesgo de alcoholismo y alcoholismo en Bucaramanga, Colombia, 2002-2003. *Colomb Med.* 2005; 36: 158-67.
- Herrán OF, Ardila MF. Validity and reproducibility of two semi-quantitative alcohol frequency questionnaires for the Colombian population. *Public Health Nutr.* 2006; 9: 763-70.
- Torres GY. Alcohol: Prevalencia de consumo y dependencia en Colombia. *Rev Med CES.* 1999; 12: 1-9.
- Presidencia de la República. Ministerio de Salud. Programa Rumbos. *Estudio Nacional Sobre Uso y Abuso de Alcohol y Consumo de Sustancias Psicoactivas.* Bogotá: Ministerio de Salud; 2000.
- Mora J, Natera G. Expectativas, consumo de alcohol y problemas asociados en estudiantes universitarios de la ciudad de México. *Salud Publica Mex.* 2001; 43: 89-96.
- Herrán OF, Ardila MF. Validez interna y reproducibilidad de la prueba CAGE en Bucaramanga, Colombia. *Biomedica.* 2005; 25: 231-41.
- Goldman MS, Brown SA, Christiansen BA. Expectancy theory: Thinking about drinking. En: Blane HT, Leonard KE, (eds.). *Psychological theories of drinking and alcoholism.* 2nd ed. New York: Guilford Press; 1999. p. 203-46.
- Mora-Ríos J, Natera G, Villatoro J, Villalvazo R. Validez factorial del cuestionario de expectativas hacia el alcohol (AEQ) en estudiantes universitarios. *Psicol Conduct.* 2000; 8: 319-28.
- Bierut LJ, Dinwiddie SH, Begleiter H, Crowe RR, Hesselbrock V, Nurnberger JI, et al. Familial transmission of substance dependence: alcohol, marijuana, cocaine and habitual smoking. *Arch Gen Psychiatry.* 1998; 55: 982-8.
- Vázquez G, Prada GE, Herrán OF. ¿Cuánto se percibe? Una aproximación a la ingesta dietaria. *Salud UIS.* 2002; 34: 89-98.
- Hair JF, Anderson RE, Tatham RL, Black WC. Modelos de ecuaciones estructurales. En: *Análisis multivariante.* 5ª ed. Bogotá: Prentice Hall; 1999. p. 611-90.
- Reynaldo J, Santos A. Cronbach's alpha: Tool for assessing the reliability of scales. *J Extens.* 1999; 37. (fecha de acceso abril 7 de 2006). Disponible en: <http://joe.org/joe/1999april/tt3.html>
- CDC. EpiInfo, version 6.04d. *Epidemiología en ordenadores.* Atlanta, Georgia. Enero, 2001.
- StataCorp. 2005. *Stata Statistical Software: Release 9.1.* College Station, TX: Stata Corporation.

15. Jöreskog K, Sörborm D. LISREL 8.70. *Scientific Software International, Inc*; 2004.
16. PROFAMILIA. Encuesta Nacional de Demografía y Salud. *Principales indicadores demográficos, socioeconómicos y de salud de la población, de las mujeres en edad fértil y de los niños menores de cinco años por zonas, regiones y departamentos, 2005*. Bogotá: PROFAMILIA; 2005.
17. Organización Panamericana de la Salud. *Escuela saludable: Una herramienta para la paz*. (fecha de acceso abril 7 de 2006). Disponible en: <http://www.col.ops-oms.org/juventudes/ESCUELASALUDABLE/>
18. Herrán OF, Ardila MF. Validez factorial del cuestionario de expectativas hacia el consumo de alcohol (AEQ-III), en adultos de Bucaramanga, Colombia. *Rev Med Chile*. 2007; 135: 488-95.