

Not to be cited without prior reference of the authors

Changes in North Sea macrofauna communities between 1986 and 2000

I. Kröncke¹, Reiss, H.¹, Bergman, M.J.N.², Cochrane, S.³, Craeymeersch, J.⁴,
Degræer, S.⁵, Desroy, M.⁶, Dewarumez, J.-M.⁶, Duineveld, G.², Eggleton, J.D.⁷,
Essink, K.⁸, Hillewaert, H.⁹, Lavaleye, M.², Moll, A.¹⁰, Nehring, S.¹¹, Newell, J.⁷,
Oug, E.¹², Pohlmann, T.¹¹, Rachor, E.¹³, Rees, H.L.⁷, Robertson, M.¹⁴, Rumohr,
H.¹⁵, Schratzberger, M.⁷, Smith, R.⁷, van den Berghe, E.¹⁶, van Dalfsen, J.¹⁷, van
Hoey, G.⁵, Vincx, M.⁵, Willems, W.⁵

¹Senckenberg Institute, Dept. for Marine Research, Wilhelmshaven, Germany

²NIOZ, Den Burg-Texel, The Netherlands

³Akvaplan-Niva, Tromsö, Norway

⁴RIVO-CSO, Yerseke, The Netherlands

⁵University of Ghent, Marine Biology Section, Ghent, Belgium

⁶University of Lille, Institute for Marine Research, Wimereux, France

⁷CEFAS, Burnham-on-Crouch, UK

⁸RIKZ, Haren, The Netherlands

⁹ILVO-Fisheries, Ostende, Belgium

¹⁰Centre of Marine and Atmospheric Science, Hamburg, Germany

¹¹Bundesamt für Gewässerkunde, Koblenz, Germany

¹²NIVA, Grimstad, Norway

¹³AWI, Bremerhaven, Germany

¹⁴Fisheries Research Services, Marine Laboratory, Aberdeen, Scotland

¹⁵Leibnitz Institute for Marine Science, Kiel, Germany

¹⁶VLIZ, Ostende, Belgium

¹⁷TNO, Den Helder, The Netherlands

Abstract

This paper is chapter 5.2 of the ICES CRR “Structure and dynamics of the North Sea benthos” (ICES 2007) compiled by the ICES Study Group on the North Sea Benthos Project 2000. The North Sea Benthos Project 2000 (NSBP) was initiated as a follow-up to the earlier 1986 ICES North Sea Benthos Survey (NSBS). One major aim of the NSBP 2000 was to compare the outcome with that of the 1986 NSBS, in order to identify any significant changes in the community structure and their likely causes.

In general, the spatial distribution of the macrofaunal communities in 2000 was rather similar to that in 1986. But changes were found within communities which were addressed to changes in the hydroclimate caused by changes in the North Atlantic Oscillation influencing changes in currents and sediment structure as well as food availability.

Keywords: macrofauna communities, long-term variability, NAOI, sediments, North Sea

Introduction

Data sets covering the whole North Sea are very limited. For the infauna, one data set available is the North Sea Benthos Survey (NSBS) in April/May 1986 initiated by the ICES Benthos Working Group (Künitzer et al. 1992, Heip & Craeymeersch 1995, Craeymeersch et al. 1997). The data from surveys during the years 1980 to 1985 in the northern North Sea by Basford & Eleftheriou (1988), Eleftheriou & Basford (1989) and Basford et al. (1990, 1993) were integrated in the NSBS. The North Sea Benthos Project (NSBP) 2000 was initiated as a follow-up to the earlier 1986 NSBS. One major aim of the NSBP 2000 was to compare the outcome with that of the 1986 NSBS, in order to identify any significant changes and their likely causes. Thus, the 1986 data were re-analysed and compared with the 2000 data and possible causal factors for observed differences are discussed.

Methods

Most stations were sampled with a 0.1 m² van Veen grab. Two replicates were taken per station. Part of the samples were taken with a Hamon grab or a box corer (see van den Berghe 2007). The samples were washed over 1 mm mesh size and fixed in 4% formalin. Although a quasi-synoptic sampling was intended for the late spring to early summer season in 2000, data e.g. for the northern parts and the British Channel were only available for the 2001 (see van den Berghe 2007).

For all analyses only NSB data for the nearest matching stations in 1986 and 2000 were used (Fig. 1). Thus, after excluding stations with a distance apart of more than 40 km, 156 matching stations were considered in the analyses. Instead of using the published results of the 1986 data (Künitzer et al. 1992) they were re-analysed since both datasets (1986 and 2000) had to be taxonomically adjusted to allow comparisons.




Fig. 1: Location of matching stations with distances apart superimposed (m).

We used the PRIMER v5 program package to perform cluster analysis and multidimensional scaling for abundance data to reveal similarities between stations (Clarke & Warwick 1994). Similarities were calculated using the Bray-Curtis coefficient. Fourth-root transformation was used prior to computation. Similarity percentage analysis (SIMPER) was used to identify species which were mainly responsible for differences in community structure. The significance of any differences in community structure at stations from 1986 and 2000 was tested with ANOSIM. The Hurlbert Index (Hurlbert 1971) was calculated as an expression of species diversity.

The comparison between the univariate parameters and community structure of 1986 and 2000 was carried out using the clusters of 1986 as a starting point. Thus, stations within each of the 1986 clusters were compared with the matching stations sampled in 2000. Additionally, the datasets of 1986 and 2000 were combined in one cluster analysis to detect differences in the cluster classification. If matching stations for both years belonged to one sub-cluster we assumed a consistent classification.

Results

Figures 2 a and b show that the spatial distribution of the macrofaunal communities in 2000 was broadly similar to that in 1986 described by Künitzer et al. (1992). The comparison of the similarity matrices revealed a significant relationship between 1986 and 2000 data (Table 1). The major divisions in the communities of the North Sea still occur at the 50 and 100 m depth contours. Also, greater heterogeneity of communities in the southern North Sea (<50 m) compared to the north is still evident.

Table 1. Correlation coefficients (Spearman rank) relating the similarity matrices of 1986 and 2000 infauna data for different transformation types (RELATE).

	R ²	p
Fourth root	0.533	0.001
Square root	0.527	0.001
Presence-absence	0.511	0.001
No transformation	0.421	0.001




Fig. 2: Spatial distribution of infaunal communities in 1986 (left) and 2000 (right) based on fourth-root transformed abundance data




Fig. 3: Cluster dendograms and groupings shown in Fig. 2 revealed with fourth-root transformed abundance data.




Fig. 4: Distribution of differences in the cluster classification between 1986 and 2000 based on fourth-root transformed abundance data. Small-sized circles represent stations which were classified identically in 1986 and 2000. The largest circles represent stations classified in different main clusters.

However, on closer inspection, there are noticeable differences between the community structure in 1986 and 2000 in some areas of the North Sea. An evaluation of differences in the composition of stations within similar clusters in 1986 and 2000 shows that these were greatest in the eastern North Sea, along the Frisian Front, towards the English Channel, and especially at depths of >50 m off the northern British coast and at >100 m in the northern North Sea (Fig. 4). These have been caused by differences in abundance and species numbers.

Deeper than 100 m, differences in community structure occurred mainly due to a decrease in abundance but an increase in species numbers (Fig. 5). In 2000, the community structure was similar to that in the central North Sea at 50-100 m depth (Fig. 1). But at some of the stations the small polychaet *Paramphinome jeffreysii* occurred in significant densities.

The differences in community structures off the northern British coast (>50 m) seem to be caused by a general decrease in species numbers in parallel with an increase in abundance in 2000 (Fig. 5). In 1986, this area was split into several clusters (mainly B, C, E4), while in 2000 the area was separated mainly into the two clusters F1 and F2 (Fig. 2). The SIMPER analyses show that the differences in community structure have been caused by an increase in the small polychaete *P. jeffreysii*, the interface-feeding polychaetes *Myriochele* spp. and *Spiophanes bombyx* (Fig. 6, Annex 1).

In the eastern North Sea, differences in 2000 occurred due to an increase in abundance of Phoronida and *S. bombyx* and of the bivalves *Fabulina fabula* and *Corbula gibba*, the amphipod *Urothoe poseidonis* and the brittle star *Acrocnida brachiata*.

In the SW North Sea towards the English Channel, both a decrease and increase in species numbers as well as an increase in abundance occurred between 1986 and 2000. At the offshore stations, differences in communities occurred due to an increase in the polychaetes *S. bombyx* and *Magelona* spp., but a decrease in e.g. *Ophelia borealis*. At the coastal stations, the abundance of the polychaetes *Lanice conchilega*, *S. bombyx* and *Lagis koreni* as well as the bivalve *Spisula* spp. increased significantly.




Fig. 5: Differences in mean abundance (left) and mean ES(100) (right) between 1986 and 2000. Green dots indicate an increase and red dots a decrease in 2000 compared to 1986.

At the western part of the Frisian Front differences in community structure could be explained by a general decrease in abundance, e.g., in the polychaete *O. borealis*, while at the eastern part and at the East Frisian coast total abundances increased due to increases in *S. bombyx* and *Magelona* spp. (Fig. 6).

The community structure in the central Oyster Ground remained rather stable over time, but a decrease in total abundance was found at some stations. Also the Dogger Bank community remained rather stable, even if the abundance of the polychaetes *O. borealis* and *Nephtys cirrosa* and the bivalve *Abra prismatica* decreased. At the Tail End abundances of *L. conchilega* and *S. bombyx* increased, while at the South West Patch *Magelona* spp. and *S. bombyx* increased but *Bathyporeia* spp. decreased.

Table 2 reveals that all of the changes in community structure evident from Figure 6 are significant ($p < 5$). An additional contribution to the apparent decline in species number at many stations in the southern North Sea (Fig. 5) may be the use of deeper-penetrating box core samples in 1986. Stations in 2000 were sampled mainly by 0.1m² van Veen grabs.




Fig. 6: MDS-plots revealed with fourth-root transformed abundance data of 1986 (\triangle) and 2000 (\blacktriangledown) for the stations of each cluster (see Fig. 3).




Fig. 6 (continued).

Detailed maps comparing species distribution in 1986 and 2000 are given by Eggleton et al. (2007). van Hoey et al. (2007) and Willems et al. (2007) present detailed correlations of faunal and environmental parameters.

Table 2: Differences between 1986 and 2000 data revealed with ANOSIM for the MDS-plots shown in Figure 6.

Cluster	R	p (%)
A	0.511	0.1
B	0.426	1.6
C	0.648	0.2
D11	0.320	0.2
D12	0.241	0.1
D13	0.190	3.8
D2	0.181	0.1
E1	0.837	0.2
E2	0.588	0.8
E3	0.372	0.1
E4	0.377	0.1
F1	0.202	4.2
F2	0.131	0.1
G	0.404	3.2

Discussion

In general, the spatial distribution of the macrofaunal communities in 2000 was rather similar to that in 1986 described by Künitzer et al. (1992). The major divisions in the structure of the communities of the North Sea still occur at the 50 and 100 m depth contours, as also described by Glemarec (1973).

The decrease in total abundance found in the northern North Sea (>100 m) was caused by the use of different mesh sizes in 1986 (0.5 mm) and 2000 (1 mm) as well as by the spatial resolution of the station grid (Fig. 1). The increase in species number in this region in 2000 might be due to improved taxonomic precision.

Thus “real” changes in community structure between 1986 and 2000 can be more confidently discussed for the southern North Sea. Changes in community structure north of the 50 m depth contour may be related to changes in the hydroclimate caused by changes in the North Atlantic Oscillation which, in positive mode, results in an increase in SST, changes in sediment structure and food availability as described by Reid & Edwards (2001) and Kröncke et al. (1998). Wiking & Kröncke (2001) described the NAO-influenced changes in hydrography especially north and south of the Dogger Bank. The increase in

inflow of Atlantic water masses through the Fair Isle channel strengthened the frontal system north of the bank, which follows the 50 m depth contour, creating a “strong” border between northern and southern water masses. High current velocities in the northern part of the Dogger Bank (Siegismund & Schrum 2001) in addition to a seasonal jet (Brown et al. 1999) limit the accumulation of particulate organic material in seabed sediments. Klein et al. (1999) showed that during storms fine sediment at the sea bed are mobilised at up to 60 m depth at the northern slope of the Dogger Bank. Thus, the decrease in total species number and the increase in species such as the small polychaete *Paramphipnoma jeffreysii*, as well as the dominance of the interface-feeding polychaetes *Myriochele* spp. and *Spiophanes bombyx* north of the 50 m depth contour, provide supporting evidence for a change in hydrodynamics affecting sediment structure and stability. Since *P. jeffreysii* and *Myriochele* spp. are considered to be cold-temperate species, their increase in abundance north of the 50 m depth contour might be a hint of colder northern water masses north of the frontal system.

Changes observed in the communities at the offshore stations in the SW North Sea towards the English Channel and the eastern part of the Frisian Front indicate similar environmental influences. The decrease in the polychaete *Ophelia borealis* in these regions (see Eggleton et al. 2007) might be due to an increase in SST, since *O. borealis* is a cold-temperate species, as well as to changes in the sediment composition as found by Wieking & Kröncke (2001) at the Dogger Bank

At the coastal stations in the SW North Sea towards the English Channel and at the western part of the Frisian Front an increase in interface-feeding polychaetes such as *S. bombyx*, *Magelona* spp. and *Lanice conchilega* as well as the bivalve *Spisula* spp. might be caused by a higher food availability at the Flamborough and Frisian Fronts due to an NAO-induced increase in SST and hydrodynamic forces. Due to frontal conditions and enhanced primary production, food supply to the benthos (quality and quantity) will be higher than in non-frontal areas at comparable depths (Duineveld & Boon 2002).

In the eastern North Sea, the increase in 2000 in phoronids and other interface-feeding species such as the polychaete *S. bombyx* and the bivalves *Fabulina fabula* and *Corbula gibba* also indicate an NAO-induced increase in food

availability due to higher primary production in the German Bight as described by Reid et al. (1998) and Reid & Edwards (2001). Current-induced changes in the sediment structure might have caused the increase in the brittle star *Amphiura brachiata* and the sea urchin *Echinocyamus pusillus*, which prefers coarser sediments, but *A. brachiata* is also a warm-temperate species (Wiekings & Kröncke 2001). Further investigations of links between benthic community structure and environmental variables, including climatic influences, can be found by van Hoey & Rees (2007) and Willems et al. (2007).

References

- Basford DJ, Eleftheriou A (1988) The benthic environment of the North Sea (56°+061 °N). J mar boil Ass UK 68: 125-141.
- Basford DJ, Eleftheriou A, Raffaelli D (1990) The infauna and epifauna of the northern North Sea. Neth J Sea Res 25 (1/2): 165-173
- Basford DJ, Eleftheriou A, Davies M, Irion G, Soltwedel T (1993) The ICES North Sea benthos survey: the sedimentary environment. ICES J mar Sci 50: 71-80.
- Brown J, Hill AE, Fernand L, Horsburgh KJ (1999) Observations of a seasonal jet-like circulation at the central North Sea cold pool margin. Estuarine, Coastal and Shelf Science 48: 343-355
- Clarke KR, Warwick RM (1994) Change in marine communities: an approach to statistical analysis and interpretation. Plymouth Marine Laboratory: 130 pp.
- Craeymeersch JA, Heip CHR, Buijs J (1997) Atlas of the North Sea benthic infauna. ICES cooperative research report 218: 1-86
- Duineveld G, Boon A (2002) Short-term variations in fluxes and composition of seston in near bottom traps of the southern North Sea. Helgoland Mar Res 150: 146-148
- Eggleton J, Smith R, Rees HL (2007) Species distributions and changes (1986-2000). ICES Cooperative Research Report No. 288: 277 pp.
- Eleftheriou A, Basford DJ (1989) The macrobenthic infauna of the offshore northern North Sea. J Mar Biol Ass UK 69: 123-143
- Glémarec M (1973) The benthic communities of the European North Atlantic continental shelf. Oceanography and Marine Biology: an Annual Review 11: 263-289.
- Heip C, Craeymeersch JA (1995) Benthic community structures in the North Sea. Helgoländer Meeresunters 49:313-328
- Hurlbert, S.H. (1971) The non-concept of species diversity: A critique and alternative parameters. Ecology, 52: 577-586
- ICES (2007) Structure and dynamics of the North Sea benthos. ICES Cooperative Research Report No. 288: 277 pp.
- Jones SE, Jago CF, Bale AJ, Chapman D, Howland RJM, Jackson J (1998) Aggregation and resuspension of suspended particulate matter at a seasonally stratified site in the southern North Sea: physical and biological controls. Continental Shelf Research 18: 1283-1309
- Klein H, König P, Frohse A (1999) Currents and Near-Bottom Suspended Matter Dynamics in the Central North Sea during stormy Weather – Results of the PIPE'98 field experiment. Deutsche Hydrografische Zeitschrift 51:1
- Kröncke I, Dippner JW, Heyen H, Zeiss B (1998) Long-term changes in macrofaunal communities off Norderney (East Frisia, Germany) in relation to climate variability. Marine Ecology Progress Series 167: 25-36

- Kröncke I, Zeiss B, Rensing C (2001) Long-term variability in macrofauna species composition off the island of Norderney (East Frisia, Germany) in relation to changes in climatic and environmental condition. *Senckenbergiana maritima* 31: 65-82
- Künitzer A, Basford D, Craeymeersch JA, Dewarumez JM, Dörjes J, Duineveld GCA, Eleftheriou A, Heip C, Herman P, Kingston P, Niermann U, Rachor E, Rumohr H, de Wilde PAJ (1992) The benthic infauna of the North Sea: species distribution and assemblages. *ICES Journal of Marine Science* 49: 127-143
- Otto L, Zimmerman JTF, Furnes GK, Mork M, Saetre R, Becker G (1990) Review of the physical oceanography of the North Sea. *Netherlands Journal of Sea Research* 26: 161-238
- Reid CP, Edwards M, Hunt HG, Warner J (1998) Phytoplankton change in the North Atlantic. *Nature* 391:546
- Reid PC, Edwards M (2001) Long-term changes in the fishery, pelagos and benthos of the North Sea. *Senckenbergiana maritima* 31: 107-115
- Siegismund F, Schrum C (2001) Decadal changes in the wind forcing over the North Sea. *Climate Research* 18:39-45
- van den Berghe, E. (2007) NSBP 2000 data management. ICES Cooperative Research Report No. 288: 277 pp.
- van Hoey G, Rees HL (2007) A comparison of indicators reflecting the status of the North Sea benthos. ICES Cooperative Research Report No. 288: 277 pp.
- Wieking G, Kröncke I (2001) Decadal changes in macrofauna communities on the Dogger Bank caused by large-scale climate variability. *Senckenbergiana maritima* 31: 125-141
- Willems W, Rees, HL, Vincx, M, Goethals P, Degraer S. (2007) Relations and interactions between environmental factors and biotic properties. ICES Cooperative Research Report No. 288: 277 pp.

Table Annex : Results of the SIMPER analysis based on fourth root transformed abundance data (Fig. 3, Chapter 5.2). The 1986 clusters were compared with the corresponding stations in 2000.

Cluster A

Average dissimilarity = 72,50

Species	Group 1986	Group 2000	Av. Abund	Av. Abund	Av. Diss	Diss/SD	Contrib%	Cum. %
Thyasira	346,29	17,53	1,18	1,59	1,62	1,62		
Lumbrineris latreilli	25,69	0,00	1,02	2,50	1,40	3,03		
Myriochele	165,65	109,06	0,98	1,27	1,35	4,38		
Paramphinome jeffreysii	29,88	97,03	0,95	1,45	1,31	5,69		
Capitella	124,88	2,33	0,92	0,97	1,27	6,96		
Pholoe baltica	74,35	4,56	0,90	1,40	1,25	8,21		
Exogone	74,86	2,47	0,84	1,26	1,16	9,37		
Aricidea catherinae	66,43	4,92	0,84	1,38	1,16	10,53		
Cirrophorus lyra	67,41	1,42	0,83	1,22	1,14	11,67		
Amphiura chiajei	2,50	55,75	0,83	1,32	1,14	12,81		
Heteromastus filiformis	0,00	45,00	0,78	1,19	1,08	13,89		
Pseudopolydora	39,85	2,11	0,76	1,27	1,05	14,94		
Prionospio	50,95	15,00	0,76	1,22	1,05	15,99		
Caulieriella	50,04	1,86	0,75	1,18	1,04	17,03		
Eriopisa elongata	42,33	15,50	0,75	1,32	1,03	18,06		
Mendicula ferruginosa	27,63	0,92	0,75	1,29	1,03	19,09		
Leucon nasica	44,15	13,50	0,73	1,09	1,01	20,10		
Diplocirrus glaucus	18,24	26,44	0,73	1,39	1,00	21,10		
Levinsenia gracilis	99,36	37,28	0,71	1,34	0,98	22,08		
Amphiura filiformis	30,06	30,17	0,69	1,31	0,95	23,03		
Lumbrineris gracilis	0,00	7,69	0,68	1,66	0,94	23,96		
Eudorella emarginata	21,05	16,92	0,68	1,20	0,93	24,90		
Pectinaria	18,33	17,50	0,67	1,44	0,92	25,82		
Glycera lapidum	57,56	2,78	0,66	1,05	0,91	26,73		
Urothoe elegans	14,04	8,39	0,62	1,40	0,85	27,58		
Spiophanes	91,23	62,69	0,61	1,08	0,84	28,43		
Laonice sarsi	33,36	5,44	0,61	1,27	0,84	29,27		
Notomastus	30,23	8,50	0,61	1,16	0,84	30,10		
Owenia fusiformis	27,60	5,53	0,61	1,19	0,84	30,94		
Goniada	19,68	9,03	0,60	1,24	0,83	31,77		
Eclysippe vanelli	20,81	1,42	0,58	1,20	0,80	32,57		
Nothria conchylega	28,25	0,75	0,58	1,05	0,80	33,37		
Aplacophora	0,00	8,56	0,57	1,26	0,79	34,16		
Eudorella truncatula	6,67	6,06	0,57	1,26	0,79	34,95		

Cluster B

Average dissimilarity = 66,44

Species	Group 1986	Group 2000	Av. Abund	Av. Abund	Av. Diss	Diss/SD	Contrib%	Cum. %
Capitella	41,56	0,00	1,20	2,25	1,81	1,81		
Ampharete lindstroemi	0,00	81,13	1,08	1,59	1,63	3,44		
Praxillella	6,02	0,00	0,78	2,94	1,18	4,62		
Paramphinome jeffreysii	0,66	25,10	0,77	1,47	1,16	5,78		
Myriochele	178,86	261,83	0,76	1,60	1,15	6,93		
Ampelisca spinipes	43,38	0,67	0,74	1,11	1,12	8,05		
Anobothrus gracilis	8,64	2,00	0,72	2,05	1,08	9,13		
Glycera rouxi	10,34	1,00	0,70	1,48	1,06	10,19		
Thyasira	35,62	16,27	0,70	1,26	1,06	11,25		
Levinsenia gracilis	38,88	16,03	0,69	1,25	1,04	12,29		
Photis	6,72	0,00	0,63	1,60	0,95	13,24		
Diplocirrus glaucus	7,94	19,63	0,63	1,41	0,94	14,18		
Cirratulus	0,34	5,83	0,61	1,49	0,92	15,11		
Heteromastus filiformis	0,00	17,40	0,61	0,98	0,92	16,03		
Nephthys hombergii	12,40	5,33	0,61	1,58	0,92	16,94		
Owenia fusiformis	5,36	26,53	0,60	2,03	0,91	17,85		
Philomedes globosus	26,44	0,00	0,59	0,87	0,89	18,74		
Echinocyamus pusillus	12,66	9,40	0,59	0,99	0,88	19,63		
Malmeagrenia arenicolae	2,68	0,00	0,57	1,80	0,86	20,48		
Harpinia antennaria	6,66	23,13	0,57	1,53	0,85	21,34		
Dipolydora flava	3,04	0,67	0,56	1,94	0,85	22,19		
Urothoe elegans	4,00	10,50	0,56	1,14	0,84	23,03		
Leucon nasica	1,66	17,47	0,56	1,07	0,84	23,87		
Nuculoma tenuis	3,68	10,77	0,54	1,38	0,81	24,68		
Scoloplos	7,70	1,33	0,53	1,15	0,80	25,48		
Terebellides stroemi	8,38	3,83	0,52	1,04	0,78	26,26		
Echinocardium flavescens	2,00	0,00	0,52	1,91	0,78	27,05		
Diastylis lucifera	0,66	16,50	0,52	0,88	0,78	27,83		
Cirrophorus lyra	11,00	7,47	0,52	0,86	0,78	28,61		
Prionospio	3,34	7,83	0,52	1,24	0,78	29,39		
Ampelisca macrocephala	1,00	6,13	0,52	1,41	0,78	30,16		
Notomastus	2,68	11,30	0,51	1,37	0,77	30,94		
Amphiura chiajei	0,00	15,50	0,51	0,79	0,77	31,71		

<i>Lucinoma borealis</i>	6,42	2,00	0,50	1,32	0,76	32,47
<i>Pectinaria</i>	5,70	26,67	0,50	1,54	0,76	33,22
<i>Scoletoma fragilis</i>	3,66	0,00	0,50	1,14	0,76	33,98
<i>Pholoe baltica</i>	8,34	4,13	0,48	1,29	0,73	34,71

Cluster C

Average dissimilarity = 65,78

Species	Group 1986	Group 2000	Av.Diss	Diss/SD	Contrib%	Cum.%
	Av.Abund	Av.Abund				
<i>Spiophanes</i>	37,30	348,06	1,55	1,22	2,36	2,36
<i>Phyllodocida groenlandica</i>	0,00	9,72	1,41	3,89	2,14	4,50
<i>Paramphithome jeffreysii</i>	3,95	23,89	1,30	1,38	1,97	6,47
<i>Myriochele</i>	13,32	60,56	1,22	1,10	1,86	8,33
<i>Ampharete lindstroemi</i>	0,00	7,78	1,18	1,97	1,79	10,12
<i>Mysella</i>	27,82	18,06	1,16	1,43	1,77	11,89
<i>Nucula nitidosa</i>	0,00	15,28	1,14	1,88	1,74	13,62
<i>Nemertina</i>	5,27	16,67	1,14	1,57	1,73	15,35
<i>Chaetozone</i>	0,00	6,67	1,08	2,01	1,64	16,99
<i>Pholoe baltica</i>	7,90	6,39	1,03	1,55	1,56	18,55
<i>Cirratulus</i>	0,00	4,17	1,01	1,96	1,54	20,09
<i>Harpinia antennaria</i>	5,27	21,94	0,96	1,19	1,47	21,56
<i>Abra prismatica</i>	2,63	8,06	0,92	1,21	1,40	22,95
<i>Bathyporeia</i>	14,72	2,78	0,90	1,16	1,36	24,31
<i>Eudorellopsis deformis</i>	39,83	16,39	0,90	1,15	1,36	25,68
<i>Polinices</i>	0,00	4,44	0,88	1,30	1,34	27,01
<i>Hippomedon</i>	0,00	4,17	0,85	1,30	1,30	28,31
<i>Goniada</i>	14,67	15,56	0,84	1,03	1,28	29,58
<i>Nuculoma tenuis</i>	3,95	15,00	0,83	1,09	1,26	30,85
<i>Ampelisca tenuicornis</i>	3,95	4,44	0,82	1,25	1,25	32,10
<i>Levinsenia gracilis</i>	9,32	0,56	0,82	1,05	1,25	33,35
<i>Nephtys hombergii</i>	15,98	6,94	0,82	1,06	1,25	34,60

Cluster D11

Average dissimilarity = 63,22

Species	Group 1986	Group 2000	Av.Diss	Diss/SD	Contrib%	Cum.%
	Av.Abund	Av.Abund				
<i>Phoronida</i>	141,62	1053,33	2,47	1,09	3,91	3,91
<i>Spiophanes</i>	15,12	164,74	1,78	1,58	2,81	6,72
<i>Chamelea gallina</i>	92,47	10,56	1,64	1,19	2,60	9,32
<i>Hexacorallia</i>	57,51	6,11	1,56	1,20	2,46	11,78
<i>Scoloplos</i>	50,48	10,93	1,45	1,16	2,29	14,08
<i>Bathyporeia</i>	45,49	62,96	1,43	1,16	2,27	16,34
<i>Pholoe baltica</i>	19,12	5,00	1,40	1,50	2,21	18,56
<i>Thracia</i>	1,89	29,67	1,40	1,34	2,21	20,77
<i>Tellina</i>	24,37	82,26	1,39	1,23	2,20	22,97
<i>Glycinde nordmanni</i>	15,10	0,56	1,36	1,44	2,15	25,13
<i>Magelona</i>	161,19	221,70	1,29	0,90	2,04	27,16
<i>Goniada</i>	1,64	26,15	1,24	1,43	1,97	29,13
<i>Urothoe poseidonis</i>	5,89	59,78	1,22	0,95	1,94	31,06
<i>Owenia fusiformis</i>	7,10	18,89	1,20	1,32	1,89	32,96
<i>Amphiura filiformis</i>	18,22	3,89	1,14	1,00	1,80	34,75

Cluster D12

Average dissimilarity = 57,08

Species	Group 1986	Group 2000	Av.Diss	Diss/SD	Contrib%	Cum.%
	Av.Abund	Av.Abund				
<i>Lanice conchilega</i>	4,66	261,58	1,14	1,23	1,99	1,99
<i>Bathyporeia</i>	245,40	229,68	0,99	1,40	1,73	3,72
<i>Amphiura filiformis</i>	70,56	12,34	0,96	1,53	1,68	5,40
<i>Mysella</i>	101,64	24,97	0,95	1,20	1,66	7,07
<i>Nephtys hombergii</i>	32,94	3,16	0,94	1,59	1,65	8,72
<i>Ophiura albida</i>	53,31	4,75	0,91	1,42	1,59	10,31
<i>Spiophanes</i>	172,64	593,01	0,90	1,52	1,58	11,89
<i>Urothoe poseidonis</i>	32,27	52,60	0,89	1,36	1,56	13,45
<i>Scoloplos</i>	31,62	28,33	0,84	1,27	1,47	14,91
<i>Ophelia borealis</i>	28,16	4,01	0,81	1,48	1,43	16,34
<i>Nephtys cirrosa</i>	38,85	6,30	0,80	1,19	1,40	17,74
<i>Abra prismatica</i>	21,04	0,67	0,80	1,60	1,39	19,14
<i>Echinocardium cordatum</i>	15,75	14,36	0,79	1,42	1,39	20,53
<i>Owenia fusiformis</i>	7,86	11,33	0,75	1,72	1,31	21,84
<i>Montacuta</i>	21,39	10,15	0,72	1,21	1,26	23,10
<i>Nephtys longisetosa</i>	10,61	0,89	0,69	1,37	1,21	24,31
<i>Acrocnida brachiatia</i>	13,51	23,65	0,69	1,32	1,21	25,53
<i>Chamelea gallina</i>	9,46	3,39	0,66	1,38	1,15	26,67
<i>Arctica islandica</i>	6,63	0,22	0,65	1,47	1,15	27,82
<i>Pectinaria</i>	3,66	23,69	0,65	1,11	1,15	28,97
<i>Dosinia lupinus</i>	6,81	1,71	0,65	1,38	1,14	30,11

Sigalion	11,28	9,39	0,65	1,26	1,14	31,25
Nucula nitidosa	13,99	10,50	0,64	0,94	1,11	32,36
Spio	18,08	11,18	0,62	1,24	1,09	33,46
Abra alba	0,94	84,05	0,62	0,67	1,08	34,54

Cluster D13

Average dissimilarity = 58,82

Species	Group 1986		Group 2000		Contrib%	Cum.%
	Av. Abund	Av. Abund	Av. Diss	Diss/SD		
Echinocardium cordatum	35,27	11,00	2,08	1,60	3,54	3,54
Urothoe poseidonis	47,64	82,17	1,90	1,47	3,24	6,78
Tellina	160,27	116,25	1,82	1,18	3,10	9,88
Magelona	112,96	296,88	1,70	1,17	2,90	12,78
Montacuta	18,24	0,91	1,65	1,61	2,81	15,58
Nephtys hombergii	12,63	1,39	1,63	1,86	2,77	18,36
Nemertina	59,36	22,14	1,50	1,15	2,55	20,91
Nephtys cirrosa	27,73	17,88	1,47	1,13	2,50	23,41
Spiophanes	38,80	146,17	1,45	1,33	2,46	25,87
Bathyporeia	325,09	114,73	1,39	1,23	2,36	28,24
Scoloplos	17,90	6,88	1,32	1,26	2,24	30,48
Donax vittatus	32,40	2,38	1,24	0,98	2,11	32,59
Spio	3,07	11,50	1,12	1,11	1,91	34,50

Cluster D2

Average dissimilarity = 59,70

Species	Group 1986		Group 2000		Contrib%	Cum.%
	Av. Abund	Av. Abund	Av. Diss	Diss/SD		
Amphiura filiformis	539,31	410,82	1,16	1,23	1,94	1,94
Mysella	215,87	106,71	1,08	1,21	1,80	3,74
Corbula gibba	4,40	103,17	1,01	1,19	1,70	5,44
Magelona	67,28	55,58	0,94	1,34	1,57	7,01
Myriochele	150,42	135,03	0,93	0,89	1,55	8,56
Ophiura albida	44,05	7,23	0,93	1,26	1,55	10,11
Scoloplos	35,52	14,31	0,88	1,28	1,48	11,59
Phoronida	74,74	64,26	0,84	1,20	1,41	12,99
Nuculoma tenuis	23,86	7,34	0,82	1,33	1,38	14,37
Chamelea gallina	32,31	4,99	0,82	1,23	1,37	15,74
Callianassa subterranea	19,62	15,71	0,80	1,34	1,34	17,08
Spiophanes	39,27	73,46	0,79	1,24	1,32	18,39
Pectinaria	25,85	13,40	0,76	1,25	1,28	19,67
Nucula nitidosa	36,79	47,63	0,73	1,11	1,23	20,90
Pholoe baltica	64,45	49,01	0,73	1,20	1,22	22,12
Montacuta	9,58	8,42	0,71	1,39	1,19	23,31
Eudorellopsis deformis	21,81	10,47	0,70	0,98	1,17	24,47
Gattyana cirrhosa	8,53	14,49	0,69	1,18	1,16	25,64
Chaetopterus	10,67	8,29	0,68	1,23	1,15	26,78
Abra alba	2,20	30,47	0,68	1,03	1,14	27,92
Polinices	13,61	10,70	0,68	1,26	1,14	29,06
Bathyporeia	10,08	9,12	0,67	1,25	1,13	30,18
Harpinia antennaria	32,79	41,90	0,67	1,19	1,12	31,30
Owenia fusiformis	9,28	5,59	0,66	1,19	1,10	32,41
Arctica islandica	7,96	0,00	0,65	1,27	1,09	33,50
Echinocardium cordatum	11,52	12,21	0,64	1,23	1,08	34,58

Cluster E1

Average dissimilarity = 65,54

Species	Group 1986		Group 2000		Contrib%	Cum.%
	Av. Abund	Av. Abund	Av. Diss	Diss/SD		
Paramphinoe jeffreysii	8,37	127,67	0,98	2,06	1,50	1,50
Echinocardium flavescentes	113,73	0,33	0,82	1,46	1,25	2,74
Exogone	55,85	1,67	0,80	1,89	1,22	3,97
Eudorellopsis deformis	105,42	2,67	0,78	1,35	1,20	5,16
Amphiura chiajei	0,00	39,50	0,76	1,66	1,15	6,32
Laonice sarsi	5,03	21,50	0,69	2,34	1,06	7,38
Thyasira	41,50	51,17	0,64	1,27	0,98	8,35
Lucinoma borealis	0,00	10,17	0,61	1,83	0,94	9,29
Philine	36,65	8,67	0,61	1,85	0,93	10,22
Glycera lapidum	23,20	17,17	0,61	1,66	0,93	11,15
Urothoe elegans	11,80	19,67	0,59	1,34	0,91	12,06
Abra prismatica	35,07	1,33	0,59	1,23	0,91	12,96
Lanice conchilega	0,00	15,17	0,59	1,88	0,90	13,86
Pholoe baltica	40,18	7,33	0,59	1,59	0,89	14,76
Pholoe pallida	0,00	12,17	0,57	1,80	0,87	15,63
Nothria conchylega	13,33	3,33	0,57	1,62	0,87	16,50
Aonides paucibranchiata	46,67	14,00	0,57	1,83	0,87	17,37
Terebellides stroemi	0,00	9,33	0,56	1,84	0,85	18,22
Aplacophora	0,00	14,83	0,55	1,21	0,84	19,07

Poecilochaetus serpens	20,17	24,83	0,55	1,12	0,84	19,91
Trichobranchus roseus	0,00	6,00	0,55	2,00	0,84	20,75
Phyllodocida mucosa	13,30	0,00	0,55	1,32	0,84	21,58
Ophiocten affinis	23,20	0,67	0,54	1,35	0,83	22,41
Pectinaria	26,77	51,67	0,54	1,42	0,82	23,24
Scolelepis	8,37	12,33	0,54	2,01	0,82	24,06
Anobothrus gracilis	8,30	11,67	0,54	1,78	0,82	24,89
Prionospio	126,92	25,67	0,54	1,52	0,82	25,71
Eudorella truncatula	28,32	8,00	0,53	1,19	0,81	26,52
Hexacorallia	26,80	21,67	0,53	1,47	0,80	27,32
Acidostoma	0,00	5,17	0,53	1,97	0,80	28,12
Aricidea catherinae	21,73	6,50	0,52	1,92	0,79	28,91
Chaetozone	28,30	12,50	0,51	1,28	0,78	29,69
Montacuta	1,67	4,33	0,51	2,78	0,77	30,46
Nephtys hombergii	6,67	11,00	0,50	1,45	0,77	31,24
Notomastus	8,37	10,00	0,50	1,58	0,77	32,00
Phoronida	10,00	23,33	0,50	1,39	0,77	32,77
Ampelisca tenuicornis	0,00	9,00	0,50	1,32	0,77	33,54
Sthenelais	16,72	3,33	0,50	1,19	0,76	34,30

Cluster E2

Average dissimilarity = 62,04

Species	Group 1986	Group 2000	Av.Diss	Diss/SD	Contrib%	Cum.%
	Av.Abrund	Av.Abrund				
Amphiura filiformis	37,74	113,20	1,45	1,54	2,34	2,34
Myriochele	3,02	232,80	1,16	0,99	1,88	4,22
Ophelia borealis	34,44	0,60	1,12	2,68	1,80	6,02
Mysella	34,56	18,40	1,00	1,71	1,62	7,64
Paramphinome jeffreysii	8,00	24,60	0,94	1,47	1,52	9,16
Phoronida	3,00	20,40	0,93	1,56	1,50	10,66
Unciola planipes	10,04	0,00	0,93	1,74	1,49	12,15
Bathyporeia	87,74	8,00	0,92	1,71	1,49	13,64
Anobothrus gracilis	0,00	4,40	0,87	5,37	1,40	15,04
Sthenelais	1,00	8,20	0,85	2,25	1,36	16,41
Tridonta montagui	24,36	0,80	0,82	1,22	1,32	17,73
Pectinaria	42,68	7,00	0,81	1,99	1,31	19,04
Timoclea ovata	10,04	0,20	0,81	1,68	1,31	20,35
Cylichna cylindracea	0,00	3,20	0,80	3,89	1,29	21,64
Diplocirrus glaucus	0,00	4,00	0,74	1,84	1,19	22,82
Magelona	1,00	6,20	0,73	1,41	1,17	23,99
Phyllodocida groenlandica	0,00	3,60	0,72	1,85	1,17	25,16
Chaetozone	34,42	3,60	0,72	1,34	1,16	26,32
Scaphopoda	7,94	6,40	0,71	1,47	1,15	27,47
Nephtys caeca	5,68	7,60	0,71	1,72	1,14	28,61
Travisia forbesii	14,62	0,40	0,70	1,21	1,12	29,74
Molgula	27,84	0,00	0,69	0,80	1,12	30,85
Nemertina	10,56	15,40	0,68	1,37	1,10	31,95
Hexacorallia	10,44	13,60	0,68	1,24	1,10	33,05
Philine	10,60	1,20	0,67	1,49	1,08	34,13

Cluster E3

Average dissimilarity = 68,78

Species	Group 1986	Group 2000	Av.Diss	Diss/SD	Contrib%	Cum.%
	Av.Abrund	Av.Abrund				
Ophelia borealis	54,80	4,18	1,37	2,08	1,99	1,99
Amphiura filiformis	48,75	65,91	0,93	1,31	1,35	3,34
Spiophanes	34,06	357,00	0,90	1,12	1,31	4,64
Echinocardium flavescens	5,63	0,00	0,85	3,21	1,24	5,88
Echinocyamus pusillus	40,98	17,48	0,84	1,49	1,22	7,10
Cochlodesma praetenuie	8,55	2,61	0,78	1,72	1,13	8,24
Exogone	29,87	0,91	0,78	1,21	1,13	9,37
Myriochele	7,20	116,36	0,77	1,15	1,11	10,48
Mysella	6,94	35,34	0,76	1,25	1,10	11,58
Ophiura albida	6,20	1,73	0,72	2,00	1,04	12,63
Owenia fusiformis	5,53	29,58	0,70	1,20	1,02	13,64
Bathyporeia	26,40	16,52	0,70	1,30	1,01	14,65
Nemertina	18,72	19,39	0,69	1,22	1,00	15,65
Amphiura chiajei	0,91	29,45	0,68	0,89	0,98	16,64
Hexacorallia	8,66	2,79	0,67	1,34	0,98	17,61
Phoronida	6,92	19,59	0,67	1,14	0,97	18,59
Tellina	17,63	2,06	0,66	1,14	0,96	19,54
Pholoe baltica	14,13	24,48	0,66	1,21	0,95	20,50
Notomastus	1,76	20,24	0,64	1,30	0,93	21,42
Thracia	14,84	4,36	0,61	0,95	0,89	22,31
Harpinia antennaria	1,82	15,70	0,60	1,02	0,87	23,19
Glycera lapidum	5,74	18,76	0,60	1,22	0,87	24,06
Nephtys caeca	2,34	7,88	0,58	1,18	0,85	24,91
Nephtys cirrosa	5,84	1,15	0,58	1,11	0,84	25,75
Chamelea gallina	0,45	13,43	0,58	1,09	0,84	26,59
Eudorellopsis deformis	4,90	5,09	0,57	1,08	0,83	27,42

Dosinia exoleta	4,96	2,00	0,56	1,19	0,82	28,24
Nephtys longosetosa	5,17	1,91	0,56	1,18	0,81	29,05
Polycirrus	2,12	7,94	0,55	1,00	0,80	29,85
Abra prismatica	6,15	5,15	0,55	1,16	0,80	30,65
Magelona	10,91	9,28	0,55	1,08	0,79	31,44
Nephtys hombergii	7,50	8,55	0,53	1,14	0,78	32,22
Aonides paucibranchiata	3,97	3,88	0,53	1,27	0,76	32,98
Lumbrineris gracilis	0,00	36,21	0,52	0,85	0,75	33,73
Sthenelais	2,34	4,04	0,51	1,11	0,75	34,48

Cluster E4

Average dissimilarity = 61,98

Species	Group 1986	Group 2000	Av. Abund	Av. Abund	Av. Diss	Diss/SD	Contrib%	Cum.%
Paramphipnoma jeffreysii			1,97	61,28	1,18	1,50	1,90	1,90
Myriochele			132,73	357,13	1,03	1,21	1,67	3,57
Nephtys longosetosa			14,08	1,10	0,87	1,64	1,40	4,97
Spiophanes			46,10	242,87	0,86	1,07	1,39	6,35
Mysella			7,91	16,21	0,85	2,00	1,36	7,72
Echinocardium flavescentes			4,65	1,92	0,79	2,28	1,28	9,00
Abra prismatica			8,06	6,67	0,78	1,70	1,25	10,25
Levinsenia gracilis			17,30	19,97	0,76	1,23	1,23	11,48
Polinices			4,02	5,00	0,74	1,83	1,20	12,67
Eudorellopsis deformis			16,10	16,82	0,74	1,21	1,19	13,86
Nephtys hombergii			8,55	1,33	0,72	1,61	1,16	15,02
Chaetozone			4,31	19,41	0,69	1,26	1,11	16,13
Pectinaria			9,18	13,44	0,67	1,17	1,09	17,21
Cirratulus			9,27	3,28	0,65	1,24	1,04	18,25
Hemilamprops rosea			4,13	2,46	0,65	1,66	1,04	19,29
Aplacophora			5,50	6,62	0,63	1,24	1,02	20,32
Montacuta			10,37	3,36	0,63	1,16	1,02	21,34
Ophiocanthes affinis			6,64	1,03	0,63	1,27	1,02	22,35
Hexacorallia			6,81	24,82	0,62	1,18	0,99	23,34
Prionospio			7,00	6,62	0,61	1,23	0,99	24,34
Phaxas pellucidus			5,80	5,82	0,61	1,22	0,99	25,32
Pholoe baltica			6,30	7,28	0,61	1,17	0,98	26,30
Ampelisca tenuicornis			9,39	5,59	0,61	1,15	0,98	27,28
Magelona			5,62	8,28	0,60	1,05	0,98	28,25
Owenia fusiformis			5,37	2,82	0,59	1,12	0,95	29,20
Westwoodilla caecula			4,45	0,87	0,58	1,26	0,94	30,14
Harpinia antennaria			2,66	6,97	0,57	1,19	0,91	31,05
Diplocirrus glaucus			3,43	5,64	0,55	1,15	0,89	31,94
Synchelidium			0,13	2,85	0,54	1,19	0,88	32,82
Trichobranchus roseus			0,88	5,33	0,54	1,12	0,87	33,69
Hippomedon			2,94	0,67	0,54	1,16	0,87	34,56

Cluster F1

Average dissimilarity = 78,73

Species	Group 1986	Group 2000	Av. Abund	Av. Abund	Av. Diss	Diss/SD	Contrib%	Cum.%
Pholoe baltica			53,42	16,67	1,29	1,23	1,64	1,64
Lanice conchilega			30,53	531,87	1,28	0,95	1,63	3,27
Capitella			31,17	0,00	1,26	1,82	1,60	4,87
Lumbrineris latreilli			26,71	0,00	1,23	1,18	1,56	6,42
Urothoe poseidonis			114,76	118,22	1,20	0,90	1,53	7,95
Mysella			111,51	39,24	1,16	0,99	1,48	9,43
Nemertina			54,67	27,96	1,14	1,07	1,45	10,88
Mya			22,24	7,14	1,11	1,37	1,41	12,29
Nicomache			122,94	1,90	1,06	1,05	1,35	13,64
Scoloplos			40,10	10,30	1,06	1,17	1,35	14,99
Spisula			1,12	739,56	1,05	0,56	1,34	16,33
Spiophanes			13,11	127,81	1,04	1,07	1,33	17,65
Pectinaria			10,09	226,12	1,04	0,89	1,33	18,98
Hexacorallia			3,46	73,81	0,99	0,95	1,26	20,23
Ophiura albida			19,77	1,90	0,98	1,53	1,24	21,48
Polycirrus			21,09	12,38	0,97	1,14	1,23	22,71
Ampelisca spinipes			18,39	1,90	0,94	1,03	1,20	23,90
Pomatoceros			10,36	77,62	0,93	1,10	1,18	25,09
Chaetozone			9,61	0,91	0,89	1,39	1,13	26,21
Scalibregma			6,76	18,10	0,88	1,16	1,11	27,33
Glycera lapidum			15,36	10,95	0,85	1,35	1,08	28,41
Urothoe brevicornis			25,36	20,16	0,82	0,93	1,04	29,45
Eusyllis blomstrandii			22,68	0,00	0,80	0,98	1,02	30,47
Eumida sanguinea			10,65	9,35	0,79	1,25	1,01	31,48
Malmgrenia arenicolae			10,52	0,00	0,78	0,91	0,99	32,46
Spio			13,95	6,63	0,77	1,10	0,98	33,44
Nephtys cirrosa			7,10	8,73	0,74	1,03	0,94	34,38

Cluster F2

Average dissimilarity = 72,82

Species	Group 1986	Group 2000	Av.Diss	Diss/SD	Contrib%	Cum.%
	Av.Abund	Av.Abund				
Spiophanes	22,55	395,85	2,08	1,21	2,85	2,85
Ophelia borealis	44,78	4,28	2,07	1,24	2,84	5,69
Magelona	72,62	118,65	2,03	1,23	2,79	8,48
Scoloplos	46,41	54,78	1,98	1,29	2,72	11,20
Urothoe poseidonis	15,72	57,60	1,73	1,04	2,38	13,58
Tellina	22,01	54,40	1,67	1,05	2,29	15,87
Bathyporeia	47,77	38,10	1,62	1,22	2,23	18,10
Spio	18,85	32,54	1,56	1,12	2,14	20,24
Lanice conchilega	9,53	48,74	1,50	0,99	2,06	22,30
Nemertina	21,02	19,76	1,45	1,17	2,00	24,30
Urothoe brevicornis	24,78	4,72	1,37	0,94	1,88	26,18
Spisula	62,82	3,92	1,35	0,96	1,85	28,03
Nephtys cirrosa	21,94	19,71	1,31	1,11	1,80	29,83
Echinocardium cordatum	11,45	4,93	1,26	1,04	1,73	31,56
Gastrosaccus spinifer	3,15	27,68	1,26	0,76	1,72	33,28
Scolelepis	5,31	6,89	1,23	1,04	1,69	34,97

Cluster G

Average dissimilarity = 80,10

Species	Group 1986	Group 2000	Av.Diss	Diss/SD	Contrib%	Cum.%
	Av.Abund	Av.Abund				
Spiophanes	14,26	53,30	1,18	1,43	1,47	1,47
Paramphinome jeffreysi	0,00	246,90	1,11	1,17	1,38	2,85
Aonides paucibranchiata	63,84	19,20	1,03	1,20	1,29	4,14
Magelona	0,00	86,67	1,03	0,65	1,28	5,42
Exogone	90,68	23,60	1,00	1,21	1,25	6,67
Glycera lapidum	76,78	24,90	0,99	1,16	1,24	7,91
Phoronida	0,00	22,90	0,97	0,94	1,21	9,12
Amphiura filiformis	3,42	64,47	0,93	1,12	1,16	10,28
Aricidea cerrutii	44,20	10,00	0,89	1,17	1,11	11,39
Myriochele	1,00	22,10	0,89	1,49	1,11	12,50
Unciola planipes	20,22	7,60	0,86	1,08	1,07	13,57
Harpinia antennaria	0,00	53,60	0,83	0,78	1,03	14,60
Hexacorallia	25,12	14,50	0,82	1,13	1,02	15,63
Scoloplos	14,68	12,33	0,82	1,32	1,02	16,64
Nephtys longosetosa	10,22	0,00	0,80	1,11	0,99	17,64
Scolelepis	12,08	18,10	0,77	1,17	0,97	18,61
Spio	2,00	23,30	0,77	1,14	0,96	19,56
Owenia fusiformis	18,88	14,60	0,76	1,16	0,95	20,51
Ophelia borealis	13,76	1,60	0,76	1,13	0,95	21,46
Corbula gibba	2,00	36,67	0,75	0,63	0,94	22,40
Echinocyamus pusillus	10,96	12,40	0,75	1,18	0,94	23,33
Bathyporeia	11,60	9,87	0,74	0,92	0,92	24,26
Goniada	41,22	21,03	0,74	1,29	0,92	25,18
Dipolydora socialis	11,96	0,00	0,73	1,13	0,92	26,10
Polinices	0,00	7,87	0,73	0,91	0,91	27,01
Pholoe baltica	0,00	9,07	0,72	1,12	0,89	27,90
Pisione remota	14,06	23,60	0,71	0,86	0,88	28,78
Nemertina	7,98	31,93	0,70	1,28	0,88	29,66
Chaetozone	9,46	17,47	0,70	1,15	0,87	30,53
Nephtys hombergii	0,00	5,30	0,69	1,06	0,86	31,39
Hesionura elongata	23,84	0,80	0,67	0,88	0,84	32,23
Spisula	6,42	0,40	0,67	1,08	0,84	33,07
Oligochaeta	0,00	26,00	0,66	0,79	0,82	33,89
Prionospio	2,00	14,10	0,65	1,17	0,81	34,70