

SUPPLEMENTARY INFORMATION

doi:10.1038/nature11338

Material and Methods

Sampling

In order to compare the commonly used $^{15}\text{N}_2$ tracer addition method to measure N_2 fixation¹ with the addition of $^{15}\text{N}_2$ -enriched water as suggested by Mohr et al. (2010)², seawater was sampled on two cruises in the Atlantic Ocean, the first on board R/V Meteor (M80/1) on a longitudinal transect (23°W) between 15°N and 5°S, the second on board R/V Polarstern (ANT-XXVI/1) on a transect between 54°N and 54°S (Bremerhaven, Germany to Punta Arenas, Chile). In total 39 triplicate incubations were conducted with both methods in parallel. On the M80/1 cruise, seawater was sampled at 11 stations from the surface (bucket), 20 m depth and the chlorophyll maximum (CTD rosette sampler) at 7:00 in the morning, whereas on the ANT-XXVI/1 cruise, seawater was sampled at 6 stations at 16:00 from the ship's clean seawater supply which is installed at 11 m depth (keel of the ship).

Incubations

Seawater samples were filled headspace-free (bubble addition method) or with a 100-150 ml headspace (dissolved method) into 4.5 L polycarbonate bottles and closed with Teflon[®] - coated butyl rubber septum caps. To determine N_2 fixation rates with the bubble-addition method, a 4.5 mL $^{15}\text{N}_2$ gas bubble (Sigma-Aldrich, ≥ 98 atom%) was injected through the septa into each of triplicate bottles (yielding a theoretical enrichment of ~ 12 atom% assuming a rapid isotopic equilibration between the added $^{15}\text{N}_2$ gas and the ambient dissolved N_2 of the water sample). After injection, bottles were gently inverted one hundred times. For comparison of N_2 fixation rates, we added $^{15}\text{N}_2$ -enriched seawater to a second set of triplicate bottles (dissolution method). In detail, the preparation of the $^{15}\text{N}_2$ -enriched seawater was started by degassing filtered seawater (0.2 μm filtered, Durapore) using a membrane flow-through system (Mini-Module, Membrana) in which the seawater flowed on the inside of the membrane and a vacuum (-960 mbar, water jet pump) was applied to the outer side of the membrane. The seawater flow rate was about 400 – 500 mL min^{-1} and seawater was recirculated for the first 10-15 min of the degassing step. Degassed seawater was then filled directly from the flow-through system into evacuated gas-tight 3L Tedlar[®] bags without a headspace. Addition of $^{15}\text{N}_2$ gas was dependent on the amount of seawater in the Tedlar[®] bag and was added at a ratio of 10 ml $^{15}\text{N}_2$ per 1L seawater. The volume of degassed seawater in

the Tedlar® bag was estimated using a balance. During the Meteor cruise, dissolution of the $^{15}\text{N}_2$ gas was achieved by ‘slapping’ the bubble with a ruler. The ‘slapping’ led to a dispersion of the large bubble into numerous small bubbles and thus an increase in the surface area to volume ratio which facilitated gas diffusion. After complete dissolution of the added $^{15}\text{N}_2$ gas ($^{15}\text{N}_2$ -enriched seawater), an aliquot of the $^{15}\text{N}_2$ enriched water was collected for each preparation of enriched seawater and stored in an Exetainer until return to the laboratory where the isotopic composition was measured by membrane-inlet mass spectrometry³. Overall, the average concentration of $^{30}\text{N}_2$ in the prepared batches of enriched water was 246 μM (standard deviation = 24.7 μM). This yielded a ^{15}N -enrichment of about 2% when 100 mL enriched seawater are added to 4.5 L of incubation volume (depending on temperature and salinity). Next, 100–150 ml of $^{15}\text{N}_2$ -enriched seawater were added to each of triplicate bottles before capping the bottles headspace-free with Teflon®-coated butyl rubber septum caps. Bottles were inverted 100 times. Primary production rates were determined in all incubation bottles by the addition of $\text{NaH}^{13}\text{CO}_3$ (~3.5 atom% final) after the addition of $^{15}\text{N}_2$ gas or $^{15}\text{N}_2$ -enriched water. All bottles were placed into on-deck incubators with a surface seawater flow-through and a 25% in situ light level established with a light foil layer (Blue Lagoon, Lee Filters). Incubations were stopped after 24 hours by filtering 2–3 liters of each incubation onto pre-combusted (450°C, 5 hours) GF/F filters (Whatman) under gentle vacuum (-200 mbar). Filters were oven-dried (50°C, 24 hours) (Meteor) and stored over desiccant until analysis or frozen at -20°C directly after filtration (Polarstern). On the Meteor, the remainder of the samples was pooled for nucleic acid sampling within each set of triplicate bottles and a total of 2 L were filtered onto Durapore filters (47 mm, 0.2 μm pore size; Millipore). Samples were shock-frozen in liquid nitrogen and stored at -80°C until analysis.

To determine the natural abundance (*NA-control*) of ^{15}N and ^{13}C isotopes in the incubations, two different controls were incubated alongside with the six experimental bottles. The control for the bubble-addition method was untreated seawater, while the control for the dissolved method received an aliquot of seawater which had previously been degassed but instead of the addition and dissolution of $^{15}\text{N}_2$ for the preparation, ambient air was used. Both controls were processed as described above for the experimental bottles.

Elemental stoichiometry and isotopic composition of particulate organic material

GF/F filters were acidified over fuming HCl overnight in a desiccator. Filters were then oven-dried for 2 hours at 50°C and pelletized in tin cups. Samples were analyzed for particulate

organic carbon and nitrogen (POC and PON) and isotopic composition using a CHN analyzer coupled to an isotope ratio monitoring mass spectrometer. Caffeine standards, calibrated against IAEA standard N1 and N2, were measured every 6 samples.

Calculation of nitrogen and carbon fixation rates

N₂ fixation rates were calculated based on the final isotopic composition of the particulate organic nitrogen after the incubation using the following equation (1):

$$N_2 \text{ fixation rate} = \frac{(A_{\text{sample}}^{PN} - A_{NA\text{-control}}^{PN})}{(A_{N_2} - A_{NA\text{-control}}^{PN})} \times \frac{[PN]}{\Delta t} \quad (1)$$

With A = atom% ¹⁵N in the particulate organic nitrogen (PN) in incubations to which ¹⁵N₂ was added (A_{sample}^{PN}), in control incubations ($A_{NA\text{-control}}^{PN}$) which were simultaneously incubated with the other bottles or in the dissolved N₂ pool (A_{N_2}). For the bubble-addition method, the atom% in the N₂ pool was calculated from the predicted ¹⁵N₂ concentrations according to Mohr *et al.* (2010)². For the dissolution method the ¹⁵N₂ concentration was calculated from the MIMS measurement value in the batch of enriched water for individual experiments and the measured volume of enriched water added to the incubation bottle. Carbon fixation rates were calculated as described for N₂ fixation rates. All rates are displayed as means of triplicate incubations with a standard error.

DNA extraction and determination of nifH phylotype gene copy number

DNA was extracted according to Langlois *et al.* 2008 using the AllPrep DNA/RNA extraction kit (QIAGEN) following the manufacturer's instructions⁴. Amplification of *nifH* genes was performed on an ABI-PRISM 7000 thermocycler, using phylotype specific probes and primers described in Langlois *et al.* 2008⁴ and Foster *et al.* 2007⁵ (Table S1). *nifH* gene copy numbers were calculated based on the attained Ct values and a linear regression from plasmid standards ranging from 10⁷ to 10¹ copies and simultaneously amplified on the same plate.

Table S1. Primers and Probes used in Taqman assays during this study. All sequences reported in 5'-3' direction.

	Reverse (5'-3')	Pos	Forward (5'-3')	Pos	Probe (5'-3')	Pos
UCYN-A	TCAGGACCACCG GACTCAAC	127-146	TAGCTGCAGAAAGA GGAAGTGTAGAAG	50-76	TAATTCCTGGCT ATAACAAC	98-117
Filamentous (Fil)	GCAAATCCACCG CAAACAAC	256-275	TGGCCGTGGTATTAT TACTGCTATC	165-189	AAGGAGCTTAT ACAGATCTA	206-225
Croco (UCYN-B)	TCAGGACCACCA GATTCTACACAC T	122-146	TGCTGAAATGGGTTC TGTTGAA	54-75	CGAAGACGTAA TGCTC	87-102
UCYN-C	GGTATCCTTCAA GTAGTACTTCGT CTAGCT	83-112	TCTACCCGTTTGATG CTACACACTAA	1-26	AAACTACCATTC TTCAGTTAGCAG	32-55
GamAO	AACAATGTAGAT TTCCTGAGCCTT ATTC	294-321	TTATGATGTTCTAGG TGATGTG	240-266	TTGCAATGCCTA TTCG	275-290
Het-1 (Rich-Rizo)	AATACCACGACC CGCACAAC	158-177	CGGTTTCCGTGGTGT ACGTT	105-124	TCCGGTGGTCCT GAGCCTGGTGT	133-155
Het-2 (Rich-Hemi)	AATGCCGCGACC AGCACAAC	158-177	TGGTTACCGTGATGT ACGTT	106-124	TCTGGTGGTCCT GAGCCTGGTGT	133-155

Statistical analysis

To test if the grouping of the Meteor dataset into Equatorial and Tropical North Atlantic stations was statistically significant, we used PRIMER (v. 6^b) to perform a principal component analysis (PCA) and an analysis of similarity (ANOSIM) with the input parameters sample depth, phosphate concentration, *Trichodesmium* dominance (filamentous *nifH* abundance relative to other *nifH* genes), temperature, oxygen, ammonium concentration, salinity and the measured N₂ fixation rates for the bubble addition and dissolution methods (Fig. S2). The PCA showed that temperature, oxygen and phosphate were driving a spread of the data according to water depth, while the dominance of *Trichodesmium*, the measured N₂ fixation rates, salinity and ammonium were responsible for the separation in latitude (Fig. S2A). The ANOSIM indicated that the division into the equatorial and tropical areas was highly significant (p<0.01).




Figure S2. A: PCA analysis showing the samples of the METEOR 23°W transect and the clustering into the tropical (green triangles) and equatorial (blue triangles) groups. The blue lines show the strength and the direction of the influence of the variable over the direction of the data distribution. B: The observed R value of 0.6 (black line) is to the right beyond the distribution function of random sampling of the variables (blue frequency bars), hence showing the statistical significance ($p < 0.01$) of the grouping in the PCA of Fig. S2A.

Areal rates

Areal rates were calculated for the Meteor cruise (15°N – 5°S) according to equation (2):

$$N_{fix} (\mu\text{mol N m}^{-2} \text{ d}^{-1}) = \frac{(r_0 + r_{20})}{2} \times 20\text{m} + \frac{(r_{20} + r_{DCM})}{2} \times (z_{DCM} - 20\text{m}) \quad (2)$$

With the areal N_2 fixation rate N_{fix} , r_0 , r_{20} and r_{DCM} the N_2 fixation rates at surface, 20 meters and the chlorophyll maximum respectively and z_{DCM} the depth of the chlorophyll maximum. Standard errors for areal N_2 fixation rates were propagated according to the propagation of errors for linear combinations⁷.

The areal N_2 fixation rates ($\mu\text{mol N m}^{-2} \text{ d}^{-1}$) calculated according to equation 2 showed a strong linear correlation (Fig. S3, $r^2 = 0.92$) with the N_2 fixation rates ($\text{nmol N l}^{-1} \text{ d}^{-1}$) measured for the surface waters during the Meteor cruise. This linear correlation (equation 3) was subsequently used to convert the N_2 fixation rates at 11m water depth from the Polarstern cruise to areal N_2 fixation rates. Note that the areal rates here only cover the surface mixed layer and may not capture the full profile of N_2 fixation, particularly sub-surface features.

$$N_{2fix} (\mu\text{mol N m}^{-2} \text{ d}^{-1}) = 35.872 \times N_{2fix(surface)} (\text{nmol N L}^{-1} \text{ d}^{-1}) \quad (3)$$




Figure S3. Correlation between N_2 fixation rates measured for the surface waters and calculated (equation 2) areal rates for the Meteor transect for both the bubble addition (red squares) and the dissolution method (blue diamonds). The black line represents the linear relationship for the entire data set with both methods, forced through zero ($r^2 = 0.92$). The slopes of the regressions for the individual regressions of the two methods separately are 36.826 ($r^2 = 0.946$) for the dissolution and 35.851 ($r^2 = 0.8624$) for the bubble-addition method.

Global N_2 fixation estimates from literature values

For the global estimates of N gain via N_2 fixation, published values of bulk water N_2 fixation rates measured with the bubble-addition method (Table S2) were weighted according to the number of stations and averaged over six ocean basins: North and South Pacific, North and South Atlantic and North and South Indian Ocean (Table S3). The basin wide average was multiplied by the dimension of the basin, whose extent was defined by the most northerly and most southerly observation of N_2 fixation mentioned. Since the Indian Ocean is poorly constrained in terms of measurements by the bubble-addition method, we used average rates from the North and South Pacific for the North and South Indian Ocean, respectively.

Table S2: Collection of publications using the bubble-addition method in the Atlantic and Pacific Oceans. S.e., standard error. † Treated as outlier, * as quoted in Mahaffey2005, ** Considering stations 11, 12, 21, 22, 31, 32, 41, 51, 52 and 71 as upwelling stations, ‡ values from supporting online material, *** used for North and South Atlantic with half the stations weighted

Publication	Ocean domain	North limit (°N)	South limit (°N)	West limit (°E)	East limit (°E)	Stations	Areal rate (μmol N m ⁻² d ⁻¹)	S.e.
Rees <i>et al.</i> 2009	Atlantic, English channel ⁸	50.2	49	-5	-4	2	350	
Fernandez <i>et al.</i> 2010	Atlantic, North ⁹	40	15	-29	-28	8	18	3
Voss <i>et al.</i> 2004	Atlantic, Western Tropical North ¹⁰	10	10	-55	-40	4	24	9
Voss <i>et al.</i> 2004	Atlantic, Eastern Tropical North ¹⁰	10	5	-25	-15	6	140	32
Voss <i>et al.</i> 2004	Atlantic, Equatorial ¹⁰	0	0	-26	-24	2	4	
Fernandez <i>et al.</i> 2010	Atlantic, Equatorial ⁹	15	-5	-29	-28	14	60	5
Mourino-Carballido <i>et al.</i> 2011	Atlantic, North ¹¹	30	16	-29	-15	6	11	4
Mourino-Carballido <i>et al.</i> 2011 ***	Atlantic, Equatorial ¹¹	16	-12	-29	-28	7	56	19
Sohm <i>et al.</i> 2011	Atlantic, South East ¹²	-14	-11	-5	0	2	24	
Sohm <i>et al.</i> 2011	Atlantic, South East ¹²	-14.75	-14.75	12	12.2	1	85	
Fernandez <i>et al.</i> 2010	Atlantic, South ⁹	-5	-30	-29	-28	12	7	1
Mourino-Carballido <i>et al.</i> 2011	Atlantic, South ¹¹	-12	-31	-36	-28	7	10	4
Shiozaki <i>et al.</i> 2009	Pacific, North ¹³	44	30	-155	-155	3	0	
Montoya <i>et al.</i> 2004	Pacific, Eastern North Gyre ¹⁴	35	25	-160	-125	10	520	160
Needoba <i>et al.</i> 2007	Pacific, North East ¹⁵	34	34	-129	-129	24	15	
Hamersley <i>et al.</i> 2011	Pacific, North East ¹⁶	34	34	-119	-119	15	150	
White <i>et al.</i> 2007	Pacific, North East ¹⁷	30	22	-122	-122	4	106	55
Shiozaki <i>et al.</i> 2009	Pacific, North ¹³	30	0	-155	-155	8	55	16
Sohm <i>et al.</i> 2011 ‡	Pacific, North ¹⁸	24	19	-160	-154	23	143	21
Sohm <i>et al.</i> 2011 ‡	Pacific, North ¹⁸	23	19	-162	-156	11	216	21
Sohm <i>et al.</i> 2011 ‡	Pacific, North ¹⁸	28	23	-180	-161	8	137	16
Church <i>et al.</i> 2009	Pacific, ALOHA ¹⁹	22.75	22.75	-158	-158	34	111	11
Zehr <i>et al.</i> 2001	Pacific, ALOHA ²⁰	22.75	22.75	-158	-158	1	95	
Montoya <i>et al.</i> 2004	Pacific, ALOHA ¹⁴	22.75	22.75	-158	-158	7	66	19
Grabowski <i>et al.</i> 2008	Pacific, ALOHA ²¹	22.75	22.75	-158	-158	9	72	4
Dore <i>et al.</i> 2002 *	Pacific, ALOHA ²²	22.75	22.75	-158	-158	4	69	13
Voss <i>et al.</i> 2006 **	Pacific, North West ²³	13.5	10	108	110.5	10	26	17
Voss <i>et al.</i> 2006	Pacific, North West ²³	13.5	10	108	110.5	18	71	16
Bombar <i>et al.</i> 2010	Pacific, North West ²⁴	13	10	108	111	4	23	3
Bombar <i>et al.</i> 2010	Pacific, North West ²⁴	13	10	108	111	7	138	45
Bombar <i>et al.</i> 2010	Pacific, North West ²⁴	13	10	108	111	4	88	49
Bombar <i>et al.</i> 2010	Pacific, North West ²⁴	13	10	108	111	4	59	23

Fernandez <i>et al.</i> 2011	Pacific, South East ²⁵	2	-18	-86	-74	8	8	2
Raimbault <i>et al.</i> 2008	Pacific, South West ²⁶	-7	-15	-141	-134	4	110	15
Fernandez <i>et al.</i> 2011	Pacific, South East ²⁵	-13	-20	-78	-70	8	190	29
Raimbault <i>et al.</i> 2008	Pacific, South West ²⁶	-15	-20	-133	-123	4	70	35
Garcia <i>et al.</i> 2007	Pacific, South West ²⁷	-20	-22	166	167	6	290	86
Raimbault <i>et al.</i> 2008	Pacific, South West ²⁶	-20	-30	-123	-101	4	60	15
Raimbault <i>et al.</i> 2008	Pacific, South West ²⁶	-30	-33	-100	-81	6	30	2
Raimbault <i>et al.</i> 2008	Pacific, South West ²⁶	-33	-35	-80	-72	2	90	47
Montoya <i>et al.</i> 2004+	Pacific, Arafura Sea ¹⁴	-10	-20	120	150	2	3955	
Montoya <i>et al.</i> 2004	Pacific, Kaneohe Bay ¹⁴	-9.5	-9.5	135	135	6	24	12
Montoya <i>et al.</i> 2004	Pacific, Arafura Sea ¹⁴	-10	-20	120	150	7	126	47

Table S3: Global rates of annual N₂ fixation. The rates are basin wide averages, obtained by station weighted averaging over the four domains (North Atlantic, South Atlantic, North Pacific and South Pacific Ocean) from table S2. * For the Indian Ocean areal rates of the Pacific Ocean were used.

Ocean Basin	North limit (°N)	South limit (°N)	Stations	Areal rate (μmol N m ⁻² d ⁻¹)	Area (10 ⁶ km ²)	Areal rate (mol N m ⁻² yr ⁻¹)	Basin (mol N yr ⁻¹)	Basin rate (tg N yr ⁻¹)
Atlantic North and Equatorial (weighted average)	50	-5	46	64	39.2	0.023	9.08E+11	12.7
Atlantic South (weighted average)	-5	-30	26	19	15.9	0.007	1.08E+11	1.5
Pacific North (weighted average)	30	0	208	115	56.3	0.042	2.36E+12	33.0
Pacific South (weighted average)	0	-35	55	103	56.1	0.038	2.11E+12	29.5
Indian North	25	0	0	115*	15.8	0.042*	6.62E+11	9.3
Indian South	0	-35	0	103*	31.8	0.038*	1.20E+12	16.7
Global Sum					215.1			102.7

Recently, the annual contribution of N_2 fixation to the Atlantic and Pacific Ocean has been estimated at 63 Tg N yr⁻¹ based on N_2 fixation rate measurements made with the bubble-addition method and the acetylene reduction assay²⁸. This estimate is comparable in magnitude to the 76 Tg N yr⁻¹ obtained in this study for Atlantic and Pacific Ocean combined.

Supplementary Online Material and Figures

Comparison of mean N_2 fixation rates measured by the bubble-addition and the dissolution methods in the tropical and equatorial regions

A pairwise comparison (t-test for dependent samples) of the mean N_2 fixation rates determined with the bubble-addition and the dissolution methods within the tropical and equatorial areas (Fig. S4) indicated that the underestimation by the bubble-addition method was statistically significant in both regions. Moreover, this underestimation was substantially higher in the equatorial region, which was dominated by diazotrophs other than *Trichodesmium*.




Figure S4: Plot of means with standard error for the dissolution method (red boxes) and bubble-addition method (green boxes). Rates were pooled over all depths for the Equatorial (4.5°N - 5°S) and the Tropical North Atlantic (15°N - 5°N) region. The mean N_2 fixation rates measured by the dissolution and the bubble-addition methods were significantly different for both the equatorial region (t-test for dependent samples, $p < 0.01$, $df = 61$), and the tropical region (t-test for dependent samples, $p < 0.01$, $df = 38$). The means of the bubble and dissolution methods over the entire dataset significantly differed by a factor of 2 (1.0 ± 0.2

nmol N L⁻¹ d⁻¹(n=116) and 2.0 +/-0.3 nmol N L⁻¹ d⁻¹ (n=115) (means and s.e.m. of the bubble-addition and the dissolution method, respectively (t-test for dependent samples, p < 0.01, df = 114)).

Table S4: Mean and standard deviation for the 39 parallel comparison experiments. Date and start time mark the beginning of an 24 hour incubation. n.d. = not detectable.

Date	Start time	Depth (m)	Lat (°N)	Lon (°E)	Bubble-addition method		dissolution method	
					(nmol N L ⁻¹ d ⁻¹)	std. dev. (n=3)	(nmol N L ⁻¹ d ⁻¹)	std. dev. (n=3)
28.10.2009	16:00	11	25.3	-16.6	3.88	0.71	3.63	0.47
21.11.2009	11:00	0	13.8	-23	6.94	0.2	9.98	0.6
21.11.2009	11:00	20	13.8	-23	7.15	0.66	12.44	0.57
21.11.2009	11:00	42	13.8	-23	n.d.	-	n.d.	-
19.11.2009	06:00	0	9	-23	3.2	0.39	4.77	2.03
19.11.2009	06:00	20	9	-23	5.74	1.2	4.09	1.72
19.11.2009	06:00	59	9	-23	0.07	0.08	0.09	0.88
17.11.2009	05:00	0	6.8	-23	0.32	0.12	n.d.	-
17.11.2009	05:00	20	6.8	-23	0.13	0.06	n.d.	-
17.11.2009	05:00	64	6.8	-23	n.d.	-	1.75	0.08
31.10.2009	08:00	0	5	-23	0.8	0.17	6.44	4.69
31.10.2009	08:00	20	5	-23	1.32	0.88	3.24	1.85
31.10.2009	08:00	71.3	5	-23	n.d.	-	1.01	0.35
03.11.2009	16:00	11	4.6	-23	0.45	0.02	0.92	0.04
15.11.2009	09:00	0	3.5	-23	0.48	0.21	1.06	0.79
15.11.2009	09:00	20	3.5	-23	0.19	0.13	n.d.	-
15.11.2009	09:00	63	3.5	-23	n.d.	-	n.d.	-
02.11.2009	06:00	0	2	-23	0.31	0.12	4.29	1.81
02.11.2009	06:00	20	2	-23	0.42	0.06	2.44	0.39
02.11.2009	06:00	76.6	2	-23	n.d.	-	1.24	1.03
05.11.2009	08:00	0	0	-23	0.31	0.11	4.23	4.4
05.11.2009	08:00	20	0	-23	0.14	0.16	n.d.	-
05.11.2009	08:00	58.5	0	-23	n.d.	-	n.d.	-
13.11.2009	06:00	0	0	-23	0.04	0.03	0.75	1.48
13.11.2009	06:00	20	0	-23	0.01	0.07	0.02	0.31
13.11.2009	06:00	55	0	-23	n.d.	-	0.75	0.3
07.11.2009	08:00	0	-1.3	-23	0.6	0.89	2.37	1.17
07.11.2009	08:00	20	-1.3	-23	0.1	0.14	1.34	0.2
07.11.2009	08:00	70.2	-1.3	-23	n.d.	-	0.89	0.1
11.11.2009	08:00	0	-2	-23	0.04	0.06	0.64	0.64
11.11.2009	08:00	20	-2	-23	0.01	0.15	0.1	0.04
11.11.2009	08:00	64.4	-2	-23	n.d.	-	n.d.	-
09.11.2009	06:00	0	-4.5	-23	0.02	0.02	0.84	0.73
09.11.2009	06:00	20	-4.5	-23	0.13	0.14	0.07	0.23
09.11.2009	06:00	68.5	-4.5	-23	n.d.	-	n.d.	-
12.11.2009	16:00	11	-24.7	-29.5	1.26	0.07	1.06	0.13
14.11.2009	16:00	11	-28.1	-35.8	4.93	1.41	5.01	0.44
18.11.2009	16:00	11	-38.1	-48.2	0.1	0.01	0.44	0.13
21.11.2009	16:00	11	-44.6	-56.4	0.18	0.06	0.54	0.21

Magnitude of the underestimation by the bubble-addition method relative to the dissolution method




Figure S5: Ratio of N₂ fixation rates measured by dissolution method relative to the bubble-addition method (note the logarithmic scale) as a function of latitude. Ratios were calculated for the paired measurements where N₂ fixation rates were detectable with both methods (n=25). Note that the ratio is particularly large (i.e. the underestimation of N₂-fixation rates by the bubble method is large) in the equatorial regions where diazotrophs other than *Trichodesmium* dominate.

Summary and implications of the results presented in Mohr et al. (2010)

Mohr et al. (2010) showed that the equilibration of a ¹⁵N₂ gas bubble in seawater is slow, taking up to 12 hours to reach ~ 80% of the calculated value based on complete equilibration of the ¹⁵N₂ label with seawater (Fig. S6, upper panel). In contrast, the ¹⁵N₂ label added as dissolved ¹⁵N₂ remained constant with time (Fig. S6, lower panel). Mohr et al. (2010) shows that the strength of agitation (their Figure 2 and 4), bottle size and bubble size (their Figure 3 and 4), affect the kinetics of the ¹⁵N₂ gas equilibration with the surrounding seawater. The consequence of the slow and variable equilibration time of the ¹⁵N₂ gas in field incubations carried out by the bubble addition method¹ is that the rates will be underestimated in most cases.

In view of the slow equilibration of ¹⁵N₂ gas with its surrounding water and the resulting underestimation of the N₂ fixation rate, we suggest that, in the future, gas exchange/equilibration in the acetylene reduction assay (ARA) may also need to be considered to ensure accuracy of the method.




Figure S6: Membrane inlet mass spectrometry measurements of $^{28}\text{N}_2$, $^{29}\text{N}_2$, $^{30}\text{N}_2$, and dissolved oxygen ($^{32}\text{O}_2$) concentrations versus equilibration time for the bubble-addition method (a) and the dissolution method (b). The calculated (expected) values for $^{15}\text{N}_2$ (mass 30) are 37.7 and 25.2 $\mu\text{mol kg}^{-1}$ for the bubble-addition and the dissolution method, respectively (red solid lines) (data redrawn from Mohr et al. (2010)).




Figure S7: Linear correlation between the $^{15}\text{N}/(^{14}\text{N}+^{15}\text{N})$ ratio for dissolved N_2 and the corresponding $^{15}\text{N}/(^{14}\text{N}+^{15}\text{N})$ ratio for particulate nitrogen from the *Crocospaera watsonii* culture after 12 h incubation. The linear relationship shows that ^{15}N -assimilation by these diazotrophs is directly proportional to the ^{15}N -labeling % for dissolved N_2 in the medium. The slope of the regression line is 0.26 indicating that the N-based doubling times for *Crocospaera* were ~ 2 days in our incubations, which is comparable to published growth rates²⁹. The different levels of ^{15}N enrichment in seawater were achieved by pre-equilibrating

a gas bubble for times varying from 1 to 24 hours prior to the inoculation with the *Crocospaera* cultures. For comparison, the values of incubations to which dissolved $^{15}\text{N}_2$ was added are displayed (filled circles). Data redrawn from Mohr et al. (2010).

Considerations about the change in gas composition in an incubation

An incubation of seawater in a bottle always presents a situation that is ideally close to, but not identical to *in situ* conditions. Since a bottle offers a very limited space, it is possible that in a 24 hour incubation some organisms or particulate matter sink to the bottom of the bottle, where oxygen will get more depleted than in the upper part of the bottle. However, such bottle effects are inherent to any kind of incubation and would not differentially affect the methods used here. In both the bubble-addition and the dissolution method a minor part of the gas composition is replaced by the added $^{15}\text{N}_2$ label. This volume needs to be kept as low as possible, but on the other hand the desire is to maximize the addition of $^{15}\text{N}_2$ label, in order to increase the detection limit. By adding 100 mL of degassed water to a 4.5 L incubation (= 2.2%), the oxygen concentration for example drops theoretically from 225 μM to 220 μM . Such a change in oxygen concentration is within the natural variability of surface oxygen concentration due to photosynthesis and respiration. In a bubble-addition type of incubation with 1 mL $^{15}\text{N}_2$ gas per L of seawater the drop in oxygen concentration would be from 225 μM to ~ 216 μM , since the bubble would hold $\sim 20\%$ oxygen after complete equilibration. The manipulation in the gas composition is therefore similar in both methods and within the range of natural variability. In Mohr et al, (2010), the concentrations of the nitrogen isotopes as well as dissolved oxygen were measured in water samples that were incubated with a $^{15}\text{N}_2$ gas bubble as well as with dissolved $^{15}\text{N}_2$ additions (see Fig. S6 above). The measured data confirm the theoretical changes mentioned here.

The strong linear correlation between the isotopic composition of dissolved N_2 in the medium and the N-isotopic composition of *Crocospaera* biomass (Fig. S7) shows that variations in $^{15}\text{N}_2$, and total dissolved N_2 concentrations themselves have no significant effect on the N_2 -fixation rate.

Examples of potential bias introduced by the species composition for the bubble-addition method

There are several ways in which the diazotrophic community composition can affect the magnitude of underestimation in a bubble-addition method incubation. One example is via

the time of N_2 fixation activity relative to the beginning of the incubation. Diazotrophic communities dominated by diazotroph species fixing at night (e.g. *Crocospaera*) will suffer less underestimation than one that is dominated by diazotrophs fixing during the day (e.g. UCYN-A) when the incubation is set up at sunrise, since the $^{15}N_2$ label had more time to equilibrate with the water phase² and vice versa.

The second way is via the relative position in the water, i.e. the “closeness” of the diazotroph to the source of label, the $^{15}N_2$ bubble. The following experiment with *Nodularia*, a buoyant diazotrophic cyanobacterium, suggests that this may also happen in natural communities. A culture of the heterocystous cyanobacterium *Nodularia spumigena* IOW-2000/1 was grown in modified artificial seawater³⁰ supplemented with phosphate and trace metals (ASW). The culture was transferred twice into fresh medium before the experiment. For the experiment, three 450 ml sub-cultures (1 week after last transfer) were transferred from culture flasks into 1.15 L sterile glass bottles. The cultures were supplemented with fresh media up to ~ 1.13 L. Cultures were left over-night to allow filaments to float to the top of the bottle. The next day, cultures were filled with fresh media from the bottom with a syringe in order to keep the floating filaments in place. The bottles were closed headspace-free with Teflon[®]-coated butyl rubber septum caps and 5 mL of $^{15}N_2$ gas were injected just underneath the septum. At the end of the ~ 6 h incubation period during the day time, the floating filaments were separated from the non-floating ones and both were separately filtered onto pre-combusted GF/F filters (Whatman). A natural abundance sample was also filtered at the end of the incubation. Filters were treated as previously described.

The nitrogen fixation rate measured in the floating fraction was significantly higher (t-test, $p < 0.05$, $n = 3$) than the measured rate of the bulk phase (Fig. S8). However, N_2 fixation rates measured with the $^{15}N_2$ isotope added as a dissolved gas (dissolution method) measured during the same experiment were approximately 2 and 6 times higher than those measured with the bubble-addition method for the floating and the non-floating fractions, respectively. Although our experimental design does not rule out that the floating and sinking fraction may have been in a different physiological state and fixing N_2 at different rates, the right panel of Figure S8 shows that a small layer of filaments close to the bubble of pure $^{15}N_2$ gas would have been exposed to very high concentration of $^{15}N_2$ gas during the incubation period. The calculated rates for the floating and non-floating fraction are in agreement with this hypothesis and suggest that such situation may also develop during calm field conditions. The culture results presented here are also congruent with the observation of less discrepancy

between the bubble-addition and dissolution method in areas which were dominated by *Trichodesmium* vs. areas not dominated by *Trichodesmium* which suffered from larger discrepancies.




Figure S8. Left: Biomass-specific N₂ fixation rates of *N.spumigena* measured with the bubble-addition method. Filaments at the very surface of the incubation bottle (floating) were analyzed separately from the bulk liquid phase (non-floating). Error bars indicate standard errors of triplicate incubations. Right: Filaments of *Nodularia* in the incubation bottles. Red box marks the floating fraction analyzed in this experiment.

Literature survey of incubation conditions used for ¹⁵N₂ fixation measurements

Since many factors besides the diazotrophic species composition (see upper section) have the potential to influence the magnitude of difference between the two methods, like bottle size, incubation time and bubble size, it is of interest to know how our experimental conditions compared to those in other studies. Table S5 gives an overview of the published literature and the experimental setup used. The global mean of ¹⁵N₂ gas added per liter of sample is 2.6 mL L⁻¹, the median is 1.0 mL L⁻¹. In this study we used 1 mL L⁻¹ of ¹⁵N₂ gas, which is a representative value of the previously published work. The incubation time and bottle size in this study were 24 h and 4.5 liters, respectively, both representative of conditions used for whole water incubations in oligotrophic regions. In addition, a 24 h incubation also ensures that the N₂ fixation measurement covers a full daily cycle. However, the table below shows the range of incubation times used and that the underestimation or rates may be highly variable especially considering the timing effect of incubation vs. the timing of N₂ fixation activity.

Table S5: Published literature about studies using the $^{15}\text{N}_2$ tracer addition (bubble-addition) method and the parameters used in their incubation setup.

Reference	Year published	Ocean	Study area or organism	Incubation time [h]	$^{15}\text{N}_2$ [ml L^{-1}]	bottle size [L]
Benavides <i>et al.</i>	2011 ³¹	Atlantic	NE Atlantic	24	1.6	1.24
Bombar <i>et al.</i>	2011 ³²	South China Sea	Mekong River Plume	6-12	0.9	2.3
Bombar <i>et al.</i>	2010 ³³	South China Sea	upwelling	6	1.1	2.3
Bonnet <i>et al.</i>	2009 ³⁴	Pacific	Equatorial W Pacific	24	0.7	4.5
Bonnet <i>et al.</i>	2011 ³⁵	Mediterranean		24	0.9	4.5
Burns <i>et al.</i>	2006 ³⁶	Atlantic / Pacific	<i>Trichodesmium</i>	4	0.4	0.25
Capone <i>et al.</i>	2005 ³⁷	Atlantic	<i>Trichodesmium</i>	2	0.3	0.31
Chen <i>et al.</i>	2008 ³⁸	South China Sea	Kuroshio and South China Sea Basin, <i>Trichodesmium</i>	3	1.7	0.12
Chen <i>et al.</i>	2011 ³⁹	Pacific	NW Pacific / Kuroshio / <i>Trichodesmium</i>	3-5	1.7	0.12
Chen <i>et al.</i>	2011 ³⁹	culture	<i>Trichodesmium</i>	24	5	1.2
Church <i>et al.</i>	2009 ¹⁹	Pacific	N Pacific subtropical gyre, station ALOHA	24	0.7	4.5
Degerholm <i>et al.</i>	2008 ⁴⁰	Baltic Sea		4	18.4	0.25
Dore <i>et al.</i>	2002 ²²	Pacific	subtropical N Pacific	24	0.2	4.7
Falcón <i>et al.</i>	2004 ⁴¹	Atlantic / Pacific	Unicellular	24	0.64	0.014
Fernandez	2010 ⁴²	Atlantic	Atlantic	24	1	2

<i>et al.</i>						
Fernandez <i>et al.</i>	2011 ²⁵	Pacific	tropical SE Pacific	24	2	2
Fong <i>et al.</i>	2008 ⁴³	Pacific	N Pacific subtropical gyre	24	0.7	4.5
Gandhi <i>et al.</i>	2011 ⁴⁴	Indian Ocean	Arabian Sea	4	1.6	1.25
Garcia <i>et al.</i>	2011 ⁴⁵	culture	<i>Trichodesmium</i>	12	1	0.16
Garcia <i>et al.</i>	2007 ²⁷	Pacific	SW Pacific	12	1.7	1.2
Goebel <i>et al.</i>	2010 ⁴⁶	Atlantic	tropical Atlantic	24	0.1	3.8
Grabowski <i>et al.</i>	2008 ²¹	Pacific	ALOHA	24	0.25	4.5*
Grosse <i>et al.</i>	2010 ⁴⁷	South China Sea	Mekong River Plume	6-12	0.9	2.3
Holl <i>et al.</i>	2007 ⁴⁸	Atlantic	W Gulf of Mexico / <i>Trichodesmium</i>	6	0.96	0.25
Ibello <i>et al.</i>	2010 ⁴⁹	Mediterranean		24	1	4.6
Konno <i>et al.</i>	2010 ⁵⁰	Pacific	NW Pacific	24-72	2-4	0.25- 0.5
Law <i>et al.</i>	2011 ⁵¹	Pacific	SW Pacific	12-36	1	2.4
Maranon <i>et al.</i>	2010 ⁵²	Atlantic	central Atlantic	24	1	2
Mills <i>et al.</i>	2004 ⁵³	Atlantic	eastern tropical N Atlantic	24	0.85	1.18
Moisander <i>et al.</i>	1996 ⁵⁴	Baltic Sea	Field and culture	2	40	0.0245
Moore <i>et al.</i>	2009 ⁵⁵	Pacific	Fiji to Hawaii	24	0.7	4.5
Mourino- Carballido	2011 ⁵⁶	Atlantic	tropical and subtropical Atlantic	24	1	2
Moutin <i>et al.</i>	2008 ⁵⁷	Pacific	tropical S Pacific	24	1.7	0.6
Mulholland	2005 ⁵⁸	culture	<i>Trichodesmium</i>	1	1	0.16

and Bernhardt						
Mulholland <i>et al.</i>	2004 ⁵⁹	culture	<i>Trichodesmium</i>	2	1	0.16
Needoba <i>et al.</i>	2007 ¹⁵	Pacific	oligotrophic N Pacific Ocean	48	2	4
Orcutt <i>et al.</i>	2001 ⁶⁰	Atlantic	BATS/ <i>Trichodesmium</i>	6	14.3	0.014
Ploug <i>et al.</i>	2010 ⁶¹	Baltic Sea	<i>Aphanizomenon</i>	3-6	8	0.25
Raimbault and Garcia	2008 ²⁶	Pacific	S Pacific Ocean	24	3.5	0.58
Rees <i>et al.</i>	2009 ⁸	Atlantic	western English Channel	24	2	0.64
Rees <i>et al.</i>	2006 ⁶²	Mediterranean	E Mediterranean Sea	24	2	10
Ridame <i>et al.</i>	2011 ⁶³	Mediterranean		24	1.1	4.5
Rijkenberg <i>et al.</i>	2011 ⁶⁴	Atlantic	subtropical NE Atlantic	≥24	1	4
Sandrone <i>et al.</i>	2007 ⁶⁵	Mediterranean	NW Mediterranean Sea	12	1.7	0.6
Shiozaki <i>et al.</i>	2009 ¹³	Pacific	NW Pacific (along 155°E)	24	0.5	4.5
Sohm <i>et al.</i> , A	2011 ¹²	Atlantic	South Atlantic Gyre	24	0.7	4.5
Sohm <i>et al.</i> , B	2011 ¹⁸	Pacific	whole water and <i>Trichodesmium</i>	24	0.7	4.5
Ternon <i>et al.</i>	2011 ⁶⁶	Mediterranean		24	1.1	4.5
Turk <i>et al.</i>	2011 ⁶⁷	Atlantic	tropical NE Atlantic	6	2	1
Twomey <i>et al.</i>	2007 ⁶⁸	Indian	SW coast of Australia	6	0.8	4
Voss <i>et al.</i>	2004 ¹⁰	Atlantic	tropical N Atlantic, transect at 10°N	6	1	1
Voss <i>et al.</i>	2006 ²³	South China Sea	Off Vietnam	6	1.1	2.3

Wannicke <i>et al.</i>	2010 ⁶⁹	Atlantic	NE Atlantic	1-20	1	2.5
Wasmund <i>et al.</i>	2005 ⁷⁰	Baltic Sea		2	2	0.25
Wasmund <i>et al.</i>	2001 ⁷¹	Baltic Sea		8	4	0.25
Watkins-Brandt <i>et al.</i>	2011 ⁷²	Pacific	North Pacific	24	0.5	4.4
White <i>et al.</i>	2007 ¹⁷	Pacific	Gulf of California	24	0.3	2
Yogev <i>et al.</i>	2011 ⁷³	Mediterranean	E Mediterranean Sea	24-30	2	4.5
Mean				15	2.6	2.2
Median				18	1.0	2.0
This study	2012	Atlantic		24	1	4.5

References

1. Montoya, J. P., Voss, M., Kahler, P. & Capone, D. G. A simple, high-precision, high-sensitivity tracer assay for N₂ fixation. *Applied and Environmental Microbiology* **62**, 986-993 (1996).
2. Mohr, W., Grosskopf, T., Wallace, D. W. R. & LaRoche, J. Methodological Underestimation of Oceanic Nitrogen Fixation Rates. *Plos One* **5(9)**, e12583. doi:10.1371/journal.pone.0012583 (2010).
3. Gao, H. et al. Aerobic denitrification in permeable Wadden Sea sediments. *Isme Journal* **4**, 417-426 (2010).
4. Langlois, R. J., Hummer, D. & LaRoche, J. Abundances and distributions of the dominant nifH phylotypes in the Northern Atlantic Ocean. *Applied and Environmental Microbiology* **74**, 1922-1931 (2008).
5. Foster, R. A. et al. Influence of the Amazon River plume on distributions of free-living and symbiotic cyanobacteria in the western tropical north Atlantic Ocean. *Limnology and Oceanography* **52**, 517-532 (2007).
6. Clarke, K. R. Nonparametric Multivariate Analyses of Changes in Community Structure. *Australian Journal of Ecology* **18**, 117-143 (1993).
7. Miller, J. N. & Miller, J. C. *Statistics and Chemometrics in Analytical Chemistry* (Pearson Education Limited, Harlow, England, 2010).

8. Rees, A. P., Gilbert, J. A. & Kelly-Gerreyn, B. A. Nitrogen fixation in the western English Channel (NE Atlantic Ocean). *Marine Ecology-Progress Series* **374**, 7-12 (2009).
9. Fernandez, A., Mourino-Carballido, B., Bode, A., Varela, M. & Maranon, E. Latitudinal distribution of *Trichodesmium* spp. and N₂ fixation in the Atlantic Ocean. *Biogeosciences* **7**, 3167-3176 (2010).
10. Voss, M., Croot, P., Lochte, K., Mills, M. & Peeken, I. Patterns of nitrogen fixation along 10N in the tropical Atlantic. *Geophysical Research Letters* **31** (2004).
11. Mourino-Carballido, B. et al. Importance of N₂ fixation vs. nitrate eddy diffusion along a latitudinal transect in the Atlantic Ocean. *Limnology and Oceanography* **56**, 999-1007 (2011).
12. Sohm, J. A. et al. Nitrogen fixation in the South Atlantic Gyre and the Benguela Upwelling System. *Geophysical Research Letters* **38** (2011).
13. Shiozaki, T., Furuya, K., Kodama, T. & Takeda, S. Contribution of N₂ fixation to new production in the western North Pacific Ocean along 155 degrees E. *Marine Ecology-Progress Series* **377**, 19-32 (2009).
14. Montoya, J. P. et al. High rates of N₂ fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean. *Nature* **430**, 1027-1031 (2004).
15. Needoba, J. A., Foster, R. A., Sakamoto, C., Zehr, J. P. & Johnson, K. S. Nitrogen fixation by unicellular diazotrophic cyanobacteria in the temperate oligotrophic North Pacific Ocean. *Limnology and Oceanography* **52**, 1317-1327 (2007).
16. Hamersley, M. R. et al. Nitrogen fixation within the water column associated with two hypoxic basins in the Southern California Bight. *Aquatic Microbial Ecology* **63**, 193-+ (2011).
17. White, A. E., Prahl, F. G., Letelier, R. M. & Popp, B. N. Summer surface waters in the Gulf of California: Prime habitat for biological N₂ fixation. *Global Biogeochemical Cycles* **21** (2007).
18. Sohm, J. A., Subramaniam, A., Gunderson, T. E., Carpenter, E. J. & Capone, D. G. Nitrogen fixation by *Trichodesmium* spp. and unicellular diazotrophs in the North Pacific Subtropical Gyre. *Journal of Geophysical Research-Biogeosciences* **116** (2011).

19. Church, M. J. et al. Physical forcing of nitrogen fixation and diazotroph community structure in the North Pacific subtropical gyre. *Global Biogeochemical Cycles* **23** (2009).
20. Zehr, J. P. et al. Unicellular cyanobacteria fix N₂ in the subtropical North Pacific Ocean. *Nature* **412**, 635-638 (2001).
21. Grabowski, M. N. W., Church, M. J. & Karl, D. M. Nitrogen fixation rates and controls at Stn ALOHA. *Aquatic Microbial Ecology* **52**, 175-183 (2008).
22. Dore, J. E., Brum, J. R., Tupas, L. M. & Karl, D. M. Seasonal and interannual variability in sources of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean. *Limnology and Oceanography* **47**, 1595-1607 (2002).
23. Voss, M., Bombar, D., Loick, N. & Dippner, J. W. Riverine influence on nitrogen fixation in the upwelling region off Vietnam, South China Sea. *Geophysical Research Letters* **33** (2006).
24. Bombar, D. et al. Sources of new nitrogen in the Vietnamese upwelling region of the South China Sea. *Journal of Geophysical Research-Oceans* **115** (2010).
25. Fernandez, C., Farias, L. & Ulloa, O. Nitrogen Fixation in Denitrified Marine Waters. *Plos One* **6**, e20539 (2011).
26. Raimbault, P. & Garcia, N. Evidence for efficient regenerated production and dinitrogen fixation in nitrogen-deficient waters of the South Pacific Ocean: impact on new and export production estimates. *Biogeosciences* **5**, 323-338 (2008).
27. Garcia, N., Raimbault, P. & Sandroni, V. Seasonal nitrogen fixation and primary production in the Southwest Pacific: nanoplankton diazotrophy and transfer of nitrogen to picoplankton organisms. *Marine Ecology-Progress Series* **343**, 25-33 (2007).
28. Luo, Y. W. et al. Database of diazotrophs in global ocean: abundances, biomass and nitrogen fixation rates. *Earth System Science Data Discussions* **5**, 47-106, doi:10.5194/essdd-5-47-2012 (2012).
29. Goebel, N. L., Edwards, C. A., Carter, B. J., Achilles, K. M. & Zehr, J. P. Growth and carbon content of three different-sized diazotrophic cyanobacteria observed in the subtropical North Pacific. *Journal of Phycology* **44**, 1212-1220 (2008).
30. Kester, D. R., Duedall, I. W., Connors, D. N. & Pytkowicz, R. M. Preparation of artificial seawater. *Limnology and Oceanography* **12**, 176-179 (1967).

31. Benavides, M., Agawin, N. S. R., Aristegui, J., Ferriol, P. & Stal, L. J. Nitrogen fixation by *Trichodesmium* and small diazotrophs in the subtropical northeast Atlantic. *Aquatic Microbial Ecology* **65**, 43-53 (2011).
32. Bombar, D. et al. Distribution of diazotrophic microorganisms and *nifH* gene expression in the Mekong River plume during intermonsoon. *Marine Ecology-Progress Series* **424**, 39-U55 (2011).
33. Bombar, D. et al. Sources of new nitrogen in the Vietnamese upwelling region of the South China Sea. *Journal of Geophysical Research-Oceans* **115** (2010).
34. Bonnet, S., Biegala, I. C., Dutrieux, P., Slemmons, L. O. & Capone, D. G. Nitrogen fixation in the western equatorial Pacific: Rates, diazotrophic cyanobacterial size class distribution, and biogeochemical significance. *Global Biogeochemical Cycles* **23** (2009).
35. Bonnet, S., Grosso, O. & Moutin, T. Planktonic dinitrogen fixation along a longitudinal gradient across the Mediterranean Sea during the stratified period (BOUM cruise). *Biogeosciences* **8**, 2257-2267 (2011).
36. Burns, J. A., Zehr, J. P., Montoya, J. P., Kustka, A. B. & Capone, D. G. Effect of EDTA additions on natural *Trichodesmium* spp. (Cyanophyta) populations. *Journal of Phycology* **42**, 900-904 (2006).
37. Capone, D. G. et al. Nitrogen fixation by *Trichodesmium* spp.: An important source of new nitrogen to the tropical and subtropical North Atlantic Ocean. *Global Biogeochemical Cycles* **19** (2005).
38. Chen, Y.-l. L., Chen, H.-Y., Tuo, S.-h. & Ohki, K. Seasonal dynamics of new production from *Trichodesmium* N₂ fixation and nitrate uptake in the upstream Kuroshio and South China Sea basin. *Limnology and Oceanography* **53**, 1705-1721 (2008).
39. Chen, Y.-l. L., Tuo, S.-h. & Chen, H.-Y. Co-occurrence and transfer of fixed nitrogen from *Trichodesmium* spp. to diatoms in the low-latitude Kuroshio Current in the NW Pacific. *Marine Ecology-Progress Series* **421**, 25-38 (2011).
40. Degerholm, J., Gundersen, K., Bergman, B. & Soderback, E. Seasonal significance of N₂ fixation in coastal and offshore waters of the northwestern Baltic Sea. *Marine Ecology-Progress Series* **360**, 73-84 (2008).

41. Falcon, L. I., Carpenter, E. J., Cipriano, F., Bergman, B. & Capone, D. G. N₂ fixation by unicellular bacterioplankton from the Atlantic and Pacific oceans: Phylogeny and in situ rates. *Applied and Environmental Microbiology* **70**, 765-770 (2004).
42. Fernandez, A., Mourino-Carballido, B., Bode, A., Varela, M. & Maranon, E. Latitudinal distribution of *Trichodesmium* spp. and N₂ fixation in the Atlantic Ocean. *Biogeosciences* **7**, 3167-3176 (2010).
43. Fong, A. A. et al. Nitrogen fixation in an anticyclonic eddy in the oligotrophic North Pacific Ocean. *Isme Journal* **2**, 663-676 (2008).
44. Gandhi, N. et al. First direct measurements of N₂ fixation during a *Trichodesmium* bloom in the eastern Arabian Sea. *Global Biogeochemical Cycles* **25** (2011).
45. Garcia, N. S. et al. Interactive Effects of Irradiance and Co(2) on Co(2) Fixation and N₂ Fixation in the Diazotroph *Trichodesmium Erythraeum* (Cyanobacteria). *Journal of Phycology* **47**, 1292-1303 (2011).
46. Goebel, N. L. et al. Abundance and distribution of major groups of diazotrophic cyanobacteria and their potential contribution to N₂ fixation in the tropical Atlantic Ocean. *Environmental Microbiology* **12**, 3272-3289 (2010).
47. Grosse, J., Bombar, D., Hai Nhu, D., Lam Ngoc, N. & Voss, M. The Mekong River plume fuels nitrogen fixation and determines phytoplankton species distribution in the South China Sea during low- and high-discharge season. *Limnology and Oceanography* **55**, 1668-1680 (2010).
48. Holl, C. M. et al. *Trichodesmium* in the western Gulf of Mexico: N-15(2)-fixation and natural abundance stable isotope evidence. *Limnology and Oceanography* **52**, 2249-2259 (2007).
49. Ibello, V., Cantoni, C., Cozzi, S. & Civitarese, G. First basin-wide experimental results on N₂ fixation in the open Mediterranean Sea. *Geophysical Research Letters* **37** (2010).
50. Konno, U. et al. Determination of total N₂ fixation rates in the ocean taking into account both the particulate and filtrate fractions. *Biogeosciences* **7**, 2369-2377 (2010).
51. Law, C. S. et al. Response of surface nutrient inventories and nitrogen fixation to a tropical cyclone in the southwest Pacific. *Limnology and Oceanography* **56**, 1372-1385 (2011).

52. Maranon, E. et al. Degree of oligotrophy controls the response of microbial plankton to Saharan dust. *Limnology and Oceanography* **55**, 2339-2352 (2010).
53. Mills, M. M., Ridame, C., Davey, M., La Roche, J. & Geider, R. J. Iron and phosphorus co-limit nitrogen fixation in the eastern tropical North Atlantic. *Nature* **429**, 292-294 (2004).
54. Moisander, P., Lehtimäki, J., Sivonen, K. & Kononen, K. Comparison of $^{15}\text{N}_2$ and acetylene reduction methods for the measurement of nitrogen fixation by Baltic Sea cyanobacteria. *Phycologia* **35**, 140-146 (1996).
55. Moore, R. M., Punshon, S., Mahaffey, C. & Karl, D. The relationship between dissolved hydrogen and nitrogen fixation in ocean waters. *Deep-Sea Research Part I-Oceanographic Research Papers* **56**, 1449-1458 (2009).
56. Mourino-Carballido, B. et al. Importance of N_2 fixation vs. nitrate eddy diffusion along a latitudinal transect in the Atlantic Ocean. *Limnology and Oceanography* **56**, 999-1007 (2011).
57. Moutin, T. et al. Phosphate availability and the ultimate control of new nitrogen input by nitrogen fixation in the tropical Pacific Ocean. *Biogeosciences* **5**, 95-109 (2008).
58. Mulholland, M. R. & Bernhardt, P. W. The effect of growth rate, phosphorus concentration, and temperature on N_2 fixation, carbon fixation, and nitrogen release in continuous cultures of *Trichodesmium* IMS101. *Limnology and Oceanography* **50**, 839-849 (2005).
59. Mulholland, M. R., Bronk, D. A. & Capone, D. G. Dinitrogen fixation and release of ammonium and dissolved organic nitrogen by *Trichodesmium* IMS101. *Aquatic Microbial Ecology* **37**, 85-94 (2004).
60. Orcutt, K. M. et al. A seasonal study of the significance of N_2 fixation by *Trichodesmium* spp. at the Bermuda Atlantic Time-series Study (BATS) site. *Deep-Sea Research Part II-Topical Studies in Oceanography* **48**, 1583-1608 (2001).
61. Ploug, H. et al. Carbon and nitrogen fluxes associated with the cyanobacterium *Aphanizomenon* sp. in the Baltic Sea. *Isme Journal* **4**, 1215-1223 (2010).
62. Rees, A. P., Law, C. S. & Woodward, E. M. S. High rates of nitrogen fixation during an in-situ phosphate release experiment in the Eastern Mediterranean Sea. *Geophysical Research Letters* **33** (2006).
63. Ridame, C. et al. Nutrient control of N_2 fixation in the oligotrophic Mediterranean Sea and the impact of Saharan dust events. *Biogeosciences* **8**, 2773-2783 (2011).

64. Rijkenberg, M. J. A. et al. Environmental Forcing of Nitrogen Fixation in the Eastern Tropical and Sub-Tropical North Atlantic Ocean. *Plos One* **6**, e28989 (2011).
65. Sandroni, V., Raimbault, P., Migon, C., Garcia, N. & Gouze, E. Dry atmospheric deposition and diazotrophy as sources of new nitrogen to northwestern Mediterranean oligotrophic surface waters. *Deep-Sea Research Part I-Oceanographic Research Papers* **54**, 1859-1870 (2007).
66. Ternon, E., Guieu, C., Ridame, C., L'Helguen, S. & Catala, P. Longitudinal variability of the biogeochemical role of Mediterranean aerosols in the Mediterranean Sea. *Biogeosciences* **8**, 1067-1080 (2011).
67. Turk, K. A. et al. Nitrogen fixation and nitrogenase (nifH) expression in tropical waters of the eastern North Atlantic. *Isme Journal* **5**, 1201-1212 (2011).
68. Twomey, L. J., Waite, A. M., Pez, V. & Pattiaratchi, C. B. Variability in nitrogen uptake and fixation in the oligotrophic waters off the south west coast of Australia. *Deep-Sea Research Part II-Topical Studies in Oceanography* **54**, 925-942 (2007).
69. Wannicke, N., Liskow, I. & Voss, M. Impact of diazotrophy on N stable isotope signatures of nitrate and particulate organic nitrogen: case studies in the north-eastern tropical Atlantic Ocean. *Isotopes in Environmental and Health Studies* **46**, 337-354 (2010).
70. Wasmund, N., Nausch, G., Schneider, B., Nagel, K. & Voss, M. Comparison of nitrogen fixation rates determined with different methods: a study in the Baltic Proper. *Marine Ecology-Progress Series* **297**, 23-31 (2005).
71. Wasmund, N., Voss, M. & Lochte, K. Evidence of nitrogen fixation by non-heterocystous cyanobacteria in the Baltic Sea and re-calculation of a budget of nitrogen fixation. *Marine Ecology-Progress Series* **214**, 1-14 (2001).
72. Watkins-Brandt, K. S. et al. Addition of inorganic or organic phosphorus enhances nitrogen and carbon fixation in the oligotrophic North Pacific. *Marine Ecology-Progress Series* **432**, 17-29 (2011).
73. Yogev, T. et al. Is dinitrogen fixation significant in the Levantine Basin, East Mediterranean Sea? *Environmental Microbiology* **13**, 854-871 (2011).