



## Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>  
Eprints ID : 4102

**To link to this article** : DOI:10.1093/jxb/erh258  
URL : <http://dx.doi.org/10.1093/jxb/erh258>

**To cite this version** : Allègre, Adeline and Silvestre, Jérôme and Morard, Philippe and Kallerhoff, Jean and Pinelli, Eric *Nitrate reductase regulation in tomato roots by exogenous nitrate: a possible role in tolerance to long-term root anoxia*. (2004) Journal of Experimental Botany, Vol. 55 (n° 408). pp. 2625-2634. ISSN 2473-2482

Any correspondance concerning this service should be sent to the repository administrator: [staff-oatao@listes-diff.inp-toulouse.fr](mailto:staff-oatao@listes-diff.inp-toulouse.fr)

# Nitrate reductase regulation in tomato roots by exogenous nitrate: a possible role in tolerance to long-term root anoxia

Adeline Allègre, Jérôme Silvestre, Philippe Morard, Jean Kallerhoff and Eric Pinelli\*

Laboratoire d'Agronomie Environnement Ecotoxicologie, ENSAT, Avenue de l'Agrobiopôle-Auzeville, Tolosane BP 107, F-31326 Castanet-Tolosan cédex, France

## Abstract

The mechanism of nitrate reductase (NR) regulation under long-term anoxia in roots of whole plants and the putative role of nitrate in anoxia tolerance have been addressed. NR activity in tomato roots increased significantly after 24 h of anaerobiosis and increased further by 48 h, with a concomitant release of nitrite into the culture medium. Anoxia promoted NR activation through dissociation of the 14-3-3 protein inhibitor and NR dephosphorylation. After 24 h of anoxia, the total amount of NR increased slightly up to 48 h. However, NR-mRNA levels remained constant between 0 h and 24 h of root anoxia and decreased after 48 h. This is probably due to the inhibition of NR degradation and the accumulation of its native form. NR was slightly dephosphorylated in the absence of oxygen and nitrate. Under anoxia, NR dephosphorylation was modulated by nitrate-controlled NR activity. In addition, the presence of nitrate prevents anoxic symptoms on leaves and delays wilting by 48 h during root anoxia. In the absence of nitrate, plants withered within 24 h, as they did with tungstate treatment, an inhibitor of NR activity. Thus, anoxia tolerance of tomato roots could be enhanced by nitrate reduction.

Key words: Anoxia, nitrate, nitrate reductase, nitrite, NR regulation, tomato roots.

## Introduction

Nitrate reductase (NR) is the key enzyme in the overall process of nitrate assimilation by plants (Botrel *et al.*, 1996; Datta and Sharma, 1999; Campbell, 2001; de la Haba *et al.*, 2001). Nitrate absorbed by roots is reduced to nitrite by nitrate reductase and then nitrite is reduced by nitrite

reductase to ammonium ion. Ammonium is thereafter incorporated into amino acids by the glutamine synthetase–glutamine-2-oxoglutarate transaminase (GS–GOGAT) enzyme system (Aslam *et al.*, 2001), providing plant autotrophy for nitrogen (Heller *et al.*, 1998). Under stress conditions, nitrite is also reduced to nitric oxide by nitrate reductase. Nitric oxide has various toxic effects and constitutes a signalling substance, so its effects should be broader. To limit the toxic effects of nitrite on root tissues, the NR activity is highly regulated through its degradation, activation, or inactivation.

Various studies indicate that nitrate reductase synthesis is promoted by nitrate, light, phytochrome, and photosynthesis (Appenroth *et al.*, 2000; de la Haba *et al.*, 2001). Nitrate reductase activity is modulated both by phosphorylation and by binding to a specific adapter protein known as 14-3-3 (Weiner and Kaiser, 1999, 2000). In spinach, phosphorylation of nitrate reductase on the 543-serine residue is the first step of enzyme inactivation (Kanamaru *et al.*, 1999). Complete inactivation occurs when the phosphorylated form is associated with the 14-3-3 protein. This binding is initiated by divalent cations such as Mg<sup>2+</sup> and Ca<sup>2+</sup> (Weiner and Kaiser, 2001). Nitrate reductase activation is related to its dissociation from the 14-3-3 protein as well as to its dephosphorylation (Huber *et al.*, 1996; Chung *et al.*, 1999). However, information on nitrate reductase regulation in roots is scarce. In the case of barley, the enzyme is activated by uncouplers, mannose feeding, or anoxia (Botrel *et al.*, 1996; de la Haba *et al.*, 2001). Anoxic root tissues have a higher level of NR activity than aerated controls (Glaab and Kaiser, 1993), but the mechanisms regulating NR in anoxic roots are not well defined and the role of NR activation remains to be unravelled.

Nitrate requirement for induction of nitrate reductase is now well established: the enhancement of NR activity is regulated by the metabolic pool of nitrate localized in the

\* To whom correspondence should be addressed. Fax: + 33 5 62 19 39 01. E-mail: pinelli@ensat.fr

cytoplasm under normoxic conditions (Ferrari *et al.*, 1973; Aslam *et al.*, 1976). The metabolic or active pool of nitrate is smaller than the large non-metabolic storage pool contained within the vacuole (Martinoia *et al.*, 1981). The concentration of nitrate in the cytoplasmic pool is lower and more constant than in the vacuolar pool (Zhen *et al.*, 1991). In barley root cells under normoxia, the exogenous concentration of nitrate determines the level of nitrate ions in the metabolic and non-metabolic pools (Aslam *et al.*, 2001). Exogenous nitrate supplied to the medium of anaerobic excised rice roots increased the adenylic energy charge and lowered the catabolic reducing charge (Reggiani *et al.*, 1985). Both factors could contribute to root survival and suggest a possible role of nitrate reduction in anoxia tolerance.

Anoxia studies were generally carried out during short periods (Botrel *et al.*, 1996; de la Haba *et al.*, 2001) on non-intact plants like maize (Xia *et al.*, 1995; Subbaiah and Sachs, 2003) or intact non-crop plants such as *Arabidopsis* (Ellis *et al.*, 1999; Paul *et al.*, 2004). The studies reported here were conducted on the effects of long-term anoxia (over 24 h) on tomato plants grown in soilless culture. In the same experimental conditions, the presence of nitrite was detected in the nutrient solution 16 h after the beginning of anoxia (Morard *et al.*, 2000). This result was corroborated on excised tomato roots (Morard *et al.*, 2004b) and also with sterile *in vitro*-cultured root tissues of tomato. Nitrate reduction is, therefore, a biochemical process linked to root cell metabolism and is not related to micro-organism activity (Morard *et al.*, 2004c).

The aim of this study was to determine the effect of long-term oxygen deprivation on NR expression and regulation. The role of nitrate on NR modulation and resistance to root anoxia was also investigated. In this paper, the following points were studied and discussed: (i) NR activity in oxygen-deprived roots, (ii) regulation of NR in anoxic conditions, and (iii) role of nitrate supplied in anoxia tolerance.

## Materials and methods

### Plant growth conditions

Experiments were conducted on 7-week-old tomato plants (*Lycopersicon esculentum* Mill.) cv. Rondello at the 10–12 leaf stage (1st flush). Plants were grown hydroponically in a culture room with a relative humidity of 50% and photoperiod of 14/10 h light/dark ( $170 \mu\text{E m}^{-2} \text{s}^{-1}$ ) provided by Philips daylight fluorescent tubes. The day/night temperature was 20/18 °C. Tomato seeds were germinated on filter paper moistened with distilled water for 1 week at 23 °C in the dark. After 7 d, plants were grown individually in a separate PVC tank containing 15 l of nutrient solution over 6 weeks. The nutrient solution contained macronutrients: 2.25 mM KNO<sub>3</sub>, 0.25 mM Ca(NO<sub>3</sub>)<sub>2</sub>, 0.35 mM KH<sub>2</sub>PO<sub>4</sub>, and 0.075 mM MgSO<sub>4</sub>; micronutrients: 268.6 μM EDTA-Fe, 8.9 μM MnSO<sub>4</sub>, 24.1 μM H<sub>3</sub>BO<sub>3</sub>, 1.7 μM ZnSO<sub>4</sub>, 3.9 μM CuSO<sub>4</sub>, and 0.1 μM Na<sub>2</sub>MoO<sub>4</sub> (Morard *et al.*, 2004a). The solution was renewed every 2 weeks or as soon as the pH

reached 7 and/or when the conductivity was reduced by half with respect to the initial value. In each experiment, the mean root fresh weight expressed per litre of nutrient solution was  $2.7 \pm 0.5$  g.

### Root anoxia

After 5 weeks, plants were transferred to 8.0 l PVC tanks containing culture medium. Shortage of oxygen was imposed progressively by sealing the PVC tank with an airtight cover, by stopping aeration, and by including an airtight seal between the root and shoot. Oxygen dissolved in the nutrient solution being thus depleted, anoxia is reached when the root system has taken up all the oxygen dissolved in the nutrient solution (Morard and Silvestre, 1996). Total depletion occurs approximately 10 h after the beginning of oxygen shortage (Morard *et al.*, 2000). By contrast, control plants were continuously bubbled with air.

Three separate sets of experiments were conducted. In the first one, whole tomato plants grown in complete medium were deprived of oxygen for 24 h and 48 h, respectively, and harvested after each period. Control plants were continuously aerated. In a second set of experiments, root anoxia was conducted in the absence of nitrate for 48 h on medium containing solely 0.22 mM K<sub>2</sub>SO<sub>4</sub>, 2.16 mM KH<sub>2</sub>PO<sub>4</sub>, 0.25 mM CaSO<sub>4</sub>, and 0.075 mM MgSO<sub>4</sub> with micronutrients. Plants were harvested after 48 h of anoxia. The aerated controls were grown on nitrate-depleted medium. In a third set of anoxia experiments, sodium molybdate was substituted by 40 μM sodium tungstate in the nutrient medium to inhibit NR activity (Deng *et al.*, 1989). In this case, ammonium was included in the medium so as to maintain nitrogen assimilation. The composition of this solution was 2 mM NH<sub>4</sub>Cl, 0.25 mM Ca(NO<sub>3</sub>)<sub>2</sub>, 2.25 mM KNO<sub>3</sub>, 0.075 mM MgSO<sub>4</sub>, 0.35 mM KH<sub>2</sub>PO<sub>4</sub>, with micronutrients without molybdate and with tungstate. Control medium contained the same macronutrients and the composition of micronutrients was with molybdate and without tungstate.

### Determination of anionic concentrations

Anion and cation concentrations in nutrient solution were periodically analysed using a Dionex high-pressure liquid ionic chromatograph DX-100 (Ionpac CS-12A and AS4A-SC, Dionex Co., Sunnyvale, CA, USA). To determine intracellular ammonium concentrations, roots were pressed at 4 °C using a hydraulic press at 27 MPa; the root juice (2 ml) was centrifuged at 10 000 g for 5 min and proteins were precipitated using 10 μl of 20 mM HCl. Supernatants were collected and ammonium concentrations determined by Dionex high pressure liquid ionic chromatography DX-100 (Ionpac CS-12A, Dionex Co., Sunnyvale, CA, USA). Nitrite concentrations were measured in the culture medium by spectrophotometry at 540 nm: a volume of 600 μl of solution was diluted with 1.4 ml of distilled water and 500 μl of 0.058 M sulphanilic acid, 0.0027 M naphthyl ethylene diamine, and 1.7 M phosphoric acid were added.

### Nitrate reductase activity assay and nitrite analysis

Nitrate reductase activity was measured in fresh root tissues after 24 h and 48 h of oxygen deprivation as well as in aerated control plants. The roots were pressed at 4 °C; 600 μl of root juice were diluted with 1 ml of 0.1 M phosphate buffer pH 7.5, 0.1 M KH<sub>2</sub>PO<sub>4</sub>, 1 mM EDTA, 14 mM mercaptoethanol containing anti-protease cocktail (Sigma, L'Isle d'Abeau Chesnes, France), and 1 mM chymostatin, in the presence of 200 μl of 1.4 mM NADH and 200 μl of 0.1 M KNO<sub>3</sub> at 30 °C in the dark. After 30 min, the reaction was stopped with 100 μl of 1 M zinc acetate, 1 ml of 1% sulphanilic acid, and 1 ml of 0.2% naphthyl ethylene diamine (Sall, 1987). Nitrite concentration was measured at 540 nm. NR activity was determined by the difference between the quantity of nitrite in root extracts after 30 min of reaction and the initial quantity. The protein content of the enzyme extracts was measured according to the method of Bradford (1976).

NR activity is expressed in  $\mu\text{mol nitrite min}^{-1} \text{g}^{-1}$  of protein in root extracts.

#### Western blot and immunoprecipitation analysis

Protein was quantified both in normal aerated plant roots (controls) and in oxygen-deprived roots over a period of 24–48 h. Two grams of fresh root were ground to a fine powder in liquid nitrogen and suspended in 2 ml of extraction buffer containing a plant anti-protease cocktail (Sigma, L'Isle d'Abeau Chesnes, France) and 1 mM chymostatin to stabilize the NR. After continued grinding in a Potter homogenizer until thawed, the suspension was centrifuged for 15 min at 4 °C at 13 500 g. Supernatants containing cytosolic proteins were then collected. For western blot analysis, an equal amount of extracts (35  $\mu\text{g}$  of total protein) was boiled in denaturing SDS (0.4%) buffer for 10 min. Proteins were separated by polyacrylamide–SDS gel electrophoresis and transferred to nitrocellulose membrane (Laemmli, 1970). Total NR was detected using monoclonal antibodies against squash (*Cucurbita maxima* L. cv. Burgess Buttercup) NR (Hyde *et al.*, 1989). The immune complex was recognized by mouse IgG antibodies coupled to peroxidase. Bound proteins were detected by enhanced chemiluminescence. Quantification was performed with a Bio-Rad scanner and Molecular Analysis Software (Bio-Rad laboratories, Richmond, CA).

For immunoprecipitation assay, 2 g of fresh roots were ground to a fine powder in liquid nitrogen and suspended in 2 ml of RIPA buffer pH 7.5 (PBS, 1 $\times$ , 1% Igepal, 0.5% sodium deoxycholate, and 0.1% SDS) and the plant anti-protease cocktail. After grinding in a Potter homogenizer until thawed, the suspension was centrifuged for 20 min at 4 °C at 13 500 g. Supernatants containing cytosolic proteins were diluted to 650  $\mu\text{g ml}^{-1}$  and 1 ml was incubated for 1 h with 1  $\mu\text{g}$  of monoclonal NR antibodies (*Cucurbita maxima* L. cv. Burgess Buttercup) and then mixed with 20  $\mu\text{l}$  of mouse IgG-protein-A-agarose beads on a rotating plate for 12 h at 4 °C. Immunoprecipitates were collected by centrifugation for 20 min at 12 000 g. The pellets were washed three times with RIPA buffer and resuspended in 15  $\mu\text{l}$  electrophoresis sample buffer, boiled and resolved by 12% SDS–PAGE. pNR protein was detected using anti-phosphoserine antibody and 14-3-3 NR protein using a 14-3-3 antibody. Bound proteins were detected by enhanced chemiluminescence.

#### RNA extraction

RNAs were extracted from roots of aerated plants (controls) and from roots following 24 h and 48 h of oxygen deprivation. RNAs were also extracted from tomato and tobacco leaves. Leaves were collected in the morning and in the evening to constitute positive controls with a high NR expression and negative controls with a low NR expression. Fresh roots (2 g) were ground to a fine powder in liquid nitrogen after which 750  $\mu\text{l}$  of hot extraction buffer (65 °C) containing 4 M LiCl, 1 M TRIS–HCl pH 8, 0.5 M EDTA, and 0.7 M SDS, was added to each sample and mixed with 750  $\mu\text{l}$  of phenol saturated with 10 mM TRIS–HCl, pH 8, and 1 mM EDTA. This mixture was homogenized for 30 s then 750  $\mu\text{l}$  chloroform-isoamylalcohol (24:1, v:v) was added and the mixture vortexed. After centrifugation at 10 000 g for 5 min, the aqueous phase was mixed with 1 vol. of 4 M LiCl. RNAs were allowed to precipitate overnight at 4 °C and collected by centrifugation at 15 000 g for 10 min. The pellets were dissolved in 250  $\mu\text{l}$  water, 25  $\mu\text{l}$  3 M sodium acetate, pH 5.6, and RNAs were precipitated with 2 vols of absolute ethanol at –20 °C for 1 h. After centrifugation, the RNA pellets were washed with 70% ethanol and dried (Verwoerd *et al.*, 1989). RNAs were then analysed by northern blot analysis.

#### Preparation of cDNA probes

The NR cDNA (1.6 kbp PT13–29 NR) probe was kindly provided by Dr C Meyer (INRA, Versailles, France). cDNA inserts for hybrid-

ization were prepared by *Eco*RI digestion and gel-purified using standard protocols (Ausubel *et al.*, 1995). DNA probes were radioactively  $^{32}\text{P}$ -labelled by random primer method using the Megaprime DNA Labelling Systems kit (Amersham), as specified by the manufacturer.

#### Northern blot hybridization

Total RNA was isolated from leaf tissue as described by Verwoerd *et al.* (1989). Total RNA (15  $\mu\text{g}$ ) was separated on 1.2% (w/v) agarose gel containing 8% (v/v) formaldehyde. RNAs were transferred to Hybond N<sup>+</sup> nylon membrane (Amersham) and fixed via UV irradiation. Hybridization was performed at 65 °C in buffer containing 7% (w/v) SDS, 0.3 M sodium phosphate (pH 7.2), and 1 mM EDTA (pH 7.5) with the addition of 50 ng of denatured probes. After hybridization, the filters were washed at room temperature, twice in 2 $\times$  SSC buffer+0.1% SDS for 10 min, and once in 0.1 $\times$  SSC buffer+0.1% SDS at 65 °C for 10 min.

#### Statistics

Experiments were replicated two to eight times (see legend of figures). Results are expressed as means  $\pm$ SD. Data were analysed by one-way analysis of variance, and multiple comparison of each treatment was calculated applying the Tukey method (Keppel, 1973).


## Results

#### Measurement of nitrate reductase activity, nitrate uptake and nitrite release under root anoxia

Oxygen deprivation appeared about 10 h after the beginning of the experiment. Nitrate and nitrite concentrations were analysed in nutrient solution during the experiment (55 h). Data were calculated by the difference between initial concentration and each value of the time-course. Positive values (nitrate) corresponded to cumulated root uptake and negative values (nitrite) corresponded to cumulated release to the nutrient medium.

In aerated conditions, cumulated nitrate uptake increased time-dependently and reached higher values after 25 h (Fig. 1A). In anoxic conditions, roots continuously absorbed nitrate and cumulated uptake increased after 30 h (Fig. 1B). So, at the end of the experiment the total uptake of nitrate was 72% higher than the aerated control. Moreover, nitrite release appeared strongly in oxygen-deprived nutrient solution (Fig. 1B). In aerated nutrient solution, no nitrite was detected (Fig. 1A).

To analyse the long-term regulation of NR under asphyxia, plant material was sampled after 24 h and 48 h of root oxygen deprivation in soilless culture. To ensure reproducibility between experiments, NR activity was measured 3 h after the beginning of the light period. Actually, NR activity is maximum in the morning and then decreases at night (Jones *et al.*, 1998; Lillo *et al.*, 2001). NR activity of tissues increased regularly during the time-course of the experiment and reached its highest value 48 h after the beginning of anoxia (Fig. 2A). NR activity was 20-fold higher than the aerated control. NR enhancement was related to a strong release of nitrite in root tissues (Fig. 2B). Anoxia induced



**Fig. 1.** Time-course of nitrate consumption with (open triangles) or without (closed triangles) aeration and cumulated nitrite released to nutrient solution with (open squares) or without (closed squares) aeration over a 55 h period. Nitrate and nitrite were expressed as cumulated quantities (mmoles of ion for 1 g fresh weight of root). Positive values correspond to root consumption and negative values to cumulated nitrite release into nutrient medium. Culture medium was not renewed during the experiment. The arrow indicates the lack of oxygen in root medium. Values represent the mean  $\pm$ SD of five separate experiments.


a non-significant increase of ammonium content in roots (Fig. 2C).

#### NR regulation under long-term root anoxia

Immunoprecipitation and western blot analysis were performed to investigate whether NR protein was phosphorylated or bound to 14-3-3 protein under anoxic stress. Anoxic conditions promoted NR dephosphorylation whereas the phosphorylated form predominated in the aerated controls. Densitometric profiles of the western blot showed a decrease to 52% of phosphorylated-NR after 24 h of anoxia and to 35% after 48 h compared with the control (Fig. 3D). The analysis of 14-3-3 protein binding to NR revealed that anoxia also decreased the total amount of the 14-3-3 NR complex (Fig. 3D). The densitometric profiles of the 14-3-3 NR complex were reduced time dependently to 75% and 67% at 24 h and 48 h, respectively. Concomitantly with the study of post-translational regulation, the total quantity of NR [NR+phosphorylated-NR (pNR) plus phosphorylated-NR 14-3-3 bound (pNR 14-3-3)] was analysed with western blotting (Fig. 3C). The total amount of NR was increased by 66% at 24 h and 83% at 48 h (Fig. 3D).

#### Regulation of NR-mRNA level during root anoxia

The level of NR-mRNA expression under root anoxia was assayed by northern blot on roots and leaves using a tobacco NR probe (Fig. 4). Tobacco leaves constitute a reference to test the NR-cDNA probe in homologous hybridization conditions. NR-mRNA of tomato leaves were also assayed as a reference for NR-mRNA levels in roots. The level of NR-mRNA hybridization of the tobacco NR-labelled probe with positive (RNA of tomato and tobacco leaves collected in the morning) and negative controls (RNA of tomato and tobacco leaves collected in the evening) was observed (Fig. 4A). In these heterologous hybridization conditions, the




**Fig. 2.** Time-dependent change in NR activity (A) and nitrite and ammonium contents (B) in root tissues under anoxia. Results are presented as the means  $\pm$ SD of five separate experiments. Two asterisks indicate significant variation at  $P < 0.01$ .

signal intensity was lower in tomato (positive control) than in tobacco. However, a strong difference in NR-transcript expression in tomato leaves was observed between the evening and the morning. The results in Fig. 4B revealed that the NR-transcript level in tomato roots remained constant between 0 h and 24 h of anoxia. Then, at 48 h of anoxia, the level of NR-transcript dramatically decreased suggesting a lack of new NR synthesis.

#### NR regulation by nitrate during root anoxia

A nitrate-depleted culture medium was used to determine the role of nitrate in NR regulation during long-term root anoxia. NR activity and NR-phosphorylation were analysed in 48 h oxygen-deprived roots with or without nitrate. There was a strong increase of NR (>500%) with anoxia in the presence of nitrate (Fig. 5A). In the absence of nitrate and in the presence of oxygen, NR activity decreased to 48% compared with aerated roots in the presence of nitrate (Fig. 5A). After 48 h, the NR activity of anoxic roots




**Fig. 3.** Time-dependent effect of whole plant-root anoxia on NR regulation. Analysis of NR phosphorylation (A) and NR binding to 14-3-3 protein (B) were performed by immunoprecipitation and western blot. NR expression (NR free+NR-P+NR-14-3-3 bound) (C) was estimated by western blot. (D) Histogram of quantitative densitometric analysis (relative units) of phosphorylated NR, NR bound to 14-3-3 and total NR protein expression. Histograms represent means of three separate experiments.

deprived of nitrate was reduced to 37% versus anoxic roots in the presence of nitrate. These results support cytoplasmic NR activity that is dependent on nitrate. In the presence of nitrate, NR was strongly dephosphorylated in oxygen-deprived roots compared with the control (Fig. 5B). In the absence of nitrate, the level of phosphorylated NR was slightly decreased in oxygen-deprived roots versus control. After 48 h of anoxia, the level of NR phosphorylation was four times higher in the absence than in the presence of nitrate (Fig. 5C).

#### Morphological state of the plant in different conditions of root anoxia

To determine the role of nitrate and nitrate reductase in tolerance to root anoxia, the morphological state of plants under oxygen deprivation was analysed (i) in the presence of nitrate, (ii) in the absence of nitrate, and (iii) in the presence of tungstate in the culture media (Fig. 6).

Tungstate is assumed to replace molybdenum in the NR enzyme complex rendering it inactive. Tungstate supply prevents nitrate reduction (Notton and Hewitt, 1971). At the



**Fig. 4.** Northern blots showing the expression of NR m-RNAs in tobacco and tomato leaves collected in the morning (positive control) and in the evening (negative control) (A), and in aerated or 24 h and 48 h oxygen-deprived roots (B). Total RNA (15 µg per lane) was hybridized to NR cDNA radiolabelled probes.

same time, plants were supplied with non-toxic concentrations of ammonium in the root medium to support nitrogen assimilation. In fact, concurrent use of tungstate and ammonium appears to be an appropriate means of restricting NR activity while maintaining nitrogen assimilation (Deng *et al.*, 1989).

The changes in leaf appearance of whole plants after 24, 48, and 72 h of root anoxia were noted (Fig. 6). Firstly, the non-toxic effects of tungstate were tested: in aerated conditions, 3 d absence of nitrate (Fig. 6A) or in the presence of tungstate with ammonium (Fig. 6B), did not generate any leaf symptoms.

After 24 h of oxygen deprivation, leaves from plants without nitrate (Fig. 6D) or with tungstate (Fig. 6E) showed the early symptoms of anoxia such as wilting and corky necrotic spots. Tomato plants with nitrate supplied did not present any leaf symptoms (Fig. 6F).

After 48 h of oxygen deprivation, necrosis was more pronounced on leaves of plants grown without nitrate (Fig. 6G) or with tungstate (Fig. 6H) than on plants with nitrate (Fig. 6I). The last one showed early wilting symptoms.


At 72 h of anoxia, leaves of plants without nitrate (Fig. 6J) or with tungstate (Fig. 6K) were curled up with necrotic symptoms. With the same oxygen deprivation, leaf necrosis of plants fed with nitrate (Fig. 6L) was slight.

Nitrate reduction by NR delayed plant wilting and leaf necrosis for about 48 h.

## Discussion


### Nitrate reductase activity in oxygen-deprived roots

In anoxic conditions only, nitrite appeared in the nutrient solution about 10 h after the beginning of the experiment



**Fig. 5.** (A) NR activity measured with nitrate in aerated conditions (CN) and after 48 h anoxic conditions (48N). NR activity measured without nitrate in aerated conditions (C) and after 48 h in anoxic conditions (48). Histograms represent mean  $\pm$ SD of three separate experiments, two asterisks indicate a significant increase of NR activity at  $P < 0.01$ . (B) Analysis of NR phosphorylation and NR expression in plants treated as described above. Each western blot is representative of three separate experiments. (C) Histogram of quantitative densitometric analysis (relative units).

(Fig. 1B). This phenomenon is related to the lack of oxygen as a result of root respiration (Morard *et al.*, 2000). Nitrate uptake was enhanced compared with the aerated control. These results suggest that NR activity was stimulated by anoxic conditions. NR activity was therefore measured in root juice pressed from whole tomato roots. The activity of nitrate reductase increased time-dependently under anoxic



**Fig. 6.** Morphological changes of tomato leaves in aerated (A–C) and oxygen-deprived (D–L) roots in different conditions. (A, D, G, J) Plants cultivated without nitrate; (B, E, H, K) plants cultivated with nitrate and tungstate supplied with ammonium; (C, F, I, L) plants cultivated with nitrate. Each photograph is representative of three separate experiments.

conditions (Fig. 2A). Concomitantly, nitrite content increased in root tissues (Fig. 2B), but not to a great extent because, in roots, nitrite is rapidly released into the nutrient medium (Fig. 1A). A constant root ammonium concentration was observed during the time-course of anoxia (Fig. 2C). These results suggest that, under anoxia, nitrite was not highly accumulated into amino-acid synthesis by the roots. Nitrite release by anaerobic tissues is not a new finding. Ferrari *et al.* (1973) and Botrel *et al.* (1996) also noticed this phenomenon. Nevertheless, the role of nitrite release by oxygen-deprived plants remains unclear. To gain an insight into the role of nitrate reduction and nitrite leakage, NR regulation must be investigated.

#### Regulation of nitrate reductase in anoxic conditions

Normally, NR is inactivated by both phosphorylation and binding to a 14-3-3 protein inhibitor (Weiner and Kaiser, 1999, 2000). However, it has been suggested that in post-translational modulation of NR, phosphorylated NR also constitutes an active form of the protein (Kaiser and Huber, 2001). These results indicate that anoxia induced NR activation by both dissociation with the 14-3-3 protein inhibitor and dephosphorylation of the NR protein (Fig. 3). After 48 h of anoxia, the respective decreases in NR phosphorylation and binding to the 14-3-3 protein inhibitor of the enzyme were very similar (26% and 36%, respectively).

These data suggest a relationship between NR phosphorylation and binding to the 14-3-3 protein inhibitor and, on the other hand, NR enzymatic activity under anoxia. The data imply that *in vivo*, in whole plant roots, the active form of the NR induced by anoxia is the native form.

Concomitantly with NR regulation, the expression of NR-mRNA was analysed by northern blots at 24 h and 48 h of root anoxia (Fig. 4). Data showed that anoxia did not increase NR-mRNA synthesis. Surprisingly, NR-mRNA was very low at 48 h of root anoxia, indicating that oxygen-deprivation strongly disturbed root metabolism. During anoxia, a general decrease in mRNA translation and an activation of the expression of anoxic genes producing anaerobically induced proteins (ANPs) has been observed (Sachs *et al.*, 1980, 1996). Normal protein synthesis was inhibited under anoxia and 10–20 ANPs appear constituting 70% of the total. The majority of the genes induced encode for enzymes involved in starch, glucose mobilization, glycolysis, and fermentation. The anaerobic response element (ARE) regulates some of these genes (Walker *et al.*, 1987). In maize, hypoxic stress rapidly induces the expression of these genes within four h. More recently Huq and Hodges (1999) identified a novel gene family called anaerobically inducible early (AIE) in a very flood-tolerant variety of rice. Tomato is not particularly adapted to long-term root anoxia. Tomato plants die under a prolonged period of root oxygen deprivation unlike tolerant species (Morard and Silvestre, 1996). However, under this study's experimental conditions, tomato was able to tolerate a prolonged period of root anoxia for up to four days. It was assumed that rapid induction of AIE genes did not occur in tomato and that NR did not constitute an ANP gene product. Nevertheless, although NR-mRNA synthesis decreased at 48 h of root anoxia, the total amount of NR remained constant between 24 h and 48 h and NR activity strongly increased. It is known that the NR association with the 14-3-3 protein inhibitor is necessary for its degradation (Kaiser and Huber, 2001). These results demonstrate that the NR–14-3-3 complex is gradually dissociated during anoxia (Fig. 3). Several factors could explain NR dissociation with the 14-3-3 protein inhibitor under anoxia. Kaiser and Brendle-Behnisch (1995) demonstrated that cellular acidification prevented its binding with NR. Under anoxia, other studies described cellular acidification (Roberts *et al.*, 1984). Both events argue for an arrest of NR degradation under anoxia. Overall, these data suggest a strong inhibition of NR degradation within 48 h and this could constitute a possible adaptive response.

Following this conclusion, the effect of nitrate on NR regulation was assayed in the presence or absence of oxygen. The results of Fig. 1 pointed out that nitrate uptake was strongly increased in oxygen-deprived roots compared with aerated roots. In this study's conditions, NR activity was maximum when nitrate was provided in the culture medium, independently of the presence of oxygen (Fig. 5).

Nevertheless, in the absence of nitrate, NR activity was much higher in oxygen-deprived than in aerated roots. Tomato plants can, therefore, tolerate nitrate shortage, but NR activity is then limited. In the absence of nitrate in the culture medium, the nitrate in cell vacuoles remains the only, although not easily available, pool for nitrate reductase activity. The nitrite content in aerated tomato roots ( $0.08 \pm 0.001 \mu\text{mol g}^{-1}$  root FW) was analogous with nitrite content in nitrate-depleted aerated and anoxic roots ( $0.1 \pm 0.01$  and  $0.12 \pm 0.05 \mu\text{mol g}^{-1}$  root FW, respectively). So, nitrate is not a limiting factor for nitrogen assimilation during the time-course of the experiment (48 h). In these conditions, the high levels of phosphorylated-NR observed suggest that nitrate plays a key role in NR post-translational regulation.

#### *Role of nitrate supplied in anoxia tolerance*

According to previous agronomic observations (Arnon, 1937; Malavolta, 1954; Guyot and Prioul, 1985), nitrate supplied to roots enhanced the plant's tolerance to anoxia. Confirmation of these findings was studied with observation of the aerial parts during tomato root anoxia, particularly the visual changes on leaves over the period from 24 h to 72 h (Fig. 6). Nitrogen nutrition can be changed in two different ways, either deprivation of nitrate supplied or blocking of NR activity. These results clearly demonstrated that the absence of nitrate accelerated the appearance of anoxic symptoms over 24–48 h (Fig. 6D, G). Likewise, when NR activity was blocked by tungstate, anoxic symptoms appeared 24–48 h earlier in anoxic plants in the presence of nitrate (Fig. 6E, H). In oxygen-deprived plants with nitrate, no symptoms appeared 24–48 h after the beginning of anoxia (Fig. 6F, I). Symptoms appeared only after 72 h (Fig. 6L). Overall, these results indicated that plant tolerance to anoxia was dramatically decreased by the absence of nitrate reduction to nitrite. Moreover, visual symptoms indicate that root anoxia is highly stressful for the aerial part of tomato plants. The principal effects of anoxia on leaves are first, stomatal closure, then epinastic curvature and wilting, and finally corky necrotic spots. Morphological changes on leaves are linked with a signal transduction pathway from roots for aerial plant parts. This phenomenon occurs under the influence of hormone signals and is related to the change of ethylene concentration interacting with gibberellins (Musgrave *et al.*, 1972), auxins (Cookson and Osborne, 1978) and cytokinins (Takei *et al.*, 2002). Stomate closing was promoted by an increase of abscisic acid in leaves (Zhang and Davies, 1987) and related to decreased water uptake (Bradford and Hsiao, 1982). In anoxia, hydraulic conductivity of root was also inhibited by gating of aquaporins, related to cytosolic acidosis (Tournaire-Roux *et al.*, 2004).

On root tissues, the first consequence of anoxia is a lowering and then a stopping of aerobic respiration. The



mitochondrial electron-transport chain is also stopped and hence the synthesis of ATP is inhibited (Blokhina, 2000). In these conditions, fermentative pathways such as lactic and ethanolic fermentation are promoted and affect cell energy charge and NADH/NAD<sup>+</sup> ratio (Chrikova and Belonogova, 1991). Too low an energy charge leads to the death of root cells by the degradation of the mitochondrial ultrastructure and this was observed in pumpkin roots after 10–15 h of anoxia (Vartapetian *et al.*, 2003). This phenomenon cannot be reversed by reaeration. Mitochondrial ultrastructure can be preserved by the hydrolysis of starch reserves and the activation of fermentation enzymes that limit the effects of anoxia. Moreover, ADH activity could be enhanced by the appearance of oxidative stress due to NADP(H) activation when oxygen disappears (Baxter-Burrell *et al.*, 2002). In these experimental conditions, root peroxidase activity, expressed in mM oxidized guaiacol  $\mu\text{g}^{-1}$  protein  $\text{min}^{-1}$ , was enhanced time-dependently from  $8.3 \pm 3.3$  to  $11 \pm 3.3$  after 24 h anoxia and to  $45 \pm 10.6$  after 72 h anoxia.

One of the major consequences of the reduction of aerobic respiration is the ATP shortage for membrane ATPases, resulting in the depolarization of the root cell membrane (Buwalda *et al.*, 1988). This cell phenomenon induced a fast K<sup>+</sup> efflux out of the roots of cucumber (Bertoni *et al.*, 1993) and of tomato (Morard *et al.*, 2000) in the same experimental conditions. Likewise, the fermentative pathway and passive leakage of H<sup>+</sup> and anions from the vacuole into the cytoplasm lead to an acidification of cell cytoplasm reducing tolerance to anoxia (Roberts *et al.*, 1984; Kaiser and Huber, 2001; Drew and Armstrong, 2002). However, in anoxic conditions, H<sup>+</sup> extrusion by maize roots increases when the cytoplasmic pH falls (Xia and Roberts, 1996), regulation of cytoplasmic pH, limiting cell acidosis, enhanced root cell survival (Drew, 1997). Moreover, cytosolic acidification promotes nitrate reductase activity, resulting in nitrite release out of root cells (Stoimenova *et al.*, 2003). One hypothesis is that nitrate reduction could prevent a pH fall by the extrusion of H<sup>+</sup> as H<sup>+</sup>NO<sub>2</sub><sup>-</sup> like nitrite transport as described in the chloroplast (Shingles *et al.*, 1996). In accordance with work on pH, these results showed a strong nitrite release over 10 h and a pH decrease from  $6 \pm 0.08$  to  $5.4 \pm 0.1$  was recorded in the nutrient solution after 48 h of anoxia.

In conclusion, these studies demonstrate that (i) root anoxia induces activation rather than new synthesis of NR and (ii) nitrate reduction improves plants' resistance to anoxia. Likewise, under oxygen deprivation, NR displayed functions other than nitrogen assimilation: it enabled NAD<sup>+</sup> regeneration and prevented the pH from dropping to life-threatening levels. To confirm these hypotheses, further studies under anoxia are necessary, in particular, investigations into the effects of the presence or absence of nitrate on the NADH/NAD<sup>+</sup> ratio and on intracellular pH variations.

## Acknowledgements

The authors thank Professor WM Kaiser, Institute of Physiology, Molecular Biology and Biophysics, Würzburg, Germany, for the generous gift of the 14-3-3 protein antibody. We also thank Dr C Meyer INRA, France, for the kind gift of cDNA NR.

## References

- Appenroth KJ, Meço R, Jourdan V, Lillo C. 2000. Phytochrome and post-translational regulation of nitrate reductase in higher plants. *Plant Science* **159**, 51–56.
- Arnon DI. 1937. Ammonium and nitrate nitrogen nutrition of barley and rice at different seasons in relation to hydrogen-ion concentrations, manganese, copper and oxygen supplied. *Soil Science* **44**, 91–121.
- Aslam M, Oaks A, Huffaker RC. 1976. Effect of light and glucose on the induction of nitrate reductase and on the distribution of nitrate in etiolated barley leaves. *Plant Physiology* **58**, 588–591.
- Aslam M, Travis RL, Rains DW. 2001. Enhancement of nitrate reductase activity and metabolic nitrate concentration by methionine sulfoximine in barley roots. *Plant Science* **161**, 133–142.
- Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K. 1995. *Current protocols in molecular biology*. New York: John Wiley.
- Baxter-Burrell A, Yang Z, Springer PS, Bailey-Serres J. 2002. RopGAP4-dependent Rop GTPase rheostat control of *Arabidopsis* oxygen deprivation tolerance. *Science* **296**, 2026–2028.
- Bertoni G, Silvestre J, Llorens JM, Morard P, Maertens C. 1993. Effect of a transient anoxia on potassium uptake in cucumber. In: Frago M, Van Beusichem M, eds. *Optimization of plant nutrition*. Dordrecht, The Netherlands: Kluwer Academic Publishers, 203–206.
- Blokhina OB. 2000. Anoxia and oxidative stress: lipid peroxidation, antioxidant status and mitochondrial functions in plants. PhD thesis, Helsinki University, Finland.
- Botrel A, Magné C, Kaiser WM. 1996. Nitrate reduction, nitrite reduction and ammonium assimilation in barley roots in response to anoxia. *Plant Physiology and Biochemistry* **34**, 645–652.
- Bradford KJ, Hsiao TC. 1982. Stomatal behavior and water relations of waterlogged of tomato plants. *Plant Physiology* **70**, 1508–1513.
- Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Analytical Biochemistry* **72**, 248–254.
- Buwalda KJ, Thomson CJ, Steiger W, Barret-Lennard EG, Gibbs J, Greenway H. 1988. Hypoxia induces membrane depolarization and potassium loss from wheat roots, but does not increase their permeability to sorbitol. *Journal of Experimental Botany* **39**, 1169–1183.
- Campbell WH. 2001. Structure and function of eukaryotic NAD(P)H: nitrate reductase. *Cell and Molecular Life Sciences* **58**, 194–204.
- Chrikova TV, Belonogova VA. 1991. Nitrate reductase activity and productivity of grain crops under waterlogged conditions. *Soviet Soil Science* **22**, 19–29.
- Chung HL, Sehnke PC, Ferl RJ. 1999. The 14-3-3 proteins: cellular regulators of plant metabolism. *Trends in Plant Science* **4**, 367–371.
- Cookson C, Osborne DJ. 1978. The stimulation of cell extension by ethylene and auxins in aquatic plants. *Planta* **144**, 39–47.
- Datta R, Sharma R. 1999. Temporal and spatial regulation of nitrate reductase and nitrite reductase in greening maize leaves. *Plant Science* **144**, 77–83.

- De la Haba P, Agüera E, Benitez L, Maldonado JM.** 2001. Modulation of nitrate reductase activity in cucumber (*Cucumis sativus*) roots. *Plant Science* **161**, 231–237.
- Deng M-D, Lamaze T, Morot-Gaudry JF.** 1989. A new experimental approach involving the simultaneous use of tungstate and ammonium for studying the physiological effects of the absence of nitrate reduction. *Plant Physiology and Biochemistry* **27**, 689–696.
- Drew MC.** 1997. Oxygen deficiency and root metabolism: injury and acclimation under hypoxia and anoxia. *Annual Review of Plant Physiology and Plant Molecular Biology* **48**, 223–250.
- Drew MC, Armstrong W.** 2002. Root growth and metabolism under oxygen deficiency. In: Waisel Y, Kafkafi U, Ezkel A, eds. *Plant root growth, the hidden half*. New York, Basel: Marcel Dekker, 1120.
- Ellis MH, Dennis ES, Peacock WJ.** 1999. *Arabidopsis* roots and shoots have different mechanisms for hypoxic stress tolerance. *Plant Physiology* **119**, 57–64.
- Ferrari TE, Yoder OC, Filner P.** 1973. Anaerobic nitrite production by plant cells and tissues: evidence for two nitrate pools. *Plant Physiology* **51**, 423–431.
- Glaab J, Kaiser WM.** 1993. Rapid modulation of nitrate reductase in pea roots. *Planta* **191**, 173–179.
- Guyot C, Prioul JL.** 1985. Correction par la fertilization minérale des effets de l'ennoyage sur le blé d'hiver. I. Expérimentation sur sol. *Agronomie* **5**, 743–750.
- Heller R, Esnault R, Lance C.** 1998. *Physiologie végétale: nutrition*, Vol. 1, 6th edn. Paris: Dunod.
- Huber SC, Bachmann M, Huber JL.** 1996. Post-translational regulation of nitrate reductase activity: a role for Ca<sup>2+</sup> and 14-3-3 proteins. *Trends in Plant Science* **1**, 432–438.
- Huq E, Hodges TK.** 1999. An anaerobically inducible early (aie) gene family from rice. *Plant Molecular Biology* **40**, 591–601.
- Hyde GE, Wilberding JA, Meyer AL, Campbell ER, Campbell WH.** 1989. Monoclonal antibody-based immunoaffinity chromatography for purifying corn and squash NADH: nitrate reductases. Evidence for an interchain disulfide bond in nitrate reductase. *Plant Molecular Biology* **13**, 233–246.
- Jones TL, Tucker DE, Ort DR.** 1998. Chilling delays circadian pattern of sucrose phosphate synthase and nitrate reductase activity in tomato. *Plant Physiology* **118**, 149–158.
- Kaiser WM, Brendle-Behnisch E.** 1995. Acid-base of nitrate reductase in leaf tissues. *Planta* **196**, 1–6.
- Kaiser WM, Huber SC.** 2001. Post-translational regulation of nitrate reductase: mechanism, physiological relevance and environmental triggers. *Journal of Experimental Botany* **52**, 1981–1989.
- Kanamaru K, Wang R, Su W, Crawford NM.** 1999. Ser-543 in the hinge-1 region of *Arabidopsis*: nitrate reductase is conditionally required for binding 14-3-3 proteins and *in vitro* inhibition. *Journal of Biological Chemistry* **274**, 4160–4165.
- Keppel G.** 1973. *Design and analysis: a researcher's handbook*. Englewood Cliffs, NJ: Prentice Hall, 658–663.
- Laemmli UK.** 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Lillo C, Meyer C, Ruoff P.** 2001. The nitrate reductase circadian system. The central clock dogma contra multiple oscillatory feedback loops. *Plant Physiology* **125**, 1554–1557.
- Malavolta E.** 1954. Studies on the nitrogenous nutrition of rice. *Plant Physiology* **29**, 98–99.
- Martinoia E, Heck U, Wiemken A.** 1981. Vacuoles as storage compartments for nitrate in barley leaves. *Nature* **289**, 292–294.
- Morard P, Silvestre J.** 1996. Plant injury due to oxygen deficiency in the root environment of soilless culture: a review. *Plant and Soil* **184**, 243–254.
- Morard P, Lacoste L, Silvestre J.** 2000. The effect of oxygen deficiency on the uptake of water and mineral nutrients by tomato plants in soilless culture. *Journal of Plant Nutrition* **23**, 1063–1078.
- Morard P, Caumes E, Silvestre J.** 2004a. Influence de la concentration de la solution nutritive sur la croissance et la nutrition minérale de la tomate. *Canadian Journal of Plant Science* **84**, 299–304.
- Morard P, Lacoste L, Silvestre J.** 2004b. Effect of oxygen deficiency on mineral nutrition of excised tomato roots. *Journal of Plant Nutrition* **27**, 611–624.
- Morard P, Silvestre J, Lacoste L, Caumes E, Lamaze T.** 2004c. Nitrate reduction and nitrite release by tomato roots in response to anoxia. *Journal of Plant Physiology* **161**, 855–866.
- Musgrave A, Jackson MB, Ling E.** 1972. *Callitriche* stem elongation is controlled by ethylene and gibberellin. *Nature New Biology* **236**, 93–96.
- Notton BA, Hewitt EJ.** 1971. The role of tungsten in the inhibition of nitrate reductase activity in spinach leaves. *Biochemical and Biophysics Research Communication* **44**, 702–710.
- Paul AL, Schuerger AC, Popp MP, Richards JT, Manak MS, Ferl RJ.** 2004. Hypobaric biology: *Arabidopsis* gene expression at low atmospheric pressure. *Plant Physiology* **134**, 215–223.
- Reggiani R, Brambilla I, Bertani A.** 1985. Effect of exogenous nitrate on anaerobic metabolism in excised rice root. *Journal of Experimental Botany* **36**, 1698–1704.
- Roberts JKM, Jardetzky O, Walbot V, Freeling M.** 1984. Cytoplasmic acidosis as a determinant of flooding intolerance in plants. *Proceedings of the National Academy of Sciences, USA* **81**, 6029–6033.
- Sachs MM, Freeling M, Okimoto R.** 1980. The anaerobic proteins of maize. *Cell* **20**, 761–767.
- Sachs MM, Subbaiah CC, Saab IN.** 1996. Anaerobic gene expression and flooding tolerance in maize. *Journal of Experimental Botany* **47**, 1–15.
- Sall K.** 1987. Influence du déficit hydrique sur les activités nitrate reductase et nitrogénase chez le soja. PhD thesis, Toulouse University, France.
- Shingles R, Roh MH, McCarty RE.** 1996. Nitrite transport in chloroplast inner envelope vesicles (direct measurement of proton-linked transport). *Plant Physiology* **112**, 1375–1381.
- Stoimenova M, Hänsch R, Mendel R, Gimmler H, Kaiser WM.** 2003. The role of nitrate reduction in the anoxic metabolism of roots. I. Characterization of root morphology and normoxic metabolism of wild type tobacco and a transformant lacking root nitrate reductase. *Plant and Soil* **253**, 145–153.
- Subbaiah CC, Sachs MM.** 2003. Molecular and cellular adaptations of maize to flooding stress. *Annals of Botany* **91**, 119–127.
- Takei K, Takahashi T, Sugiyama T, Yamaya T, Sakakibara H.** 2002. Multiple routes communicating nitrogen availability from roots to shoots: a signal transduction pathway mediated by cytokinin. *Journal of Experimental Botany* **53**, 971–977.
- Tournaire-Roux C, Suktra M, Javot H, Gout E, Gerbeau P, Luu DT, Bligny R, Maurel C.** 2004. Cytosolic pH regulates root water transport during anoxic stress through gating of aquaporins. *Nature* **425**, 393–397.
- Vartapetian BB, Andreeva IN, Generozova IP, Polyakova LI, Maslova IP, Dolgikh YI, Stepanova AY.** 2003. Review: functional electron microscopy in studies of plant response and adaptation to anaerobic stress. *Annals of Botany* **91**, 155–172.
- Verwoerd TC, Dekker BMM, Hoekema A.** 1989. A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Research* **17**, 2362.
- Walker JC, Howard EA, Dennis ES, Peacock WJ.** 1987. DNA sequences required for anaerobic expression of the maize *alcohol dehydrogenase-1* gene. *Proceedings of the National Academy of Sciences, USA* **84**, 6624–6628.
- Weiner H, Kaiser WM.** 1999. 14-3-3 proteins control proteolysis of nitrate reductase in spinach leaves. *FEBS Letters* **455**, 75–78.

- Weiner H, Kaiser WM.** 2000. Binding to 14-3-3 protein is not sufficient to inhibit nitrate reductase in spinach leaves. *FEBS Letters* **480**, 217–220.
- Weiner H, Kaiser WM.** 2001. Antibodies to assess phosphorylation of spinach leaf nitrate reductase on serine 543 and its binding to 14-3-3 proteins. *Journal of Experimental Botany* **52**, 1165–1172.
- Xia JH, Saglio P, Roberts JKM.** 1995. Nucleotide levels do not critically determine survival of maize root tips acclimated to a low-oxygen environment. *Plant Physiology* **108**, 589–595.
- Xia JH, Roberts JKM.** 1996. Regulation of H<sup>+</sup> extrusion and cytoplasmic pH in maize root tips acclimated to a low-oxygen environment. *Plant Physiology* **111**, 227–233.
- Zhang J, Davies WJ.** 1987. ABA in root and leaves of flooded pea plants. *Journal of Experimental Botany* **38**, 649–659.
- Zhen RG, Koyro HW, Leigh RA, Tomos AD, Miller AJ.** 1991. Compartmental nitrate concentrations in barley root cells measured with nitrate-selective microelectrodes and by single-cell sap sampling. *Planta* **185**, 356–361.