



Progress in Photosynthesis Research

Volume 1

*Proceedings of the VIIth International Congress on Photosynthesis
Providence, Rhode Island, USA, August 10–15, 1986*

edited by

J. BIGGINS

*Division of Biology and Medicine, Brown University
Providence, RI 02912, USA*

1987 **MARTINUS NIJHOFF PUBLISHERS**
a member of the KLUWER ACADEMIC PUBLISHERS GROUP
DORDRECHT / BOSTON / LANCASTER



GENERAL CONTENTS

Volume I

1. Excitation Energy Transfer	1
2. Primary Electron Transfer Reactions	151
3. Chlorophylls and Model Systems	297
4. Structure of Molecular Complexes: Crystallographic and Physical Studies	353
5. Oxygen Evolution	453

Volume II

1. Components and Pigment Protein Complexes a) b) c) d)	1
2. Stoichiometry of Photosystem I and Photosystem II	233
3. Lateral Heterogeneity and Stacking	269
4. Localization of Membrane Components	293
5. Effects of Environmental Changes and Growth Conditions	339
6. Quinone Acceptors	387
7. Cytochromes (b-f) Complex	441
8. Lateral Electron Transport, Connectivity Between Photosystems	501
9. Plastocyanin	521
10. PSI-cyclic Electron Transport	537
11. Inhibition and Environmental Effects	553
12. General Aspects	613
13. Higher Plants	669
14. Organisms Containing Phycobilins	757
15. Membrane Protein Kinases	797

VI

Volume III

1. Proton ATP-ase	1
2. Electrochemical Proton Gradients and ATP Synthesis	127
3. Light-dark Regulation of Carbon Metabolism	233
4. Metabolite Regulation of Carbon Metabolism	273
5. Ribulose 1,5 Bisphosphate Carboxylase Oxygenase	371
6. Other Chloroplast Enzymes	435
7. Metabolism of C ₄ and CAM Plants	503
8. Integration of Carbon and Nitrogen Metabolism	535
9. Photorespiration	549
10. Carbon partitioning	675
11. Herbicide Action	763

Volume IV

1. Photoinhibition	1
2. Temperature	99
3. Water Potentials	147
4. Salinity and Nutrient Deficiency	185
5. Regulation of Gas Exchange	209
6. Mechanisms of CO ₂ Concentration	289
7. Crop Productivity	361
8. Biosynthesis of Photosynthetic Pigments	423
9. Photoregulation of Chloroplast Development	511
10. Chloroplast Molecular Genetics	617
11. Photosynthetic Bacteria	691
12. Cyanobacteria	749

CONTENTS TO VOLUME I

General Contents	V
Contents Volume I	VII
Preface	XXI
Acknowledgments	XXIII
Local Organizing Committee	XXV
1. Excitation Energy Transfer	
Picosecond Absorption and Fluorescence Spectroscopy of Energy Transfer and Trapping in Photosynthetic Bacteria R. van Grondelle	1
Excitation Energy Transport in the Antenna Systems of Purple Bacteria, Studied by Low-intensity Picosecond Absorption Spectroscopy V. Sundström, R. van Grondelle, H. Bergström, E. Åkesson, T. Gillbro	9
The Organization of the Light Harvesting Antenna of Purple Bacteria M. Vos, R.J. van Dorssen, R. van Grondelle, C.N. Hunter, J. Amesz, L.N.M. Duysens	13
Photochemical and Non-photochemical Holeburning Studies of Energy and Electron Transfer in Photosynthetic Reaction Centers and Model Systems Steven G. Boxer, Thomas R. Middendorf, David J. Lockhart, David S. Gottfried	17
The Temperature Dependence of Electron Back-transfer from the Primary Radical Pair of Bacterial Photosynthesis David E. Budil, Stephen V. Kolaczkowski, James R. Norris	25
Supramolecular Organisation of Light-harvesting Pigment-protein Complexes of <i>Rhodobacter Sphaeroides</i> Studied by Excitation Energy Transfer and Singlet-singlet Annihilation at Low Temperature in Phospholipid-enriched Membranes Willem H.J. Westerhuis, Marcel Vos, Rob J. van Dorssen, Rienk van Grondelle, Jan Amesz, Robert A. Niederman	29

VIII

Correlation between the Efficiency of Energy Transfer and the Polyene Chain Structure of Carotenoids in Purple Photosynthetic Bacteria	33
H. Hayashi, K. Iwata, T. Noguchi, M. Tasumi	
Triplet Energy Transfer between Photosynthetic Pigments: An ESR Study of B800-850 Light-harvesting Complexes and Synthetic Carotenoporhyrin Molecules	37
Harry A. Frank, Barry W. Chadwick, Chaoying Zhang, Jung Jin Oh	
Picosecond Excitation Energy Transfer between Different Light-harvesting Complexes and Reaction Centres in Purple Bacteria	41
V.I. Godik, A. Freiberg, K. Timpmann, A.Yu. Borisov, K.K. Rebane	
Spectral Dependence of the Fluorescence Lifetime of <i>Rhodospirillum Rubrum</i> . Evidence for Inhomogeneity of B880 Absorption Band	45
A. Freiberg, V.I. Godik, K. Timpmann	
Protein Phosphorylation: A Mechanism for Control of Excitation Energy Distribution in Purple Photosynthetic Bacteria	49
Nigel G. Holmes, John F. Allen	
A Model for the Functional Antenna Organization and Energy Distribution in the Photosynthetic Apparatus of Higher Plants and Green Algae	53
Alfred R. Holzwarth	
Picosecond Transient Absorbance Spectra and Fluorescence Decay Kinetics in Photosystem II Particles	61
A.R. Holzwarth, H. Brock, G.H. Schatz	
Picosecond Time Resolved Chlorophyll Fluorescence Spectra from Pea Chloroplast Thylakoids	67
G.H. Schatz, A.R. Holzwarth	
Picosecond Fluorescence Spectra of Synchronous Cultures of the Green Alga <i>Scenedesmus Obliquus</i>	71
E. Bittersmann, H. Senger, A.R. Holzwarth	
Measurements and Kinetic Modeling of Picosecond Time-resolved Fluorescence from Photosystem I and Chloroplasts	75
Bruce P. Wittmershaus	
Time-resolved Fluorescence Decay Kinetics in Photosystem I. Experimental Estimates of Charge Separation and Energy Transfer Rates	83
T.G. Owens, S.P. Webb, D.D. Eads, R.S. Alberte, L. Mets, G.R. Fleming	
Spectral Properties of Photosystem I Fluorescence at Low Temperatures	87
J. Wachtveitl, H. Krause	

Analysis of Pigment System I Chl <i>a</i> Fluorescence at Room Temperature by the Steady State Spectrum and the Time Resolved-spectrum in Picosecond Time Range	91
Mamoru Mimuro, Iwao Yamazaki, Naoto Tamai, Tomoko Yamazaki, Yoshihiko Fujita	
Spectral Shifts in Picosecond Transient Absorption Spectra Due to Stimulated Emission from Chlorophyll <i>in vitro</i> and in Protein Complexes	95
D.R. Klug, B.L. Gore, L.B. Giorgi, G. Porter	
Fast Fluorescence and Absorption Measurements of Photosystem I from a Cyanobacterium	99
E. Hilary Evans, Raymond Sparrow, Robert G. Brown, David Shaw, John Barr, Martin Smith and William Toner	
Anomalous Fluorescence Induction on Subnanosecond Time Scales and Exciton-exciton Annihilations in PSII	103
A. Dobek, J. Deprez, N.E. Geacintov, J. Breton	
Laser Flash-induced Non-sigmoidal Fluorescence Induction Curves in Chloroplasts	107
Nicholas E. Geacintov, Jacques Breton, Lee France, Jean Deprez, Andrzej Dobek	
Is Variable Fluorescence Due to Charge Recombination?	111
I. Moya, M. Hodges, J-M. Briantais	
Time Resolved Chlorophyll Fluorescence Studies of Photosynthetic Pigment Protein Complexes: Characterisation of Five Kinetic Components	115
M. Hodges, I. Moya, J-M. Briantais, R. Remy	
Multivariate Analysis of Photosystem II Chlorophyll Fluorescence Quenching by Quinones	119
K.K. Karukstis, S.C. Boegeman, S.M. Gruber, C.R. Monell, J.A. Fruetel, M.H. Terris	
Energy Transfer in Chlorophyll Antennae of Isolated PSII Particles	123
Tomas Gillbro, Åke Sandström, Villy Sundström, Michael Spangfort, Bertil Andersson, Göran Lagenfelt	
Polarized Spectra of PS2 Particles in PVA Films	127
D. Frackowiak, W. Hendrich, M. Romanowski, A. Szczepaniak, R.M. Leblanc	
The Dependence of the Energy Transfer Kinetics of the Higher Plant Light Harvesting Chlorophyll-protein Complex on Chlorophyll/Detergent Resolubilisation Ratios	131
J.P. Ide, D.R. Klug, B. Crystall, B.L. Gore, L.B. Giorgi, W. Kuhlbrandt, J. Barber, G. Porter	

X

Characterization of the Fluorescence Decays of the Chlorophyll <i>a/b</i> Protein D.D. Eads, S.P. Webb, T.G. Owens, L. Mets, R.S. Alberte, G.R. Fleming	135
Fluorescence Decay and Depolarization Kinetics Calculated Using Förster Inductive Resonance and the Molecular Coordinates for C-phycoyanin Kenneth Sauer, Hugo Scheer	139
Photochemistry and Photophysics of C-phycoyanin Hugo Scheer	143
2. Primary Electron Transfer Reactions	
Primary Reactions of Photosynthesis: Discussion of Current Issues Paul Mathis	151
Selective Reduction and Modification of Bacteriochlorophylls and Bacteriopheophytins in Reaction Centers from <i>Rhodospseudomonas Viridis</i> V.A. Shuvalov, A.Ya. Shkuropatov, M.A. Ismailov	161
Spectroscopic and Primary Photochemical Properties of Modified <i>Rhodospseudomonas Sphaeroides</i> Reaction Centers Dewey Holten, Christine Kirmaier, Leanna Levine	169
Fourier Transform Infrared (FTIR) Spectroscopic Investigations of the Primary Reactions in Purple Photosynthetic Bacteria E. Nabedryk, B.A. Tavitian, W. Mäntele, W. Kreutz, J. Breton	177
Picosecond Characterization of Primary Events in <i>Rhodospseudomonas Viridis</i> Whole Cells by Transmembrane Potential Measurements J. Deprez, H.-W. Trissl, J. Breton	181
Excitation of Antenna Pigments and Electron Transfer upon Picosecond Flash Illumination of Membranes of <i>Chloroflexus Aurantiacus</i> A.M. Nuijs, H. Vasmel, L.N.M. Duysens, J. Amesz	185
Electron Transport in <i>Heliobacterium Chlorum</i> H.W.J. Smit, J. Amesz, M.F.R. van der Hoeven, L.N.M. Duysens	189
A Possible Mechanism for Electron Transfer in the Diquinone Acceptor Complex of Photosynthetic Reaction Centers S.K. Buchanan, K. Ferris, G.C. Dismukes	193
Triplet-minus-singlet Absorption Difference Spectra of Some Bacterial Photosynthetic Reaction Centers with and without Carotenoids Recorded by Magneto-optical Difference Spectroscopy (MODS) at 290 and 20 K E.J. Lous, A.J. Hoff	197

An E.P.R. Signal Arising from Q_B^- -Fe in <i>Chromatium Vinosum</i> Strain D P. Heathcote, A.W. Rutherford	201
Photochemical Reduction of either of the Two Bacteriopheophytins in Bacterial Photosynthetic Reaction Centers Sandra Florin, David M. Tiede	205
Reconstitution of Reaction Centers in Planar Bilayer Lipid Membranes (BLM) H. Ti Tien	209
$^3(P^+I^-)$ Lifetime as Measured by B_1 Field Dependent RYDMR Triplet Yield Stephen Kolaczowski, David Budil, James R. Norris	213
Electron Transfer in Reaction Center Protein from <i>R. Sphaeroides</i> : Generation of a Spin Polarized Bacterio-chlorophyll Dimer EPR Signal Whose Formation is Modulated by the Electron Transfer Rate from Bacteriopheophytin to Q_A M.R. Gunner, D.E. Robertson, R.L. LoBrutto, A.C. McLaughlin, P.L. Dutton	217
Electric Field Dependence of Electron Transfer in Photosynthetic Reaction Centers from <i>Rhodospseudomonas Sphaeroides</i> G.A. Alegria, P.L. Dutton	221
Hydrocarbon Tail Structure and its Effect on the Affinity and Kinetic Performance of Quinones at the Q_A Site in Reaction Centers of <i>Rhodobacter Sphaeroides</i> R26 K. Warncke, M.R. Gunner, B.S. Braun, C.-A. Yu, P.L. Dutton	225
Excited States and Primary Photochemical Reactions in Photosystem I A.M. Nuijs, V.A. Shuvalov, H.W.J. Smit, H.J. van Gorkom, L.N.M. Duysens	229
Characterization of the Electron Acceptor A_1 in Photosystem I by Flash-absorption Spectroscopy at Low Temperature: Evidence that A_1 is Vitamin K_1 K. Brettel, P. Setif, P. Mathis	233
EPR Evidence that the Photosystem I Acceptor A_1 is a Quinone Molecule M.C. Thurnauer, P. Gast, J. Petersen, D. Stehlik	237
Investigation of the Chemical Nature of Electron Acceptor A_1 in Photosystem I of Higher Plants R.W. Mansfield, J.H.A. Nugent, M.C.W. Evans	241
Evidence for the Existence of Electron Acceptors A_0 and A_1 in Cyanobacterial Photosystem I N.S. Smith, R.W. Mansfield, J.H.A. Nugent, M.C.W. Evans	245

XII

Iron X-ray Absorption Spectra of Acceptors in PS I Ann E. McDermott, Vittal K. Yachandra, R.D. Guiles, R. David Britt, S.L. Dexheimer, Kenneth Sauer, Melvin P. Klein	249
Photosystem I Charge Separation in the Absence of Centers A & B: Biochemical Characterization of the Stabilized P700 A2(X) Reaction Center John H. Golbeck, Kevin G. Parrett, Leslie L. Root	253
Picosecond Transient Absorption Spectroscopy of Photosystem 1 Reaction Centres from Higher Plants L.B. Giorgi, B.L. Gore, D.R. Klug, J.P. Ide, J. Barber, G. Porter	257
Light-induced Fourier Transform Infrared (FTIR) Spectroscopic Investigations of Primary Reactions in Photosystem I and Photosystem II B.A. Tavitian, E. Nabedryk, W. Mäntele, J. Breton	261
Chlorophyll Organization in Photosystem-I Reaction-center of Spinach Chloroplasts Isamu Ikegami, Shigeru Itoh	265
Bound Quinones in the Reaction Centres of Bacteria and Plants M.C.W. Evans	269
How Close is the Analogy between the Reaction Centre of PSII and that of Purple Bacteria? 2. The Electron Acceptor Side A.W. Rutherford	277
Depletion and Reconstitution of the Quinone at the Q _B Site in Photosystem II: A Thermoluminescence Study T. Wydrzynski, Y. Inoue	285
Chemically-induced Dynamic Electron Polarization in Photosystem 2 Reaction Centers Joseph T. Warden, Nathan M. Lacoﬀ, Károly Csatorday	289
The Mechanism of Fatty Acid Inhibition in Photosystem 2 Károly Csatorday, Claire Walczak, Joseph T. Warden	293
 3. Chlorophylls and Model Systems	
<i>In vivo</i> Spectral Peaks Related to New Chemical Species of Chlorophylls: 4-Vinyl-4-Desethyl Maarib B. Bazzaz	297
Chlorophyll <i>a'</i> in Photosynthetic Apparatus: Reinvestigation Tadashi Watanabe, Masami Kobayashi, Masataka Nakazato, Isamu Ikegami, Tetsuo Hiyama	303

Are Chlorinated Chlorophylls Components of Photosystem I Reaction Centers?	307
J. Fajer, E. Fujita, H.A. Frank, B. Chadwick, D. Simpson, K.M. Smith	
Environmental Effects on the Properties of Chlorophylls <i>in vivo</i> Theoretical Models	311
L.K. Hanson, M.A. Thompson, J. Fajer	
Effects of Structure and Geometry of Pigment-Protein Complexes on Experimental Quantities in Primary Processes of Photosynthesis	315
K. Vacek, M. Ambroz, O. Bilek, J. Hala, V. Kapsa, P. Pancoska, I. Pelant, L. Skala, L. Souckova	
Infrared Study of Solid Chlorophyll <i>a</i> Absorbing Near 700 nm at Room Temperature	321
Camille Chapados	
Borohydride Reduction of Bacteriochlorophyll <i>a</i> in the Light Harvesting Protein of <i>Rhodospirillum Rubrum</i>	325
Patricia M. Callahan, Therese M. Cotton, Paul A. Loach	
Fourier-transform Infrared (FTIR) Spectroelectrochemistry of Bacteriochlorophylls	329
W. Mäntele, A. Wollenweber, E. Nabedryk, J. Breton, F. Rashwan, J. Heinze, W. Kreutz	
Solvent Effects on the Transfer Kinetics of Bacteriochlorophyll Oxidation	333
Therese M. Cotton, Randall L. Heald	
X- and Y-polarized Absorptions of Chlorophyll <i>a</i> and Pheophytin <i>a</i> Oriented in a Lamellar Phase of Glycerylmonooctanoate/water	337
M. Fragata, T. Kurucsev, B. Nordén	
The Bacteriochlorophyll <i>c</i> Dimer in Carbon Tetrachloride	341
J.M. Olson, G.H. van Brakel, P.D. Gerola, J.P. Pedersen	
Superoxide Photogeneration by Chlorophyll <i>a</i> in Water/Acetone Solutions. Electron Spin Resonance Studies of Radical Intermediates in Chlorophyll <i>a</i> Photoreactions <i>in vitro</i>	345
Jun-Lin You, Karen S. Butcher, Angela Agostiano, Francis K. Fong	
Resonant Energy Transfer between Bulk Chlorophyll <i>a</i> and Chlorophyll <i>a</i> Dihydrate Dimers in Water/Acetone Mixtures. A Model of Sensitized Excitation in Plant Photosynthesis	349
Angela Agostiano, Karen A. Butcher, Michael S. Showell, Jun-Lin You, Albert J. Goth, Michael S. Showell	
4. Structure of Molecular Complexes: Crystallographic and Physical Studies	
The Structural Organization of Photosynthetic Reaction Centers	353
Hartmut Michel, Johann Deisenhofer	

XIV

Relating Structure to Function in Bacterial Photoreaction Centers J.R. Norris, D.E. Budil, D.M. Tiede, J. Tang, S.V. Kolaczowski, C.H. Chang, M. Schiffer	363
Crystallographic Studies of the Photosynthetic Reaction Center from <i>R. Sphaeroides</i> C.-H. Chang, D. Tiede, J. Tang, J. Norris, M. Schiffer	371
Structure Analysis of the Reaction Center from <i>Rhodopseudomonas</i> <i>Sphaeroides</i> : Electron Density Map at 3.5Å Resolution J.P. Allen, G. Feher, T.O. Yeates, D.C. Rees	375
Evidence of the Primary Charge Separation in the D ₁ D ₂ Complex of Photosystem II from Spinach: EPR of the Triplet State M.Y. Okamura, K. Satoh, R.A. Isaacson, G. Feher	379
Crystallization and Spectroscopic Investigations of the Pigment- protein Complexes of <i>Rhodopseudomonas Palustris</i> T. Wacker, K. Steck, A. Becker, G. Drews, N. Gad'on, W. Kreutz, W. Mäntele, W. Welte	383
Spectroscopy, Structure and Dynamics in the Reaction Center of <i>Rhodopseudomonas Viridis</i> J. Breton, J. Deprez, B. Tavitian, E. Nabedryk	387
Interspecific Structural Variations of the Primary Donor in Bacterial Reaction Centers Qing Zhou, Bruno Robert, Marc Lutz	395
Linear-Dichroic Absorbance Detected Magnetic Resonance (LD-ADMR) Spectroscopy of the Photosynthetic Reaction Center of <i>Rhodopseudomonas Viridis</i> . Spectral Analysis by Exciton Theory E.J. Lous, A.J. Hoff	399
Optical Properties of the Reaction Center of <i>Chloroflexus Aurantiacus</i> at Low Temperature. Analysis by Exciton Theory H. Vasmel, R.F. Meiburg, J. Amesz, A.J. Hoff	403
The Photochemical Reaction Center of <i>Chloroflexus Aurantiacus</i> : Isolation and Protein Chemistry of the Purified Complex Judith A. Shiozawa, Friedrich Lottspeich, Reiner Feick	407
Structures of Antenna Complexes and Reaction Centers from Bacteriochlorophyll b-containing Bacteria: Resonance Raman Studies Bruno Robert, Robert Steiner, Qing Zhou, Hugo Scheer, Marc Lutz	411
Strong Orientational Ordering of the Near-infrared Transition Moment Vectors of Light-harvesting Antenna Bacterioviridin in Chromatophores of the Green Photosynthetic Bacterium <i>Chlorobium</i> <i>Limicola</i> , Strain c Z.G. Fetisova, S.G. Kharchenko, I.A. Abdourakchmanov	415

Light Absorption and Fluorescence of BChl <i>c</i> in Chlorosomes from <i>Chloroflexus Aurantiacus</i> and in an <i>in vitro</i> Model Daniel C. Brune, Robert E. Blankenship	419
Serrins as a Probe for Pigments Located near the Surfaces of Bacterial Photosynthetic Membranes Rafael Picorel, Randall E. Holt, Therese M. Cotton, Michael Seibert	423
Optical Excited Triplet States in Antenna Complexes of the Photosynthetic Bacterium <i>Rhodospseudomonas Capsulata</i> A1a ⁺ Detected by Magnetic Resonance in Zero-field A. Angerhofer, J.U. von Schütz, H.C. Wolf	427
Singlet Energy Transfer in Photosynthetic Bacteria: Absorption and Fluorescence Excitation of B800-850 Complexes Barry W. Chadwick, Harry A. Frank, Chaoying Zhang, Shahriar S. Taremi, Richard J. Cogdell	431
Properties of the Core Complex of Photosystem II J.J. Plijter, R.J. van Dorssen, J.P. Dekker, F.T.M. Zonneveld, H.J. van Gorkom, J. Amesz	435
Pigment Arrangement in Photosystem II R.J. van Dorssen, J.J. Plijter, A. den Ouden, J. Amesz, H.J. van Gorkom	439
Three-dimensional Crystals of the Light-harvesting Chlorophyll a/b Protein Complex from Pea Thylakoids W. Kuehlbrandt	443
Interpretation of Transient Linear Dichroism Spectra of LHC Particles Robert S. Knox, Su Lin	445
Resonance Raman Spectroscopy of Chlorophylls and the Light-harvesting Chlorophyll-a/b Protein H.N. Fonda, G.T. Babcock	449

5. Oxygen Evolution

Oxygen-evolving Complex of Photosystem II in Higher Plants Norio Murata, Mitsue Miyao	453
Kinetics and Structure on the High Potential Side of Photosystem II G.T. Babcock, T.K. Chandrashekar, D.F. Ghanotakis, C.W. Hoganson, P.J. O'Malley, I.D. Rodriguez, C.F. Yocum	463
Endor Characterization of H ₂ O/D ₂ O Exchange in the D ⁺ Z ⁺ Radical in Photosynthesis I.D. Rodriguez, T.K. Chandrashekar, G.T. Babcock	471

XVI

Endor Characterization of the Z ⁺ /D ⁺ Species in Photosystem II and Relevant Model Compounds	475
T.K. Chandrashekar, P.J. O'Malley, I.D. Rodriguez, G.T. Babcock	
Time-resolved ESR Spectrum of Z ⁺ in Oxygen-evolving Photosystem II Membranes	479
C.W. Hoganson, Y. Demetriou, G.T. Babcock	
Spatial Relationship between the Intramembrane Components (D ⁺ , Z ⁺) which Give Rise to Signal II and the Membrane Peripheral Proteins Working in Photosystem II Oxygen Evolution Studied by the Effect of Spin-relaxing Reagent Dysprosium	483
Shigeru Itoh, Yasuhiro Isogai, Xiao-Song Tang, Kimiyuki Satoh	
The Effects of Chemical Oxidants on the Electron Transport Components of Photosystem II and the Water-oxidizing Complex	487
J. Tso, D. Hunziker, G.C. Dismukes	
On the Mechanism of Photosynthetic Water Oxidation	491
Gary W. Brudvig, Julio C. de Paula	
Coordination of Ammonia, but not Larger Amines, to the Manganese Site of the O ₂ -evolving Center in the S ₂ State	499
Warren F. Beck, Gary W. Brudvig	
EPR Studies of the Oxygen-evolving system. The Interaction with Amines	503
Lars-Erik Andreasson, Örjan Hansson	
Cooperative Binding of Hydroxylamine and Hydrazine to the Water-oxidizing Complex	511
Verena Förster, Wolfgang Junge	
Reaction Mechanisms of H ₂ O Substrate Analogues at the PS II-donor Side in Thylakoids and PS II-particles	515
B. Hanssum, G. Renger	
Proton Release by Photosynthetic Water Oxidation	519
Ralf Diedrich-Glaubitz, Manfred Völker, Gernot Renger, Peter Gräber	
On the Cleavage of Water Pattern of Charges and Protons. States of Water and Manganese. Routes and Rate of Intermediates	523
H.T. Witt, Ö. Saygin, K. Brettel, E. Schlodder	
Absorption Changes with Periodicity Four. Associated with Photosynthetic Oxygen Evolution	533
Jan P. Dekker, Johan J. Plijter, Hans J. van Gorkom	
State of Manganese During Water Splitting	539
Ö. Saygin, H.T. Witt	
New Results about the Molecular Mechanism of Photosynthetic Water Oxidation	541
G. Renger, B. Hanssum, W. Weiss	

The Modification of the Donor Side Reaction Pattern in PS II Membrane Fragments by Trypsin and CaCl ₂ M. Völker, H.J. Eckert, G. Renger	545
Studies on Water Oxidation by Mass Spectrometry in the Filamentous Cyanobacterium <i>Oscillatoria Chalybea</i> Klaus P. Bader, Pierre Thibault, Georg H. Schmid	549
Flash-induced Enhancements in the ¹ H-relaxation Rate of Photosystem II Particles A.N. Srinivasan, R.R. Sharp	553
The State of Manganese in the Photosynthetic Apparatus: An X-ray Absorption Spectroscopy Study Vittal K. Yachandra, R.D. Guiles, Ann McDermott, James Cole, R. David Britt, S.L. Dexheimer, Kenneth Sauer, Melvin P. Klein	557
Structural Features of the Manganese Cluster in Different States of the Oxygen Evolving Complex of Photosystem II: An X-ray Absorption Spectroscopy Study R.D. Guiles, Vittal K. Yachandra, Ann E. McDermott, R. David Britt, S.L. Dexheimer, Kenneth Sauer, Melvin P. Klein	561
Characterization of the Mn-containing O ₂ Evolving Complex from the Cyanobacterium <i>Synechococcus</i> Using EPR and X-ray Absorption Spectroscopy Ann McDermott, Vittal K. Yachandra, R.D. Guiles, R. David Britt, S.L. Dexheimer, Kenneth Sauer, Melvin P. Klein	565
The Flash Number Dependence of EPR Signal II Decay As a Probe for Charge Accumulation in Photosystem II James Cole, Kenneth Sauer	569
Electron Spin Echo Studies of PSII Membranes R. David Britt, Kenneth Sauer, Melvin P. Klein	573
EPR Studies at 9 and 34 GHz of the Multiline and <i>g</i> = 4.1 S ₂ Signals Roland Åasa, Örjan Hansson, Tore Vänngård	577
Structural and Functional Aspects of Electron Transfer in Photosystem 2 of Oxygen-evolving Organisms V.V. Klimov, I.B. Ganago, S.I. Allakhverdiev, M.A. Shafiev, G.M. Ananyev	581
The Study of Effects on Strongly-bound Manganese of Oxygen Evolving Complex in Wheat Chloroplasts by EPR Sun Qi, Luo Chang-Mei, Zhang Li-Li, Fang Zhao-Xi, Mei Zhen-An	585
Evidence for the Role of Functional Manganese in Hydrogen-peroxide-stimulated Oxygen Production of the First Flash in CaCl ₂ -washed Photosystem II Membranes Steven P. Berg, Michael Seibert	589

XVIII

Interaction between Manganese and the 33-kilodalton Protein in Spinach PS II Yasusi Yamamoto	593
Manganese and Calcium Binding Properties of the Extrinsic 33 kDa Protein and of Photosystem II Membranes D. Hunziker, D.A. Abramowicz, R. Damoder, G.C. Dismukes	597
The 33 kDa Extrinsic Polypeptide of Photosystem II is not a Ligand to Manganese in the O ₂ Evolving Complex Anne-Frances Miller, Julio C. de Paula, Gary W. Brudvig	601
Effect of Release of the 17 and 23 kDa Polypeptides of Photosystem II on Cytochrome b ₅₅₉ Julio C. de Paula, Brian W. Wu, Gary W. Brudvig	605
Cytochrome b ₅₅₉ Plays a Structural Role in the Oxygen Evolving Complex of Photosystem II Lynmarie K. Thompson, Julian M. Sturtevant, Gary W. Brudvig	609
Effect of the 33-kDa Protein on the S-state Transition in the Oxygen-evolving Complex M. Miyao, N. Murata, B. Maison-Peteri, A. Boussac, A.-L. Etienne, J. Lavorel	613
PSII Ca Abundance and Interaction of the 17,24 kD Proteins with the Cl ⁻ /Ca ²⁺ Essential for Oxygen Evolution Kirk Cammarata, George Cheniae	617
Photoactivation of the Water Oxidizing Complex by Photosystem 2 Membranes N. Tamura, G. Cheniae	621
Numbers of Calcium Ions Associated with Oxygen Evolving Photosystem II Preparations with Different Affinities Sakae Katoh, Kazuhiko Satoh, Takashi Ohno, Jian-Ren Chen, Yasuhiro Kasino	625
Involvement of Ca ²⁺ and the 33 kD Polypeptide in Cl ⁻ Binding to the Oxygen Evolving Complex of Photosystem II W.J. Coleman, Govindjee, H.S. Gutowsky	629
Inhibition at the CA ²⁺ Sensitive Site of the Oxygen Evolving Center by Ruthenium Red Sylvie Lemieux, Robert Carpentier	633
Thermoluminescence Studies of the Abnormal S-states Formed in Cl ⁻ -depleted or 33 kDa Extrinsic Protein-depleted PSII Yorinao Inoue	637
Temperature Dependence of the S-state Transition in a Thermophilic Cyano-bacterium Measured by Thermoluminescence Hiroyuki Koike, Yorinao Inoue	645

- Depletion of Cl^- or 33 kDa Extrinsic Protein Modifies the Stability of $\text{S}_2\text{Q}_\text{A}^-$ and $\text{S}_2\text{Q}_\text{B}^-$ Charge Separation States in PS II 649
Imre Vass, Taka-aki Ono, Peter H. Homann, Hermann Gleiter, Yorinao Inoue
- Abnormal S_2 State Formed in Chloride Depleted Photosystem II as Revealed by Manganese EPR Multiline Signal 653
T. Ono, J.L. Zimmermann, Y. Inoue, A.W. Rutherford
- Cl^- Dependent Binding of the Extrinsic 23 kDa Polypeptide at the Water Oxidizing Site of Chloroplast Photosystem II 657
Peter H. Homann
- Effects of Chloride on Paramagnetic Coupling of Manganese in Calcium Chloride-washed Photosystem II Preparations 661
Gopinath Mavankal, Douglas C. McCain, Terry M. Bricker
- Accessibility for, and Production of H_2O_2 Related to PS-II 665
Wolfgang P. Schröder, Hans-Erik Åkerlund
- Reversible Inhibition of Photosystem Two Electron Transfer Reactions and Specific Removal of the Extrinsic 23 kDa Polypeptide by Alkaline pH 669
David J. Chapman, James Barber
- O_2 Flash Yield Sequences of Photosystem II Membranes—Sequential Extraction of the Extrinsic Proteins 673
Michael Seibert, Brigitta Maison-Peteri, Jean Lavorel
- Comparative Study of Period 4 Oscillations of the Oxygen and Fluorescence Yield Induced by a Flash Series in Inside out Thylakoids 677
M.J. Delrieu, F. Rosengard
- Purification of an Oxygen Evolving Photosystem II Reaction Center Core Preparation 681
D.F. Ghanotakis, D.M. Demetriou, C.F. Yocum
- Selective Depletion of Water-soluble Polypeptides Associated with Photosystem II 685
Charlene M. Waggoner, Charles F. Yocum
- Binding of the 17 and 23 kDa Water-soluble Polypeptides to a Highly-resolved PSII Reaction Center Complex 689
Stewart Merritt, Patrik Ernfors, Demetrios Ghanotakis, Charles Yocum
- A Manganese Containing Protein Complex Isolated from Photosystem II Preparations of Spinach 693
Neil R. Bowlby, Wayne D. Frasch
- Purification and Proteinchemical Characterization of the Extrinsic Membrane Proteins in the Water Splitting System of Spinach 697
Joachim Vater, Johann Salnikow, Ricci Zepmeusel, Christer Jansson

XX

Partial Amino Acid Sequences of the Proteins of Pea and Spinach Photosystem II Complex	701
N. Murata, H. Kajiura, Y. Fujimura, M. Miyao, T. Murata, A. Watanabe, K. Shinozaki	
Proline-rich Structure at Amino-terminal Region of the 18-kDa Protein of Photosynthetic Oxygen-evolving Complex	705
Tomohiko Kuwabara, Teruyo Murata, Mitsue Miyao, Norio Murata	
Topographical Studies on Subunit Polypeptides of Oxygen-evolving Photosystem II Preparations by Reversible Crosslinking: Functions of Two Chlorophyll-carrying Subunits	709
Isao Enami, Takeshi Miyaoka, Sahoko Igarashi, Kazuhiko Satoh, Sakae Katoh	
Tenacious Association of the 33kDa Extrinsic Polypeptide (Water Splitting) with PS II Particles	713
Edith L. Camm, Beverley R. Green	
Thermodynamic Constraints to Photosynthetic Water Oxidation	717
Lee Spencer, Donald T. Sawyer, Andrew N. Webber, Robert L. Heath	
Binuclear and Tetranuclear Manganese Complexes: As Models for the Site for Photosynthetic Water Oxidation	721
J.E. Sheats, B.C. UnniNair, V. Petrouleas, S. Artandi, R.S. Czernuszewicz, G.C. Dismukes	
Models for Manganese Centers in Metalloenzymes	725
Vincent L. Pecoraro, Dimitris P. Kessissoglou, Xinhua Li, William M. Butler	
Molecular Orbital Study (IV) on the 'Microsurface' Model of Catalytic Binuclear Manganese Complex in Photosynthetic Water-splitting and Oxygen-evolving Reaction	729
Masami Kusunoki	
Dynamic Linearity of the Bare Platinum Electrode for Oxygen Exchange Measurements in Marine Algae	733
S.I. Swenson, C.P. Meunier, K. Colbow	
A Dynamic Model for the Bare Platinum Electrode	737
C.P. Meunier, S.I. Swenson, K. Colbow	
Index of names	741

PHOTOCHEMISTRY AND PHOTOPHYSICS OF C-PHYCOCYANIN

Hugo Scheer, Botanisches Institut der Universität, Menzinger Str.67
D-8000 München 19, Federal Republic of Germany

Introduction

Phycocyanin (PC) belongs to a group of pigments functional for light-harvesting in cyanobacteria, red and cryptophyte algae. In the former two classes of organisms, it is a major constituent of phycobilisomes, the light-harvesting complexes located at the outer surface of the photosynthetic membrane. There, it absorbs light energy in the spectral range between 580 and 640 nm, and transfers it via a second biliprotein, allophycocyanin (APC), to the chlorophyll within the membrane. In many species, PC also accepts energy from a third type of biliprotein, e.g. phycoerythrin (PE), thus acting as an intermediate carrier in the energy transfer from the latter to APC.

The simplest PC, which is found in cyanobacteria (C-PC), contains three chromophores of the dihydrobilindion type, each of them being attached covalently to the apoprotein via a single thioether bond to cysteine. The same chromophore is present in APC, and a chromophore differing only in one of the β -pyrrolic substituents is found in the plant photomorphogenetic pigment, phytochrome. According to the different functions of these three pigments, the properties of the chromophores in each of them are quite different from each other, and they all differ considerably from the properties of free pigments bearing this type chromophore (1). The factors responsible for the different adaptations of these structurally so similar chromophores are still only partly understood. From reversible denaturation studies in C-PC, it appears that they are mostly due to non-covalent protein-chromophore interactions.

The recent elucidation of the x-ray structures (2,3) of C-PCs from two different organisms, has greatly advanced our knowledge of these pigments. It has for the first time in any photosynthetic antenna system become possible to 'look' at the native chromophore structures on a molecular level, and to obtain direct information on their conformations and relative orientations. This renders it possible to test the viability of theoretical models applied in the calculation of the their spectral properties, of the energy transfer pathways, the kinetics among them, etc., by using the structural data as input parameters.

This report summarizes recent work carried out along these lines in München. It contains data on the photochemistry of C-PC from the cyanobacterium, Mastigocladus (M.) laminosus, which are compared to the respective properties of the photochromic plant photoreceptor, phytochrome, as well as theoretical and experimental results on the energy transfer in aggregates of different sizes from this chromoprotein.

1.1.144

Chromophore structure

The three chromophores of **C-PC** are bound to the protein via single thioether bonds to cysteine 84 on the α -subunit (cys α -84), and to cys β -84 and cys β -155 on the β -subunit. A linkage to the C-3 ethyl-substituent of the hydrogenated ring A had been established for chromophores of all type of plant biliproteins including phytochrome (1,4). A more complex binding pattern has recently been proposed, which involves a different linkage and structure for the chromophore β -155 (5). Instead of being hydrogenated at ring A, this chromophore is hydrogenated at ring D and bound via the 'exo' 18-thioethyl substituent.

All chromophores are present in more or less extended conformations (2,3), which account for the observed absorption increase of the red bands, and a concomitant decrease of the near-uv bands (1). It is also likely, that the chromophores have a reduced conformational mobility, which accounts for their high fluorescence quantum yields, and for their inertness to a variety of chemical reagents (metal ions, reducing agents) known to react readily with the free chromophores (see 1 for leading references). The confirmation of this general structural principle in the crystal structure of C-PCs from two different organisms (*M. laminosus* and *Agmenellum (A.) quadruplicatum*), makes it likely that the spectrally similar chromophores of **APC** and phytochrome have similar native structures as well.

Chromophore assignment

While the x-ray results supported this general structure principle, they show on the other hand pronounced differences among the details of the chromophore conformations, of the binding sites and of likely interactions with the apoprotein. Such differences account for a variety well documented spectroscopic and chemical results indicating the presence of a set of distinct chromophores in almost any phycobiliprotein (1). In **C-PC**, these allowed the definition of three distinct chromophores, e.g. α -1, β -1 and β -2.

In the case of **C-PC** from *M. laminosus*, the following data were combined for the spectral resolution: The integral pigment can be separated into two subunits, the α -subunit bearing only one chromophore, and the β -subunit bearing two chromophores. Since the absorption spectra of the two subunits -weighted properly according to the subunit stoichiometry- add up to the spectrum of the monomeric pigment, it is likely that the states of the chromophores remain unchanged during subunit separation, and that strong inter-subunit chromophore-chromophore interactions are absent. This yields directly the required absorption and fluorescence spectra of the α -subunit, and reduces the problem to the resolution of the β -subunit spectrum. The presence of two spectrally distinct chromophores in this subunit is derived from several lines of evidence:

Reversible photochemistry: Native **PC** has a high fluorescence quantum yield. Its photochemistry is characterized by an irreversible bleaching, which has a low quantum yield (0.4%) and proceeds probably via the triplet state because it is slowed down in the presence of oxygen. This irreversible reaction occurs with a similar quantum yield also in phytochrome (Scheer, unpublished results). Addition of urea to **PC** causes a gradual, reversible unfolding of the protein and a concomitant loss of its interactions with the chromophore. At 8 M urea, the protein is completely denatured, and the chromophores then attain the properties characteristic of free bile pigments (1). Here, the fluorescence is greatly reduced ($\phi < 10^{-3}$). The photochemical reactivity is increased, but it is again irreversible, leading to a variety of tri- and tetrapyrroles absorbing at shorter wave lengths. The onset of unfolding at intermediate urea concentrations is characterized by a reduction of fluorescence and the concomitant occurrence of a reversible photochemical reaction, which is maximum at about 5 M urea. Similar reactions have been observed as

well with other denaturants at moderate concentrations, and they have been related to the reversible Z,E-isomerization of the phytochrome chromophore, and to the primary reactions of the less well understood phytochromes (see 1). If the isolated subunits of C-PC from *M. laminosus* are titrated with urea this reaction is negligibly in the α -subunit, but much more pronounced in the β -subunit. The absorption difference spectrum of the latter shows a single negative band in the visible spectral range peaking at 624 nm, which is considerably to the red of the absorption maximum at 606 nm, and its shape is similar to that of a typical bile pigment (6). This suggests, that only one of the two chromophores on the β -subunit is susceptible to this reaction, and that this chromophore absorbs at longer wavelengths than the second, inactive one.

Fluorescence polarization: The fluorescence polarization spectra of nearly all phycobiliproteins show distinct discontinuities, and the anisotropy rises in discrete steps towards longer wavelengths (13, 14, see also 1). Since the red absorption band of bile pigments corresponds to a single electronic transition, this has been interpreted as the result of several distinct chromophores being present, with different orientations and different absorption spectra, among which energy transfer occurs. The fluorescence polarization spectrum of the β -subunit of C-PC from *M. laminosus* shows two distinct regions of anisotropy (7). Below 600 nm, it is nearly wavelength independent about 0.2, and then rises sharply to 0.4. There are, therefore, at least two chromophores present absorbing below and above this threshold wavelength. Assuming a similar Stokes' shift for the fluorescing chromophore of the β -subunit ($\lambda_{\max}^{\text{fluor}} = 643$ nm) and the one of the α -subunit ($\lambda_{\max}^{\text{abs}} = 616$, $\lambda_{\max}^{\text{fluor}} = 641$ nm), an absorption around 620 nm can be estimated for the former.

Circular dichroism: The cd spectrum of the α -subunit shows a single positive band in the visible spectral region peaking close to its absorption maximum. The cd-spectrum of the β -subunit also shows a single positive peak. Its intensity is decreased by 40% on a molar basis, increased by 20% on a chromophore basis, and centered well to the blue (590nm) of the absorption maximum. At longer wavelengths, the band trails slightly and indicates the presence of a smaller, much less intense band. This result, which has been reported independently by Mimuro et al. (7), is again, best interpreted as to arise from two different chromophores, one of them is strongly optically active and absorbs around 595 nm, the second one is much less active and absorbs above 610 nm.

Curve resolution of absorption spectrum: To better define the absorption bands of these chromophores, the spectrum of the β -subunit was resolved by computer analysis. It was assumed, that the shape and width of the bands were identical to that of the alpha chromophore, and the starting wavelengths for the analysis were estimated from the aforementioned data. The absorption band was fit best with two bands peaking at 598 and 622 nm, with molar absorptivities of 92 and 60%, respectively, of that of the α -subunit. For an estimation of the individual fluorescence spectra, similar Stokes' shifts and fluorescence lifetime was furthermore assumed. The resulting spectral data are given in table 1 and compared to a similar analysis by Mimuro et al. (7).

Chemical reactivity and assignment to binding sites

Whereas the aforementioned results allowed the distinction of two spectrally defined chromophores on the β -subunit, the correlation between the different chromophores defined above (α -1, β -1 and β -1), and the ones defined by their binding sites (α -84, β -84, β -155), respectively, was still lacking. Following a suggestion by Schirmer, Bode and Huber (2,3), we have been able to make this assignment by treatment of C-PC with organic mercurials (8). There is only a single free cysteine present in C-PC located at position β -111 (9), which is very close to the chromophore β -84, but more than 22A from chromophores β -155 and β -84 (2,3). This cysteine is the only site to which mercurials were

I.1.146

bound in heavy-atom derivatives of C-PC crystals, and it was expected that a binding of the bulky reagent at this position would have a discernible effect on the spectral properties of the neighboring chromophore, and only on this one. In solution, titration of C-PC from M. laminosus leads to a decrease of absorption at about 620nm, and a concomitant, albeit smaller, increase around 650nm, and a general absorption increase in the near-uv spectral region (Fig. 1). In the β -subunit, the same spectral changes are observed, but the relative amplitudes are increased. Titration experiments showed, that the reaction is complete after the addition of 1 ± 0.2 moles of the mercurial (p-chloro-mercuribenzenesulfonate, PCMS) per mole of C-PC or β -subunit, respectively, and that the reaction can be reversed to more than 80% by addition of thiols. In the α -subunit, the reaction is negligible (Fig.1).

These findings are interpreted in the following way:

- 1: The reaction site is the single free cys-111 on the β -subunit, similar to the situation in the crystal.
- 2: No irreversible reaction occurs between mercurials and the chromophores, because the α -subunit is inert.
- 3: Since the difference absorption maximum is on the red side of the absorption maximum in the β -subunit, it must be related to a spectral change of the long-wavelength absorbing chromophore, e.g. the one defined above spectroscopically as β -1.
- 4: Due to the spatial relationships, this chromophore is the one bound to cys- β -84, e.g. chromophore β -84 is identical with chromophore β -1.
- 5: The indirect effect of the mercurial binding on the absorption of chromophore β -84 involves probably a conformational change from the native, extended to a more denatured, cyclic-helical conformation, as indicated by the overall decrease of absorption in the red, and an increase in the near-uv spectral region (1).
- 6: The resulting spectral data for the three chromophores are summarized in table 1.
- 7: Due to the inertness of the chromophores to a direct reaction with mercurials, these reagents are suitable to test the accessibility of cys-111 in higher aggregates, and at the same time to identify the absorption of the β -84 chromophores.

	α -84	β -1 = β -84	β -2 = β -155
This work	616(120)	622(72)	598(106)
Mimuro et al. (7)	618(108)	624(103)	594(113)

Table 1: Absorption maxima [λ_{\max} ($\epsilon \times 10^{-3}$)] of the individual chromophores of C-phycoyanin from Mastigocladus laminosus.

The assignment (β -1 = β -84, β -2 = β -155) agrees with the ad-hoc assignment of by Mimuro et al. (7), and there is also a reasonable agreement of the spectral data of the individual chromophores. The most pronounced differences are the position of the β -155 absorption, which is displaced to the blue by appx. 4nm by these authors, and a lower absorptivity of the β -84 absorption in our calculations.




Fig.1: Reaction of trimeric C-phycocyanin from *Mastigocladus laminosus* (A), its α - (B) and β -subunit (C). Figure (A) shows both the absorption spectra before (a) and after (b) reaction with PCMS (1 mole / mole C-PC), and the difference spectra during the titration (c,d,e). The other figures show the absorption spectrum before addition of PCMS, and the absorption difference spectrum after its addition (1 mole / mole subunit). All reactions in potassium phosphate buffer (50mM, pH 7.5).

1.1.148

Kinetics of energy transfer

Energy transfer in phycobiliproteins has been suggested to occur mainly via the induced dipole or Förster type mechanism. The main arguments to this have come originally from the absence in most biliproteins, including C-PC from *M. laminosus*, of strong, s-shaped signals indicative of strong exciton couplings, and from polarization spectroscopy (see 1). Over the past 5 years, the amount of kinetic data on energy transfer has tremendously increased. It has been found in particular, that the fastest energy transfer processes take place on a time scale in the order of tens of picoseconds or even less (10,11). With all the necessary information at hand, it was then intriguing to test if such fast kinetics could be matched by theory on the basis of a pure Förster transfer mechanism.

The rate constant for energy transfer is described by:

$$k \approx \frac{\kappa^2}{R^6} \int F_{\text{Donor}} \cdot A_{\text{Acceptor}}$$

The major variables in this equation are the distance R between the transition dipoles of the donor and acceptor chromophores, the relative orientations factors κ of the two, and the overlap of the fluorescence of the donor (F_{Donor}) with the absorption of the acceptor (A_{Acceptor}). The only other variable in the equation, e.g., the refractive index of the medium, is hidden in the proportionality constant. A value of 1.56 has been used throughout. The first two pieces of information have been taken from the x-ray data of Schirmer et al. for C-PC from *A. quadruplicatum* (3). This pigment has a very similar chromophore arrangement as C-PC from *M. laminosus* (2), but it has been resolved to higher accuracy. Moreover, it crystallizes as stacks of hexamers rather than trimers, so that orientations and distances between chromophores on different trimers and hexamers are available. These authors have tabulated the distances of the centers of gravity of the π -systems for all chromophores, as well as the relative orientations of chromophores as defined by the masses of the atoms present in the chromophore π -systems projected on a straight line. These distances and orientations do, therefore, not strictly correspond to the transition dipoles, but rather to their reduced masses. In view of the elongated structure of the chromophores, it is likely, however, that the deviations are reasonable. The overlap integral was finally calculated from the individual absorption and fluorescence bands of the three different chromophores as given in the top row of table 1.

The details of these calculations, which are the result of a continuing cooperation between K. Sauer and our group, are being published elsewhere (12), and a summary is presented in the poster abstracts of this conference. There is a good agreement with most of the currently available experimental data. The calculations show an increased transfer rate with increasing aggregate size. They support, in particular, a preferential energy transfer along the rods of trimer-stacks, as compared to energy transfer within trimers. Such a preferential transfer would greatly facilitate the funneling of energy towards the reaction centers. According to these results, the energy transfer in C-PC can be accounted for well by the Förster mechanism, and the flow of excitation energy in these moderately complex aggregates can be analyzed on a molecular basis. Since the data can be transformed readily to mimic a variety of experimental conditions (different excitation and emission wavelengths, static and dynamic depolarization), a comparison with new data and/or assistance in the choice of experimental conditions are expected to further evaluate this conclusion critically.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft, Bonn (SFB 143 and Forschergruppe "Pflanzliche Tetrapyrrole"). The cooperation with K. Sauer, Berkeley, was rendered possible by an Alexander-von-Humboldt Award. The following coworkers were involved in this work as detailed in the references: R. Fischer, W. John, G. Schmidt, G. Schoy, S. Siebzehrübl (all Botanisches Institut der Universität München) and P.Sauer (Carlton College).

References

1. Scheer, H. (1982) in *Light Reaction Path of Photosynthesis* (Fong, F.K., ed.) pp. 7-45, Springer-Verlag, Berlin.
2. Schirmer, T., Bode, W., Huber, R., Sidler, W. and Zuber, H. (1985) *J. Mol. Biol.* 184, 257-277.
3. Schirmer, T., Huber, R., Schneider, M., Bode, W., Miller, M. and Hackert, M. L. (1986) *J. Mol. Biol.* 188, 651-676.
4. Rüdiger, W. and Scheer, H. (1983) in *Handbook of Plant Physiology*, Vol. 16 (Shropshire, W. and Mohr, H., eds), pp. 119-151, Springer Verlag, Berlin
5. Bishop, J. E., Lagarias, J. C., Nagy, J. O., Schoenleber, R. W., Rapoport, H., Klotz, A. V. and Glazer, A. N. (1986) *J. Biol. Chem.* 261, 6790-6796.
6. Siebzehrübl, S., Fischer, R. and Scheer, H. (1985) *Proc. Int. Symp. Energy Transfer*, Charles University Press, Praha.
7. Mimuro, M., Füglistaller, P., Rübeli, R. and Zuber, H. (1986) *Biochim. Biophys. Acta* 848, 155-166.
8. Siebzehrübl, S., Fischer, R. and Scheer, H. (1986) *FEBS Lett.*, submitted.
9. Zuber, H. (1985) in *Antennas and Reaction Centers of Photosynthetic Bacteria* (Michel-Beyerle, M. E., ed.) pp. 3-25, Springer Verlag, Berlin.
10. Scheer, H. (1986) in *Handbook of Plant Physiology*, Vol. 19 (Staehelein, L. A. and Arntzen, C. J., eds.) pp. 327-337, Springer Verlag, Berlin.
11. Earlier and related work has been reviewed in: Holzwarth, A. R. (1986) *Photochem. Photobiol.* 43, yearly review.
12. Sauer, K. and Scheer, H., this volume, p.
13. Dale, R. E. and Teale, F. J. W. (1970) *Photochem. Photobiol.* 12, 99-117
14. Grabowski, J. and Gantt, E. (1978) *Photochem. Photobiol.* 28, 39-46 and 47-56