



---

# PERSPECTIVES IN PHOTOSYNTHESIS

PROCEEDINGS OF THE TWENTY-SECOND JERUSALEM SYMPOSIUM ON  
QUANTUM CHEMISTRY AND BIOCHEMISTRY HELD IN  
JERUSALEM, ISRAEL, MAY 15-18, 1989

*Edited by*

J. JORTNER

*Department of Chemistry,  
University of Tel-Aviv, Israel*

and

B. PULLMAN

*Institut de Biologie Physico-Chimique  
(Fondation Edmond de Rothschild), Paris, France*



KLUWER ACADEMIC PUBLISHERS

DORDRECHT / BOSTON / LONDON

## TABLE OF CONTENTS

PREFACE	ix
H. MICHEL and J. DEISENHOFER / The Photosynthetic Reaction Center from the Purple Bacterium <i>Rhodospseudomonas Viridis</i> : Relation of Structure and Function	1
J.R. NORRIS, T.J. DiMAGNO, A. ANGERHOFER, C.H. CHANG, O. EL-KABBANI and M. SCHIFFER / A Structural Basis for Electron Transfer in Bacterial Photosynthesis	11
J. BRETON / Orientation of the Pheophytin Primary Electron Acceptor and of the Cytochrome B559 in the D1D2 Photosystem II Reaction Center	23
S.G. BOXER, D.J. LOCKHART, C. KIRMAIER and D. HOLTEN / Mechanism of Charge Separation in Photosynthetic Reaction Centers : Electric Field Effects on the Initial Electron Transfer Kinetics	39
M.R. GUNNER and B. HONIG / Electrostatic Analysis of the Midpoints of the Four Hemes in the Bound Cytochrome of the Reaction Center of <u>RP. Viridis</u>	53
C. KIRMAIER, L.M. McDOWELL, E.J. BYLINA, D.C. YOVAN and D. HOLTEN / Primary Photochemistry in Reaction Centers from the His <sup>M200</sup> Leu Mutant of <i>Rhodobacter Capsulatus</i>	61
A.J. HOFF, E.J.J. GROENEN and E.J. LOUS / A Note on Singlet-Singlet and Singlet-Triplet Exciton Interactions in the Bacterial Photosynthetic Reaction Center	71
W. LERSCH, E. LANG, R. FEICK, W.J. COLEMAN, D.C. YOVAN and M.E. MICHEL-BEYERLE / Determination	

of the Exchange Interaction in the Primary Radical Ion Pair in Reaction Centers	81
H. LEVANON / Photoelectron Transfer Reactions : Porphyrins-Quinone. FT- and CW-Time Resolved-EPR Spectroscopies	91
D. TANG, S.G. JOHNSON, R. JANKOWIAK, J.M. HAYES, G.J. SMALL and D.M. TIEDE / Structure and Marker Mode of the Primary Electron Donor State Absorption of Photosynthetic Bacteria : Hole Burned Spectra	99
R. FISCHER, J. GOTTSTEIN, S. SIEBZEHRÜBL and H. SCHEER / Influence of Chromophores on Quarternary Structure of Phycobiliproteins from the Cyanobacterium, <i>Mastigocladus laminosus</i>	121
J.T.HYNES , E.A. CARTER, G. CICCOTTI, H.J. KIM, D.A. ZICHI, M. FERRARIO, R. KAPRAL / Environmental Dynamics and Electron Transfer Reactions	133
R. SILBEY / Relaxation and Coherence in Simple Model Systems	149
M.D. NEWTON / Superexchange Coupling Mechanisms for Electron Transfer Processes	157
Y. HU and S. MUKAMEL / Sequential Versus Superexchange Electron Transfer in the Photosynthetic Reaction Center	171
M.A. RATNER / Non-Adiabatic Electron Transfer : Some Dynamical and Electronic Extensions of Standard Rate Expressions	185
J.N. ONUCHIC, R.F. GOLDSTEIN and W. BIALEK / Biomolecular Dynamics - Quantum or Classical? Results for Photosynthetic Electron Transfer	211
N. MATAGA / Femtosecond-Picosecond Laser Photolysis Studies on Electron Transfer Dynamics and Mechanisms in Some Model Systems	227
A.M. KUZNETSOV, J. ULSTRUP and M.G. ZAKARAYA / Protein Configurational Fluctuation Dependence of the Electronic Tunnel Factor of Modified Metalloprotein Electron Transfer Systems and in	

Fast Direct and Superexchange Separation and Recombination in Bacterial Photosynthesis	241
H. HEITEL, S. WEEREN, F. POLLINGER and M.E. MICHEL-BEYERLE / Solvent Polarity Effects on Thermodynamics and Kinetics of Intramolecular Electron Transfer Reactions	261
P.F. BARBARA, Tai Jong KANG, W. JARZEBKA and T. FONSECA / Solvation Dynamics and Ultrafast Electron Transfer	273
I. RIPS and J. JORTNER / Frequency Factor for the Outer-Sphere Electron Transfer. Bridging Between the Diabatic, Adiabatic and Solvent-Controlled Limits	293
D. HUPPERT and V. ITTAH / Static and Dynamic Electrolyte Effects on Excited State Intramolecular Electron Transfer and Excited-State Solvation	301
J. KLAFTER, A. BLUMEN and G. ZUMOFEN / Hierarchical Relaxation : Ultrametric Spaces	317
M. BIXON, J. JORTNER and M.E. MICHEL-BEYERLE / The Superexchange Model for the Primary Charge Separation in Bacterial Photosynthesis	325
R.A. FRIESNER and Y. WON / Role of Charge Transfer States of the Special Pair in Primary Photosynthetic Charge Separation	337
M.R. WASIELEWSKI, M.P. NIEMCZYK, D.G. JOHNSON, W.A. SVEC and D.W. MINSEK / Photosynthetic Model Systems that Address the Role of Superexchange in Electron Transfer Reactions	347
P.O.J. SCHERER and S.F. FISCHER / Electronic Excitations and Electron Transfer Coupling within the Bacterial Reaction Center Based on an INDO/S-CI Supermolecule Approach including 615 Atoms	361
A. SCHERZ, V. ROSENBAACH-BELKIN and J.R.E. FISHER / Self Assembly of Bacteriochlorophyll a and Bacteriopheophytin a in Micellar and Non-Micellar Aqueous Solutions ; Application to the Pigment-Protein Organization in Light-Harvesting Complexes and Reaction Centers	371

E.W. KNAPP and L. NILSSON / Protein Dynamics in the Photosynthetic Reaction Center : the Electron Transfer from Cytochrome c to the Special Pair	389
F. PARAK, A. BIRK, E. FROLOV, V. GOLDANSKII, I. SINNING and H. MICHEL : Mössbauer Spectroscopy on Photosynthetic Bacteria : Investigation of Reaction Centers of Rhodopseudomonas Viridis.	413
M. PLATO, K. MOBIUS, W. LUBITZ, J.P. ALLEN and G. FEHER / Magnetic Resonance and Molecular Orbital Studies of the Primary Donor States in Bacterial Reaction Centers	423
Index.	435

INFLUENCE OF CHROMOPHORES ON QUARternary STRUCTURE OF  
PHYCOBILIPROTEINS FROM THE CYANOBACTERIUM, *Mastigocladus*  
*lam.nosus*

R. FISCHER, J. GOTTSTEIN, S. SIEBZEHRÜBL,  
H. SCHEER  
Botanisches Institut der Universität  
Menzinger Str. 67  
D - 8000 München 19  
FRG

ABSTRACT. Chromophores of C-phycoerythrin and phycoerythrocyanin have been chemically modified by reduction to rubins, bleaching, photoisomerization, or perturbation with bulky substituents. Pigments containing modified chromophores, or hybrids containing modified and unmodified chromophores in individual protomers have been prepared. All modifications inhibit the association of the ( $\alpha\beta$ )-protomers of these pigments to higher aggregates. The results demonstrate a pronounced effect of the state of the chromophores on biliprotein quaternary structure. It may be important in phycobilisome assembly, and also in the dual function of biliproteins as (i) antenna pigments for photosynthesis and (ii) reaction centers for photomorphogenesis.

## 1. INTRODUCTION

Cyanobacteria, red algae and cryptophytes have specialized antenna pigments which enable them to harvest light efficiently in the green spectral region where chlorophylls have only poor absorption. These are the phycobiliproteins, extra-membraneous proteins carrying covalently bound open-chain tetrapyrrolic chromophores (Gantt, 1986; Glazer, 1983; MacColl and Guard-Friar, 1983; Scheer, 1982; Schirmer *et al.*, 1987; Wehrmeyer, 1983; Zuber, 1986). They are characterized by a remarkable degree of spectral adaptations covering the range from 480 to 670 nm. The spectral variation is based on the usage of only four different chromophore types, each of them being further modulated in its absorption properties by non-covalent interactions with the apoproteins.

In cyanobacteria and red algae, the phycobiliproteins show a high degree of organisation: Together with the linker

polypeptides, which are often devoid of chromophores, they form self-organizing microscopic structures, the phycobilisomes, which in turn form ordered arrays on the outer thylakoid membrane surface (Gantt, 1986; Glazer, 1983; Mörschel and Schatz, 1988; Wehrmeyer, 1983). By a combination of energetic and spatial ordering, and low internal conversion rates of the chromophores, the phycobilisomes are an antenna system working with quantum efficiencies approaching 100%.

The factors controlling tuning of the absorption, photochemistry and aggregation in biliproteins are presently becoming understood in considerable detail. This is based on an increasing body of structural data (Bishop *et al.*, 1987; Bryant, 1988; Düring and Huber, 1989; Glazer, 1983; Schirmer *et al.*, 1987; Tandeau de Marsac, Zuber, 1986) with theoretical (Sauer and Scheer, 1988; Schneider *et al.*, 1988, Scheer, 1987) and spectroscopic studies (Fischer *et al.*, 1988; Glazer, 1983; MacColl and Guard-Friar, 1987; Mimuro *et al.*, 1986a,b, Scheer, 1982; Schirmer *et al.*, 1987). They involve conformational changes of the chromophores imposed by non-covalent interactions with the apoproteins (e.g., point charges, and  $\pi$ - $\pi$ -interactions, and -in higher aggregates- excitonic interactions among chromophores (MacColl and Guard-Friar, 1983; Sauer and Scheer, 1988; Schneider *et al.*, 1988).

The assembly of phycobilisomes is a complex, and to a large extent autonomous process believed to be controlled by the presence of appropriate components. The factors controlling the aggregation of monomers ( $\alpha\beta$ ) to oligomers ( $\alpha\beta$ ) have hitherto mainly been located on the proteins. Here we want to present data which show a pronounced influence of the chromophores, too, on the aggregation. The results may also be important for an understanding of a the second function of biliprotein, e.g. as photomorphogenic reaction center pigments (Björn and Björn, 1980; Rüdiger and Scheer, 1983; Scheer, 1982; Song, 1988).

## 2. MATERIALS AND METHODS

C-phycoerythrin (PC) and phycoerythrocyanin (PEC) were isolated from the cyanobacterium, *Mastigocladus laminosus*, as described before (Fischer *et al.*, 1988). Subunits were prepared by isoelectric focusing (Schmidt *et al.*, 1988).

Chromophores were bleached by irradiation with 350 nm light (Scheer, 1987).

Reduction of cyanin to rubin chromophores was done by a modification of the method of Kufer and Scheer, 1982). Denatured PC or isolated subunits (8-25  $\mu$ M) in potassium phosphate buffer (0.9 M, pH = 7) were treated with NaBH<sub>4</sub> (final concentration  $\approx$ 170 mM) for 45 min at ambient temperature. The reaction was followed spectrophotometrically (decrease at  $\approx$ 600, increase at  $\approx$ 420 nm). After completion,

excess reagent was destroyed by addition of glucose. Modified PC was renatured by dialysis against phosphate buffer (100 mM, pH = 7) at ambient temperature. Hybridization to native protomers with modified chromophores on only one of the subunits, was done by combining stoichiometric amount of the appropriate subunits in buffer containing urea (8M), followed by dialysis against decreasing amounts of urea and finally urea-free buffer.

Cystein-111 of PC was modified by treatment with p-chloro-mercuribenzenesulfonate (PCMS) (Siebzehrübl *et al.*, 1987).

Photochemistry was induced with a cold light source (150W) equipped with light-guide and suitable interference filters ( $\approx 10$  nm fwhh). Spectra were recorded with lambda2 (Perkin-Elmer) or ZWSII (sigma) spectrophotometers. Aggregation state of samples was determined by ultracentrifugation (Martin and Ames, 1961) with myoglobin and trimeric phycocyanin as reference.

### 3. RESULTS AND DISCUSSION

#### 3.1. Effect of reduced chromophores on aggregation

Chromophores of PC were completely reduced with NaBH in the presence of urea (8M) to rubins, which show no absorption at wavelengths  $>500$ nm. After renaturation, the modified chromophores are unstable in the native protein environment and are slowly oxidized to cyanin chromophores within several days. The process can be followed quantitatively by monitoring the absorption increase at 620nm.

The aggregation state of phycocyanin containing reduced chromophores on both subunits is monomeric. There is only a single, yellow band present after centrifugation of freshly prepared "phycorubin". Under uv-light, this band is non-fluorescent. After prolonged standing of the samples, they become slowly green due to re-oxidation of the chromophores. Ultracentrifugation of such samples shows generally two bands: a greenish one at the position of monomers, and an additional one at the position of the PC trimer. The latter exhibits the characteristic absorption and fluorescence of PC trimer, and lacks the 420 nm band of the rubin chromophore(s) ( $\lambda_{max} \approx 420$  nm). It is at first only visible by its red fluorescence, and at increasing re-oxidation times becomes concentrated enough to become visible to the eye too. We, therefore, conclude that the lower band contains trimeric PC containing the re-oxidized native chromophores. If the aggregation pattern is followed through the reoxidation, the trimer band is always fluorescent and blue (if visible), whereas the monomer band is yellow or green. This shows clearly that modified PC bearing rubin chromophores can no longer aggregate.



In order to test if the inhibition of aggregation is due to specific chromophore(s), similar experiments were carried out with hybrids in which the  $\alpha$ -subunit was bearing a modified rubin chromophore and the  $\beta$ -subunit cyanin chromophores. For this purpose, isolated  $\alpha$ -subunits were modified, and then hybridized with the respective complementary unmodified subunits. The results can be summarized in one sentence: Whenever there was a rubin chromophores present in the hybrid, the hybrids were monomeric; and they did only aggregate when the chromophores became re-oxidized.

### 3.2. Effect of cystein-111 modification on aggregation

The single free cystein residue in PC, e.g.  $\beta$ -111 in the immediate vicinity of chromophore  $\beta$ -84, was modified with PCMS, and the aggregation studied at different protein concentrations. At low concentrations ( $<0.3 \mu\text{M}$ ), both PCMS-modified and unmodified PC were monomeric. Increasing concentrations shifted the equilibrium to trimers in unmodified PC, which at concentrations  $\geq 3 \mu\text{M}$  PC was mostly present as trimer. The PCMS-modified PC remained monomeric up to the highest concentrations investigated, e.g.  $10 \mu\text{M}$ .

This inhibition of aggregation was reversible, too. Treatment of PCMS-modified PC with an excess of dithiothreitol, removes the mercurial reagent from the chromoprotein as shown by the blue-shift of the spectrum. The difference spectra are mirror-images of the reaction with the mercurial, and the final spectrum of the recovered PC showed again the spectral characteristics of trimeric PC.

### 3.3. Effect of aggregation on photochemistry

#### 3.3.1. Phycocyanin

Long-lived photoproducts have been observed in several phycobiliproteins. In phycocyanin, a small but measurable photochemistry is found only upon partial denaturation and/or disaggregation of the native trimers. This photochemistry was originally studied in phycocyanin treated with urea at concentrations of 3-5M, which are known to change aggregation and to some extent also the tertiary structure. The difference spectra are characterized by a decrease around 622nm, and only a minor increase in the 500nm range (John *et al.*, 1985). A similar photochemistry (if judged from absorption difference spectra) has now been observed under a variety of mildly denaturing conditions. Its magnitude (defined by the ratio of the amplitude of the difference spectrum, to the maximum absorption) can be as high as 60% (in the presence of 20% mercaptoethanol, Schmidt *et al.*, 1988). From comparison with the photoreactions of phycocyanin in which the native 15Z-configured (Schirmer *et al.*, 1987) chromophores have been partially converted to the 15E-isomers (Schmidt *et al.*, 1988), it is most likely due to a Z/E interconversion of the chromophore(s) at the C-15,16 double-bond. Under such condi-

Figure 1: Difference spectrum of phycoerythrocyanin (pre-irradiated with green light, 500 nm), and the same sample after orange irradiation (600 nm).




Table 1 : Absorption difference extrema and amplitudes of different PEC samples or samples subjected to different pre-treatments.

Sample	C <sub>urea</sub> <sup>a)</sup> [M]	pH	Difference extrema [nm]		ΔΔA <sup>b)</sup> [%]
			orange	green	
PEC	0	7.0	570	503	18
PEC	1	7.1	569	503	32
PEC	4	7.3	567	502	35
PEC	6	7.5	600	510	12
PEC	8	7.0	599	515	6
PEC	8	3.0	598	507	13
PEC-monomer	c)	7.0	565	504	36
α-Subunit	0	7.0	569	504	50
phycobilisome	d)	7.0	567	502	0.31
phycobilisome, diss	e)	7.0	567	504	2.82

a) Samples in potassium phosphate buffer (100 mM, pH = 7), if not indicated otherwise.

b) ΔΔA is the amplitude of the s-shaped difference spectrum from minimum to maximum, divided by the maximum absorbance of the sample (see Fig.1).

c) 1 M KSCN, no urea.

d) Coupled phycobilisomes, in phosphate buffer (0.9 M).

e) Dissociated phycobilisomes, in distilled water.

tions, the chromophores are apparently capable to perform the same type of photochemistry as that of phytochrome, but the subsequent changes -if there are any- must of course be different (as are the proteins).

### 3.3.2. Phycoerythrocyanin

Phycocyanin shows photochemistry only under partially denatured conditions and is photochemically "silent" in the native as well as in the fully denatured state. A closely related pigment, phycoerythrocyanin (PEC), shows a pronounced photochemistry under all conditions from fully native (Fig. 1) to fully denatured (Siebzehrübl et al., 1989). There is nonetheless a pronounced effect of aggregation on the amplitude of the photochemical difference spectrum. It is very small in phycobilisomes, increases upon dissociation to trimers and further to monomers, and then decreases again in denatured PEC. Thus, the basic pattern is similar in PC and PEC, but the amplitudes of the difference spectra are always much larger in PEC (Table 1) than in PC (John et al., 1985).

The  $\alpha$ -subunit of PEC had been linked previously to photochromic activities in cyanobacterial extracts, and possibly to photomorphogenesis (Björn and Björn, 1980; Kufer, 1988). This pigment, which is structurally very similar to PC (Bryant, 1982; Düring and Huber, 1989) carries a rare phycoviolobin chromophore (alternatively also called PXB or cryptoviolin) at cystein  $\alpha$ -84 (Bishop et al., 1987), which replaces the common phycocyanobilin chromophore present at the same location in PC. Being a component of the phycobilisome, it is commonly regarded a light-harvesting pigment. A distinct difference from other phycobiliproteins, is however its pronounced photochemistry in the native state. It involves probably a Z/E-isomerization at the C-15 methine bridge similar to the photoreaction of the chromophore in phytochrome (Thümmler et al., 1983), but this still has to be demonstrated. This reaction would require a decreased rigidity in the environment of  $\alpha$ -84, which (contrary to preliminary indications) is not obvious in the crystal structure of (trimeric) PEC (Düring and Huber, 1989).

### 3.4. Effect of chromophore $\alpha$ -84 photochemistry in PEC on aggregation

During photochemical studies with PEC, we noticed that it shows not only increased photochemistry upon disaggregation, but that there is also a reciprocal dependence of biliprotein aggregation on photochemistry. When PEC is alternately irradiated with orange (600nm) and green light (500nm), the  $\alpha$ -subunit becomes enriched in one (15E) and the other form (15Z), respectively. Ultracentrifugation analysis showed, that at the same time there occurs a photoreversible change in aggregation: The amount of trimer increased each time the last irradiation was performed with green light, and decreased each time it was

performed with orange light. This means, that the configuration of  $\alpha$ -84 influences aggregation, a fact which can be rationalized from the x-ray structure:  $\alpha$ -84 is located very close to the contact surface of monomers in trimers (Düring and Huber, 1989).

#### 4. Discussion

In many cyanobacteria, profound alterations in phycobilisome structure (and other cyanobacterial activities) can be induced by changes in the environmental light conditions. The photomorphogenetic receptors (adaptachromes, photomorphochromes) are hitherto unknown, but action spectra suggest that they also belong to the biliproteins (Björn and Björn, 1980; Scheer, 1982). Pigment(s) of the biliprotein type, albeit with very different apoproteins, are functional in higher plants for the same purpose. The green-plant photomorphogenetic reaction center pigment, phytochrome, has a chromophore which is very similar to the ones of the light harvesting pigment, phycocyanin (Rüdiger and Scheer, 1983). This multiple function of bile pigment chromophores is an aspect of biliproteins which is still poorly understood. The first event in photomorphogenetic pigments, is generally believed to be a structural change of the chromophore(s). This signal can then be propagated by induced structural changes of the protein, and beyond. In phytochrome, the primary reaction has been shown to involve a photoreversible Z/E-isomerization and possibly a protonation/deprotonation of the chromophore, while the induced changes on the protein are less understood (Rüdiger and Scheer, 1983; Song, 1988; Thümmler *et al.*, 1983).

The multiple function of biliproteins containing very similar chromophores, is reminiscent of the multiple functions performed by chlorophylls of the same molecular structure in photosynthesis. In both cases, the photochemical and photophysical properties of the chromophores are diverse. The relative contributions of different de-excitation pathways must be regulated by specific interactions between the chromophores and the proteins. In analogy to interactions among living systems, these interactions can be termed as "molecular ecology".

#### 5. SUMMARY

The results obtained by different modifications of the chromophores or their immediate vicinity, demonstrate an involvement of the biliprotein chromophores not only in energy transfer and photochemistry of biliproteins, but also in their quaternary structure. This effect may well be at the origin of a signal chain leading eventually to photomorphogenesis. In a more general context, it is an example for the intricate interplay of proteins with their cofactors, which leads to the stunning variety of properties of pigments with the same or very similar molecular structures.

## References:

- Bishop, J. E., H. Rapoport, A. V. Klotz, C. F. Chan, A. N. Glazer, P. Füglistaller, and H. Zuber, Chromopeptides from phycoerythrocyanin. Structure and linkage of the three bilin groups. *J. Am. Chem. Soc.* **109**, 875-881.
- Björn, G. S. (1979) Action spectra for *in vivo* and *in vitro* conversions of Phycochrome b, a reversibly photochromic pigment in a blue-green alga, and its separation from other pigments. *Physiol. Plant.* **46**, 281-286.
- Björn, G. S., N. Ekelund, and L. O. Björn (1984) Light-induced linear dichroism in photoreversibly photochromic sensor pigments. 4. Lack of chromophore rotation in phycochrome-B immobilized *in vitro*. *Physiol. Plant.* **60**, 253-256.
- Björn, L.O., and Björn, G.S. (1980), Photochromic Pigments and photoregulation in blue-green algae, *Photochem. Photobiol.*, **32**, 849-852
- Braslavsky, S. E., A. R. Holzwarth, and K. Schaffner (1983) Solution conformations, photophysics and photochemistry of bili-pigments - bilirubin and biliverdin dimethyl esters and related linear tetrapyrroles. *Angew. Chem. Int. Ed.* **22**, 656-674.
- Bryant, D. A. (1982) Phycoerythrocyanin and Phycoerythrin - properties and occurrence in Cyanobacteria. *J. Gen. Microbiol.* **128**, 835-844.
- Bryant, D. A. (1988) Phycobilisomes of *Synechococcus* sp. PCC 7002, *Pseudoanabaena* sp. PCC 7409, and *Cyanophora paradoxa*: An analysis by molecular genetics. In *Photosynthetic Light Harvesting Systems*. (Eds. H. Scheer and S. Schneider), pp. 217-233, De Gruyter, Berlin.
- Düring, M. and R. Huber (1989) private communication
- Fischer, R., S. Siebzehrübl, and H. Scheer (1988) C-phycoyanin from *Mastigocladus laminosus*: Chromophore Assignment in higher Aggregates by Cystein Modification. In *Photosynthetic Light Harvesting Systems: Organization and Function* (Eds. H. Scheer and S. Schneider), pp. 71-76, De Gruyter, Berlin.
- Gantt, E. (1986) Phycobilisomes. In *Photosynthesis III* (Eds. L. A. Staehelin and C. J. Arntzen), pp. 260-268, Springer-Verlag, Berlin.
- Glazer, A. N. (1983) Comparative biochemistry of photosyn-

thetic light-harvesting systems. *Ann. Rev. Biochem.* 52, 125-157.

John, W., R. Fischer, S. Siebzehnrübl, and H. Scheer (1985) C-Phycocyanin from *Mastigocladus laminosus*. Isolation and properties of subunits and small aggregates. In *Antenna and Reaction Centers of Photosynthetic Bacteria - Structure, Interactions and Dynamics* (Ed. M. E. Michel-Beyerle), pp. 17-25, Springer, Berlin.

Kußer, W. (1988) Concerning the relationship of light harvesting Biliproteins to Phycochromes in cyanobacteria. In *Photosynthetic Light Harvesting Systems*. (Eds. H. Scheer and S. Schneider), pp. 89-93, De Gruyter, Berlin.

Kußer, W. and H. Scheer (1982) Rubins and rubinoid addition products from phycocyanin. *Z. Naturforsch.* 37c, 179-192.

MacColl, R. and D. Guard-Friar (1987) *Phycobiliproteins*. CRC Press, Boca Raton.

MacColl, R., K. Csatorday, and D. S. Berns (1981) The relationship of the quaternary structure of Allophycocyanin to its spectrum. *Arch. Biochem. Biophys.* 208, 42-48.

Martin, R. G. and B. N. Ames (1961) A method for determining the sedimentation behavior of enzymes: Application to protein mixtures. *J. Biol. Chem.* 236, 1372-1379.

Mimuro, M., P. Füglistaller, R. Rübli, and H. Zuber (1986 a) Functional assignment of chromophores and energy transfer in C-phycocyanin isolated from the thermophilic cyanobacterium *Mastigocladus laminosus*. *Biochim. Biophys. Acta* 848, 155-166.

Mimuro, M., R. Rübli, P. Füglistaller, and H. Zuber (1986 b) The microenvironment around the chromophores and its changes due to the association states in C-phycocyanin isolated from the cyanobacterium *Mastigocladus laminosus*. *Biochim. Biophys. Acta* 851, 447-456.

Mörschel, E. and G.-H. Schatz (1988) On the structure of photosystem II-Phycobilisome complexes of cyanobacteria. In *Photosynthetic Light Harvesting Systems*. (Eds. H. Scheer and S. Schneider), pp. 21-35, De Gruyter, Berlin.

Rüdiger, W. and H. Scheer (1983) Chromophores in Photomorphogenesis. In *Photomorphogenesis - Encyclopedia of Plant Physiology*, New Series. (Eds. W. Shropshire jr. and H. Mohr), Vol. 16, pp. 119-151, Springer, Heidelberg.

Sauer, K., and H. Scheer (1988) Excitation transfer in C-phycocyanin: Förster transfer rate and exciton calculations based on new crystal structure data for C-phycocyanins from *Agmenellum quadruplicatum* and *Mastigocladus laminosus*, *Biochim. Biophys. Acta*.

- Scheer, H. (1982) Light reaction Path of Photosynthesis. In *Phycobiliproteins: Molecular Aspects of Photosynthetic Antenna Systems* (Ed. F. K. Fong), pp. 7-45, Springer, Springer.
- Scheer, H. (1987) Photochemistry and Photophysics of C-phycocyanin. In *Progress in Photosynthesis Research* (Ed. J. Biggins), pp. 1.1.143-149, Martinus Nijhoff, Dordrecht.
- Schirmer, T. and M. G. Vincent (1986) Polarized absorption and fluorescence spectra of single crystals of C-phycocyanin. *Biochim. Biophys. Acta* **893**, 379-385.
- Schirmer, T., W. Bode, and R. Huber (1987) Refined Three-dimensional structures of 2 cyanobacterial C-phycocyanins at 2.1. and 2.5 Å resolution - A common principle of phycobilinprotein interaction. *J. Mol. Biol.* **196**, 677-695.
- Schmidt, G., S. Siebzehrübl, R. Fischer, and H. Scheer (1988) Photochromic properties of C-phycocyanin. In *Photosynthetic Light Harvesting Systems. Organization and Function* (Eds. H. Scheer and S. Schneider), pp. 77-89, De Gruyter, Berlin.
- Schneider, S., C. Scharnagl, M. Düring, T. Schirmer, and W. Bode (1988) Effect of protein environment and the excitonic coupling on the excited-state properties of the bilinchromophores in C-phycocyanin. In *Photosynthetic Light Harvesting Systems*, (Eds. H. Scheer and S. Schneider), pp. 483-491, De Gruyter, Berlin.
- Schneider, S., P. Geiselhart, F. Baumann, S. Siebzehrübl, R. Fischer, and H. Scheer (1988) Energy transfer in "native" and chemically modified C-phycocyanin trimers and the constituent subunits. In *Photosynthetic Light Harvesting Systems. Organization and Function* (Eds. H. Scheer and S. Schneider), pp. 469-483, De Gruyter, Berlin.
- Siebzehrübl, S., R. Fischer, and H. Scheer (1987) Chromophore Assignment in C-phycocyanin from *Mastigocladus laminosus*. *Z. Naturforsch.* **42c**, 258-262.
- Siebzehrübl, S., R. Fischer, W. Kufer, and H. Scheer (1989) Photochemistry of phycobiliproteins: reciprocity of reversible photochemistry and aggregation in phycoerythrocyanin from *Mastigocladus laminosus*. *Photochem. Photobiol.* **49**, 753 - 762.
- Song, P. S. (1988) The molecular topography of Phytochrome -Chromophore and Apoprotein. *J. Photochem. Photobiol.* **2**, 43-57.
- Tandeau de Marsac, N. and G. Cohen-Bazire (1977) Molecular composition of bacterial Phycobilisomes. *Proc. Natl. Acad.*

*Sci. USA* **74**, 1635.

Thümmeler, F., W. Rüdiger, E. Cmiel, and S. Schneider (1983) Chromopeptides from phytochrome and phycocyanin. Nuclear magnetic resonance studies of the P<sup>+</sup> and P<sup>-</sup> chromophore of phytochrome and E,Z isomeric<sup>+</sup> chromophores of phycocyanin. *Z. Naturforsch.* **38 c**, 359-368.

Wehrmeyer, W. (1983) Phycobiliproteins and phycobiliprotein organization in the photosynthetic apparatus of cyanobacteria, red algae, and cryptophytes. In *Proteins and nucleic acids in plant systematics* (Eds. U. Jensen and D. E. Fairbrother), Springer, Berlin.

Zuber, H. (1986) Primary structure and function of the light-harvesting polypeptides from cyanobacteria, red algae and purple photosynthetic bacteria. In *Photosynthesis III* (Eds. L. A. Staehelin and C. J. Arntzen), pp. 238-251, Springer, Berlin.