

85-1

ZEITSCHRIFT FÜR NATURFORSCHUNG

SECTION C BIOSCIENCES

Council

E. BÜNNING, Tübingen
A. BUTENANDT, München
M. EIGEN, Göttingen

Editorial Board

A. HAGER, Tübingen
W. HASSELBACH, Heidelberg
P. KARLSON, Marburg
F. KAUDEWITZ, München
E. WECKER, Würzburg

Advisory Editorial Board

P. BÖGER, Konstanz
D. BÜCKMANN, Ulm
K. G. GÖTZ, Tübingen
G. GOTTSCHALK, Göttingen
H. HOFFMANN-BERLING, Heidelberg
R. JAENICKE, Regensburg
G. F. MEYER, Tübingen
M. RAJEWSKY, Essen

H. SCHIMASSEK, Heidelberg
D. SCHULTE-FROHLINDE, Mülheim/R
F. F. SEELIG, Tübingen
J. SEELIG, Basel
H. SIMON, München
W. STEGLICH, Bonn
A. TREBST, Bochum

EDITED IN COLLABORATION
WITH THE INSTITUTES OF THE MAX-PLANCK-GESELLSCHAFT

VOLUME 37c

NUMBER 1/2

JANUARY/FEBRUARY 1982

VERLAG DER ZEITSCHRIFT FÜR NATURFORSCHUNG
TÜBINGEN

37c, 1-2
1982

3 20

WL

Anschrift des Verlages: Postfach 2645, D-74 Tübingen
Satz und Druck: Konrad Triltsch, Würzburg

Nachdruck — auch auszugsweise — nur mit schriftlicher Genehmigung des Verlags



Section a
Physics, Physical Chemistry, Cosmic Physics

Section b
Inorganic and Organic Chemistry

Contents

Contents of Number 1/2

Original Communications

- Antibiotics from Basidiomycetes. XV[1]. 1-Hydroxy-2-nonyl-4-one, an Antifungal and Cytotoxic Metabolite from *Ischnoderma benzoinum* (Wahl.) Karst. (In German)
T. ANKE, B.-M. GIANNETTI, and W. STEGLICH 1
- Vinyl-Olefines and Sesquiterpenes in the Root-Oil of *Senecio isatideus*
W. BOLAND, L. JAENICKE, and A. BRAUNER 5
- Carotenoid Composition of Chlorophyll-Carotenoid-Proteins from Radish Chloroplasts
H. K. LICHTENTHALER, U. PRENZEL, and G. KUHN 10
- Carotenoid Biogenesis in the Stick Insect, *Carausius morosus*, during a Larval Instar
H. KAYSER 13
- Hydroxylation of the B-Ring of Flavonoids in the 3'- and 5'-Position with Enzyme Extracts from Flowers of *Verbena hybrida*
G. STOTZ and G. FORKMANN 19
- Dissociation of FAD from The NAD(P)H-Nitrate Reductase Complex from *Ankistrodesmus braunii* and Role of Flavin in Catalysis
M. A. DE LA ROSA, A. J. MÁRQUEZ, and J. M. VEGA 24
- Spectrophotometric Isolation of Kinetically Different Pools of P-700 and Their Correlation to the Reduction of NADP by Isolated Chloroplasts. I. The Effect of Light Quality and Intensity
H. J. RURAINSKI, R. GERHARDT, and G. MADER 31
- Chemical Aspects of the Mutagenic Activity of the Ascorbic Acid Autoxidation System
W. H. KALUS, W. G. FILBY, and R. MÜNZNER 40
- Mevinolin: A Highly Specific Inhibitor of Microsomal 3-Hydroxy-3-methylglutaryl-Coenzyme A Reductase of Radish Plants
T. J. BACH and H. K. LICHTENTHALER 46
- Free Activation Energies and Activation Volumes for the Amide Rotation in Some Peptides Studied by High Pressure ¹H-High Resolution NMR
H. HAUER, H.-D. LÜDEMANN, and R. JAENICKE 51
- Investigation of Large Intramolecular Movements within Metmyoglobin by Rayleigh Scattering of Mössbauer Radiation (RSMR)
Y. F. KRUPYANSKII, F. PARAK, V. I. GOLDANSKII, R. L. MÖSSBAUER, E. E. GAUBMAN, H. ENGELMANN, and I. P. SUZDALEV 57
- On the Free Radical-Induced Aggregation of Ribonuclease - A Pulse Radiolysis Study Using the Light Scattering Detection Method
H. SEKI and W. SCHNABEL 63
- Dark Oxidation of Unsaturated Lipids by the Photoxidized 8-Methoxypsoralen
A. Ya. POTAPENKO, M. V. MOSHNIN, A. A. KRASNOVSKY JR., and V. L. SUKHORUKOV 70
- Growth Specific Thylakoid Differentiation of the Endocyanelle *Cyanocyta korschikoffiana* in the Symbiotic Association with *Cyanophora paradoxa* (In German)
P. BRANDT, S. MARTEN, B. MÜLLER, and W. WIESSNER 75
- Differential Fluorescence and Kinetic Studies on the Template-Binding of RNA Polymerase from Parsley and *Escherichia coli*
K. GROSSMANN, H. BISSWANGER, and H. U. SEITZ 81
- Sensitivity of *Escherichia coli* to Viral Nucleic Acid. XVI. Temperature Conditions for Ca²⁺ Dependent DNA Uptake in *Escherichia coli*
A. TAKETO 87
- Studies on ¹⁸O₂-Uptake in the Light by Entire Plants of Different Tobacco Mutants
R. ISHII and G. H. SCHMID 93
- Immobilization of Yeast Cells by Radiation-Induced Polymerization
T. FUJIMURA and I. KAETSU 102

<i>Nanochlorum eucaryotum</i> : a Very Reduced Coccoid Species of Marine Chlorophyceae C. WILHELM, G. EISENBEIS, A. WILD, and R. ZAHN	107	A New Catechin Glycoside from <i>Polypodium vulgare</i> (In German) C. KARL, G. MÜLLER, and P. A. PEDERSEN	148
Growth and Photosynthesis of <i>Nanochlorum eucaryotum</i> , a New and Extremely Small Eucaryotic Green Alga C. WILHELM and A. WILD	115	On the Essential Oil Components from Blossoms of <i>Artemisia vulgaris</i> L. (In German) K. MICHAELIS, O. VOSTROWSKY, H. PAULINI, R. ZINTL, and K. KNOBLOCH	152
Kinetics of Proton-Hydroxyl Transport across Lecithin Vesicle Membranes as Measured with a Lipoid pH-Indicator W. G. POHL	120	Phenazine and Phenoxazinone Biosynthesis in <i>Brevibacterium iodinum</i> R. B. HERBERT, J. MANN, and A. RÖMER	159
<i>Notes</i>		Separation, Partial Purification and Characterization of a Fatty Acid Hydroperoxide Cleaving Enzyme from Apple and Tomato Fruits P. SCHREIER and G. LORENZ	165
¹ H-NMR- and ESR-Investigations on the Structures of Dehydroascorbic Acid and the Semidehydroascorbate Radical (In German) H. SAPPER, A. PLEYER-WEBER, and W. LOHMANN	129	The Site of Indole-3-acetic Acid Synthesis in Mesophyll Cells of <i>Spinacia oleracea</i> B. HEILMANN, W. HARTUNG, and H. GIMMLER	174
Hydroxyrutacridone Epoxide, a New Acridone Alkaloid from <i>Ruta graveolens</i> (In German) U. EILERT, B. WOLTERS, A. NAHRSTEDT, and V. WRAY	132	Rubins and Rubinoid Addition Products from Phycocyanin W. KUFER and H. SCHEER	179
Flavonol 3-O-Methyltransferase in Plant Tissues V. DE LUCA, G. BRUNET, H. KHOURI, R. IBRAHIM, and G. HRAZDINA	134	Isolation of a <i>b</i> -Type Cytochrome Oxidase from Membranes of the Phototrophic Bacterium <i>Rhodospseudomonas capsulata</i> H. HÜDIG and G. DREWS	193
Enzymatic Hydroxylation of Ruthenocen and Osmocene (In German) M. SCHNEIDER, M. WENZEL, and G. SCHACHSCHNEIDER	136	Tetrapyrrol Derivatives Shown by Fluorescence Emission and Excitation Spectroscopy in Cells of <i>Rhodospseudomonas capsulata</i> Adapting to Phototrophic Conditions J. BECK and G. DREWS	199
Neutral Red Fluorescence of Chromatin: Specificity and Binding Mechanism R. H. ESPELOSIN and J. C. STOCKERT	139	Formation of Hydrocarbons by Photobleaching Cyanobacterium, <i>Anacystis nidulans</i> G. SCHMETTERER	205
Proteinase-Inhibitors in Albumin Glands of <i>Achatina fulica</i> R. O. OKOTORE and G. UHLENBRUCK	142	Peptidoglycan and Protein, the Major Cell Wall Constituents of the Obligate Halophilic Bacterium <i>Rhodospirillum salexigens</i> M. H. TADROS, G. DREWS, and D. EVERS	210
Erratum	144	CO ₂ Fixation in <i>Anabaena cylindrica</i> G. DÖHLER	213
Contents of Number 3/4		On the Occurrence of Monoacylglycerol Derivatives in Lipid Metabolism of Chloroplasts A. SAUER and K.-P. HEISE	218
<i>Original Communications</i>			
Structure Elucidation of Pterosupin from <i>Pterocarpus marsupium</i> , the First Naturally Occurring C-Glycosyl- β -hydroxy-dihydrochalcone D. ADINARAYANA, K. V. SYAMASUNDAR, O. SELIGMANN, and H. WAGNER	145		

Possible Models of Murein and Their Fourier Transforms H. FORMANEK	226	The Influence of Detergents on the Ca ²⁺ - and Mg ²⁺ -Dependent Adenosine Triphosphatase of the Sarcoplasmic Reticulum H. LÜDI, B. RAUCH, and W. HASSELBACH	299
Binding of Antibodies onto the Thylakoid Membrane. VII. Localization of Coupling Factor of Photophosphorylation in the Lamellar System of Chloroplasts from <i>Antirrhinum majus</i> A. RADUNZ and R. MEIER	236	Inhibition of Acetyl CoA Carboxylase by a High Molecular Weight Protein in Rat Liver M. N. ABDEL-HALIM and S. Y. K. YOUSUFZAI	308
Photophosphorylation by Chloroplasts: Effects of Low Concentrations of Ammonia and Methylamine C. GIERSCH	242	Subcellular Fractionation of Bone Marrow-Derived Macrophages: Localization of Phospholipase A ₁ and A ₂ and Acyl-CoA:1-Acylglycero-3-phosphorylcholine-0-acyltransferase E. E. KRÖNER, H. FISCHER, and E. FERBER	314
Digitonin Precipitable Sterols Inducing the Aggregation of Prolamellar Body-Like Structures from Completely Dissolved PLB Components A. NISIUS, M. MÜLLER, and H. G. RUPPEL	251	Periodicities of Dinucleotide Self-Information Values in Φ X174 DNA N. B. FURLONG and C. F. BECKNER	321
Isolation of PS II-Particels with <i>in vivo</i> Characteristics from <i>Euglena gracilis</i> , Stamm Z (In German) F. SCHULER, P. BRANDT and W. WIESSNER	256	Growth Characteristics of <i>in vitro</i> Cultured Ehrlich Ascites Tumor Cells under Anaerobic Conditions and after Rearation R. MERZ and F. SCHNEIDER	326
Absorption and Picosecond Fluorescence Characteristics of Chlorophyll Vesicles as a Function of Temperature S. S. BRODY	260		
Herbicides which Inhibit Electron Transport or Produce Chlorosis and Their Effect on Chloroplast Development in Radish Seedlings. I. Chlorophyll a Fluorescence Transients and Photosystem II Activity K. H. GRUMBACH	268	<i>Notes</i>	
Use of Isolated Leaf Cells of <i>Abutilon theophrasti</i> to Localize the Action of Two Aminotriazinone Herbicidal Derivatives K. K. HATZIOS	276	3-O-Methyl, 8-C-methyl Quercetin a New Flavonoid from <i>Dasylirion acrotrichum</i> and <i>Xanthorrhoea hastilis</i> , Liliales (In German) C. LARACINE, J. FAVRE-BONVIN, and P. LEBRETON	335
Structure Investigations of Agonists of the Natural Neurotransmitter Acetylcholine II [1]. X-ray Structure Analysis of Trimethyl(4-oxopentyl)-ammonium-chloride A. GIEREN and M. KOKKINIDIS	282	New Flavonoids from the Farina of <i>Pityrogramma</i> Species C. HITZ, K. MANN, and E. WOLLENWEBER	337
Bile Salt Delipidation, Residual Phospholipids and Reactivation of the Ca ²⁺ -ATPase from Sarcoplasmic Reticulum G. SWOBODA and W. HASSELBACH	289	Actinioerythrin in <i>Mullus barbatus</i> L. (Pisces; Teleostei; Mullidae) (In German) F.-C. CZYGAN and A. KRÜGER	340
		The Importance of the Position of the Adenosine Linkage in NAD ⁺ for Its Coenzyme Activity <i>via</i> Lactate Dehydrogenase (In German) E. SCHLIMME, R. MATTERN, and E. SCHOTT	342
		Characterization of the Nuclear Lamina in an Insect by Differential Staining R. MARX	345

Trail Orientation of Wood Ants (<i>Formica polyctena</i> Förster) (In German) K. HORSTMANN	348	Proton NMR-Relaxation Dispersion in Meconium Solutions and Healthy Amniotic Fluid: Possible Applications to Medical Diagnosis G. J. BÉNÉ, B. BORCARD, V. GRAF, E. HILTBRAND, P. MAGNIN, and F. NOACK	493
Erratum	350	Monospecific Antibody against 5-Methyl-Cytidine for the Structural Analysis of Nucleic Acids S. K. MASTRONICOLIS, V. M. KAPOULAS, and H. KRÖGER	399
Contents of Number 5/6			
<i>Original Communications</i>			
Structure of Echinacoside (In German) H. BECKER, W. CH. HSIEH, R. WYLDE, C. LAFFITE, and C. ANDARY	351	On the Oxidative Main-Chain Scission of Nucleic Acids — Pulse Radiolysis Studies (In German) O. DENK and W. SCHNABEL	405
Synthesis of Naturally Occurring Prenylated Naphthalene Derivates. Isolation of a New Prenyl-naphthoquinone from <i>Galium mollugo</i> (In German) L. HEIDE and E. LEISTNER	354	Binding Constants, Binding Enthalpies and Entropies of the Non-Competitive and the Competitive Binding of Acriflavine, Tetramethylacriflavine and Acridine Orange to DNA (In German) H. WILLE, J. PAULUHN, and H. W. ZIMMERMANN	413
Biosynthesis of Isoflavonoid Phytoalexins: Incorporation of Sodium [1,2- ¹³ C ₂]Acetate into Phaseollin and Kievitone P. M. DEWICK, M. J. STEELE, R. A. DIXON, and I. M. WHITEHEAD	363	Possible Orientation of the Fatty Acid Chains in Lipopolysaccharide H. FORMANEK	428
Diurnal Fluctuation of Quinolizidine Alkaloid Accumulation in Legume Plants and Photomixotrophic Cell Suspension Cultures M. WINK and T. HARTMANN	369	Screening for Highly Active Plasmid Promoters via Fusion to β -Galactosidase Gene A. ROSNER, M. GORECKI, and H. AVIV	441
Volatile Compounds from Cephalic Secretions of Females in two Cleptoparasite Bee Genera, <i>Epeolus</i> (Hym., Anthophoridae) and <i>Coelioxys</i> (Hym., Megachilidae) J. TENGÖ, G. BERGSTRÖM, A.-K. BORG-KARLSON, I. GROTH, and W. FRANCKE	376	Anomalous Reduction of Cytochrome <i>b</i> in Highly Purified Complex III from Baker's Yeast F. F. DE LA ROSA and G. PALMER	445
Purification and Properties of Glucosaminephosphate Isomerase of <i>Proteus mirabilis</i> B. CIFUENTES and C. VICENTE	381	Changes in the Fluorescence Emission Spectrum of <i>Chlorella emersonii</i> Induced by Cold Treatment; a Possible Regulative Feature of Energy Uptake G. HARNISCHFEGGER and H. JARRY	448
Effects of Trypan Blue and Related Compounds on Production and Activity of Streptolysin S Y. TAKEO and A. TAKEO	385	Studies of Photosynthesis Inhibition by Phytoluminography R. BLAICH, O. BACHMANN, and I. BAUMBERGER	452
Antiviral Activity of 2-(α -Hydroxybenzyl)-benzimidazole and other 2-Substituted Benzimidazoles against TMV in Tomato Leaf Discs A. C. CASSELLS and F. M. COCKER	390	A Study of State Changes in <i>Chlorella</i> : The Effect of Uncoupler and Energy Transfer Inhibitors P. V. SANE, D. FURTADO, T. S. DESAI, and V. G. TATAKE	458
		Adaptation of Chloroplast-Ultrastructure and of Chlorophyll-Protein Levels to High-Light and Low-Light Growth Conditions H. K. LICHTENTHALER, G. KUHN, U. PRENZEL, C. BUSCHMANN, and D. MEIER	464

Protochlorophyllide Photoconversion Initiates the Transformation of Reaggregated Prolamellar Body Tubules <i>in vitro</i> M. MÜLLER and A. NISIUS	476	Hyperglycemia in the Fresh Water Field Crab (<i>Oziotelphusa senex senex</i>) Produced by a Pesticide (BHC) P. SREENIVASULA REDDY, S. B. RAMESH BABU, and R. RAMAMURTHI	545
The Sarcoplasmic Reticulum of Smooth Muscle Fibers L. RAEYMAEKERS	481	Coenzyme Binding at Different Ionization States of Cytoplasmic and Mitochondrial Malate Dehydrogenase K. SCHWERDTFEGER, C. WOENCKHAUS, D. M. PARKER, and J. J. HOLBROOK	547
Histochemical Demonstration of an ATP-Dependent Ca ²⁺ -Pump in Bullfrog Myocardial Cells R. MEYER, W. STOCKEM, M. SCHMITZ, and H. G. HAAS	489	A New Method to Prepare Membrane Fractions Containing Ionophore-Stimulated ATPase from Pumpkin Hypocotyls (<i>Cucurbita maxima</i> , L.) G. F. E. SCHERER	550
Ca Entry and Contraction as Studied in Isolated Bovine Ventricular Myocytes G. ISENBERG	502		
Fluorescence Energy Transfer between ATPase Monomers in Sarcoplasmic Reticulum Reconstituted Vesicles, in the Presence of Low Concentrations of a Nonionic Detergent P. CHAMPEIL, J.-L. RIGAUD, and M. P. GINGOLD	513		
Functional Significance of Quaternary Organization of the Sarcoplasmic Reticulum Ca ²⁺ -ATPase J. V. MØLLER, T. S. MAHROUS, J. P. ANDERSEN, and M. LE MAIRE	517		
Equilibrium Constants for Some Steps of the Reaction Cycle of the Sarcoplasmic Reticulum Calcium Pump C. TANFORD and D. W. MARTIN	522		
Phosphorylation of Ca ²⁺ -ATPase by Inorganic Phosphate in Water-Organic Solvent Media: Dielectric Constant and Solvent Hydrophobicity Contribution A. DE SOUZA OTERO and L. DE MEIS	527		
ATPases of the Cat Carotid Body and of the Neighbouring Ganglia H. STARLINGER	532		
<i>Notes</i>			
Synthesis of 6-Chlorogenistein A. LÉVAI and A. L. TÖKÉS	540		
Effect of Penicillins on the Level of Inorganic Pyrophosphatase in <i>Escherichia coli</i> K 12 E. I. KUKKO and J. K. HEINONEN	542		
		Contents of Number 7/8	
		<i>Original Communications</i>	
		ESR Studies on Platinum Uracil Blue H. NEUBACHER, M. SEUL, and W. LOHMANN	553
		Roseanolone: A New Diterpene from <i>Hypoestes rosea</i> J. I. OKOGUN, A. A. ADESOMOJU, G. A. ADESIDA, H. J. LINDNER, and G. HABERMEHL	558
		6-C- α -L-Rhamnopyranosylapigenin 7-O- β -D-glucopyranoside (Isofurcatain 7-O- β -D-glucoside), a New Flavone Glycoside from <i>Metzgeria furcata</i> K. R. MARKHAM, R. THEODOR, R. MUES, and H. D. ZINSMEISTER	562
		On the Essential Oil Components from <i>Humulus lupulus</i> L. var. <i>neomexicanus</i> Nels. & Cockerell. I. Contribution K. KNOBLOCH, H. PAULINI, C. ELEY, J. H. ELEY, E. ZIEGLER, H. BRANDAUER, K. MICHAELIS, and O. VOSTROWSKY	565
		Demethylation, Methylation and 3'-Hydroxylation of Isoflavones by <i>Fusarium</i> Fungi K.-M. WELTRING, K. MACKENBROCK, and W. BARZ	570
		Accumulation of Acridone-Epoxides in Callus Cultures of <i>Ruta graveolens</i> Increased by Coculture with Not Host-Specific Fungi (In German) B. WOLTERS and U. EILERT	575

- Precursors of Cyclic Polysulphides in Seeds of *Parikia speciosa* Hassk. (In German)
R. SUSILO and R. GMELIN 584
- Formation of Flavonol 3-O-Diglycosides and Flavonol 3-O-Triglycosides by Enzyme Extracts from Anthers of *Tulipa* cv. Apeldoorn
G. KLEINEHOLLENHORST, H. BEHRENS, G. PEGELS, N. SRUNK, and R. WIERMANN 587
- Polynucleotide Phosphorylase from a Cyanobacterium (*Synechococcus* sp.): Subunit Composition and Properties
W.-T. NOLDEN and G. RICHTER 600
- DEAE-Dextran Induced Increase of Membrane Permeability and Inhibition of Photosynthesis in *Dunaliella parva*
H. GIMMLER and G. LOTTER 609
- High- and Low-Affinity Binding of Photosystem II Herbicides to Isolated Thylakoid Membranes and Intact Algal Cells
H. LAASCH, K. PFISTER, and W. URBACH 620
- Herbicides which Inhibit Electron Transport or Produce Chlorosis and Their Effect on Chloroplast Development in Radish Seedlings. II. Pigment Excitation, Chlorophyll Fluorescence and Pigment-Protein Complexes
C. BUSCHMANN and K. H. GRUMBACH 632
- Herbicides which Inhibit Electron Transport or Produce Chlorosis and Their Effect on Chloroplast Development in Radish Seedlings. III. Plastid Pigment and Quinone Composition
K. H. GRUMBACH 642
- Effects of Substituted 2-Phenylamino-1,4,5,6-tetrahydropyrimidines on ATP Formation in Isolated Spinach Chloroplasts
G. VAN DEN BERG, M. BRANDSE, and J. TIPKER 651
- A Cyanobacterial ATPase Distinct from the Coupling Factor of Photophosphorylation
W. LOCKAU and S. PFEFFER 658
- The Anatomy of the Sarcoplasmic Reticulum in Vertebrate Skeletal Muscle: Its Implications for Excitation Contraction Coupling
J. R. SOMMER, M. D. 665
- Sodium-Potassium Movement and the Regulation of Cardiac Muscle Activity
J. DAUT and R. RÜDEL 679
- Regulation of the Sarcoplasmic Reticular Ca²⁺ Transport ATPase by Phosphorylation and Dephosphorylation
L. M. G. HEILMEYER, Jr. and M. VARSANYI 682
- Equilibrium and Kinetic Studies of Calcium Transport and ATPase Activity in Sarcoplasmic Reticulum
G. INESI, M. KURZMACK, D. KOSK-KOSICKA, D. LEWIS, H. SCOFANO, and H. GUIMARAES-MOTTA 685
- Comparative Studies on the ATP-Binding Sites in Ca²⁺-ATPase and (Na⁺ + K⁺)-ATPase by the Use of ATP-Analogues
W. SCHONER, E. H. SERPERSU, H. PAULS, R. PATZELT-WENZLER, H. KREICKMANN, and G. REMPETERS 692
- Electromechanical Coupling I. Introduction
H. CH. LÜTTGAU 706
- Electromechanical Coupling II. The Effect of Perchlorate Upon Excitation-Contraction Coupling in Frog Skeletal Muscle Fibres
M. GOMOLLA, G. GOTTSCHALK, and H.-CH. LÜTTGAU 707
- Electromechanical Coupling III. Estimation of the Ca Storage Capacity of the SR by Analysing the Time Course of Caffeine-Induced Tension Transients of Skinned Muscle Fibres
R. THIELECZEK 709
- Intracellular Membranes as Boundaries for Ionic Distribution. *In Situ* Elemental Distribution in Guinea Pig Heart Muscle in Different Defined Electro-Mechanical Coupling States
M. F. WENDT-GALLITELLI, R. JACOB, and H. WOLBURG 712
- Notes
- New Flavonol Acetates from the Frond Exudate of the Fern *Notholaena aschenborniana*
M. JAY, M.-R. VIRICEL, J. FAVRE-BONVIN, B. VOIRIN, and E. WOLLENWEBER 721

Isolation and Identification of Lathycarpin, a New Pterocarpin Phytoalexin from <i>Lathyrus sativus</i> J. L. INGHAM and K. R. MARKHAM	724	Synthesis of Galloyl-Coenzyme A Thioester G. G. GROSS	778
Irregularities in the Circular Dichroism of Oligoribonucleotides W. ZACHARIAS and H. FOLLMANN	727	Nitrite Inhibition of Bacterial Dinitrogen Fixation F. CASTILLO and J. CÁRDENAS	784
Adult T-Cell Leukemia-Associated Antigen (ATLA): Detection of a Glycoprotein in Cell- and Virus-Free Supernatant N. YAMAMOTO, J. SCHNEIDER, Y. HINUMA, and G. HUNSMANN	731	Energy Transfer Inhibition Induced by Nitrofen B. HUCHZERMEYER	787
Contents of Number 9		Natural Inhibitors of Germination and Growth, I. Development of a Quantitative Biotest and Application upon Extracts from Husks of <i>Avena sativa</i> L. (In German) R. KARL and W. RÜDIGER	793
<i>Original Communications</i>		Natural Inhibitors of Germination and Growth II. Isolation and Structure of Inhibitors from <i>Avena sativa</i> L. (In German) E. LOHAUS, I. BLOS, W. SCHÄFER, and W. RÜDIGER	802
Nonlinear Polarisationoscillations in a Biophysical Model. System II: External Dynamics Z. SZABO and F. KAISER	733	5-Methylcytosine Content in Erythroleucemic Cells of Rats after Induction with Dimethylsulfoxid (In German) R. BAUR, N. KLUGE, H. KRÖGER, and H. WOHLERT	812
The Phytochemistry of South American <i>Nephroma</i> -Species (In German) B. RENNEN, A. HENSSEN, and E. GERSTNER	739	Catabolic Properties of 5',5''-Linked Dinucleoside-phosphates in Rat Liver Nuclei (In German) S. BORNEMANN and E. SCHLIMME	818
Detection of Glyceollin on the Cellular Level in Infected Soybean by Laser Microprobe Mass Analysis P. MOESTA, U. SEYDEL, B. LINDNER, and H. GRISEBACH	748	The Interaction of Calf Thymus DNA with Mercuric Acetate and 3,6-Bis-(acetatomercurimethyl)-dioxane. Small-Angle X-Ray Scattering and Viscosity Studies P. ZIPPER, G. RIBITSCH, J. SCHURZ, and H. BÜNEMANN	824
Oxygen-Isotope Effect in Enzymatic Cleavage Reaction of 13-L-Hydroperoxylinoleic Acid to Hexanal and 11-Formyl- <i>cis</i> -9-undecenoic Acid A. HATANAKA, T. KAJIWARA, J. SEKIYA, and T. FUKUMOTO	752	Growth Characteristics of Anaerobically Treated Ehrlich Ascites Tumor Cells after Reaeration as Studied by Combination of Flow Cytometry and Centrifugal Elutriation R. MERZ and F. SCHNEIDER	833
On the Biosynthesis of C ₃₀ Carotenoid Acid Glucosyl Esters in <i>Pseudomonas rhodos</i> . Analysis of <i>car</i> -Mutants H. KLEINIG and R. SCHMITT	758	Inverse Relationships between Ecdysteroid Titres and Total Body Metabolism in Insects K. SLÁMA	839
Biosynthesis of Photosynthates and Taxonomy of Algae B. P. KREMER and G. O. KIRST	761	<i>Notes</i>	
Specific Localization of β -D-Glucoside Conjugates of 2,4-Dichlorophenoxyacetic Acid in Soybean Vacuoles R. SCHMITT and H. SANDERMANN, Jr.	772	Yeast Aminopeptidase II: Rapid Purification Using Affinity Chromatography of the Periplasmic Enzyme J. KNÜVER and K.-H. RÖHM	845

Contents of Number 10

Original Communications

- On the Biogenesis of Aroma Compounds in Plants and Fruits, XVII. Anabolic Properties of Strawberry Fruit Tissue for the Biosynthesis of Aroma Compounds (In German)
F. DRAWERT and R. G. BERGER 849
- Indole Alkaloids from Cell Suspension Cultures of *Stemmadenia tomentosa* and *Voacanga africana*
J. STÖCKIGT, K.-H. PAWELKA, A. ROTHER, and B. DEUS 857
- Degradation of the Isoflavone Biochanin A by *Fusarium javanicum*
U. WILLEKE and W. BARZ 861
- Changes of C-Phycocyanin in *Synechococcus* 6301 in Relation to Growth on various Sulfur Compounds
A. SCHMIDT, I. ERDLÉ, and H.-P. KÖST 870
- State of Iron in the Archaeobacterium *Methanosarcina barkeri* Grown on Different Carbon Sources as Studied by Mössbauer Spectroscopy
P. SCHERER and CH. SAUER 877
- Effect of Aging on the Fluorescence Lifetime of Chloroplasts
S. S. BRODY 881
- On the Role of Hydrogen Peroxide in Peroxidase Catalyzed Metabolism of Indole-3-acetic Acid
H. J. GRAMBOW 884
- Distribution and Effects of Bentazon in Crop Plants and Weeds
H. K. LICHTENTHALER, D. MEIER, G. RETZLAFF, and R. HAMM 889
- Mechanism of Bleaching in Leaves Treated with Chlorosis-Inducing Herbicides
J. FEIERABEND, TH. WINKELHÜSENER, P. KEMMERICH, and U. SCHULZ 898
- Rotating-Field-Induced Rotation and Measurement of the Membrane Capacitance of Single Mesophyll Cells of *Avena sativa*
W. M. ARNOLD and U. ZIMMERMANN 908
- The Triton X-100 and High Salt Resistant Residue of *Saccharomyces cerevisiae* Nuclear Membranes
K. MANN and D. MECKE 916
- Malonate and Krebs Cycle Intermediates Utilization in the Presence of other Carbon Sources by *Rhizobium japonicum* and Soybean Bacteroids
D. WERNER, W. DITTRICH, and H. THIERFELDER 921
- Alanine Dehydrogenase from Bacteroids and Free Living Cells of *Rhizobium japonicum*
P. MÜLLER and D. WERNER 927
- The Tautomerism of Cytosine and Hydroxycytosine. A Quantum-Mechanical Study
J. S. KWIATKOWSKI, B. LESYNG, M. H. PALMER, and W. SAENGER 937
- The Reversibility of the Vitamin C Redox System: Electrochemical Reasons and Biological Aspects
H. SAPPER, S.-O. KANG, H.-H. PAUL, and W. LOHMANN 942
- Coupled Gas Chromatography — Single Cell Recording: a New Technique for Use in the Analysis of Insect Pheromones
L. J. WADHAMS 947
- Evaluation of (Z)-5-Decen-1-ol as an Attractant for Male Larch Casebearer Moths, *Coleophora laricella*
E. PRIESNER, W. ALTENKIRCH, W. BALTENSWEILER, and H. BOGENSCHÜTZ 953
- A Steroid-Binding Protein from Insect Haemolymph Isolated by Photoaffinity Labelling and Immuno-adsorption
L. REUM, G. KÄUSER, U. ENDERLE, and J. KOOLMANN 967
- Structure Investigations of Agonists of the Natural Neurotransmitter Acetylcholine, III[1]. X-Ray Structure Analysis of (2-Ethoxyethyl)trimethylammonium Chloride
A. GIEREN and M. KOKKINIDIS 977
- Polycation-Bacterium Interactions and Wall Subunits as Endocytosis Factors. Topoisomeraselike Action of Basic Polypeptides Suggesting a 7th Class of Enzymes: The Stereases
S. ANTOHI 985
- Tl⁺-Ions: Influence on Cardiac Contractility
R. ZISKOVEN, C. ACHENBACH, J. WIEMER, and U. WINTER 995

<p>Tl⁺-Ions: Effects on the Automaticity of Sinoatrial Tissue and on dV/dt_{\max} and i_{K2} of Cardiac Purkinje Fibres C. ACHENBACH, J. WIEMER, R. ZISKOVEN, and U. WINTER 1006</p> <p>Tl⁺-Ions: Comparison of the Effects on the Slow Inward Current and Contractility of Ventricular Tissues J. WIEMER, R. ZISKOVEN, and C. ACHENBACH 1015</p> <p>Differentiation in <i>Volvox carteri</i>: Study of Pattern Variation of Reproductive Cells R. GILLES and L. JAENICKE 1023</p> <p>Thermodynamic Approach to a Possible Theory of the Evolution of a Genetic Code M. F. MACCHIATO and A. TRAMONTANO 1031</p> <p><i>Notes</i></p> <p>B-Decachloro-<i>o</i>-carborane Derivatives Suitable for the Preparation of Boron-Labeled Biological Macromolecules D. GABEL and R. WALCSZYNA 1038</p> <p>Temperature-Dependent Colour Change in Larvae of the Green Spider <i>Micromata rosea</i> (Sparassidae) (In German) A. HOLL 1040</p> <p>"Memory" of First Interaction with Physiological or Biologically Active Foreign Molecules (Benzopyrene, Gibberelline) in a Unicellular (<i>Tetrahymena</i>) Model System G. CSABA, G. NÉMETH, and P. VARGHA 1042</p> <p>On the Nerve Growth Factor (NGF) from Human Glandula submandibularis (In German) R. RIEMSCHEIDER, F. DEMBÉLÉ, and I. A. K. GHOURI 1045</p> <p>Spatio-Temporal Visual Receptive Fields as Revealed by Spatio-Temporal Random Noise E. HIDA and KEN-ICHI NAKA 1048</p> <p>On the Taxonomic Position of Hypocomidae (Ciliata) (In German) K. G. GRELL and A. MEISTER 1050</p>	<p>Contents of Number 11/12</p> <p><i>Original Communications</i></p> <p>Composition of Epicuticular Waxes from Fruits of <i>Jobba</i> (<i>Simmondsia chinensis</i> [Link] Schneider) P.-G. GÜLZ 1053</p> <p>Biliverdin IXα, Intermediate and End Product of Tetrapyrrole Biosynthesis H.-P. KÖST and E. BENEDIKT 1057</p> <p>The Oxidative Degradation of L-Ascorbic Acid via an α-Ketoaldehyde S.-O. KANG, H. SAPPER, and W. LOHMANN 1064</p> <p>Further Observations on the Source of Nitrogen in Phenazine Biosynthesis A. RÖMER and R. B. HERBERT 1070</p> <p>Photosynthetic Adaptation in <i>Synechococcus</i> Cells G. DÖHLER and J.-C. LÉCLERC 1075</p> <p>Interrelationship between Quinolizidine Alkaloid Producing Legumes and Infesting Insects: Exploitation of the Alkaloid-Containing Phloem Sap of <i>Cytisus scoparius</i> by the Broom Aphid <i>Aphis cytisorum</i> M. WINK, T. HARTMANN, L. WITTE and J. RHEINHEIMER 1081</p> <p>Biological Activities of Sesquiterpene Lactones from <i>Helianthus annuus</i>: Antimicrobial and Cytotoxic Properties; Influence on DNA, RNA, and Protein Synthesis O. SPRING, J. KUPKA, B. MAIER, and A. HAGER 1087</p> <p>Bleaching Activity of New 2-Phenylpyridazinones: Structure-Activity Relationship G. SANDMANN and P. BÖGER 1092</p> <p>Some Observations on the Saponin Accumulation in Oat Seedlings and on the Transformation of the Avenacosides to the Antibiotic 26-Desglucoavenacosides U. LAUDENBACH and J. KESSELMBIER 1095</p> <p>Naramycin B, an Antibiotic from <i>Streptomyces griseus</i> Strain 587 with Herbicidal Properties-Fermentation, Isolation, and Identification D. BERG, M. SCHEDEL, R. R. SCHMIDT and K. DITGENS and H. WEYLAND 1100</p>
--	---

The Intracellular Distribution of Enzymes of the Glycerol Cycle in the Unicellular Alga <i>Dunaliella parva</i> H. GIMMLER and G. LOTTER	1107	Release of K ⁺ and H ⁺ from Poly U in Aqueous Solution upon γ and Electron Irradiation. Rate of Strand Break Formation in Poly U E. BOTHE and D. SCHULTE-FROHLINDE	1191
Osmoregulation in <i>Dunaliella</i> . Intracellular Distribution of Enzymes of Glycerol Metabolism A. D. BROWN, R. MCC. LILLEY, and T. MARENGO	1115	Model Reactions for the Degradation of DNA-4' Radicals in Aqueous Solution. Fast Hydrolysis of α Alkoxyalkyl Radicals with a Leaving Group in β -Position Followed by Radical Rearrangement and Elimination Reactions G. BEHRENS, G. KOLTZENBURG, and D. SCHULTE-FROHLINDE	1205
Multicomponent Mandibular Gland Secretions in Three Species of <i>Andrena</i> Bees (Hym., Apoidea) G. BERGSTRÖM, J. TENGÖ, W. REITH, and W. FRANCKE	1124	Induction of Single- and Double-Strand Breaks in Linear and Superhelical DNA by Phleomycin (In German) F. SCHMÜLLING and W. KÖHNLEIN	1228
The Sex Pheromone of the Silver Y Moth <i>Chryso-deixis eriosoma</i> (Doubleday) in New Zealand M. H. BENN, R. A. GALBREATH, V. A. HOLT, H. YOUNG, G. DOWN, and E. PRIESNER	1130	Effect of Water Soluble Polymer, Polyethylene-glycol, and Glass-Forming Compounds on Cell Fusion F. YOSHII and I. KAETSU	1234
Structure-Activity-Relation of Analogues and Homologues of Blepharismone, the Low-Molecular Conjugation Hormone of <i>Blepharisma japonicum</i> (In German) M. ENTZEROTH and L. JAENICKE	1136	System Analysis of the Circadian Rhythm of <i>Euglena gracilis</i> , I. Linearities and Non-Linearities in the Response to Temperature Signals W. LORK, T. KREUELS, W. MARTIN, and K. BRINKMANN	1240
Conformational Analysis of Benzyloxycarbonyl-Protected Peptides I. KNACK	1141	Exogenous and Endogenous Control of Swimming Activity in <i>Astyanax mexicanus</i> (Characidae, Pisces) by Direct Light Response and by a Circadian Oscillator. I. Analyses of the Time-Control Systems of an Epigeal River Population W. ERCKENS and W. MARTIN	1253
Immunochemical Characterization of <i>Porphyridium cruentum</i> B-Phycoerythrin: Proof of Cross-Reaction between Chromophore-Free Apoprotein and Holoprotein-Specific Antibodies E. SEFF, G. WANNER, J. EDER, and H.-P. KÖST	1146	Exogenous and Endogenous Control of Swimming Activity in <i>Astyanax mexicanus</i> (Characidae, Pisces) by Direct Light Response and by a Circadian Oscillator. II. Features of Time-Controlled Behaviour of a Cave Population and their Comparison to a Epigeal Ancestral Form W. ERCKENS and W. MARTIN	1266
X-Ray Studies on Phospholipid Bilayers. II. Polymorphic Forms of Dipalmitoyl Phosphatidylethanolamine M. SUWALSKY and E. KNIGHT	1157	Functional Regeneration of the Visual System in Teleosts. Comparative Investigations after Optic Nerve Crush and Damage of the Retina R. KÄSTNER and H. WOLBURG	1274
Statistical Description of Isotope Exchange Processes: A Multisite Model for the ¹⁸ O Exchange P. RÖSCH	1161		
Fluorescence Studies on N-(3-Pyrene)Maleinimide-Labeled Sarcoplasmic Reticulum ATPase in Native and Solubilized Membranes H. LÜDI and W. HASSELBACH	1170		
Localization of Ca ²⁺ at the Plasma Membrane of Bullfrog Myocardial Cells R. MEYER, M. SCHMITZ, W. STOCKEM, and H. G. HAAS	1180	Notes	
		Monoterpenes from the True Bug <i>Harpocera thoracica</i> (Hemiptera) H.-P. HANSEN and J. JACOB	1281

Diterpenes of <i>Cheilanthes argentea</i> , a Fern from Asia E. WOLLENWEBER, P. RÜEDI, and D. S. SEIGLER	1283	Isolation and Synthesis of Hemoregulatory Peptide W. R. Paukovits and O. D. Laerum	1297
Composition of Phospholipids in Seed Oil of <i>Jajoba</i> (<i>Simmondsia chinensis</i> [Link], Schneider) P. G. GÜLZ and C. EICH	1286	Erratum To N. Yamamoto, J. Schneider, Y. Hinuma, and G. Hunsmann (37c, 731—732 [1982])	1300
Adenine Phosphoribosyltransferase Activity in Mitochondria of <i>Catharanthus roseus</i> Cells F. HIROSE and H. ASHIHARA	1288	Subject Index	1301
Inactivation of the Sarcoplasmic Reticulum Cal- cium-Transport-ATPase by Lasoloid in Com- bination with Triton X-100 W. HASSELBACH, H. LÜDI, and A. MIGALA	1290	Authors Index	1327
Small- and Large-Angle X-Ray Scattering Studies of Counter Ion Influence on tRNA Conformation P. WILHELM, I. PILZ, G. DEGOVICS, and F. VON DER HAAR	1293	Errata in this volume To A. Hager and M. Helmle (36c, 997 [1981]) To J. Kesselmeier and D. Strack (36c, 1072 [1981]) To N. Yamamoto, J. Schneider, Y. Hinuma, and G. Hunsmann (37c, 731 [1982])	144 350 1300

Rubins and Rubinoid Addition Products from Phycocyanin *

W. Kufer and H. Scheer

Botanisches Institut der Universität München, Menzinger Straße 67, D-8000 München 19

Z. Naturforsch. 37 c, 179–192 (1982); received December 15, 1981

Phycocyanin, Phycorubin, Bile Pigments, Biliverdin, Bilirubin

The verdin-type chromophore of denatured C-phycocyanin (**1**) from *Spirulina platensis* is reduced to the corresponding rubin (**2a**) by sodium borohydride. The structure assigned is in agreement with the uv-vis spectroscopic properties of the product and was deduced from model studies with free bile pigments.

Analogous model studies using sodium dithionite demonstrated a two-fold reactivity for this reagent, leading to products which are both of the rubin spectral type under the conditions tested. True rubins (10,22-dihydrobilindions) are formed in low yield only if an excess of reagent is used in methanol/water mixtures. It is accompanied by polar addition product(s) of the same spectral type, which are generally formed exclusively. In particular, no bilirubin was formed under the reaction conditions previously applied for the chemical modification of phycobiliproteins and phytochrome. From this finding and from the strikingly different properties of the borohydride and dithionite products, of phycocyanin upon renaturation, the dithionite product is suggested to be a rubinoid addition product (**2b**) rather than a hydrogenation product.

In contrast to the dithionite addition product **2b** of phycocyanin, the chromophore of the true phycorubin (**2a**) remains stable upon renaturation. The uv-vis spectral properties of the chromophore are not markedly different whether the apoprotein is in its native or denatured state. The different electrophoretic mobilities of native (renatured) phycocyanin compared to the renatured borohydride product suggest that these two have different protein conformations.

The preparation of these phycorubins renders the extensive techniques of bilirubin chemistry applicable in the study of biliproteins.

Introduction

Bile pigments, open-chain tetrapyrrolic compounds, serve important functions in plants. In contrast to mammalian bile pigments derived from heme degradation, they are biosynthesized as photoreceptors which are covalently bound to apoproteins, thus forming "biliproteins" (for a recent review, see [1]). Two functionally different groups can be distinguished: Firstly the phycobiliproteins as light harvesting pigments of photosynthesis in cyanobacteria, red and cryptophyten algae, and secondly phytochrome and the phycochromes as photomorphogenetic receptor pigments of higher plants and some algae.


The physico-chemical properties of the bile pigment chromophores are profoundly influenced by the native proteins. This is mainly due to non-covalent protein chromophore interactions which are essential in biliproteins to optimize the proper-

Abbreviations: PC, phycocyanin; tlc, thin layer chromatography; mob, electrophoretic mobility relative to standard.



* Studies on Plant Bile Pigments 12; part 11 of this series: W. Kufer and H. Scheer, *Angew. Chem.*, in press.

Reprint requests to Prof. Dr. H. Scheer.

0341-0382/82/0300-0179 \$ 01.30/0



1



ties of the chromophores in their function as photo-receptors. This can be easily seen from a comparison of native and (reversibly) denatured pigments (see ref. [1] for a survey). Recently, we developed a technique for the selective chemical modification of biliprotein chromophores [2, 3]. The experiments reported here have been performed to yield more detailed information about the structure of the chemically modified products, and the non-covalent protein chromophore interactions in biliproteins. The previous experiments of this type were carried out with the reducing agents dithionite, sulfite and 2-mercaptoethanol, and pigments of the rubinoid spectral type were obtained [3]. Here, we wish to report the reaction of C-phycocyanin (1) (C-PC) and some free bilins as model compounds with sodium borohydride. This reagent has been used [4, 5] for selective reduction of bilindiones ("biliverdin" type chromophores like (3)) to 10,22-dihydrobilindiones ("bilirubin" type chromophores like (4))*.

Results

Reaction of phycocyanin with sodium borohydride

The reactions are summarized in Scheme 1. When PC (concentration range 30–130 μM chromophores**) denatured by 8 M urea in 50 mM sodium phosphate buffer, pH 7.5, was treated at 0 °C with solid sodium borohydride (25–40 mM***), the bands in the uv-vis spectrum at 602 nm ($\epsilon = 15400$ per chromophore) and 355 nm ($\epsilon = 37000$) [3] disappeared completely after 20–30 min and a new band at 416 nm arose simultaneously. The pH of the solution increased to about 9, which could be avoided without significant changes in reactivity by using a buffered solution of sodium borohydride. Using PC (48 μM chromophores) and 13 mM NaBH_4 , a 90% decrease of the 602 nm band was observed within 30 min. The reaction went to completion

within additional 30 min after a further addition of the same amount of borohydride (Fig. 1)*. Assuming a 1:1 stoichiometry with no by-product formed, $\epsilon_{416} = 24200$ (per chromophore) was calculated from the spectral changes, and with $\epsilon_{602} = 15400$ [3] for the educt chromophores**. The vis-spectrum remained unaltered when the reducing agent was removed by gel filtration on a column equilibrated with 8 M urea. The yellow colour of the borohydride product also remained unchanged after simultaneous removal of urea and borohydride by gel filtration ($\lambda_{\text{max}} = 418$ nm). The extinction coefficient of the pigment thus obtained was determined by unfolding the protein again, by addition of solid urea to a final concentration of 8 M. With $\epsilon = 24200$ as reference value for the denatured pigment (see above) and allowing for the dilution caused by addition of urea, $\epsilon_{418} = 21900$ was obtained for "native" phycorubin.

The electrophoretic mobility of the product obtained after removal of urea and borohydride differed from that of native PC in tris-glycine electrophoresis. With equal protein concentrations applied, the yellow product had a higher mobility (mob = 0.72 vs. bromophenolblue as reference compound) as compared to PC (mob = 0.59) (Fig. 2). They were identical, however, in SDS-gel electrophoresis having a single band corresponding to MW = 19900 in coelectrophoresis using the system of Weber and Osborn [6] (a separation into subunits was generally not obtained with PC from *Spirulina platensis*). Reaction of the yellow pigments with diazotized ethyl anthranilate (5–10 fold molar excess) yielded uv-vis spectroscopically identical pigments with a long wavelength band at 485 nm, irrespective of the removal of only NaBH_4 or both NaBH_4 and urea before the reaction.

Addition of solid sodium borohydride (43 mM) to native PC (6.6 μM chromophores, calculated from $\epsilon_{620} = 98700/\text{chromophore}$ [3]) led to only partial conversion to the yellow pigment. After 10 min reaction




* See ref. [43] for the nomenclature of bile pigments, and formulas for the numbering system. In addition, the non-systematic terms "rubin" and "verdin" are used to characterize pigments bearing the conjugation systems of bilirubin and biliverdin, respectively.

** Concentrations throughout are given for chromophores. Since monomeric ($\alpha_1 \beta_1$) C-PC contains three chromophores, the protein concentration is one third of the chromophore concentration.


*** NaBH_4 is unstable in protic solvents. All concentrations given are the initial concentrations of the reagent.

* Addition of a larger excess of NaBH_4 results in a further reaction of the product ($\lambda_{\text{max}} = 416$ nm) to products not absorbing in the visible spectral range. It is thus recommended to carry out the reaction with a minimum amount of borohydride added in successive portions, and to follow the reactions spectrophotometrically.


** The extinction coefficients for the reaction products of denatured C-PC with sodium dithionite, sodium sulfite and 2-mercaptoethanol have been determined by the same method to 25500, 27900 and 23600 $\text{l} \cdot \text{M}^{-1}$, respectively [3].



Type A



Type B

Type C: R = SO₂⁻
 Type D: R = H

Scheme 1. Reaction scheme of the reversible renaturation of PC from *Spirulina platensis*, and of the reactions with sodium borohydride (A) and sodium dithionite (B). The chromophore structures A–D are schematic representations of the chromophore geometry. Structure A is representative of an extended conformation without major steric hindrance of the β -pyrrolic substituents. Structure B is the cyclic-helical conformation found for biliverdin in solution [22, 23]. Structures C and D have been drawn with the two dipyrromethone units nearly perpendicular to each other, similar to the crystal structure of bilirubin [46, 47]. The reactions of PC with sodium sulfite and 2-mercaptoethanol principally follow Scheme 1B as well but different concentrations of the reagents are needed.

Purification of cytochrome oxidase

All operations were performed at 4 °C unless stated otherwise. The membrane fraction was re-suspended in TRIPE buffer at a concentration of about 6 mg protein per ml. Triton X-100 (10% stock solution) in TRIPE buffer was added dropwise to the suspension under stirring to a final concentration of 1.3% (w/v, protein to detergent ratio of 1 : 2). After incubation for 30 min insoluble material was precipitated at $145,000 \times g$, 90 min, 4 °C. The supernatant contained about 90% of the cytochrome oxidase. The enzyme extract was diluted with TRIPE buffer to a final concentration of 1% Triton X-100.

10 ml of the supernatant (4 mg protein/ml) were applied to a column (1 × 8 cm) of DEAE Sepharose CL-6B (Pharmacia Freiburg) equilibrated with TRIPE buffer plus 1% Triton. The column was washed with 30 ml of TRIPE buffer plus 1% Triton and eluted with a linear gradient of 0–1 M KCl (2 × 40 ml). Fractions of 3 ml were collected and tested for cytochrome oxidase activity.

The peak fractions eluted at 0.3 M KCl were pooled and desalted as described in [10]. 9 ml of the desalted enzyme solution (5.6 mg protein) was then applied to a cytochrome *c*-thiol-activated Sepharose 4B column (1 × 4 cm; Pharmacia) equilibrated with TRIPE buffer plus 1% Triton X-100. The coupling of cytochrome *c* from *Saccharomyces cerevisiae* (Sigma, München, type VIII) and the preparation of the cysteine inactivated column were performed as described in [12]. The binding capacity of the column was 7.7 mg cytochrome *c* per ml of packed material. The column was washed with 20 ml of TRIPE buffer and then eluted with a linear gradient of 0–0.5 M KCl solution (2 × 40 ml).

Determination of enzymatic activity

Cytochrome oxidase was assayed at 30 °C by following the decrease of absorption at 546 nm in an Eppendorf photometer (type 1101 M) using dithionite reduced horse heart cytochrome *c* as electron donor ($\epsilon = 21 \times 10^3 \text{ cm}^2/\text{mmol}$). The reaction mixture contained 50 mM TRIPE buffer (pH 8), 1 mg ferrocyanochrome *c* and enzyme solution in a final volume of 1 ml. The reaction was started by addition of 10 to 100 μl enzyme solution. When inhibitors were used, the enzyme preparation was preincubated for 5 min with the inhibitor. The reaction was then started by addition of reduced cytochrome *c*. Horse

heart ferrocyanochrome *c* was prepared as described in [10] with the modification that the pH of the ferrocyanochrome *c* solution was adjusted to pH 8 with 1 M Tris.

Native polyacrylamide gel electrophoresis (charge-shift-electrophoresis)

Electrophoresis in 5% polyacrylamide slab gels containing 0.1% Triton X-100 was carried out according to the method of Simons *et al.* [13] with the modification that DTT was added to the gel and the buffers in 1 mM concentration. The samples were preincubated in the sample buffer for 15 min at 4 °C. Electrophoresis was performed at 4 °C with 2 mA/gel track. After the run one track was cut into 2 mm slices. The slices were incubated over night with 200 μl of TRIPE buffer plus 0.1% Triton and tested afterwards for activity in the enzyme assay. The slices of the second track were extracted for 4 h with 50 μl of sample buffer (4% SDS) at 60 °C. The extracts were then subjected to SDS-gel electrophoresis. The third track was stained with 0.04% Coomassie brilliant blue in 25% isopropanol and 10% acetic acid (v/v) and destained with 10% acetic acid and 10% methanol.

Sodium dodecylsulfate polyacrylamide gel electrophoresis

Slab-SDS-gel electrophoresis was performed according to the method of Laemmli [14] using 11.5–16.5% acrylamide gradient gels of 1 mm thickness. The samples were dialyzed for two days, lyophilized and resuspended in sample buffer. All samples were preincubated at 60 °C for 30 min or at 100 °C for 10 min, respectively.

Determination of protein

Protein concentration was determined by the method of Lowry *et al.* [15]. The presence of Triton X-100 was corrected by addition of 0.5% (w/v) sodium dodecylsulfate to the alkali solution of the Lowry reagent [16]. Bovine serum albumin was used as standard.

Measurement of difference absorption spectra

Difference spectra were obtained with a Perkin-Elmer split beam spectrophotometer model 330 (Überlingen) using 1 cm light-path cuvettes at room

Table I. Purification of cytochrome oxidase from *Rhodospseudomonas capsulata* strain 37 b4. The purification procedure and the enzyme assay are described under Materials and Methods.

	Activity [$\mu\text{mol}/\text{min} \cdot \text{ml}$]	Protein [mg/ml]	Specific Activity [$\mu\text{mol}/\text{min}$ mg prot.]	Purification (= fold)	Yield [%]
Membranes	2.3	6.15	0.37	1	100
Triton extract	3.4	3.9	0.87	2.4	148
DEAE-Sepharose eluate; peak fraction	1.19	0.19	6.7	17	51
Cytochrome <i>c</i> -thiol- activated Sepharose eluate; peak fraction	0.63	0.011	59.7	161	29

temperature. 150 μM potassium ferricyanide was added to the reference cuvette and 100 μM sodium dithionite to the sample. Before recording the spectrum the cuvettes were kept for 2 min in the dark.

Results and Discussion

Solubilization of cytochrome *c* oxidase

Ninety seven percent of the enzyme was solubilized at a Triton concentration of 1.3% and a protein to detergent ratio of 1 : 2. The activity of the enzyme was enhanced about 30–50%. Lowering the Tris concentration from 50 mM to 20 mM reduced the solubilization of the enzyme. An increase of detergent concentration to more than 1.3% resulted in a loss of activity. Addition of KCl up to 1 M stimulated the activity about 10% (in contrast to NaCl) but had no effect on the solubilization of the cytochrome oxidase activity.

The detergent *N,N*-dimethylaurylamineoxide (LDAO) solubilized the enzyme activity at a concentration of 1.3% nearly 100% but the half life of the activity dropped to 3–5 h. A mixture of LDAO and other detergents or increasing the ionic strength by addition of KCl or NaCl caused a complete loss of the activity. The detergents Brij-58, Tween 20, Lubrol-PX and SDS applied at 4 °C and at room temperature solubilized the enzyme only partially or resulted in inactivation of cytochrome oxidase. Sodium cholate solubilized cytochrome oxidase activity up to 90% but with a lower yield than Triton. In the presence of 1.3% Triton X-100 the cytochrome oxidase had a half-life of 24 h at 4 °C. Freezing of the solubilized enzyme even in the presence of 50% glycerol destroyed the activity. The crude Triton-extract lost approx. 20% of activity at 4 °C in 24 h.

Purification

Triton-extract from membranes was applied to a DEAE-Sepharose CL 6B column. Enzyme activity eluted shortly after a dark red band at 300 mM KCl (Fig. 1). The specific activity was increased 17-fold (Table I).

The enzyme fraction which eluted from the DEAE-Sepharose column contained considerable amounts of cytochrome *c* besides cytochrome *b*. The desalted fraction was loaded on a cytochrome *c*-thiol activated Sepharose 4B column. The cytochrome *c* was covalently linked through its cysteine residue located close to the N-terminus [17] thus leaving free the important lysine residues for binding cytochrome *c* oxidase and reductase [12]. The enzyme was not bound to the column when cystein was inactivated.




Fig. 1. Ion exchange chromatography on DEAE-Sepharose Cl-6B-column (1 × 8 cm; Pharmacia) in presence of 1% Triton X-100 and standard buffer (0.05 M Tris-HCl pH 8, 0.1 mM PMSF, 0.1 mM EDTA). —■—■— Cytochrome oxidase-activity; --○--○-- A_{290} ; ——— KCl-gradient 0–1 M KCl, 2 × 40 ml; fractions containing 3 ml.

Table I (continued).

Compound and reagent	Absorptions λ_{\max} [nm]	λ_{\max} after extraction in CHCl_3	tlc: R_F -values $\times 100$								
			a	b	c	d	e	f	g	h	
Mesobilirubin, 4b	424, 398 sh ¹⁾ , 419 ^{m)}		66–74					60–69	31		65
Mixture of III, IX and XIII α -isomers of mesobilirubin (5b , 4b , 6b)			68					60	31		
			74								
			78								
Octaethylbilindion, 7	646 ¹⁾										47
7 + NaBH_4 ¹⁾	400, 420 sh										11
7 + 2-mercaptoethanol ^{1, s)}	426										
2,3-Dihydrooctaethyl- bilindion, 8	596 ¹⁾										65
8 + NaBH_4 ¹⁾	400										05
8 + dithionite ¹⁾	398 ^{c)}										
8 + 2-mercaptoethanol ^{1, s)}	410										

The octaethylbilindion **7** and its 2,3-dihydro-derivate **8** similarly yielded yellow products upon treatment with solid borohydride in methanol. **7** reacted smoothly at ambient temperature, **8** only at elevated temperatures (50 °C, under nitrogen). The conjugation system of **8** is identical to that of C-PC.

Using sodium dithionite instead of borohydride, the results depended critically on the reaction conditions. In an experiment in methanol/water (1:1, v/v) analogous to the one described above, a yellow product different from **4a** was obtained from **3a** with 1 mM dithionite. It migrated with an R_F -value of or close to zero on silica gel-tlc, developed with neutral or acidic solvent systems (a, e, f in Table I). On polyamide-tlc with a basic solvent system, it migrated close to the front (g in Table I). No bilirubin (**4a**) could be detected under these conditions. Upon rising the dithionite concentration to 24 mM, **4a** and its isomers (**5**, **6**) could be identified as by-products besides the aforementioned polar product(s). The product mixture was analyzed quantitatively after reaction on the preparative scale (**3a** = 10^{-4} M, dithionite = 2×10^{-2} M). After work up, esterification with diazomethane and preparative tlc, **4c** was isolated in 4% yield and identified by uv-vis and tlc comparison with authentic bilirubin dimethylester (**4c**). The results obtained with mesobiliverdin (**3b**) were analogous (see Table I).

In another series of experiments, biliverdin was treated with dithionite in the absence of organic solvents. In particular, the reaction of **3a** (9×10^{-6} M) was studied in phosphate buffer (50 mM sodium phosphate, pH 7.5) containing urea (8 M), e. g. under

conditions identical with those used for the reaction of PC [3]. With dithionite concentrations ranging from 2×10^{-4} to 2×10^{-2} M, no bilirubin could be identified in the yellow product mixtures. Chromatography revealed only polar product(s) ($R_F = 0$ in system a, as compared to 0.85 for **4a**). The same results were obtained using buffer without urea.

The conformationally restricted pigments phorbacilin (**9**) and isophorbacilin (**10**) reacted smoothly, too, with dithionite (see Table II). Treatment of **9** (13×10^{-6} M) with sodium dithionite (5×10^{-5} M) in a buffer/methanol mixture led to decrease of the

Table II. UV-vis spectral data of the biliverdin **11**, the verdins **9** and **10** with an extended conformation similar to the one suggested for the chromophores of native PC [25], and of their reaction products with reducing agents.

Compound	λ_{\max} [nm] in methanol
Phorbacilin-dimethylester, 9	555 562 ^{b)}
9 + NaBH_4	417
9 + dithionite (0.05 mM)	412 ^{b)}
Isophorbacilin-dimethylester, 10	606, 690 (sh) 614 ^{b)}
10 + NaBH_4	417
10 + dithionite (0.5 mM)	435, 455 (sh), 405 (sh) ^{b)}
Biliverdin IX γ -dimethylester, 11	645
11 + NaBH_4	408
11 + dithionite ^{b)}	c)

a) See footnote (s), Table I.

b) 50 mM sodium phosphate buffer, pH 7.5/methanol = 1:1

c) Not determined due to precipitation.




Fig. 3. Titration of isophorbocabilindimethylester (**10**) (9.6×10^{-6} M) in a 1:1 mixture of methanol and phosphate buffer (50 mM, pH 7.5) with sodium dithionite. The absorption spectra were recorded at the following concentrations of dithionite curve 1: 0 M, 2: 5×10^{-6} M, 3: 5×10^{-5} M, 4: 5×10^{-4} M, and have been recorded each 5 min after the respective additions of the reagent.

long-wavelength band ($\lambda_{\max} = 562$ nm) by 95%, and the appearance of only one product band ($\lambda_{\max} = 412$ nm). Titration of **10** with dithionite (Fig. 3) led to 96% bleaching with 5×10^{-4} M dithionite, again forming a yellow product ($\lambda_{\max} = 435$, shoulders at 435 and 405 nm).

In addition to the results described here, some earlier results [3] of the reaction of model compounds for the PC chromophore with sodium sulfite and 2-mercaptoethanol have been incorporated in Table 1. They have already been published, together with preliminary results obtained with sodium dithionite.


Discussion

The reaction product of denatured PC with sodium borohydride has an absorption maximum at 416 nm. Spectroscopically similar products were found with other reducing agents, viz. sodium dithionite, sodium sulfite and 2-mercaptoethanol [2, 3] (see also Scheme 1). This suggests for all these products an interruption of the conjugated system of the PC chromophore (**1a**) at the C-10 position. For the rings C and D, this leads to the chromophoric system of mesobilirubin (**4c**, $\lambda_{\max} = 419$ nm in phosphate buffer containing 8 M urea). All four products have similar extinction coefficients: $\epsilon_{416} = 24200$ with borohydride, $\epsilon_{418} = 25500$ with dithionite [3], $\epsilon_{418} = 27900$ with sulfite and $\epsilon_{418} = 23600$ with 2-mercaptoethanol. They amount to roughly 50% of the value reported for mesobilirubin (**4c**) (ϵ_{433} (CHCl₃) = 54600 [10]); thus supporting the general structure **2**. For rings A and B, the interruption of


the conjugation at C-10 leads to a vinylpyrrole chromophore not absorbing in the visible range (e.g. 313 nm for a series of 3,4-dihydro-6(1H)-pyromethenons [8, 9] closely related to the ring A, B-fragment of **2a**). The borohydride product has an absorption in this spectral range ($\lambda_{\max} = 310$ nm) which appears as a shoulder of the protein band ($\lambda_{\max} = 280$ nm), the other products could not be investigated in this region due to strong absorption of the reagents.

Although the phycocyanin reaction products with these four reductants are uv-vis spectroscopically nearly identical, they differ in their structure concerning the residue R in formula **2**. From uv-vis and tlc-data of the reaction products of model compounds with sulfite, and from the acid reversible product formation with 2-mercaptoethanol, the structures **2c** and **2d** were suggested [3] for these pigments arising from addition of the respective reagents. The pronounced reactivity of C-10 in bilindiones towards nucleophiles had been inferred first from MO calculations [12]. After circumstantial evidence [3, 13a, b], NMR-data of adducts of thioles with bilindiones in agreement with the proposed structure **2d** have been reported [14, 15], and the thermodynamic data for related equilibria have been obtained [15].


Reaction of the same model compounds with sodium borohydride led via addition of hydride at C-10 to true reduction products, e.g. bilirubins, as revealed by comparison of their uv-vis and tlc-data with those of the authentic corresponding rubins (see Table I). Reduction of verdins to rubins with borohydride in methanol has been reported in the




3a R = C₂H₃ R' = H
3b R = C₂H₅ R' = H
3c R = C₂H₃ R' = CH₃




R = R' =
4a C₂H₃ H
4b C₂H₅ H
4c C₂H₃ CH₃




5a R = C₂H₃
5b R = C₂H₅




6a R = C₂H₃
6b R = C₂H₅




7




8



9



10



11

literature [4, 5]. The appearance of III- and XIII- α isomers **5** and **6** besides bilirubin (**4a**) and mesobilirubin (**4c**) in methanol/water can be explained by the isomerisation ("scrambling") reaction, which has been observed with bilirubin in moderately basic aqueous solutions [11]. Structure **2a** was thus assigned to the reaction product of phycocyanin with borohydride, and since it is a true hydrogenation product it should be named "phycorubin".

Conflicting results have been reported for the reaction of verdins with dithionite. Bilirubin has been isolated in low yield from the reaction of sodium dithionite with biliverdin, in aqueous alkaline solution [16], and the reduction of a biliverdin-iodine complex to bilirubin has been reported [17]. On the other hand, a product different from bilirubin has been identified as well [18]. The model studies reported here support this differential reactivity of bilindiones towards dithionite. In the neutral pH-range studied, rubins have only been identified in low yield, and only in solutions containing methanol. In particular, rubins have in no case been observed under the conditions used for the reactions of PC with sodium dithionite. A product with similar chromatographic mobilities as the sulfite product (see Table I) arose instead. Derivatives of nicotinamide are reduced by dithionite in a complex mechanism. Addition of sulfinate, the homolytic fission product of dithionite, has been suggested as the first step, followed by a slower and pH-dependent elimination and a final true reduction step with HSO_2^- rather than $\text{S}_2\text{O}_4^{2-}$ as the reagent proper for hydride transfer [19]. Whereas this sequence eventually leads to reduction of nicotinamide in good yields, this is apparently not the case with bilindiones, possibly due to the increased stability of the first addition products*. Even prolonged treatment of the product at pH 2.7 during workup did not result in bilirubin formation. Structure **2b** is thus suggested for the dithionite reaction products of PC and APC, and analogous structures

for the reaction products with other biliproteins, e.g. phycoerythrin, and phytochrome P_r. They had earlier been tentatively identified as true reduction products [2, 3], which has to be corrected in view of the comparison with the rubins proper obtained with borohydride. Thus, the stability of phycorubin (**2a**) upon renaturation (removal of urea e.g. by gel filtration), was strikingly different as compared to the products obtained with sulfite (**2c**), 2-mercaptoethanol (**2d**) and dithionite (**2b**) [3] see Scheme 1. The yellow colour of the borohydride product remained unchanged, while the products with sulfite, 2-mercaptoethanol and dithionite were converted back to native PC.

The reactions summarized in Scheme 1 B, can readily be explained by a reversible addition to the central methine bridge, with the equilibrium strongly dependent on the state of the protein. In the denatured state with the chromophores still bound but uncoupled from the protein, the addition-elimination equilibrium is shifted to the addition product, whereas the educt is favored in the native or renatured pigments even in the presence of a large excess of dithionite. Also, the equilibrium is differentially affected by the different environments of the chromophores of identical molecular structure (e.g. three in PC). Reversible addition reactions are thus a useful tool to probe these different environments [3].

The different stabilities of the native and denatured pigments may originate from a different conformational freedom of the chromophores in the two states. Biliverdin dimethylester (**3b**) and related bilindiones are rather flexible molecules, which exist in solution in several rapidly interconverting forms [21] of predominantly cyclic conformation [22, 23]. The nearly identical spectroscopic properties of denatured biliproteins make a similar mobility likely for their chromophores. The comparably narrow absorption bands of native biliproteins, as well as their high fluorescence yield indicate that the chromophores are rigidly fixed in (an extended [24, 25]) conformation. The nucleophilic addition results in the conversion from an sp^2 to sp^3 hybridised C-10, and thus necessarily a conformational change of the chromophores. A conformational fixation could then shift the equilibrium between the biliprotein chromophore proper, and its addition product. We have attempted to test this hypothesis by studying the reaction of phorbabilin-dimethylester (**9**) and

* Caughey and Schellenberg [20] have identified sulfinate addition products of nicotinamides by ^1Hmr spectroscopy. Attempts to identify the corresponding product of biliverdin failed. **3c** in C^2HCl_3 (0.02 M) was treated with an aqueous solution of sodium dithionite, and the yellow solution formed was thoroughly washed with water and dried. The solution exhibited only broad signals. When the reaction was carried out with sodium borohydride instead, the spectrum of the product was identical to that of bilirubin dimethylester (**4c**) (see experimental part).

isophorcabilin dimethylester (**10**), bilins held in an extended conformation by additional bridges between the pyrrole rings. Unfortunately, however, the central seven membered ring is still flexible enough to allow both sp^2 - and sp^3 -hybridization of C-10 without steric restrictions. Thus, **9** and **10** react with both dithionite and borohydride as well as does biliverdin-IX γ -dimethylester **11** (see Table II).

The slow formation of denatured PC from the dithionite product [3] after removal of the reducing agent (gel filtration, see Scheme 1a) is then explained by a shift in the equilibrium to the side of phycocyanin.

According to the structure of the phycorubins obtained with NaBH_4 as true reduction products, their properties are quite different. The major difference between the addition of nucleophiles to and the hydrogenation of biliverdin is, that the former reaction is in thermodynamic equilibrium, whereas the latter is essentially irreversible under the reaction conditions. Biliverdin can be reduced to bilirubin only with rather strong reductants (e.g. NaBH_4), whereas the oxidation of bilirubin to biliverdin requires high-potential quinones* [4, 14]. Recent electrochemical studies have shown, too, that the biliverdin-bilirubin redox couple is not an equilibrium system [26]. The striking difference between phycorubin and the addition products of phycocyanin summarized in Scheme 1 can readily be explained on this basis. The yellow phycorubin chromophore remains stable upon renaturation of the protein and removal of the reducing agent, sodium borohydride. Moreover, the absorption maximum remained roughly in the same position and the extinction coefficient was the same within experimental error. The geometry of the phycocyanin chromophores is profoundly changed when the protein is renatured and *vice versa*, which is reflected by pronounced spectral changes [25]. The absence of similar effects in phycorubin could then indicate, that the altered chromophores do no longer fit the binding sites, and remain – although covalently attached – only loosely coupled to the protein. It should be noted, though, that the uv-vis-

spectra of bilirubins [28] appear less conformation dependent than those of biliverdins [29].

The different binding situation of the chromophores in native phycocyanin and phycorubin can also be the reason for their different electrophoretic mobilities (Fig. 2). Phycorubin migrates faster towards the anode, which could be due to an increase of the net negative charge by the free carboxylic acid side chain(s) at or close to the surface. The ready reaction of renatured phycorubin with diazotized ethyl anthranilate leading to a product which cannot be distinguished uv-vis spectroscopically ($\lambda_{\text{max}} = 485 \text{ nm}$) from the denatured form, is compatible with this proposal. In SDS-gel electrophoresis, PC and phycorubin migrated identically, indicating the absence of proteolytic artefacts during the preparation of the latter pigment.

In view of the irreversible reactions of NaBH_4 , the treatment of native PC may be used to discriminate between thermodynamic and kinetic effects in the reactions of biliprotein chromophores. In first order, one may assume that (i) the chromophores react with NaBH_4 if only their methine bridges are accessible to the reagent, and that (ii) this reaction may be slowed down by the influence of the protein, but is irreversible once the chromophore has been reduced. The influence of the state of the protein on the reaction kinetics with NaBH_4 has already been studied by Crespi *et al.* [45], and interpreted as a change in accessibility of the chromophores to the reagent. An alternative mechanism of the protein to slow down the reaction with sodium borohydride would be a conformational change of the chromophore. The similar reactivity of biliverdin and isophorcabilin does not prove this assumption, but indicates that a mere conformational change without the coplanar fixation suggested for PC has no dominant influence.

The reaction of native PC with dithionite is incomplete even at high concentrations of the reagent. In this case the reaction is determined by thermodynamics. Sodium borohydride reduces native PC slowly, but completely. This demonstrates, that C-10 of all chromophores is accessible by the reagent dissolved in the aqueous phase. Similar conclusions have been reached by Troxler [30] for the terminal O-1 and O-19, and hence practically the entire chromophore must be accessible to the aqueous phase. It requires further work to determine, whether the chromophores are actually at the

* We have achieved the reoxidation of phycorubin with benzoquinone to products identical with PC if judged from absorption, fluorescence and electrophoretic mobilities. Details of this procedure and the reaction of PC with quinones [27] are to be published separately.

surface, or rather in interior hydrophilic parts of the protein.

The possibility of the preparation of a phycobiliprotein with a rubin-type chromophore described here makes further modifications feasible, since bilirubins are accessible to a variety of reactions. A first example is the use of the diazoreaction of bilirubin as a novel degradation technique for biliproteins [31].

Materials and Methods

PC was isolated from *Spirulina platensis* as described previously [3].

Preparation of Phycorubin

10.0 ml of a solution of denatured PC (5×10^{-5} M chromophores) in 50 mM sodium phosphate/8 M urea-buffer (containing 5 mM EDTA and 5 mM NaN_3), pH 7.5, were cooled with ice and 500 μl of a solution of 10 mg/ml sodium borohydride in the same buffer were added. The solution was left standing for 1/2 h and then again 500 μl of the borohydride solution were added (2.4×10^{-2} M final concentration). Within again 1/2 h the reaction followed spectrophotometrically went to completion.

Renaturation of phycorubin

For removal of urea (and borohydride) 1.0 ml of the solution obtained by the reaction described above was passed through a 1.7×12 cm biogel P 2 column (Bio-Rad, Richmond, California) which was previously equilibrated with 0.05 M sodium phosphate buffer, pH 7.5. The procedure was carried out in a cold room at 4 °C.

Preparation of model bilins for the PC-chromophore and of authentic rubins

Bilirubin (**4a**), biochemical grade, was purchased from Merck, Darmstadt. Mesobilirubin (**4b**) was prepared from 200 mg of **4a** by catalytic hydrogenation [4, 32] (40 mg of 10% Pd on charcoal) in 20 ml 0.1 N NaOH. The reaction was followed spectrophotometrically, adding aliquots from the reaction mixture to methanol (λ_{max} (**4a**) = 452 nm, λ_{max} (**4b**) = 428 nm). The mixture was worked up with chloroform after addition of glycine/HCl-buffer, pH 2.7, saturated with ammonium sulfate. Yield: 98%. Biliverdin (**3a**) was prepared from 50 mg bili-

rubin (**4a**) dissolved in 50 ml dimethylsulfoxide by oxidation with a solution of 50 mg 2,3-dichloro-5,6-dicyanobenzoquinone in 10 ml of the same solvent in a nitrogen atmosphere [4]. After work-up with chloroform and water (yield: 38%), the reaction product was purified by preparative tlc on silica using the upper phase of a mixture of toluene/acetic acid/water 5:5:1 (v/v/v) as solvent system. The green band (R_F -value 0.1) was eluted with acetic acid and worked up by partitioning between chloroform and water. The chloroform phase was washed with water and 1% aqueous NaHCO_3 -solution until neutral (yield: 26%). For crystallization the product was dissolved in 3 ml chloroform (containing a small amount of methanol). 12 ml of *n*-hexane were added and the mixture was left standing at -20 °C over night (yield: 9%).

Mesobiliverdin (**3b**) was prepared by an analogous procedure from mesobilirubin (**4b**).

10 mg of crude (**3a**) dissolved in 150 ml methanol were esterified to the corresponding dimethylester (**3c**) by treatment with 60 ml methanol containing 20% (w/w) BF_3 [4] under reflux for 10 min in a nitrogen atmosphere. After work-up with chloroform and water, the crude product was purified by tlc on silica with chloroform/acetone 95:5 (v/v) as solvent [33]. The green zone was eluted with acetone (yield: 85%).

Bilirubin dimethylester (**4c**) was prepared from 30 mg (**4a**) suspended in 10 ml CHCl_3 by esterification with diazomethane over night [34, 35]. The chloroform phase was washed with an aqueous 10% Na_2CO_3 solution and the product subsequently chromatographed on neutral Al_2O_3 (2×6 cm; super activity I, Woelm, Eschwege) [36]. By-products were eluted with CHCl_3 , (**4c**) with $\text{CHCl}_3/\text{MeOH}$ 9:1 (yield: 65%).

Octaethylbilindiones (**7**) and (**8**) were synthesized from octaethylporphyrin [37, 38]. The products were purified by tlc on silica with $\text{CCl}_4/\text{acetone}$ 9:1 (v/v) as solvent.

Biliverdin-IX γ -dimethylester (**11**): a mixture of the isomeric biliverdin-IX dimethylesters was obtained by coupled oxidation ($\text{O}_2/\text{hydrazine}$) of hemin with subsequent esterification [33]. Biliverdin-IX dimethylester was freed from the β and δ -isomers by tlc on silica with chloroform/acetone 97:3 (v/v) [39], and subsequently from the α -isomer by tlc on silica with toluene/ethylmethylketone/acetic acid 10:5:0.5 (v/v/v) [40].

Phorcabilindimethylester (**9**) was prepared by heating biliverdin-IX γ -dimethylester (**11**) for 1 h at 100 °C in dimethylsulfoxide under nitrogen [39]. Isophorcabilindimethylester (**10**) was obtained from phorcabilindimethylester (**9**) in 20% MeOH/H₂SO₄, the mixture being kept under reflux for 1/2 h [39]. The products were purified by tlc on silica with chloroform/acetone 8:2 (v/v) as solvent system and crystallized from CHCl₃/*n*-pentane 1:25 at -20 °C.

Mixture of the III-, IX- and XIII- α -isomers of bilirubin (**4a**, **5a**, **6a**) and mesobilirubin (**4b**, **5b**, **6b**): bilirubin (**4a**) or mesobilirubin (**4b**) were dissolved in 0.5 ml 0.1 N NaOH and added to 5.5 ml 50 mM sodium phosphate buffer, pH 7.5, containing 8 M urea. The samples were purged with nitrogen and kept at 34–38 °C for 2 h. The rubins were extracted with chloroform after addition of glycine/HCl buffer, pH 2.7, saturated with (NH₄)₂SO₄. For preparation of pure isomers of bilirubin, the mixture was chromatographed on silica with CHCl₃/HOAc 97:3 (v/v) [7c] and the products eluted with chloroform.

The model compounds were reacted with sodium borohydride (Merck-Schuchardt, Hohenbrunn, p.s.) and sodium dithionite (technical grade, Merck, Darmstadt). The reaction mixtures were worked up by extraction with 1.5 vol of chloroform after addition of 1.5 vol. glycine/HCl-buffer, pH 2.7, saturated with ammonium sulfate in the case of free acids [5]. The chloroform phase was briefly dried over NaCl and after filtration on cotton, uv-vis spectra were taken. The samples were dried in a stream of nitrogen for tlc.

Preparative scale reaction of biliverdin dimethylester (3c) with dithionite

A solution of 10 mg **3c** in 10 ml methanol was added to a solution of 50 mg sodium dithionite in 10 ml doubly glass distilled water under a nitrogen atmosphere. After 10 min, 20 ml glycine/HCl-buffer, pH 2.7 [5] were added and nitrogen was bubbled through the solution for further 30 min. The mixture was extracted twice with 10 ml chloroform, with some yellow pigment remaining in the aqueous phase. Preparative tlc of the organic extracts on silica gel (0.75 mm) with benzene/ethanol = 25:2 (v/v) yielded four zones: I, yellow, $R_F = 0$; II, yellow, $R_F = 0.3-0.5$; III, yellow with greenish and brownish components, $R_F = 0.65$; IV, blue, $R_F = 0.70$ (identified as the starting material by analytical tlc

with system "a" of Table I). Bands II–IV were eluted with acetone, elution of zone I was impossible. Zones II and III were each fractionated by column chromatography (2 × 5 cm valumina, neutral, activity super 1; Woelm, Eschwege): zone II: the main fraction was eluted with chloroform, a minor fraction with chloroform/methanol 9:1, zone III: two fractions were separated with chloroform, one with chloroform/methanol 9:1, a further zone could not be eluted with methanol.

Preparative scale reaction of biliverdin (3a) with dithionite

7.8 mg biliverdin were dissolved in 50 ml hot methanol and after cooling, added to a solution of 500 mg sodium dithionite in 50 ml water under a continuous stream of nitrogen. 400 ml glycine/HCl-buffer, saturated with ammonium sulfate, pH 2.7 [5], were immediately added and the aqueous phase was extracted four times with a total volume of 300 ml of chloroform. The chloroform phase was concentrated in vacuo to 30 ml and fractionated by tlc on silica gel plates (0.25 mm) with benzene/ethanol = 25:2 (v/v). A zone with green, brown and violet coloured substances with a yellow margin remained at the start. The only major zone (orange $R_F = 0.4$) was eluted with 130 ml chloroform to yield 0.8 mg crude bilirubin (**4a**); it was esterified with CH₂N₂ in CHCl₃. Chromatography on Al₂O₃ yielded the following fractions: I, yellow substance eluted with chloroform; II, main fraction, yellow substance eluted with CHCl₃/MeOH 9:1 = yield 0.3 mg; λ_{\max} (CHCl₃) = 403 nm, shoulder at 435 nm; identical in cochromatography with bilirubin-dimethylester (**4c**) with system a of Table I; III, yellow-greenish substance, eluted with CHCl₃ acetic acid = 1:1.

¹HMR-spectra of biliverdin dimethylester (**3c**), its reaction products with borohydride and dithionite, and of authentic bilirubin dimethylester (**4c**):

Preparation of the reactions products: **3c**/NaBH₄ and **3c**/dithionite: A solution of 6 mg biliverdin dimethylester (**3c**) in 2 ml CHCl₃ was shaken with 1.0 ml of an aqueous solution of 50 mg/ml dithionite until the colour changed to yellow-green. The CHCl₃ phase was washed with water, dried on NaCl and evaporated under a stream of nitrogen. An analogous experiment was carried out using sodium borohydride instead of dithionite.

Biliverdin dimethylester (**3c**): 1.80 (*exo*-CH₃); 2.01, 2.04, 2.10 (*endo*-CH₃); 2.56 t, 2.93 t (8.12-CH₂-CH₂); 3.61 (OCH₃); 5.28–6.61 (vinyl-protons); 5.94, 6.00 (5.15-CH); 6.73 (10-CH).

Bilirubin dimethylester (**4c**): 1.68 (*exo*-CH₃); 1.91, 2.02 (*endo*-CH); 2.39 t, 2.84 t (8.12-CH₂-CH₂); 3.63 (OCH₃); 4.10 (10-CH₂); 4.65–6.66 (vinyl protons); 5.84, 6.13 (5.15-CH); 10.04, 10.12, 10.42, 11.14 (NH).

3c/NaBH₄: 1.68 (*exo*-CH₃); 1.91, 2.01 (*endo*-CH₃); 2.38 t, 2.83 t (8.12-CH₂-CH₂); 3.63 (OCH₃); 4.09 (10-CH₂); 4.65–6.66 (vinyl protons); 5.84, 6.12 (5.15-CH); 10.03, 10.12, 10.41, 11.11 (NH).

3c/dithionite: Broad signals in the ranges 1.5–3.6 ppm and 5.1–6.4 ppm; no signal detectable in the range 3.7–5.0 ppm.

General methods:

UV-vis spectra were taken with a model DMR 22 spectrophotometer (Zeiss, Oberkochen), the spectra of compounds 9–12 with a model Superscan spectrophotometer (Varian, Palo Alto). ¹Hmr spectra were recorded in C ²HCl₃ with a model HFX 90 spectrometer (Bruker, Karlsruhe). Chemical shifts in δ (ppm) are given relative to Si(CH₃)₄ as internal standard. Signals appeared as singlets, unless otherwise noted. t = triplet. The vinyl protons appeared as ABX system.

Tris-glycine electrophoresis was done by the method of Davies [41] as modified by Wagenmann [42], SDS-electrophoresis with the system of Weber and Osborn [6]. Densitograms were obtained with a type TLD 100 densitometer (Vitatron). Analytical tlc was carried out on silica HPTLC plates and polyamide 11 F 254 plates (both Merck, Darmstadt). Preparative separations were performed on self-made 20 × 20 cm plates covered with 0.75 mm silica H (Merck, Darmstadt).

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft, Bonn-Bad Godesberg. We are indebted to Prof. W. Rüdiger for continuing support. We thank Prof. R. Gautron (Grenoble) for stimulating discussions and Dr. C. Petrier (Grenoble) for acquainting one of us (W.K.) with the preparation of isophorocobilin. The visit to Grenoble was made possible by a travel grant from the European Photochemistry Association. We thank Mrs. G. Schild for the measurement of the ¹HMR spectra, Mrs. H. Wieschoff and Mrs. C. Bubenzer for valuable technical assistance in the preparation of the phorocobilins and C-PC, respectively.

- [1] H. Scheer, *Angew. Chem.* **93**, 230 (1981); *Angew. Chem. Int. Ed.* **20**, 241 (1981).
- [2] W. Kufer and H. Scheer, *Z. Naturforsch.* **34 c**, 776 (1979).
- [3] W. Kufer and H. Scheer, *Hoppe-Seyler's Z. Physiol. Chem.* **360**, 935 (1979).
- [4] M. S. Stoll and C. H. Gray, *Biochem. J.* **163**, 59 (1977).
- [5] N. Blanckaert, K. P. M. Heirwegh, and F. Compernelle, *Biochem. J.* **155**, 405 (1976).
- [6] K. Weber and M. Osborn, *J. Biol. Chem.* **244**, 4406 (1969).
- [7] a) A. F. McDonagh and F. Assisi, *FEBS Lett.* **18**, 315 (1971). b) D. Manitto and P. Monti, *Experientia* **29**, 137 (1973). c) A. F. McDonagh and L. A. Palma, unpublished, cited in A. F. McDonagh in D. Dolphin (ed.), *The Porphyrins*, Vol. VI, p. 441, Academic Press, New York, 1979. d) Z. J. Petryka and C. J. Watson, *J. Chromatogr.* **37**, 76 (1968).
- [8] A. Gossauer and M. Blacha-Puller, *Liebigs Ann. Chem.* **1981**, 1492.
- [9] A. Gossauer and H. Plieninger, in: D. Dolphin (ed.), *The Porphyrins*, Vol. VI, p. 585, Academic Press, New York 1979.
- [10] C. H. Gray, A. Kulzycka, and D. C. Nicholson, *J. Chem. Soc.* **1961**, 2268.
- [11] A. F. McDonagh and F. Assisi, *Biochem. J.* **129**, 797 (1972).
- [12] B. Pullman and A. Pullman, *Quantum Biochemistry* p. 247, Wiley-Interscience, New York 1963.
- [13] a) A. R. Holzwarth, H. Lehner, S. E. Braslavsky, and K. Schaffner, *Liebigs Ann. Chem.* **1978**, 2002. b) H. Falk and T. Schlederer, *Monatsh. Chem.* **109**, 1013 (1978).
- [14] P. Manitto and D. Monti, *Experientia* **35**, 1418 (1979).
- [15] H. Falk, N. Müller, and T. Schlederer, *Monatsh. Chem.* **111**, 159 (1980).
- [16] H. Fischer and H. Plieninger, *Hoppe-Seyler's Z. Physiol. Chem.* **274**, 231 (1942).
- [17] A. J. Fatiadi and R. Schaffer, *Experientia* **27**, 1139 (1971).
- [18] E. C. Foulkes, R. Lemberg, and P. Purdom, *Proc. Roy. Soc. London B* **138**, 386 (1951).
- [19] G. Blankenhorn and E. G. Moore, *J. Amer. Chem. Soc.* **102**, 1092 (1980).
- [20] W. S. Caughey and K. A. Schellenberg, *J. Org. Chem.* **31**, 1978 (1966).
- [21] H. Lehner, W. Riemer, and K. Schaffner, *Liebigs Ann. Chem.* **1979**, 1798; S. E. Braslavsky, A. R. Holzwarth, E. Langer, H. Lehner, J. I. Matthews, and K. Schaffner, *Isr. J. Chem.* **20**, 196 (1980).
- [22] H. Lehner, S. E. Braslavsky, and K. Schaffner, *Liebigs Ann. Chem.* **1978**, 1990.
- [23] H. Falk, K. Grubmayr, and K. Thirring, *Z. Naturforsch.* **33 b**, 924 (1978).

- [24] C. Petrier, P. Jardon, C. Dupuy, and R. Gautron, *J. Chim. Phys.* **78**, 519 (1981).
- [25] H. Scheer and W. Kufer, *Z. Naturforsch.* **32 c**, 513 (1977).
- [26] P. Longhi, P. Manitto, D. Monti, T. Mussini, and S. Rondini, *Electrochim. Acta* **26**, 541 (1981).
- [27] H.-P. Köst, K. Rath, G. Wanner, and H. Scheer, *Photochem. Photobiol.* **30**, 139 (1981).
- [28] G. Blauer and G. Wagnière, *J. Am. Chem. Soc.* **97**, 1949 (1975).
- [29] G. Wagnière and G. Blauer, *J. Am. Chem. Soc.* **98**, 7806 (1976).
- [30] R. Troxler, private communication (1980).
- [31] W. Kufer and H. Scheer, *Angew. Chem.*, in press.
- [32] H. Fischer and H. W. Haberland, *Hoppe-Seyler's Z. Physiol. Chem.* **232**, 236 (1935).
- [33] R. Bonnett and A. F. McDonagh, *J. Chem. Soc. Perkin I* **1973**, 881 (1973).
- [34] W. Küster, *Hoppe-Seyler's Z. Physiol. Chem.* **141**, 40 (1924).
- [35] H. Fischer, H. Plieninger, and O. Weissbarth, *Hoppe-Seyler's Z. Physiol. Chem.* **268**, 197 (1941).
- [36] C. C. Kuenzle, *Biochem. J.* **119**, 395 (1970).
- [37] H. W. Whitlock jr., R. Hanauer, M. Y. Oester, and B. K. Bower, *J. Am. Chem. Soc.* **91**, 7485 (1969).
- [38] J. A. S. Cavaleiro and K. M. Smith, *J. Chem. Soc. Perkin I* **1973**, 2149 (1973).
- [39] C. Petrier, Dissertation, Université Scientifique et Médicale de Grenoble, 1978.
- [40] P. O'Carra and E. Colleran, *J. Chromatogr.* **50**, 458 (1970).
- [41] B. J. Davis, *Ann. N. Y. Acad. Sci.* **121**, 404 (1964).
- [42] R. Wagenmann, Dissertation, Universität München 1977.
- [43] IUPAC-IUB Joint Commission on Biochemical Nomenclature, *Pure Appl. Chem.* **51**, 2251 (1979).
- [44] T. A. Wooldridge and D. A. Lightner, *J. Liquid Chromatogr.* **1**, 653 (1978).
- [45] H. L. Crespi, U. Smith, and J. J. Katz, *Biochemistry* **7**, 2232 (1968).
- [46] R. Bonnett, J. E. Davies, M. B. Hursthouse, and G. M. Sheldrick, *Proc. Roy. Soc. (London) B* **202**, 249 (1978).
- [47] W. Becker and W. S. Sheldrick, *Acta Crystallogr. B* **34**, 1298 (1979).