

STEREOCHEMISTRY OF C-METHYLATION IN THE BIOSYNTHESIS OF RHODODENDRIN IN *ALNUS* AND *BETULA*

MARTINA KLISCHIES and MEINHART H. ZENK

Lehrstuhl für Pflanzenphysiologie, Ruhr-Universität Bochum, D 4630 Bochum, West Germany

(Received 12 December 1977)

Key Word Index—*Betula alba*; *Alnus glutinosa*; Betulaceae; cinnamic acid pathway; C-methylation; stereochemistry; rhododendrin; phenylpropanoid.

Abstract—Using differently labelled precursors, it was established that rhododendrin (3-(4-hydroxyphenyl)-1-methylpropyl- β -D-glucopyranoside) is formed through the phenylpropane pathway via *p*-coumaryl alcohol, dihydro-*p*-coumaryl alcohol and C-methylation of the γ -C-atom of the C₆-C₃ unit with methionine supplying the methyl group. It was demonstrated that the *pro*-(S)-hydrogen atom of dihydro-*p*-coumaryl alcohol is replaced stereospecifically by the methyl group.

INTRODUCTION

Rhododendrin (3-(4-hydroxyphenyl)-1-methylpropyl- β -D-glucopyranoside) was discovered for the first time in leaves of *Rhododendron chrysanthemum* [1] and later in the bark of *Betula* [2, 3]. More recently its distribution in the genus *Rhododendron* (Ericaceae) has been surveyed [4, 5] and it has been found in *Bergenia* (Saxifragaceae) [4]. No information regarding the biosynthesis of this interesting phenolic glucoside is available. Rhododendrin and its aglycone rhododendrol may be assumed to be derived from *p*-coumaric acid via dihydro-*p*-coumaryl alcohol and subsequent C-methylation of the γ -C-atom of the phenylpropyl alcohol. This latter reaction would create a chiral center with unknown stereochemical configuration. The availability of prochiral tritium labelled cinnamyl alcohols [6] of known stereochemistry [7] and high configurational purities make it possible to attempt to study the steric course of the assumed C-methylation and furthermore to clarify the relative configuration of rhododendrol and its glucoside.

RESULTS AND DISCUSSION


A survey of the biosynthetic capacity of different rhododendrin-containing species was made in order to select the most suitable experimental material. *Rhododendron chrysanthemum* as well as *Bergenia speciosa* proved to be unsuitable for incorporation experiments with labelled precursors due to low metabolic rate or seasonal variations. However, 9-month-old seedlings of *Betula alba* proved to be well suited and material was available all the year round. Another member of the Betulaceae, *Alnus glutinosa*, was found to contain rhododendrin and this plant also proved to be suitable for incorporation experiments at the 9 month seedling stage. The occurrence of rhododendrin in *Alnus* and *Betula* was established by TLC on 5 solvents, UV spectroscopy and conversion to rhododendrol by β -glucosidase followed by subsequent GC/MS of the aglycone. Rhododendrin was present

Table 1. Incorporation of potential ¹⁴C and ³H labelled precursors into rhododendrin of *Alnus glutinosa* and *Betula alba*

Precursor applied	(μ mol)	Total act. (dpm $\times 10^6$)	Isolated (μ mol)	Rhododendrin Sp. act (dpm/ μ mol)	Incorporation (%)
<i>p</i> -coumaric acid [U- ¹⁴ C]*	0.05	5.55	1.24	8309	0.34
<i>p</i> -coumaric acid [U- ¹⁴ C]†	0.05	5.55	1.04	10673	0.22
dihydro- <i>p</i> -coumaric acid [U- ¹⁴ C]*	0.04	4.22	1.90	13089	1.32
dihydro- <i>p</i> -coumaric acid [U- ¹⁴ C]†	0.03	2.55	1.60	20558	0.78
<i>p</i> -coumaryl alcohol [U- ¹⁴ C]*	0.07	0.15	1.28	941	1.25
dihydro- <i>p</i> -coumaryl alcohol [U- ¹⁴ C]*	0.20	0.19	1.27	1466	0.98
dihydro- <i>p</i> -coumaryl alcohol [U- ¹⁴ C]†	0.07	0.15	1.40	300	0.28
L-methionine [methyl- ³ H]*	1.00	2.22 $\times 10^2$	2.00	85665	0.08

* *Alnus glutinosa* † *Betula alba*

at 0.15% in *A. glutinosa* and 0.30% in *B. alba* based on dry matter. In order to establish the pathway for the biosynthesis of rhododendrin, a series of ¹⁴C- or ³H-labelled precursors were fed to young terminal shoots of both *A. glutinosa* and *B. alba* and the degree of incorporation was measured. As shown in Table 1, both *p*-coumaric acid and its dihydro derivative were satisfactorily incorporated into the glucoside. The same holds true for both coumaryl alcohol and dihydrocoumaryl alcohol. Thus the phenylpropane pathway is involved in the biosynthesis of rhododendrin. From the incorporation data it is, however, not possible to distinguish between pathway A or B in the Scheme 1. Surprisingly both the acid and the alcohol containing either a double bond between the α - and β -C-atom of the side chain or the fully hydrogenated derivatives are incorporated approximately equally. This shows that the enzymes involved in the hydrogenation of the double bond of either *p*-coumaric acid or its alcohol exhibit a considerable lack of specificity. The assumed precursor of rhododendrin, dihydro-*p*-coumaryl alcohol, may



Scheme 1. Proposed pathway for the biosynthesis of rhododendrin.

therefore be formed *in vivo* either by reduction of *p*-coumaric acid to dihydro-*p*-coumaric acid followed by activation and reduction of the carboxyl group [8] or by reduction of *p*-coumaryl alcohol. Thus the exact mechanism can only be decided by using cell-free systems. The incorporation of the C₆-C₃ units into rhododendrin accounts for the biosynthetic origin of six of the total of seven required carbon atoms. In order to determine the origin of the missing carbon, L-methionine (—C³H₃) was supplied to the shoot of *A. glutinosa* and an incorporation of 0.08% was observed (Table 1). In order to prove that the methyl group of methionine was incorporated into the methyl group of the glucoside, rigorously purified rhododendrin (51 339 dpm) was diluted and subjected to Kuhn-Roth oxidation. The resulting acetate, which is derived from the methyl group and the adjacent carbon atom carrying the alcohol function (corresponding to γ -C of dihydro-*p*-coumaryl alcohol), contained 506 dpm/ μmol (theoretically 555 dpm/ μmol) and thus this compound carried 91% of the required radioactivity. This proves that the methyl group of rhododendrin is derived from the methyl group of methionine. Thus the original hypothesis on the biosynthesis of rhododendrin was confirmed in that dihydro-*p*-coumaryl alcohol is C-methylated at the alcohol function of the phenylpropyl derivative. The possibility that the C-methylation occurs at the level of the *p*-coumaryl alcohol and that the methylated propene derivative is subsequently reduced is unlikely but cannot be excluded.


If there is a stereospecific step involved in the course of the methylation, clearly one hydrogen atom of the alcohol group, either the *pro-R* or the *pro-S*, has to be removed and replaced by the methyl group derived from S-adenosylmethionine. In order to prove this

reaction and to establish its stereochemical course, if any, tritium labelled *R*- and *S*-*p*-coumaryl alcohol and dihydro-*p*-coumaryl alcohol were synthesized according to known methods [6]. The introduction of the label was achieved using the highly stereospecific horse liver alcohol dehydrogenase system [7].

In a preliminary experiment doubly labelled (*R,S*)-(γ -³H)-[U-¹⁴C]-*p*-coumaryl alcohol was fed to a rigorously growing shoot of *A. glutinosa*. The ³H/¹⁴C ratio of the precursor was 18.31:1. The labelled rhododendrin formed in this experiment was isolated, purified and was found to have a ³H/¹⁴C ratio of 10.30:1. This result corresponds to a loss of about 45% of the tritium label (no isotope effects were taken into consideration) which is in accordance with the assumption that one of the hydrogen atoms of the γ -C-atom is replaced by the methyl group introduced from methionine. The crucial experiment consisted in feeding the two pure labelled enantiomers to the plants. It was observed in both *Alnus* as well as in *Betula* that the (*R*)-*p*-coumaryl alcohol was incorporated into rhododendrin to an extent of 2.7%

Table 2. Incorporation of (*R*)-(γ -³H)- and [U-¹⁴C]- as well as (*S*)-(γ -³H)- and [U-¹⁴C]-dihydro-*p*-coumaryl alcohol into rhododendrin in *Alnus glutinosa* and *Betula alba*

Plant species	Precursor	Rhododendrin				
		³ H/ ¹⁴ C ratio	Incorporation (%) ³ H	¹⁴ C	Loss of ³ H (%)	
<i>A. glutinosa</i>	{ (<i>R</i>)-(γ - ³ H)/(U- ¹⁴ C)	9 86 1	0 96	0 98	9 59 1	2 7
	{ (<i>S</i>)-(γ - ³ H)/(U- ¹⁴ C)	24 00 1	0 08	0 96	2 35 1	90 2
<i>B. alba</i>	{ (<i>R</i>)-(γ - ³ H)/(U- ¹⁴ C)	6 18 1	0 23	0 22	6 26 1	0
	{ (<i>S</i>)-(γ - ³ H)/(U- ¹⁴ C)	12 88 1	0 09	0 28	3 17 1	75 39



Scheme 2. Biosynthetic incorporation of (*R*)-(γ - ^3H)- and (*S*)-(γ - ^3H)-dihydro-*p*-coumaryl alcohol into rhododendrin in *Alnus glutinosa* and *Betula alba*.

and 5.4% respectively, while the (*S*)-enantiomer was incorporated only to an extent of 0.08% and 0.8% respectively. The much lower incorporation of the (*S*)-enantiomer suggested that it is the *pro*-(*S*) hydrogen atom which is replaced by methylation. However, final proof that this assumption is correct came from an experiment using $^3\text{H}/^{14}\text{C}$ doubly labelled pure enantiomers of both *p*-coumaryl alcohol and dihydro-*p*-coumaryl alcohol.

The results of the double label experiment are shown in Table 2. This clearly shows which of the γ -hydrogen atoms is replaced. Invariably in both plants the *pro*-*R*-hydrogen atom of dihydroxy-coumaryl alcohol was completely retained and the *pro*-*S*-hydrogen atom was largely lost.

These experiments prove that during the enzymatically catalyzed methylation of the γ -C-atom of either *p*-coumaryl alcohol or dihydro-*p*-coumaryl alcohol the *pro*-*S* hydrogen atom is stereospecifically removed according to Scheme 2.

EXPERIMENTAL

Plant material. *A. glutinosa* and *B. alba* was purchased from the firm Borntäger (Offstein, Germany). The plants were grown in a Phytotron chamber with a photoperiod of 16 hr day (2500 lx), 8 hr night and a 23°–18° day–night temperature cycle at 70–75% rel. hum. The plants were used at an age of 9 months.

Feeding and isolation procedure. The labelled compounds were fed through the cut end of an excised young terminal shoot (ca 4 cm in height). The tracer was usually allowed to be metabolized for 24 hr. Subsequently the shoot was cut into 0.5 cm pieces and exhaustively extracted with boiling 80% EtOH. The extract was purified first by preparative layer (1.5 mm)

chromatography in $\text{CHCl}_3\text{-MeOH}$ (4:1) and then by TLC in $\text{C}_6\text{H}_6\text{-EtOH}$ (8:2); $\text{CHCl}_3\text{-EtOH}$ (7.7:1); $\text{EtOAc-MeCOEt-HCOOH-H}_2\text{O}$ (5:3:1:1:1), rhododendrin R_f values 0.4; 0.60; 0.25; 0.72, respectively. The ϵ_{278} (MeOH) was $1.59 \times 10^6 \text{ M}^{-1} \text{ cm}^{-1}$. Rhododendrin was converted to rhododendrol by incubation with β -glucosidase (Boehringer), the aglycone was then ether extracted from the incubation mixture and purified by TLC using $\text{CHCl}_3\text{-EtOH}$ (7.7:1). The R_f was 0.6. Both the glycoside and the aglycone gave a red coloration on spraying with conc H_2SO_4 . Rhododendrol was further characterised by GLC (3% SE 30 on supelcoport, He 36 ml/min, programme 100–200°, 10°/min). The rhododendrol retention time was 6.7 min and MS m/e : 166 (M^+), 148 ($\text{M-H}_2\text{O}$), 133, 107.

Synthesis and degradation of labelled compounds. [$\text{U-}^{14}\text{C}$]-L-tyrosine was deaminated using PAL to *p*-coumaric acid which was then hydrogenated (Pd/H_2) to yield dihydro-*p*-coumaric acid. [$\text{U-}^{14}\text{C}$]-*p*-coumaric acid was transformed by standard methods into *p*-coumaryl alcohol via acetyl-*p*-coumaric acid, acetyl-*p*-coumaryl chloride reduction, and alkaline hydrolysis. The labelled alcohol was further hydrogenated to dihydro-*p*-coumaryl alcohol (Pd/H_2). The stereospecifically labelled alcohols were synthesized to the corresponding coniferyl alcohols [6]. (*R,S*)-(γ - ^3H)-*p*-coumaryl alcohol was synthesized from *p*-coumaryl aldehydes by reduction with NaBH_3T (yield via ^3H 22.4%; sp. act. 239 $\mu\text{Ci}/\mu\text{mol}$). (*S*)-(γ - ^3H)-*p*-coumaryl alcohol (25% yield, sp. act. after dilution 28 $\mu\text{Ci}/\mu\text{mol}$) *p*-coumaryl aldehyde (yield 45%, sp. act. 119.5 $\mu\text{Ci}/\mu\text{mol}$). (*R*)-(γ - ^3H)-*p*-coumaryl alcohol via ^3H -cyclohexanol [7] (yield 9.2% sp. act. 1.12 $\text{mCi}/\mu\text{mol}$). All labelled *p*-coumaryl alcohol species were hydrogenated ($\text{Pd}:\text{H}_2$) at 70–80% yield into the correspondingly labelled dihydro-*p*-coumaryl alcohol with unchanged sp. act. (MS m/e : 152 (M^+), 134 ($\text{M-H}_2\text{O}$), 107, 77). Kuhn-Roth degradation was conducted under conditions given in [9].

Acknowledgements—We thank Prof. H. Thieme, Leipzig, for a generous gift of pure rhododendrin, Dr. G. Wilson, Dublin,

for his help in the preparation of the English version of this manuscript and the 'Deutsche Forschungsgemeinschaft' for financial support.

REFERENCES

1. Archangelsky, K. (1901) *Arch. Exp. Path. Pharmacol.* **46**, 313.
2. Sosa, A. (1933) *Chem. Zentralblatt* **II**, 1193.
3. Kim, K.-U. (1942) *J. Pharm. Soc. Japan* **62**, 455.
4. Thieme, H. and Winkler, H. J. (1969) *Pharmazie* **24**, 703.
5. Thieme, H., Walewska, E. and Winkler, H. J. (1969) *Pharmazie* **24**, 648.
6. Harborne, J. B. and Williams, C. A. (1971) *Phytochemistry* **10**, 2727.
7. Klischies, M., Stöckigt, J. and Zenk, M. H. (1978) *Phytochemistry* (in press).
8. Battersby, A. R., Staunton, J. and Wiltshire, H. R. (1975) *J. Chem. Soc. Perkin Trans.* **1** 1156.
9. Gross, G. G., Stöckigt, J., Mansell, R. L. and Zenk, M. H. (1973) *FEBS Lett.* **31**, 283.
10. Durand, R. and Zenk, M. H. (1974) *Phytochemistry* **14**, 1483.