

Chemische Berichte

Inhalt

Contents

A

Teil A: Anorganische und elementorganische Chemie / Part A: Inorganic and Organometallic Chemistry

- Bromm, D., Seibold, U., Noltemeyer, M., Meller, A. 2645 ▶ 1,2,3-Triborolane aus Norbornenderivaten, Allylbenzol und Dehalogenierungsprodukten von Dichlor(diisopropylamino)-boran; 2,4,5-Tris(diisopropylamino)-2,4,5-triborabicyclo[1.1.1]pentan — Bicyclus oder Carboran?
- Klingebiel, U., Pauer, F., Sheldrick, G. M., Stalke, D. 2651 ▶ 1,3-Bis[*tert*-butyl(di-*tert*-butylfluorsilyl)-amino]trisulfan — eine Verbindung mit FSiN₃SSNSiF-Helix
- Olms, P., Roesky, H. W., Keller, K., Noltemeyer, M., Bohra, R., Schmidt, H.-G., Stalke, D. 2655 ▶ Synthesen und Strukturen von cyclischen und acyclischen Vanadium(V)- und Molybdän(VI)-haltigen Verbindungen
- Herberich, G. E., Englert, U., Roos, P. 2663 ▶ Neue Cyclooctatetraen-Komplexe des Niobiums
- Proft, B., Pörschke, K.-R., Lutz, F., Krüger, C. 2667 ▶ 1,6-Heptadien-Nickel(0)-Komplexe: *rac/meso*-($\mu\text{-}\eta^2,\eta^2\text{-C}_7\text{H}_{12}$) $\{\text{Ni}(\eta^2,\eta^2\text{-C}_7\text{H}_{12})\}_2$ und L-Ni($\eta^2,\eta^2\text{-C}_7\text{H}_{12}$)
- Goede, S. J., Bickelhaupt, F. 2677 ▶ Synthese und Reaktionen von *P*-Supermesityl-C-halophosphaalkenen
- Nagy-Magos, Z., Feßenbecker, A., Pritzkow, H., Siebert, W., Hyla-Kryspin, I., Gleiter, R. 2685 ▶ Komplexierung von 1,3-Dihydro-1,3,10,10-tetramethyl-1,3-dibora-4,5-benzopentafulven mit Fe(CO)₅- und CpCo-Fragmenten: Synthese, Kristall- und Elektronenstruktur von Tricarbonyl[(η^5 -cyclopentadienyl)cobalt][$\mu,\eta^5\text{:}\eta^4$ -(1,3-dihydro-1,3,10,10-tetramethyl-1,3-dibora-4,5-benzopentafulven)]jiron
- Lindner, E., Fawzi, R., Hiller, W., Carvill, A., McCann, M. 2691 ▶ Synthese, Kristallstruktur und Beweglichkeit von *fac*-[Ru(CH₃CO₂)₂(PPh₃)₃] · 2 CH₃CO₂H
- Hansert, B., Powell, A. K., Vahrenkamp, H. 2697 ▶ Azoalkan- und Nitren-verbrückte Carbonylmétall-Cluster des Eisens und Rutheiniums
- Vogt, R., Jones, P. G., Kolbe, A., Schmutzler, R. 2705 ▶ Verbindungen mit dem 2,4-Dimethyl-2,4-diaza-1,5-diphosphapentan-3-on-Grundgerüst: Synthese, Struktur und Eigenschaften von 1,5-Di-*tert*-butyl-2,4-dimethyl-1,5-diphenyl-2,4-diaza-1,5-diphosphapentan-3-on und Derivaten
- Köster, R., Seidel, G., Wrackmeyer, B., Bläser, D., Boese, R., Bühl, M., von Raguer Schleyer, P. 2715 ▶ Decaethyl-2,6,8,10-tetracarba-*nido*-decarboran(10) — Herstellung, Struktur im festen Zustand und Stabilität
- Jones, P. G., Thöne, C. 2725 ▶ Gold-Komplexe mit Selen-Liganden, IV. — Herstellung, Kristallstrukturen und Reaktionen von Phosphan(selenoharnstoff)-gold(I)-Komplexen
- Davies, S. J., Compton, N. A., Huttner, G., Zsolnai, L., Garner, S. E. 2731 ▶ Synthese und Reaktivität von „Bismuthiniden“-Verbindungen und die Bildung von Bi¹-Chelat-Komplexen
- ▶ Publikationssprache
- 1,2,3-Triborolanes from Norbornene Derivatives, Allylbenzene and Dehalogenation Products of Dichlorodiisopropylamino-boran; 2,4,5-Tris(diisopropylamino)-2,4,5-triborabicyclo[1.1.1]pentane — Bicyclic or Carborane?
- 1,3-Bis[*tert*-butyl(di-*tert*-butylfluorosilyl)-amino]trisulfane — a Compound with FSiN₃SSNSiF Helix
- Syntheses and Structures of Cyclic and Acyclic Compounds Containing Vanadium(V) and Molybdenum(VI)
- New Cyclooctatetraene Complexes of Niobium
- 1,6-Heptadiene Nickel(0) Complexes: *rac/meso*-($\mu\text{-}\eta^2,\eta^2\text{-C}_7\text{H}_{12}$) $\{\text{Ni}(\eta^2,\eta^2\text{-C}_7\text{H}_{12})\}_2$ and L-Ni($\eta^2,\eta^2\text{-C}_7\text{H}_{12}$)
- Synthesis and Reactions of *P*-Supermesityl-C-halophosphaalkenes
- Complexation of 1,3-Dihydro-1,3,10,10-tetramethyl-1,3-dibora-4,5-benzopentafulven with Fe(CO)₅ and CpCo Fragments: Synthesis, Crystal and Electronic Structure of Tricarbonyl[(η^5 -cyclopentadienyl)cobalt][$\mu,\eta^5\text{:}\eta^4$ -(1,3-dihydro-1,3,10,10-tetramethyl-1,3-dibora-4,5-benzopentafulven)]jiron
- Synthesis, X-Ray Crystal Structure, and Fluxional Behaviour of *fac*-[Ru(CH₃CO₂)₂(PPh₃)₃] · 2 CH₃CO₂H
- Azoalkane- and Nitrene-Bridged Carbonyl Metal Clusters of Iron and Ruthenium
- Compounds with the 2,4-Dimethyl-2,4-diaza-1,5-diphosphapentan-3-one Skeleton: Synthesis, Structure, and Properties of 1,5-Di-*tert*-butyl-2,4-dimethyl-1,5-diphenyl-2,4-diaza-1,5-diphosphapentan-3-one and Derivatives
- Decaethyl-2,6,8,10-tetracarba-*nido*-decarborane(10) — Preparation, Structure in the Solid State, and Stability
- Gold Complexes with Selenium Ligands, IV. — Preparation, Crystal Structures and Reactions of Phosphine(selenourea)gold(I) Complexes
- Synthesis and Reactivity of “Bismuthinidene” Compounds and the Formation of Bi-Chelate Complexes
- Language of Publication

Kumberger, O., Riede, J., Schmidbaur, H.	2739	Orotat-Komplexe, II. – Herstellung und Kristallstrukturen von Calcium- und Zink-orotat(2-)hydraten	► Orotate Complexes, II. – Preparation and Crystal Structures of Calcium and Zinc Orotate(2-) Hydrates
--	------	--	--

Notizen / Notes

Mikulcik, P., Riede, J., Schmidbaur, H.	2743	Metall-Ionen-Fixierung durch Dipeptide: Struktur des durch „Aspartam“-Hydrolyse gebildeten Produkts, Natrium- <i>cyclo(L-α-aspartyl-L-phenylalanin)-tetrahydrats, Na[c(L-Asp-L-Phe)] · 4 H₂O</i>	► Metal Ion Binding by Dipeptides: Structure of the Product Generated in “Aspartame” Hydrolysis, Sodium <i>cyclo(L-α-Aspartyl-L-phenylalanine)</i> Tetrahydrate, Na[c(L-Asp-L-Phe)] · 4 H ₂ O
Kreißl, F. R., Stegmair, C. M.	2747	► Carbonyl(η^5 -cyclopentadienyl)(trimethylphosphane)[η^2 -(methylthio)phenylcarben]-molybdän-tetrafluoroborat – ein amphiphiler Carbenkomplex	Carbonyl(η^5 -cyclopentadienyl)(trimethylphosphane)[η^2 -(methylthio)phenylcarbene]molybdenum Tetrafluoroborate – an Amphiphilic Carbene Complex

Querverweise / Cross References

Regitz, M. et al.	2843	► Organophosphorverbindungen, 51	Organophosphorus Compounds, 51
Grevels, F.-W., Özkar, S. et al.	2857	► Photoinduzierte Reaktionen von Cr(CO) ₃ -koordiniertem 1,3,5-Cycloheptatrien	Photoinduced Reactions of Cr(CO) ₃ -Coordinated 1,3,5-Cycloheptatriene

B

Teil B: Organische Chemie / Part B: Organic Chemistry

Roth, W. R., Langer, R., Ebbrecht, T., Beitat, A., Lennartz, H.-W.	2751	► Zur Energie-Delle von Diradikalen, III. – 2,3,5,6-Tetramethylen-1,4-cyclohexadiyl	Energy Well of Diradicals, III. – 2,3,5,6-Tetramethylene-1,4-cyclohexadiyl
Bestmann, H. J., Schaper, W., Ruppert, D., Zimmermann, R., Burzlaff, H.	2761	► Phosphinalkylen, 49. – Chemie und Stereochemie des 9,10,19,20-Tetrahydrotetrabenzo[<i>a,c,g,i</i>]cyclododecens und einiger seiner Derivate	Phosphine Alkylidenes, 49. – Chemistry and Stereochemistry of 9,10,19,20-Tetrahydrotetrabenzo[<i>a,c,g,i</i>]cyclododecene and of Some Derivatives
Bestmann, H. J., Schaper, W., Holzmann, H. G., Zimmermann, R.	2773	► Synthese und Reaktionen der Stereoisomeren von Tetrabenzo- und Tetranaphtho[<i>a,c,g,i</i>]cyclododecen	Synthesis and Reactions of the Stereoisomers of Tetrabenzo- and Tetranaphtho[<i>a,c,g,i</i>]cyclododecene
Liebscher, J., Riemer, B., Hassoun, A., Stößer, R., Grubert, L.	2781	► Synthese neuartiger Bispyrimido[1,2- <i>b</i> :1',2'- <i>e</i>][1,2,4,5]tetrazine	Synthesis of Novel Bispyrimido[1,2- <i>b</i> :1',2'- <i>e</i>][1,2,4,5]tetrazines
Ohm, S., Bäuml, E., Mayr, H.	2785	► Eine vielseitige Synthese von 1,4-Dienen: Verwendung von Vinylether als Vinyl-Kation-Äquivalente	A Versatile Synthesis of 1,4-Dienes: Use of Vinyl Ethers as Vinyl Cation Equivalents
Reichardt, C., Schäfer, G., Allmann, R., Krestel, M., Knecht, J., Gärtner, R.	2791	► Über die Konformation eines trinuclearen [1.1.1]Tetramethinium-Cyaninfarbstoffs mit Indolin-Endgruppen	Conformation of a Trinuclear [1.1.1]Tetramethinium Cyanine Dye with Indoline End Groups
Schulz, J., Nieger, M., Vögtle, F.	2797	► Regioselektive Komplexierung von [2.2]-Metacyclophanen mit Triaminetricarbonylchrom	Regioselective Complexation of [2.2]Metacyclophanes with Triaminetricarbonylchromium
Sustmann, R., Korth, H.-G., Nüchter, U., Siangouri-Feulner, I., Sicking, W.	2811	► Sind Charge-Transfer-Komplexe Zwischenstufen bei Diels-Alder-Reaktionen? Eine Fallstudie über die Reaktion von 1,2-Dimethylenyclopentan mit Tetracyanoethylene	Are Charge-Transfer Complexes Intermediates in Diels-Alder Reactions? A Case Study of the Reaction of 1,2-Dimethylenecyclopentane with Tetracyanoethylene
Auchter-Krummel, P., Krummel, G., Lex, J., Müllen, K.	2819	► Reduktive Umwandlungen, 18. – Cycloannellierung und Überbrückung von Cyclooctatetraen	Reductive Transformations, 18. – Cycloannulation and Bridging of Cyclooctatetraene
Schade, P., Schäfer, T., Müllen, K., Bender, D., Knoll, K., Bronstert, K.	2833	► Reduktive Umwandlungen, 19. – Elektron-Transfer-induzierte Dimerisierung von β -Alkylstyrolen und die Strukturen der resultierenden 1,4-Dilithiobutane	Reductive Transformations, 19. – Electron-Transfer-Induced Dimerization of β -Alkylstyrenes and the Structures of the Resulting 1,4-Dilithiobutanes
Fuchs, E., Breit, B., Heydt, H., Schoeller, W., Busch, T., Krüger, C., Betz, P., Regitz, M.	2843	► Organophosphorverbindungen, 51. – Phosphatriafulvene – Phosphaalkene mit inverser Elektronendichte	Organophosphorus Compounds, 51. – Phosphatriafulvenes – Phosphaalkenes with Inverse Electron Density

Inhalt (Fortsetzung)

Contents (Continued)

Fischler, I., Grevels, F.-W., Leitich, J., Özkar, S.	2857	Photoinduzierte Reaktionen von Cr(CO) ₃ -koordiniertem 1,3,5-Cycloheptatrien: [6 + 2]-Cycloaddition mit einem Alkin und katalytische 1,6-Hydrierung	Photoinduced Reactions of Cr(CO) ₃ -Coordinated 1,3,5-Cycloheptatriene: [6 + 2] Cycloaddition with an Alkyne and Catalytic 1,6-Hydrogenation
Elsässer, D., Hassenrück, K., Martin, H.-D., Mayer, B., Lutz, G., Prinzbach, H.	2863	Kleine und mittlere Ringe, 77. — Sich der klassischen π,π -Konjugation nähernende nichtkonjugierte π,π -Wechselwirkungen — PE-spektroskopische Untersuchungen des 7,8;12,19-Diseco-1,16-dodecahedradien-System	Small and Medium Rings, 77. — Nonconjugated π,π Interactions Approaching Classical π,π Conjugation — PE-Spectroscopic Investigation of the 7,8;12,19-Diseco-1,16-dodecahedradiene System
Albert, B., Elsässer, D., Martin, H.-D., Mayer, B., Chow, T. J., Marchand, A. P., Ren, C.-T., Paddon-Row, M. N.	2871	Kleine und mittlere Ringe, 78. — Käfig-Dimere von Norbornadien mit rechtwinkliger Anordnung von Subchromophoren: Orbital-Wechselwirkung im Heptacyclo-[6.6.0 ^{2,6} .0 ^{3,13} .0 ^{4,11} .0 ^{5,9} .0 ^{10,14}]tetradecan-System	Small and Medium Rings, 78. — Cage Dimers of Norbornadiene with Perpendicular Arrangement of Subchromophores: Orbital Interaction in the Heptacyclo-[6.6.0 ^{2,6} .0 ^{3,13} .0 ^{4,11} .0 ^{5,9} .0 ^{10,14}]tetradecane System
Huda, E., Martin, H.-D., Mayer, B., Somnitz, K.-H., Steigel, A., Haddad, H., Distefano, G., Modelli, G.	2879	Kleine und mittlere Ringe, 79. — 4-Aza-tetracyclo[3.3.0.0 ^{2,8} .0 ^{3,6}]octane: Neuer Heterocyclen durch Addition von Sulfonylaziden an 7-substituierte Norbornadiene — Experimentelle Bestätigung von σ^* -Wechselwirkungen im Norbornan-Gerüst	Small and Medium Rings, 79. — 4-Aza-tetracyclo[3.3.0.0 ^{2,8} .0 ^{3,6}]octanes: New Heterocycles by Addition of Sulfonyl Azides to 7-Substituted Norbornadienes — Experimental Verification of σ^* Interactions in the Norbornane Skeleton
Beck, A. K., Seebach, D.	2897	► Aldol- und Michael-Additionen fluorierter Nitroalkane an Aldehyde, Ketone und α,β -ungesättigte Carbonylverbindungen	Aldol and Michael Additions of Fluorinated Nitroalkanes to Aldehydes, Ketones, and α,β -Unsaturated Carbonyl Compounds

Notiz / Note

Drewes, S. E., Malissar, D. G. S., Roos, G. H. P.	2913	Kristalline, enantiomerenreine Aldole aus einem von (–)-Ephedrin abgeleiteten <i>N</i> -Acylimidazolidin-2-on	► Crystalline, Enantiomerically Pure Aldols from a (–)-Ephedrine-Derived <i>N</i> -Acylimidazolidin-2-one
---	------	---	---

Querverweis / Cross Reference

Schmidbaur, H. et al.	2743	Metall-Ionen-Fixierung durch Dipeptide	► Metal Ion Binding by Dipeptides
		► Publikationssprache	► Language of Publication

A Versatile Synthesis of 1,4-Dienes: Use of Vinyl Ethers as Vinyl Cation Equivalents*

Stefan Ohm, Engbert Bäuml, and Herbert Mayr*

Institut für Chemie, Medizinische Universität zu Lübeck,
Ratzeburger Allee 160, W-2400 Lübeck 1, F.R.G.

Received June 3, 1991


Key Words: Enol ether / Allylsilane / Bromo ether / 1,4-Diene / Alkoxyalkylation

The α,β -dibromo ethers **2**, in situ generated from the vinyl ethers **1** and bromine, react with the allylsilanes **3** in the presence of $ZnCl_2$ to give the β -bromo ethers **4**. Treatment of **4**

with Na in diethyl ether yields 1,4-dienes **5** in 62–79% overall yield.


Most syntheses of 1,4-dienes proceed by combination of vinylic nucleophiles with allylic electrophiles¹⁾, and examples for the coupling of a vinylic electrophile with an allylic nucleophile are rare¹⁾. A general method for the synthesis of 1,4-dienes using the latter approach has recently been described by Lipshutz (Scheme 1)²⁾.

Scheme 1



We now describe another example for constructing 1,4-dienes from vinyl cation and allyl anion equivalents, where the readily available vinyl ethers are used as vinyl cation equivalents (Scheme 2).

Scheme 2




For this purpose, the reaction sequence depicted in Scheme 3 has been elaborated. A comparison of structures **1** and **5** shows that, overall, the alkoxy group of vinyl ethers is replaced by an allylic group.

Formation of the β -Bromo Ethers **4**

In view of the high tendency of vinyl ethers to undergo cationic polymerization, the 1,2-dibromo ethers **2** have been prepared by dropwise addition of the vinyl ethers **1** to so-

Scheme 3



lutions of bromine in CH_2Cl_2 at $-78^\circ C$. Even under these conditions a small excess of **1** over bromine is usually needed to achieve complete decolorization of the bromine solution (see Experimental). The highly reactive and *presumably toxic* dibromo ethers **2** have not been isolated, and the β -bromo ethers **4** have been produced in a one-pot reaction by adding $ZnCl_2$ –Et₂O³⁾ and the allylsilanes **3** to the dichloromethane solutions obtained by bromination of the vinyl ethers⁴⁾. Compounds **4** have either been purified by distillation or immediately converted into the 1,4-dienes **5**.

All combinations of enol ethers and allylsilanes examined in this work gave the β -bromo ethers **4** in good yield. Contaminations observed in some runs were due to the use of moist bromine, leading to HBr adducts of the vinyl ethers, which then reacted with the allylsilanes **3** to give debrominated analogs of **4** (e.g. **7**). As the stereochemistry of the intermediate bromo ethers **4** is usually not important for the structure of the elimination products **5**, the diastereoselectivities of the C–C bond-forming reactions have generally not been investigated. Only in the case of **4c**, a ¹³C-NMR spectrum has been taken, which showed the highly selective formation of one diastereomer ($\approx 95:5$) from a 2:1 mixture of diastereomeric dibromo ethers **2c**.

Elimination Reactions

The production of alkenes by Zn-promoted elimination of halogen and alkoxy from β -halogeno ethers has long been known⁵⁾. The reaction conditions were optimized for the conversion $4d \rightarrow 5d$, and it was found that a mixture

of $4d$ and 5 equivalents of zinc powder in ethanol or methanol had to be refluxed for approximately 20 hours to achieve complete consumption of the bromo ether. A slight retardation of the elimination was observed, when 4 mol-% of $ZnCl_2$ was added to the reaction mixture.


Table 1. 1,4-Dienes 5 from vinyl ethers 1 and allylsilanes 3

Vinyl Ethers	Allylsilanes	β -Bromo ethers (% Yield)	1,4-Dienes	(% Yield) $Zn/EtOH^a$) Na/Et_2O^a)
				(79)
				(40)
				(38) ^{c)}
				(65)
				(74)
				(72)
				(62)
				(64)
				(75)
				(54)
				(67)

^{a)} Yield with respect to vinyl ethers 1 and allylsilanes 3. — ^{b)} Yield of crude material. — ^{c)} Ref.¹⁰⁾.

While the elimination with zinc in alcohols gave satisfactory results in several cases (Table 1), there were systems, where generally high amounts of side products were observed under these conditions. Treatment of **4o** with zinc dust (Riedel-de Haen) in ethanol, for example, gave **5o**, **6**, and **7** (Scheme 4) in a 58:33:9 ratio (GC), and zinc powder (Merck) in ethanol yielded these compounds in a 72:22:6 ratio. Heating of **4o** with a Zn/Cu couple (Merck) even led to the predominant formation of **6** (**5o**:**6**:**7** = 38:46:16). While no reaction was observed when **4o** was treated with magnesium turnings in diethyl ether, the analogous treatment with sodium wire led to the selective formation of compound **5o**⁶.

Scheme 4




As this method gave good yields of 1,4-dienes in all cases examined, the elimination with Na in Et₂O is the method of choice for further applications.

Conclusion

Scheme 3 provides a straightforward synthesis for 1,4-dienes with variable substitution pattern. This method may be considered as a variant of the Boord olefin synthesis (Scheme 5)^{5,7}, which has also been employed for the synthesis of 1,4-dienes by using allylic Grignard reagents⁸. As unsymmetrical allylmagnesium halides often do not react regioselectively, the advantage of the method presented in this article is not only the higher overall yield but also the greater regioselectivity of the C–C bond-forming reaction.

Scheme 5



We thank R. Wendt for orientating experiments, the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for financial support. We also would like to express our sincere thanks to Profs. Tochtermann and Hoppe for accepting this work as a part of a Diplomarbeit at the Universität Kiel.

Experimental

¹H NMR: EM 390 (Varian). — ¹³C NMR: XL 200 (Varian). — MS: 70–250 E (VG-Instruments).

The enol ethers were commercially available (**1a**–**c**), or were synthesized from the corresponding acetals by H₃PO₄-catalyzed alcohol elimination using literature procedures (**1d**–**f**)⁹. The allylsilanes **3b** and **3e** were obtained by magnesium-promoted coupling of the corresponding allyl chlorides with chlorotrimethylsilane^{10,11}, **3c** was prepared by photochemically induced hydrosilation of 2,3-dimethyl-1,3-butadiene in the presence of Cr(CO)₆¹² and **3d** from (trimethylsilyl)methylmagnesium chloride and 1-(trimethylsiloxy)-1-cyclohexene in the presence of catalytic amounts of nickel acetyl-acetonate¹³.

*Typical Procedure for the Synthesis of the β-Bromo Ethers **4**:* Bromine (1.60 g, 10.0 mmol) was dissolved in CH₂Cl₂ (20 ml) in a 100-ml flask and the solution cooled at –78 °C. A solution of the enol ether **1e** (1.39 g, 11.0 mmol) in CH₂Cl₂ (20 ml) was added dropwise with stirring, leading to complete decolorization of the solution. Allylsilane **3a** (1.26 g, 11.0 mmol) and ZnCl₂–Et₂O [4.8 mmol; 2.5 ml of a solution obtained by dissolving ZnCl₂ (50 g) in diethyl ether (60 ml) and dichloromethane (120 ml)]³ were added, and the mixture was kept at –78 °C for 2 h. The solution was washed with two 35-ml portions of 25% aqueous NH₄Cl and dried with MgSO₄. After evaporation of the solvent, spectroscopically pure *1-allyl-2-bromo-1-ethoxycyclohexane* (**4e**) was obtained in 84% yield (2.29 g). — ¹H NMR (90 MHz, CDCl₃): δ = 1.05–2.78 (m, 13 H), 3.20–3.70 (m, 2 H, OCH₂CH₃), 4.10 (m, 1 H, CHBr), 5.0–5.4 (m, 2 H, =CH₂), 5.73 (m, 1 H, CH=).

Note: This procedure is recommended for all compounds **4**, though in the early phase of this project different amounts of the reactants have been used (see below).

5-Bromo-4-ethoxy-1-pentene (**4a**): was obtained in 87% yield (1.17 g) from Br₂ (1.12 g, 7.01 mmol), **1a** (500 mg, 6.93 mmol), ZnCl₂–Et₂O (1.75 ml, 3.36 mmol), and **3a** (1.20 g, 10.5 mmol) after a reaction time of 15 h; b.p. 40–45 °C (bath)/2–5 mbar [ref.¹⁴] 81.8–84.0 °C (33 mbar). — ¹H NMR (CCl₄): δ = 1.20 (t, *J* = 6.0 Hz, 3 H, OCH₂CH₃), 2.37 (m, 2 H, 3-H), 3.23–3.75 (m, 5 H, 4-H, 5-H, OCH₂CH₃), 4.97–5.27 (m, 2 H, 1-H), 5.80 (m, 1 H, 2-H).

5-Bromo-4-methoxy-4-methyl-1-pentene (**4b**): Vinyl ether **1b** (720 mg, 10.0 mmol), Br₂ (1.60 g, 10.0 mmol), ZnCl₂–Et₂O (2.5 ml, 4.84 mmol), and **3a** (1.71 g, 15.0 mmol) gave **4b** (1.40 g, 73%) with b.p. 25–30 °C (bath)/0.4 mbar (reaction time 15 h). — ¹H NMR (CCl₄): δ = 1.22 (s, 3 H, CH₃), 2.35 (br. d, *J* = 6 Hz, 2 H, 3-H), 3.21 (s, 3 H, OCH₃), 3.27 (s, 2 H, 5-H), 4.93–5.22 (m, 2 H, 1-H), 5.47–5.97 (m, 1 H, 2-H).

5-Bromo-4-ethoxy-1-hexene (**4c**): The dibromo ether prepared from Br₂ (1.12 g, 7.01 mmol) and (*E,Z*)-**1c** (600 mg, 7.00 mmol, *E*-**1c**:*Z*-**1c** = 30:70) reacted within 15 h with **3a** (1.20 g, 10.5 mmol) in the presence of ZnCl₂–Et₂O (1.8 ml, 3.46 mmol) to give **4c** (1.33 g, 92%) with b.p. 40–45 °C (bath)/4 mbar. — ¹H NMR (CDCl₃): δ = 1.21 (t, *J* = 7.0 Hz, 3 H, OCH₂CH₃), 1.70 (d, *J* = 6.8 Hz, 3 H, 6-H), 2.42 (br. t, *J* = 6.2 Hz, 2 H, 3-H), 3.41 (q, *J* = 5.4 Hz, 1 H, 4-H), 3.63 (m, 2 H, OCH₂CH₃), 4.14 (qd, *J* = 6.8/5.4 Hz, 1 H, 5-H), 5.06–5.20 (m, 2 H, 1-H), 5.85 (m, 1 H, 2-H). — ¹³C NMR (CDCl₃): δ = 15.48 (q, OCH₂CH₃), 21.35 (q, C-6), 36.49 (t, C-3), 51.31 (d, C-5), 66.52 (t, OCH₂CH₃), 83.02 (d, C-4), 117.58 (t, C-1), 134.13 (d, C-2). — The ¹³C-NMR spectrum indicates a 95:5 ratio of diastereomers. — MS (70 eV): *m/z* (%) = 165, 167 (94, 92) [$M^+ - C_3H_5$], 137, 139 (84, 82) [$M^+ - C_3H_4 - C_2H_4$], 57 (100) [$M^+ - C_3H_5 - C_2H_4 - HBr$].

5-Bromo-4-methoxy-4-phenyl-1-pentene (**4d**): 5.00 g of a 83:17 mixture of **1d** and acetophenone dimethyl acetal (total 37.0 mmol), bromine (5.92 g, 37.0 mmol), ZnCl₂–Et₂O (9.3 ml, 17.9 mmol), and **3a** (6.30 g, 55.1 mmol) gave 8.67 g (92%) of spectroscopically pure

4d. — ^1H NMR (CDCl_3): $\delta = 2.74$ (br. d, $J = 6.5$ Hz, 2H, 3-H), 3.10 (s, 3H, OCH_3), 3.69 (s, 2H, 5-H), 4.93–5.28 (m, 2H, 1-H), 5.29–5.82 (m, 1H, 2-H), 7.28 (m_c, 5H, Ph).

4e: See typical procedure.

5-Bromo-4-ethoxy-5-methyl-1-hexene (4f) was prepared in 81% yield (3.92 g) from **1f** (2.20 g, 22.0 mmol), bromine (3.20 g, 20.0 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (5.0 ml, 9.60 mmol), and **3a** (2.51 g, 22.0 mmol); b.p. 62–73°C/2–5 mbar. — ^1H NMR (CDCl_3): $\delta = 1.17$ (t, $J = 6.0$ Hz, 3H, OCH_2CH_3), 1.72, 1.77 (2 s, 6H, 5- CH_3 , 6-H), 1.03–2.83 (m, 2H, 3-H), 3.30 (m_c, 1H, 4-H), 3.63 (q, $J = 6.0$ Hz, 2H, OCH_2CH_3), 4.87–5.27 (m, 2H, 1-H), 5.60–6.20 (m, 1H, 2-H).

5-Bromo-4-methoxy-2,4-dimethyl-1-pentene (4g): Vinyl ether **1b** (1.52 g, 21.1 mmol), bromine (3.07 g, 19.2 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (4.8 ml, 9.22 mmol), and **3b** (2.70 g, 21.0 mmol) gave 3.77 g (86%) of **4g**; b.p. 50–70°C (bath)/1 mbar (ref.¹⁰) 30–40°C (bath)/0.02 mbar. — ^1H NMR (CDCl_3): $\delta = 1.27$ (s, 3H, 4- CH_3), 1.80 (br. s, 3H, 2- CH_3), 2.40 (br. s, 2H, 3-H), 3.23 (s, 3H, OCH_3), 3.40 (s, 2H, 5-H), 4.80, 4.90 (2 m_c, 2H, 1-H).

5-Bromo-4-methoxy-2-methyl-4-phenyl-1-pentene (4h): 1.47 g of a 83:17 mixture of **1d** and acetophenone dimethyl acetal (total 11.0 mmol), bromine (1.60 g, 10.0 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (2.5 ml, 4.84 mmol), and **3b** (1.41 g, 11.0 mmol) gave **4h** (2.86 g, 97%) as an orange, spectroscopically pure liquid. — ^1H NMR (CCl_4): $\delta = 1.36$ (s, 3H, 2- CH_3), 2.50, 2.75 (AB system, $J = 13.5$ Hz, 2H, 3-H), 3.08 (s, 3H, OCH_3), 3.72, 3.90 (AB system, $J = 9.0$ Hz, 2H, BrCH_2), 4.73 (m_c, 2H, $\text{C}=\text{CH}_2$), 7.28 (m_c, 5H, Ph).

2-Bromo-1-ethoxy-1-(2-methylallyl)cyclohexane (4i): Compound **1e** (2.77 g, 22.0 mmol), bromine (3.20 g, 20.0 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (5.0 ml, 9.6 mmol), and **3b** (2.82 g, 22.0 mmol) gave 3.93 g (68%) of crude **4i**. — ^1H NMR (CDCl_3): $\delta = 1.05$ –2.68 (m, 16H, 3-, 4-, 5-, 6-H, OCH_2CH_3 , CH_3 , = CCH_2), 3.53 (m_c, 2H, OCH_2CH_3), 4.26 (m_c, 1H, 2-H), 4.92 (br. s, 2H, $\text{C}=\text{CH}_2$).

5-Bromo-4-ethoxy-2,5-dimethyl-1-hexene (4j) was obtained in 92% yield (1.19 g) from **1f** (550 mg, 5.49 mmol), bromine (800 mg, 5.00 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (1.3 ml, 2.52 mmol), and **3b** (705 mg, 5.50 mmol). — ^1H NMR (CDCl_3): $\delta = 1.18$ (t, $J = 6$ Hz, 3H, OCH_2CH_3), 1.71, 1.78, 1.85 [3 s, 9H, $\text{BrC}(\text{CH}_3)_2$, 2- CH_3], 2.02–2.70 (m, 2H, 3-H), 3.31–3.84 (m, 3H, OCH_2CH_3 , 4-H), 4.83 (br. s, 2H, $\text{C}=\text{CH}_2$).

5-Bromo-4-ethoxy-2,3,3-trimethyl-1-pentene (4k): Vinyl ether **1a** (1.59 g, 22.0 mmol), bromine (3.20 g, 20.0 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (5.0 ml, 9.60 mmol), and **3c** (4.37 g, 22.0 mmol) gave 8.52 g of a mixture of **4k** and of hexaethyldisiloxane which was not separated but subjected to treatment with sodium (see below). The spectral data of **4k** were taken from this mixture. — ^1H NMR (CDCl_3): $\delta = 1.03$, 1.10 (2 s, 6H, 3- CH_3), 1.20 (t, $J = 6.0$ Hz, 3H, OCH_2CH_3), 1.77 (s, 3H, 2- CH_3), 3.10–4.06 (m, 5H, OCH_2CH_3 , 4-, 5-H), 4.82 (br. s, 2H, $\text{C}=\text{CH}_2$).

1-(2-Bromo-1-ethoxypropyl)-2-methylenecyclohexane (4l): Compound **1c** (*E,Z* mixture, 1.57 g, 18.2 mmol), bromine (2.64 g, 16.5 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (3.6 ml, 6.90 mmol), and **3d** (3.10 g, 18.4 mmol) gave 4.70 g (99%) of **4l**. — ^1H NMR (CDCl_3): $\delta = 1.08$ –2.73 (m, 14H, 2 CH_3 and cyclohexane H), 3.33–4.56 (m, 5H, OCH_2CH_3 , BrCH , OCH , CH), 4.60–4.93 (m, 2H, $\text{C}=\text{CH}_2$). On the basis of ^1H -NMR spectral data it could not be decided whether this was a mixture of two diastereoisomers or a single isomer.

2-Bromo-1-(2-cyclopenten-1-yl)-1-methoxy-1-phenylethane (4m) was obtained in 92% yield (2.83 g) from a 83:17 mixture of **1d** and acetophenone dimethyl acetal (1.47 g, total 11.0 mmol), bromine (1.60 g, 10.0 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (2.5 ml, 4.80 mmol), and **3e**

(1.54 g, 11.0 mmol). — ^1H NMR (CCl_4): $\delta = 1.66$ –1.93 (m, 4H, CH_2CH_2), 3.29 (s, 3H, OCH_3), 3.38–4.06 (m, 3H, $\text{CH}-\text{C}\text{H}=$, BrCH_2), 5.68 (m_c, 2H, vinyl H), 7.21 (m_c, 5H, Ph).

2-Bromo-1-(2-cyclopenten-1-yl)-2-ethoxycyclohexane (4n): Enol ether **1e** (2.77 g, 22.0 mmol), bromine (3.20 g, 20.0 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (5.0 ml, 9.60 mmol), and **3e** (3.08 g, 22.0 mmol) gave 5.29 g (88%) of **4n**. — ^1H NMR (CDCl_3): $\delta = 0.99$ –2.57 (m, 15H, 6 CH_2 , OCH_2CH_3), 3.28–4.04 (m, 3H, OCH_2CH_3 , $\text{CH}-\text{C}\text{H}=$), 4.18 (m_c, 1H, BrCH), 5.78 (m_c, 2H, vinyl H).

2-Bromo-1-(2-cyclopenten-1-yl)-1-ethoxy-2-methylpropane (4o): was synthesized in 88% yield (1.20 g) from enol ether **1f** (550 mg, 5.49 mmol), bromine (800 mg, 5.00 mmol), $\text{ZnCl}_2-\text{Et}_2\text{O}$ (1.3 ml, 2.52 mmol), and **3e** (770 mg, 5.50 mmol). — ^1H NMR (CDCl_3): $\delta = 0.95$ –1.32 (m, 4H, 5'-H_A, OCH_2CH_3), 1.52–2.50 [m, 9H, 4'-H, 5'-H_B, $\text{C}(\text{CH}_3)_2$], 3.11–3.75 (m, 4H, 1'-H, $\text{CHOCH}_2\text{CH}_3$), 5.69 (m_c, 2H, vinyl H).

Synthesis of the 1,4-Dienes

Elimination with Zinc/Ethanol (Procedure A): Crude **4h**, obtained from 11.0 mmol of **1d** and **3b**, was dissolved in absolute ethanol (28 ml). After addition of zinc powder (6.50 g, 100 mmol), the mixture was heated at reflux for 15 h. Pentane (40 ml) was added, the solvent mixture was decanted, and the remaining zinc powder was washed with four 20-ml portions of pentane. The combined solvents were successively washed with 10% aqueous NH_4Cl solution (40 ml) and with four 20-ml portions of water. The solvent was evaporated, and the residue was distilled to give 1.30 g (75%) of **5h** with b.p. 100°C (bath)/10–20 mbar (ref.¹⁵) 90–92°C/16 mbar. For spectral data see below.

Elimination with Sodium in Diethyl Ether (Procedure B): Dry diethyl ether (40–50 ml) and sodium wire (3.79 g, 165 mmol) were placed into a 250-ml two-necked flask under nitrogen. Crude **4i**, prepared from 22.0 mmol of **1e** and **3b**, was added, and the mixture was heated under reflux for 4 h. The solvent was decanted, and the solid residue was washed with pentane (20 ml, 10 ml, and 5 ml). The solutions were combined, washed with two 40-ml portions of water (caution: small pieces of Na may be in the solution!), and dried with Na_2SO_4 . After careful evaporation of the solvents, the residue was distilled to give 2.20 g (73%) of **5i** with b.p. 68°C (bath)/5–20 mbar. For spectral data see below.

2-Phenyl-1,4-pentadiene (5d): Compound **4d**, obtained from 22.0 mmol **1d** and **3a**, gave 2.50 g (79%) of **5d** according to procedure A. b.p. 80°C (bath)/22–30 mbar (ref.¹⁵) 82–84°C/16 mbar. — ^1H NMR (CDCl_3): $\delta = 3.22$ (br. d, $J = 6.7$ Hz, 2H, 3-H), 4.93–5.21 (m, 3H, 1-H_A, 5-H), 5.38 (s, 1H, 1-H_B), 5.88 (m_c, 1H, 4-H), 7.21–7.45 (m, 5H, Ph). — ^{13}C NMR (CDCl_3): $\delta = 39.49$ (t, C-3), 113.10 (t, C-1), 116.45 (t, C-5), 125.95 (d, C_o), 127.41 (d, C_p), 128.22 (d, C_m), 136.14 (d, C-4), 140.88 (s, C_i), 146.25 (s, C-2). — MS (70 eV): m/z (%) = 144 (100) [M^+], 129 (73) [$\text{M}^+ - \text{CH}_3$], 115 (10) [$\text{M}^+ - \text{C}_2\text{H}_5$], 103 (57) [$\text{M}^+ - \text{C}_3\text{H}_5$], 77 (16) [Ph^+].

$\text{C}_{11}\text{H}_{12}$ Calcd. 144.0939 Found 144.0932 (MS)

1-Allyl-1-cyclohexene (5e): Procedure A was used to prepare 272 mg (40% with respect to **1e**) from crude **4e**. b.p. 55°C (bath)/5–20 mbar (ref.¹⁶) 154–158°C. — ^1H NMR (CDCl_3): $\delta = 1.54$ –1.64 (m, 4H, 4-, 5-H), 1.89–2.01 (m, 4H, 3-, 6-H), 2.67 (d, $J = 6.9$ Hz, 2H, allyl CH₂), 4.97–5.08 (m, 2H, $\text{CH}=\text{CH}_2$), 5.44 (m_c, 1H, 2-H), 5.70–5.91 (m, 1H, $\text{CH}=\text{CH}_2$). — ^{13}C NMR (CDCl_3): $\delta = 22.50$, 22.97, 25.31, 28.34 (4 t, C-3, -4, -5, -6), 42.57 (t, CH_2CH), 115.44 (t, $\text{CH}=\text{CH}_2$), 121.87 (d, C-2), 136.30 (s, C-1), 136.97 (d, $\text{CH}=\text{CH}_2$). — MS (70 eV): m/z (%) = 122 (28) [M^+], 107 (14) [$\text{M}^+ - \text{CH}_3$], 93 (16) [$\text{M}^+ - \text{C}_2\text{H}_5$], 81 (100) [$\text{M}^+ - \text{C}_3\text{H}_5$].

C_9H_{14} Calcd. 122.1096 Found 122.1091 (MS)

2-Methyl-4-phenyl-1,4-pentadiene (5h) was synthesized according to procedure A (yield 75%, details see above) and procedure B (yield 2.26 g = 65% with respect to 1d). — ^1H NMR (CDCl_3): δ = 1.72 (s, 3 H, CH_3), 3.21 (s, 2 H, 3-H), 4.77, 4.81 (2 m, 2 H, 1-H), 5.11 (d, J = 1.4 Hz, 5-H), 5.43 (d, J = 1.5 Hz, 1 H, 5-H), 7.27–7.46 (m, 5 H, Ph). — ^{13}C NMR (CDCl_3): δ = 22.28 (q, CH_3), 44.00 (t, C-3), 112.58, 114.27 (2 t, C-1, -5), 126.07 (d, C_o), 127.34 (d, C_p), 128.15 (d, C_m), 140.96, 143.39, 145.59 (3 s, C_i , C-2, -4). — MS (70 eV): m/z (%) = 158 (14) [M^+], 143 (100) [$\text{M}^+ - \text{CH}_3$], 128 (30) [$\text{M}^+ - 2\text{CH}_3$], 103 (33) [$\text{M}^+ - \text{C}_4\text{H}_7$], 77 (22) [Ph^+].

$\text{C}_{12}\text{H}_{14}$ Calcd. 158.1096 Found 158.1095 (MS)

1-(2-Methylallyl)-1-cyclohexene (5i) was prepared following procedure A (yield 459 mg = 61% with respect to 1e) and B (yield 74%, details see above). — ^1H NMR (CDCl_3): δ = 1.50–1.64 (m, 7 H, CH_3 , 4-, 5-H), 1.81–2.10 (m, 4 H, 3-, 6-H), 2.63 (s, 2 H, allyl CH_2), 4.70, 4.75 (2 m, 2 H, $\text{C}=\text{CH}_2$), 5.46 (m, 1 H, 2-H). — ^{13}C NMR (CDCl_3): δ = 21.84 (q, CH_3), 22.49, 23.00, 25.35, 27.70 (4 t, C-3, -4, -5, -6), 47.14 (t, CH_2), 111.28 (t, $\text{C}=\text{CH}_2$), 123.00 (d, C-2), 135.51 (s, C-1), 144.28 (s, $\text{C}=\text{CH}_2$). — MS (70 eV): m/z (%) = 136 (30) [M^+], 121 (64) [$\text{M}^+ - \text{CH}_3$], 107 (21) [$\text{M}^+ - \text{C}_2\text{H}_5$], 93 (49) [$\text{M}^+ - \text{C}_3\text{H}_7$], 81 (100) [$\text{M}^+ - \text{C}_4\text{H}_7$].

$\text{C}_{10}\text{H}_{16}$ Calcd. 136.1252 Found 136.1257 (MS)

2,5-Dimethyl-1,4-hexadiene (5j) has been synthesized according to procedure A (yield 122 mg = 20% with respect to 1f) and procedure B (1.52 g = 72% with respect to 1f). B.p. 89 °C (bath)/150–190 mbar (ref.¹⁷) does not report the b.p.). — ^1H NMR (CDCl_3): δ = 1.63, 1.72 [2 br. s, 9 H, 2- CH_3 , $\text{C}(\text{CH}_3)_2$], 2.68 (br. d, J = 7.1 Hz, 2 H, 3-H), 4.69 (m, 2 H, 1-H), 5.19 (m, 1 H, 4-H). — ^{13}C NMR (CDCl_3): δ = 17.62, 22.55, 25.77 (3 q, 3 CH_3), 36.59 (t, C-3), 109.71 (t, C-1), 121.94 (d, C-4), 132.92 (s, C-5), 145.46 (s, C-2). — MS (70 eV): m/z (%) = 110 (52) [M^+], 95 (100) [$\text{M}^+ - \text{CH}_3$], 67 (47) [$\text{M}^+ - \text{C}_3\text{H}_7$].

C_8H_{14} Calcd. 110.1096 Found 110.1098 (MS)

2,3,3-Trimethyl-1,4-pentadiene (5k): The mixture of 4k and hexaethyldisiloxane described above was treated with Na (procedure B) to give 1.50 g (62% with respect to 1a) of 5k. B.p.¹⁸ 102 °C. — ^1H NMR (CDCl_3): δ = 1.15 [s, 6 H, $\text{C}(\text{CH}_3)_2$], 1.70 (m, 3 H, 2- CH_3), 4.74–4.82 (m, 2 H, 5-H), 4.93–5.03 (m, 2 H, 1-H), 5.82 (m, 1 H, 4-H). — ^{13}C NMR (CDCl_3): δ = 19.74 (q, 2- CH_3), 26.01 [q, $\text{C}(\text{CH}_3)_2$], 42.18 (s, C-3), 109.30, 110.87 (2 t, C-1, -5), 147.14 (d, C-4), 151.61 (s, C-2). — MS (70 eV): m/z (%) = 110 (20) [M^+], 95 (100) [$\text{M}^+ - \text{CH}_3$], 67 (44) [$\text{M}^+ - \text{C}_3\text{H}_7$], 41 (47) [C_3H_5^+].

C_8H_{14} Calcd. 110.1096 Found 110.1098 (MS)

1-Methylene-2-((E,Z)-1-propenyl)cyclohexane (5l): Treatment of crude 4l (4.70 g) with Na according to procedure B gave 1.60 g (64% with respect to 1c) of 5l as a mixture of (E)- and (Z)-5l. B.p. 90 °C (bath)/16–90 mbar. — ^1H NMR (CDCl_3): δ = 1.20–1.90 (m, 9 H, 3-, 4-, 5-H, CH_3), 2.02–2.22, 2.32–2.50 (2 m, 2 H, 6-H), 2.66–2.80, 2.98–3.12 (2 m, 1 H, 2-H), 4.64–4.80 (m, 2 H, $\text{C}=\text{CH}_2$), 5.40–5.72 (m, 2 H, $\text{CH}=\text{CH}$). — ^{13}C NMR (CDCl_3): δ = 12.96 (q, CH_3 , Z), 18.08 (q, CH_3 , E), 25.07, 25.42, 28.16, 28.27, 34.45, 34.79, 35.29, 35.54 (8 t, C-3, -4, -5, -6, E,Z), 41.14, 46.59 (2 d, C-2, E,Z), 106.10, 106.34 (2 t, $\text{C}=\text{CH}_2$, E,Z), 123.95, 124.86 (2 d, $\text{CH}_3\text{CH}=\text{CH}$, E,Z), 132.77, 133.69 (2 d, $\text{CH}_3\text{CH}=\text{CH}$, E,Z), 151.00, 152.54 (2 s, C-1, E,Z). — MS (70 eV): m/z (%) = 136 (64) [M^+], 121 (60) [$\text{M}^+ - \text{CH}_3$], 107 (50) [$\text{M}^+ - \text{C}_2\text{H}_5$], 93 (76) [$\text{M}^+ - \text{C}_3\text{H}_7$], 79 (100) [$\text{M}^+ - \text{C}_2\text{H}_5 - \text{C}_2\text{H}_4$].

$\text{C}_{10}\text{H}_{16}$ Calcd. 136.1252 Found 136.1256 (MS)

3-(1-Phenylvinyl)-1-cyclopentene (5m): 1.41 g of compound 5m was obtained from crude 4m and Zn (procedure A). Yield: 75% (with respect to 1d). B.p. 87–110 °C (bath)/2–4 mbar. — ^1H NMR

(CDCl_3): δ = 1.46–1.68 (m, 1 H, 4-H_A), 2.10–2.48 (m, 3 H, 4-H_B, 5-H), 3.85 (m, 1 H, 3-H), 5.03, 5.25 (2 m, 2 H, $\text{C}=\text{CH}_2$), 5.70–5.92 (m, 2 H, 1-, 2-H), 7.25–7.47 (m, 5 H, Ph). — ^{13}C NMR (CDCl_3): δ = 31.14, 32.00 (2 t, C-4, -5), 49.77 (d, C-3), 110.81 (t, $\text{C}=\text{CH}_2$), 126.38 (d, C_o), 127.19 (d, C_p), 128.17 (d, C_m), 131.90, 133.29 (2 d, C-1, -2), 141.95 (s, C_i), 152.24 (s, $\text{C}=\text{CH}_2$). — MS (70 eV): m/z (%) = 170 (100) [M^+], 155 (57) [$\text{M}^+ - \text{CH}_3$], 103 (75) [$\text{M}^+ - \text{cyclopentenyl}$], 77 (52) [Ph^+], 67 (40) [cyclopentenyl^+].

$\text{C}_{13}\text{H}_{14}$ Calcd. 170.1096 Found 170.1086 (MS)

1-(2-Cyclopenten-1-yl)-1-cyclohexene (5n): Procedure B yielded 1.77 g (54% with respect to 1e) of 5n from crude 4n. B.p. 100 °C/16–20 mbar (ref.¹⁹) does not report the b.p.). — ^1H NMR (CDCl_3): δ = 1.50–1.78 (m, 5 H), 1.89–2.22 (m, 5 H), 2.37 (m, 2 H), 3.27 (m, 1 H, 3-H), 5.43 (m, 1 H), 5.64–5.90 (m, 2 H, 1-, 2-H). — ^{13}C NMR (CDCl_3): δ = 22.73 (t), 23.09 (t), 25.24 (t), 26.40 (t), 29.30 (t), 32.38 (t), 52.93 (d), 119.73 (d), 131.23 (d), 133.65 (d), 140.97 (s). — MS (70 eV): m/z (%) = 148 (86) [M^+], 133 (15) [$\text{M}^+ - \text{CH}_3$], 119 (36) [$\text{M}^+ - \text{C}_2\text{H}_5$], 105 (33) [$\text{M}^+ - \text{C}_3\text{H}_7$], 91 (65) [$\text{M}^+ - \text{C}_4\text{H}_9$], 80 (100), 77 (25), 67 (53) [cyclopentenyl^+].

$\text{C}_{11}\text{H}_{16}$ Calcd. 148.1252 Found 148.1258 (MS)

3-(2-Methyl-1-propenyl)-1-cyclopentene (5o) was prepared according to procedure A (yield 200 mg = 30% with respect to 1f) and procedure B (1.80 g = 67% with respect to 1f). B.p. 73 °C (bath)/30–45 mbar (ref.¹⁹) does not report the b.p.). — ^1H NMR (CDCl_3): δ = 1.36–1.55 (m, 1 H, 4-H_A), 1.67, 1.69 [2 br. s, 6 H, $\text{C}(\text{CH}_3)_2$], 2.02–2.40 (m, 3 H, 4-H_B, 5-H), 3.49 (m, 1 H, 3-H), 5.00 (br. d, J = 9.1 Hz, 1 H, $\text{C}=\text{CH}$), 5.50–5.80 (m, 2 H, 1-, 2-H). — ^{13}C NMR (CDCl_3): δ = 17.93, 25.73 [2 q, $\text{C}(\text{CH}_3)_2$], 31.19, 32.22 (2 t, C-4, -5), 44.25 (d, C-3), 129.13, 130.45 (2 d, C-1, -2), 130.60 [s, $(\text{CH}_3)_2\text{C}=\text{CH}$], 134.72 [d, $(\text{CH}_3)\text{C}=\text{CH}$]. — MS (70 eV): m/z (%) = 122 (50) [M^+], 107 (100) [$\text{M}^+ - \text{CH}_3$], 79 (64) [$\text{M}^+ - \text{C}_3\text{H}_7$], 67 (17) [cyclopentenyl^+].

C_9H_{14} Calcd. 122.1096 Found 122.1098 (MS)

The reaction with Zn in ethanol gave compounds 6 and 7 in addition to 5o. These two ethers (6, 7) were adsorbed, when the product mixture was passed through silica with pentane as eluent, and eluted with CH_2Cl_2 . The 2:1 mixture of 6 and 7 was identified by ^{13}C -NMR spectroscopy. — **3-(2-Cyclopenten-1-yl)-3-ethoxy-2-methyl-1-propene (6)**: ^{13}C NMR (CDCl_3): δ = 15.29 (q, OCH_2CH_3), 16.76 (q, CH_3), 27.30, 31.96 (2 t, C-3, -4), 48.59 (d, C-3), 63.37 (t, OCH_2CH_3), 88.18 (d, CH), 114.11 (t, $\text{C}=\text{CH}_2$), 131.51, 132.08 (2 d, $\text{CH}=\text{CH}$), 144.45 (s, $\text{C}=\text{CH}_2$). — **1-(2-Cyclopenten-1-yl)-1-ethoxy-2-methylpropane (7)**: ^{13}C NMR (CDCl_3): δ = 15.83 (q, OCH_2CH_3), 18.11, 20.21 [2 q, $\text{CH}(\text{CH}_3)_2$], 25.15, 32.10 (2 t, C-4, -5), 31.87 [d, $\text{CH}(\text{CH}_3)_2$], 48.90 (d, C-3), 68.27 (t, OCH_2CH_3), 87.84 (d, CH), 131.62, 132.87 (2 d, C-1, -2).

CAS Registry Numbers

1a: 109-92-2 / **1b:** 116-11-0 / **1c (E):** 4696-26-8 / **1c (Z):** 4696-25-7 / **1d:** 4747-13-1 / **1e:** 1122-84-5 / **1f:** 927-61-7 / **3a:** 762-72-1 / **3b:** 18292-38-1 / **3c:** 64545-12-6 / **3d:** 58541-14-3 / **3e:** 14579-08-9 / **4a:** 22089-55-0 / **4b:** 135312-62-8 / **4c (R*, R*):** 135312-63-9 / **4c (R*, S*):** 135312-79-7 / **4d:** 135312-64-0 / **4e:** 135312-65-1 / **4f:** 135312-66-2 / **4g:** 135312-67-3 / **4h:** 135312-68-4 / **4i:** 135312-69-5 / **4j:** 135312-70-8 / **4k:** 135312-71-9 / **4l:** 135312-72-0 / **4m:** 135312-73-1 / **4n:** 135312-74-2 / **4o:** 135312-75-3 / **5a:** 591-93-5 / **5b:** 763-30-4 / **5c (E):** 7319-00-8 / **5c (Z):** 7318-67-4 / **5d:** 35342-69-9 / **5e:** 13511-13-2 / **5f:** 763-88-2 / **5g:** 4161-65-3 / **5h:** 52713-62-9 / **5i:** 135312-76-4 / **5j:** 927-97-9 / **5k:** 756-02-5 / **5l (E):** 135312-77-5 / **5l (Z):** 135312-80-0 / **5m:** 135312-78-6 / **5n:** 119946-62-2 / **5o:** 16839-59-1

* Dedicated to Professor Karl Heinz Büchel on the occasion of his 60th birthday.

¹⁾ See references 2–6 in ref.²⁾.

- ²⁾ B. H. Lipshutz, T. R. Elworthy, *J. Org. Chem.* **55** (1990) 1695.
³⁾ H. Mayr, W. Striepe, *J. Org. Chem.* **50** (1985) 2995.
⁴⁾ For a comprehensive literature report on Lewis acid-catalyzed reactions of α -halo ethers with allylsilanes see: I. Fleming, J. Dunoguès, R. Smithers, *Org. React.* **57** (1989) 57.
⁵⁾ H. B. Dykstra, J. F. Lewis, C. E. Board, *J. Am. Chem. Soc.* **52** (1930) 3396.
⁶⁾ For related eliminations with Na in ether solvents see: ^{6a)} H. O. House, R. S. Ro, *J. Am. Chem. Soc.* **80** (1958) 182. — ^{6b)} R. Paul, O. Riobé, M. Maumy, *Org. Synth.*, Coll. Vol. VI (1988) 675.
⁷⁾ M. Schlosser in *Methoden der organischen Chemie* (Houben-Weyl-Müller) Bd. 5/1b, 4. Aufl., p. 212, Georg Thieme Verlag Stuttgart 1972.
⁸⁾ B. H. Shoemaker, C. E. Board, *J. Am. Chem. Soc.* **53** (1931) 1505.
⁹⁾ H. Meerwein in *Methoden der organischen Chemie* (Houben-Weyl-Müller), Bd. 6/3, 4. Aufl., p. 95, Georg Thieme Verlag Stuttgart 1965.
¹⁰⁾ B. Irrgang, H. Mayr, *Tetrahedron* **47** (1991) 219.
¹¹⁾ G. Hagen, H. Mayr, *J. Am. Chem. Soc.* **113** (1991) 4954.
¹²⁾ M. S. Wrighton, M. A. Schroeder, *J. Am. Chem. Soc.* **96** (1974) 6235.
¹³⁾ T. Hayashi, Y. Katsuro, M. Kumada, *Tetrahedron Lett.* **21** (1980) 3915.
¹⁴⁾ D. C. Rowlands, K. W. Greenlee, J. M. Derfer, C. E. Board, *J. Org. Chem.* **17** (1952) 807.
¹⁵⁾ Y. Frangin, M. Gaudemar, *Bull. Soc. Chim. Fr.* **1976**, 41.
¹⁶⁾ J. Barluenga, M. Yus, J. M. Concellon, P. Bernad, *J. Chem. Res. (M)* **1980**, 677.
¹⁷⁾ Y. Tajima, E. Kunioka, *J. Chem. Soc., Chem. Commun.* **1968**, 603.
¹⁸⁾ N. F. Cywinski, *J. Org. Chem.* **30** (1965) 361.
¹⁹⁾ R. C. Larock, W. H. Gong, *J. Org. Chem.* **54** (1989) 2047.

[220/91]