


## BROMINATION OF OCTAMETHYLCYCLOPENTENE - THE IRREGULAR REACTIVITY OF A STERICALLY HINDERED CYCLOALKENE

Herbert Mayr\*, Elfriede Will<sup>1</sup>, Ulrich W. Heigl and Christian SchadeInstitut für Chemie der Medizinischen Universität zu Lübeck  
Ratzeburger Allee 160, D-2400 Lübeck 1, Federal Republic of Germany


(Received in Germany 28 December 1985)

**Abstract:** The title compound **7** reacts with bromine under mild conditions (CCl<sub>4</sub>, 20°C) to give the substitution products **9** and/or **10** via an electrophilic process. The tetrabromo compound **11** is formed from **7** and bromine in refluxing CCl<sub>4</sub>, via a radical mechanism.


The addition of bromine is one of the best known reactions of unsaturated nonaromatic hydrocarbons. While most alkenes react smoothly with formation of addition products, a different behaviour is encountered with sterically hindered double bonds. Adamantylidene adamantane, for example, was reported to yield a crystalline bromonium tribromide when treated with Br<sub>2</sub>.<sup>2</sup> The bromination of di- and polyarylethylenes gave charge transfer complexes or bromonium ions, sometimes in equilibrium with the reactants and regular addition products.<sup>3</sup> Meisenheimer reported that an allylic substitution product **3** was formed from **1** and bromine.<sup>4</sup> This reaction was later demonstrated by Ziegler and Bähr to proceed via bromine addition (→ **2**), HBr elimination and allylic rearrangement.<sup>5</sup>




Lenoir reported the formation of 2,3-di-tert.butyl-1,3-butadiene (**5**) by treatment of **4** with one equivalent of bromine and the formation of **6** from **4** and two equivalents of bromine.<sup>6</sup>



In this paper we report the course of the bromination of **7**, which we observed during an attempt to convert the readily available octamethylcyclopentene **7** into **8** by a bromination elimination sequence. The latter compound was required for the mechanistic study of cycloaddition reactions of 2,3-shielded cisoid 1,3-dienes.<sup>8</sup>



**Products.** When 7 was treated with 1 equivalent of Br<sub>2</sub> in CCl<sub>4</sub> at room temperature, bromine was consumed instantaneously, and a mixture of 7, 9 and 10 (1 : 2.4 : 1) was observed in the <sup>1</sup>H NMR spectrum. An addition product could not be detected. The selective monobromination was achieved by treating 7 with 1 equivalent of pyridinium tribromide in acetic acid.



The crude material, obtained from 7 and 5 equivalents of Br<sub>2</sub> in CCl<sub>4</sub> at room temperature showed a <sup>1</sup>H-NMR spectrum with only 3 signals (6 : 12 : 4), indicating the presence of 10. When a 5 : 1 mixture of 7 and Br<sub>2</sub> was heated in CCl<sub>4</sub> at reflux, preferably in the presence of a radical starter like azoisobutyronitrile, 11 was formed in fair yield. The temperature dependent <sup>1</sup>H and <sup>13</sup>C NMR spectra of 11 indicate hindered rotation of the -CHBr<sub>2</sub> groups. At low temperature C-1, C-2 as well as C-3, C-5 and the attached groups are nonequivalent. The rotational barrier ( $\Delta G^\ddagger$ ) of the dibromomethyl groups was calculated<sup>9</sup> to be 57 kJ/mol from the coalescence of the methine protons in the 60 MHz NMR spectrum at 12°C. The conformational equilibria of 11 are presently under investigation.

Table: <sup>13</sup>C NMR Spectra of Some Octamethylcyclopentene Derivatives (CDCl<sub>3</sub>)


R	R'	C-1	1-R	C-2	2-R'	C-3	3-CH <sub>2</sub>	C-4	4-CH <sub>2</sub>	C-5	5-CH <sub>2</sub>	
7	CH <sub>3</sub>	CH <sub>3</sub>	135.0	10.0	135.0	10.0	49.8	24.2	45.7	21.7	49.8	24.2
9	CH <sub>2</sub> Br	CH <sub>3</sub>	146.7	26.7	135.7	10.6	49.7	23.9	46.3	21.3	50.1	24.8
10	CH <sub>2</sub> Br	CH <sub>2</sub> Br	144.7	23.4	144.7	23.4	50.1	24.5	46.9	21.0	50.1	24.5
11 a)	CHBr <sub>2</sub>	CHBr <sub>2</sub>	143.8	29.7	146.7	34.7	50.2	23.5	47.9	20.3	52.8	25.0

a) -20°C; the relative assignments of C-1,2, C-3,5 and the attached groups are uncertain.


**Reaction Mechanism.** The formation of 9 and 10 is best explained by an ionic pathway, comparable to the mechanism suggested by Wynberg for the halogenation of tetraalkylethylenes with N-halo-succinimides.<sup>10</sup> The alternative radical mechanism is excluded since the reaction proceeds fast at low temperature and was not inhibited by 2,4,6-tri-*tert.*butyl phenol.

The initial formation of the bromonium ion 12, which is suggested in the Scheme is in accord with the behaviour of other sterically hindered alkenes.<sup>9</sup> Successive deprotonation yields the tertiary allylic bromide 13, which rearranges to the allylic isomer 9. An analogous mechanism

9 → 15 → 10 may explain the production of 10 from 9 and bromine in CCl<sub>4</sub> at 20°C. In accord with the relative electron releasing effects of CH<sub>3</sub> and CH<sub>2</sub>Br, 9 was observed to consume Br<sub>2</sub> less rapidly than 7.



However, there must be an additional route from 7 to 10, as the dibromo compound 10 was also formed during the treatment of 7 with 0.2 equivalents of Br<sub>2</sub> in CCl<sub>4</sub> at 20°C (10 : 9 = 1 : 3). Since 9 is less reactive than 7, a pathway is required, which avoids the intermediacy of 9. As suggested in the Scheme, deprotonation of 14 may yield 8, which was found to add bromine instantaneously with formation of 10. This route is analogous to that observed by Lenoir.<sup>6</sup> When the HBr elimination from 14 was suppressed by saturating the CCl<sub>4</sub> solution of 7 with HBr, prior to adding 0.5 equivalents of bromine at 0°C, the exclusive formation of 9 was found.



Bromination of 7 in more polar solvents (CH<sub>2</sub>Cl<sub>2</sub>, CHCl<sub>3</sub>, CH<sub>3</sub>CO<sub>2</sub>H) gave several side products, probably because of rearrangements of intermediate carbenium ions. When 7 was allowed to stand with 5 equivalents of bromine in CHCl<sub>3</sub> at 20°C for 1 hour, a complex mixture containing 9, 10 and 4% of the tetrabromo compound 16 was isolated.

Carbenium ion rearrangements are unfavourable in CCl<sub>4</sub>. Even under reflux conditions the tetrabromo compound 11 was the only product isolated from 7 and excess bromine. Since this reaction was accelerated by azobisisobutyronitrile, the formation of 11 is explained by a radical mechanism.

The conversion of 9 - 11 into 2,3 shielded 1,3-dienes will be reported in a subsequent paper.

We thank Dr. H. Klein for exploratory experiments and the Fonds der Chemischen Industrie for financial support.

## EXPERIMENTAL

<sup>1</sup>H NMR spectra: JNM-C-60-HL (JEOL). <sup>13</sup>C NMR spectra: JNM-PS-100 (JEOL). TMS as internal standard. Mass spectra: CH 4 B MAT (Varian) and 311 A MAT (Varian); only the most intensive mass peaks are listed.

**Octamethylcyclopentene (7)** was prepared via [3+2] cycloaddition of the 1,1,2,3-tetramethylallyl cation to 2,3-dimethyl-2-butene.<sup>7</sup> The published procedure<sup>7</sup> was modified by reducing the alkene/alkyl chloride ratio from 3.2 to 1.3 (yield: 74%).

**1-Bromomethyl-2,3,3,4,4,5,5-heptamethyl-1-cyclopentene (9)**. Pyridinium tribromide (11.6 g, 36 mmol) was added in small portions to a stirred solution of **7** (5.0 g, 28 mmol) in 50 ml glacial acetic acid while the temperature was kept below 15°C. After 30 min, the solution was poured onto 300 mL of water. The product was extracted with ether, and the ethereal layer was washed with dilute aqueous HCl, water, and aqueous NaHCO<sub>3</sub> solution. After drying with MgSO<sub>4</sub>, the solvent was evaporated and the residue was distilled to give **9** (5.3 g, 73%) as a colourless liquid with b.p. 55 - 65°C (bath) / 0.1 mmHg. - <sup>1</sup>H NMR (CCl<sub>4</sub>): δ 0.79 (s, 6 H), 0.92 (s, 6 H), 1.03 (s, 6 H), 1.63 (s, 3 H), 3.92 (s, 2 H). - Mass spectrum (96 eV): m/e = 260, 258 (13%, 13%, M<sup>+</sup>), 245, 243 (41, 43, C<sub>12</sub>H<sub>22</sub>Br), 179 (100, C<sub>11</sub>H<sub>22</sub>), 163 (94, C<sub>11</sub>H<sub>19</sub>), 135 (C<sub>10</sub>H<sub>19</sub>), 123 (44), 121 (53), 99 (21), 97 (40). Anal. Calcd. for C<sub>11</sub>H<sub>22</sub>Br: C, 60.23; H, 8.94. Found: C, 60.57; H, 9.15.

**1,2-Di(bromomethyl)-3,3,4,4,5,5-hexamethyl-1-cyclopentene (10)**. A solution of bromine (17.6 g, 110 mmol) in 30 mL of CCl<sub>4</sub> was added dropwise to a solution of **7** (9.00 g, 50.0 mmol) in 50 mL of CCl<sub>4</sub> at 20°C. After 15 h, a nitrogen stream was passed through the solution to remove HBr and the solvent was evaporated. Distillation gave **10** (14.3 g, 84%), a colourless liquid with bp. 95 - 100°C (bath) / 0.05 mmHg. - <sup>1</sup>H NMR (CCl<sub>4</sub>): δ 0.83 (s, 6 H), 1.09 (s, 12 H), 4.05 (s, 4 H). - Mass spectrum (70 eV): m/e = 340, 338, 336 (3%, 7%, 3%, M<sup>+</sup>), 325, 323, 321 (19, 41, 20, C<sub>11</sub>H<sub>19</sub>Br<sub>2</sub>), 259, 257 (21, 21, C<sub>11</sub>H<sub>22</sub>Br), 245, 243 (10, 13, C<sub>12</sub>H<sub>22</sub>Br), 178 (24, C<sub>11</sub>H<sub>22</sub>), 177 (37, C<sub>11</sub>H<sub>21</sub>), 163 (100, C<sub>12</sub>H<sub>19</sub>). - Anal. Calcd. for C<sub>11</sub>H<sub>22</sub>Br<sub>2</sub>: C, 46.18; H, 6.56. Found: C, 46.17; H, 6.33.

**1,2-Bis(dibromomethyl)-3,3,4,4,5,5-hexamethyl-1-cyclopentene (11)**. A solution of **7** (5.0 g, 28 mmol) and bromine (24 g, 0.15 mol) in 600 mL CCl<sub>4</sub> was heated at reflux for 6 h. During this period small amounts (~ 50 mg) of azoisobutyronitrile were added every hour. When the solvent was evaporated, **11** instantaneously crystallized to yield colourless prisms (8.2 g, 59%) which were recrystallized from hexane. mp. 150 - 152°C. - <sup>1</sup>H NMR (CDCl<sub>3</sub>, +60°C): 0.83 (s, 6 H), 1.23 (s, 12 H), 6.83 (br. s, 2 H); (-60°C): 0.87 (s, 6 H), 1.07 (s, 3 H), 1.40 (s, 3 H), 6.06 (s, 1 H), 7.56 (s, 1 H). - Mass spectrum (70 eV): m/e = 500, 498, 496, 494, 492 (3%, 11%, 17%, 13%, 3%, M<sup>+</sup>), 485, 483, 481, 479, 477 (4, 18, 27, 19, 5, C<sub>12</sub>H<sub>19</sub>Br<sub>4</sub>), 419, 417, 415, 413 (12, 35, 36, 12, C<sub>11</sub>H<sub>22</sub>Br<sub>3</sub>), 338, 337, 336, 335, 334, 333 (34, 44, 68, 54, 34, 21, C<sub>11</sub>H<sub>22</sub>Br<sub>2</sub> und C<sub>11</sub>H<sub>19</sub>Br<sub>2</sub>), 297, 295, 293, 291 (20, 54, 47, 17, C<sub>10</sub>H<sub>19</sub>Br<sub>2</sub>, C<sub>10</sub>H<sub>17</sub>Br<sub>2</sub>), 281, 279, 277 (20, 43, 23, C<sub>9</sub>H<sub>17</sub>Br<sub>2</sub>), 257, 255 (C<sub>11</sub>H<sub>19</sub>Br), 215, 213 (79, 85, C<sub>10</sub>H<sub>17</sub>Br), 176 (85, C<sub>11</sub>H<sub>20</sub>), 161 (85, C<sub>12</sub>H<sub>17</sub>), 57 (100, C<sub>8</sub>H<sub>9</sub>). - Anal. Calcd. for C<sub>11</sub>H<sub>20</sub>Br<sub>4</sub>: C, 31.49; H, 4.06. Found: C, 31.38; H, 3.84.

**Bromination of 7 in CHCl<sub>3</sub>**. A 2M solution of bromine in CHCl<sub>3</sub> (12.5 mL, 25 mmol) was added dropwise to a solution of **7** (1.00 g, 5.5 mmol) in 12.5 mL CHCl<sub>3</sub> at 20°C. The mixture was washed with aqueous NaHSO<sub>3</sub> solution, dried, and the solvent evaporated. Chromatographic separation (Silica 60, petroleum ether/40 - 60°C) yielded **16** (80 mg, 4%) besides **9**, **10** and nonidentified products.

**r-1,c-2,c or t-4-tri(bromomethyl)-1,2,4-trimethyl-3,5-dimethylene-cyclopentane (16)**. Colourless viscous oil. - <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.28 (s, 6 H), 1.41 (s, 3 H), 3.40 (s, 2 H), 3.43 and 3.62 (AB system of 1,2-CH<sub>2</sub>Br with J = 10.5 Hz), 5.16 (d, J = 0.8 Hz, 2 H), 5.30 (d, J = 0.8 Hz, 2 H). - <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 22.6 (q, 1,2-CH<sub>2</sub>), 30.0 (q, 4-CH<sub>2</sub>), 39.6 (t, 1,2-CH<sub>2</sub>Br), 44.2 (t, 4-CH<sub>2</sub>Br), 47.0 (s, C-4), 51.9 (s, C-1,2), 111.7 (t, = CH<sub>2</sub>), 158.4 (s, C-3,5). - Mass spectrum (70 eV): m/e = 337, 335, 333 (13%, 24%, 11%, M<sup>+</sup>-Br), 323, 321, 319 (50, 100, 56, C<sub>12</sub>H<sub>17</sub>Br<sub>2</sub>), 133 (85, C<sub>11</sub>H<sub>19</sub>). - No UV-absorption for λ > 200 nm. - Anal. Calcd. for C<sub>11</sub>H<sub>19</sub>Br<sub>3</sub>: C, 37.62; H, 4.61. Found: C, 36.96; H, 4.55.

## REFERENCES

- 1) New address: Max-Planck-Institut für Med. Forschung, Abt. Organ. Chemie, Heidelberg.
- 2) H. Slebocka-Tilk, R.G. Ball, and R.S. Brown, *J. Am. Chem. Soc.* **107**, 4504 (1985).
- 3) P.B.D. De la Mare and R. Bolton, *Electrophilic Additions to Unsaturated Systems*, Elsevier, Amsterdam, 1982, pp 150.
- 4) J. Meisenheimer, *Liebigs Ann. Chem.* **456**, 142 (1927).
- 5) K. Ziegler and K. Bähr, *Ber. Deut. Chem. Ges.* **62**, 1695 (1929).
- 6) D. Lenoir, *Chem. Ber.* **111**, 411 (1978); D. Lenoir and B. Meyer, Abstract A44, Chemiedozententagung 1983, Dortmund.
- 7) H. Klein and H. Mayr, *Angew. Chem.* **93**, 1069 (1981); *ibid.* *Int. Ed. Engl.* **20**, 1027 (1981).
- 8) H. Mayr and U.W. Heigl, *Angew. Chem.* **97**, 567 (1985); *ibid.* *Int. Ed. Engl.* **24**, 579 (1985).
- 9) H. Günther, *NMR-Spektroskopie*, 2. Aufl., Thieme, Stuttgart, 1983, p. 229.
- 10) E.W. Meijer, R.M. Kellogg and H. Wynberg, *J. Org. Chem.* **47**, 2005 (1982).