

## Inverted cusps in electron spectra near zero electron velocity in inelastic ion-atom collisions

H. Böckl, R. Spies, and F. Bell

*Physics Section, University of Munich, D-8046 Garching, Federal Republic of Germany*

D. H. Jakubassa-Amundsen

*Physics Department, Technical University of Munich, D-8046 Garching, Federal Republic of Germany*

(Received 27 September 1983)

Expressions for inverted cusps in electron spectra near zero velocity from inelastic ion-atom collisions are derived by the plane-wave Born approximation. In addition, it is shown that for  $2p_0$  ionization not only the cusp-shaped electron-loss peak is inverted but the binary-encounter peak too.

Recently, Burgdörfer<sup>1</sup> showed that the usual cusp-shaped electron spectrum from inelastic ion-atom collisions is inverted for  $2p_0$  ionization of the projectile. Here it is assumed that the cusp is dominated by electron loss to the continuum (ELC).<sup>2</sup> Thus the cusp shows a dip at an electron velocity  $v_e = v_p$  which gets more pronounced with increasing projectile velocity  $v_p$ . Burgdörfer<sup>1</sup> obtained his results by an algebraic  $O(4,2)$  approach and showed a few graphical examples of electron spectra. In this Rapid Communication we will give a formula for the inverted cusp which shows a general scaling for arbitrary projectile-target combinations at any projectile velocity. This scaling is correct for hydrogenic wave functions of the initial and final electron state and for an unscreened Coulomb interaction. In the first-order Born approximation the double-differential cross section for the ejection of a projectile electron from an arbitrary state  $|nlm\rangle$  is easily obtained by means of partial derivatives of the ionization matrix element of the  $1s$  state. In the limit of vanishing electron velocity  $u$  (in the projectile frame), this matrix element acquires a simple form, such that the cross section reduces to a series of Legendre polynomials:

$$\lim_{u \rightarrow 0} \left( u \frac{d\sigma}{d\bar{u}} \right) = \sigma_0 \left( 1 + \sum_{\lambda=1}^n a_{2\lambda} P_{2\lambda}(\cos\theta) \right). \quad (1)$$

The target is thereby left in its ground state.  $\theta$  is the electron ejection angle relative to  $-\hat{v}_p$  and the beam direction is taken as quantization axis of the magnetic substates.  $\sigma_0$  and  $a_{2\lambda}$  are independent of  $\theta$ . The coefficients  $a_{2\lambda}$  scale with  $v_p/v_{or}$ , where  $v_{or} = Z_p/n$  is the orbital velocity of the electronic initial state and  $Z_p$  the projectile charge, but their structure depends on  $n$ ,  $l$ , and  $m$ . An equivalent expression for the double-differential cross section for the special case of  $1s$  ionization has been given by Day<sup>3,4</sup> and Briggs and Day.<sup>5</sup> For the  $2p_0$  ionization the angular distribution is determined by the coefficients  $a_2$  and  $a_4$ . Figure 1 shows  $a_2$  and  $a_4$  as a function of  $v_p/v_{or}$ . Transforming Eq. (1) to the laboratory frame and integrating over the angular acceptance  $\theta_0$  of the electron spectrometer (neglecting finite energy resolution) gives the scaled cusp yield  $I_{\text{cusp}}$  as a function of

the laboratory frame electron velocity  $v_e$ ,

$$I_{\text{cusp}}(\eta) = \text{const} \left[ (A + B - C) |\eta| - (1 + \eta^2)^{1/2} \left( \frac{A\eta^4}{(1 + \eta^2)^2} + \frac{B\eta^2}{1 + \eta^2} - C \right) \right], \quad (2)$$

with

$$\begin{aligned} A &= 35a_4/24; \quad B = 3a_2/2 - 15a_4/4, \\ C &= 1 - a_2/2 + 3a_4/8; \quad \eta = (v_e - v_p)/(v_e v_p)^{1/2} \theta_0. \end{aligned} \quad (3)$$

For  $v_p \rightarrow 0$  one obtains  $a_2 = a_4 = 0$  and for  $v_p \rightarrow \infty$  the coefficients are  $a_2 = \frac{5}{7}$ ,  $a_4 = -\frac{12}{7}$ . The latter values hold for an




FIG. 1. Angular distribution parameters  $a_2$  and  $a_4$  as a function of  $v_p/v_{or}$ . Solid lines: unscreened Coulomb interaction; dashed lines: exponentially screened Coulomb interaction. The Thomas-Fermi screening parameter  $\lambda = 1.13Z_T^{1/3}$  ( $Z_T$  is the target atomic number) was chosen to be that of an argon target (Ref. 6).




FIG. 2. The cusp yield  $I_{\text{cusp}}$  as a function of the scaled electron velocity (see text). The curves belong to different projectile velocities  $v_p$  ( $v_{\text{or}}=1$ ).

unscreened Coulomb interaction. Inspection of Fig. 1 reveals that  $a_2$  and  $a_4$  approach these limiting values logarithmically only. It follows that for small projectile velocities  $v_p$  one gets the usual cusp-shaped electron peak ( $A=B=0$ ,  $C=1$ ), but for  $v_p \rightarrow \infty$  the dip within the cusp reaches even zero electron yield ( $C=0$ ). Figure 2 shows the cusp yield as a function of the scaled electron velocity  $\eta$  for different projectile velocities. The curve for  $v_p=0$  might be somewhat unrealistic since the plane-wave Born approximation ceases to be valid in this limit. Clearly, the development of the cusp inversion with increasing projectile velocity can be seen.

In general, an inversion occurs for  $2a_2 - 8a_4/3 - 1 \geq 0$ . This condition holds not only for  $2p_0$ —but also for  $1s$  ionization, where  $a_4=0$ .<sup>3,4</sup> Thus even the cusp from  $1s$  ionization could be inverted for  $a_2 \geq \frac{1}{2}$ ,<sup>4</sup> at least in principle. It is interesting to note that, within the first Born approximation,  $a_2$  always stays below 0.5.<sup>3</sup> In addition, we have investigated  $2s$  ionization where both  $a_2$  and  $a_4$  do not vanish identical but again the condition for cusp inversion cannot be fulfilled within the whole  $v_p$  range. In our treatment of cusp inversion we have neglected the simultaneous excitation of the target atom. For a  $\text{He}^+/\text{Ar}$  interaction Burgdörfer<sup>1</sup> has found that this excitation did not seriously affect the dip structure since the doubly inelastic contribution was a small fraction of the total cusp cross section only.

Whereas Eqs. (1) and (2) hold only for zero electron velocity  $u$  in the projectile reference frame, Fig. 3 displays an isometric plot of  $d^2\sigma/dE d\Omega$  as a function of the scaled




FIG. 3. An isometric plot of the double-differential cross section for  $2p_0$  ionization (in units of  $b/\text{keV sr}$ ) of a Ne target by protons with  $v_p=20$  a.u.

electron velocities  $u_{\perp}$  and  $u_{\parallel}$ , perpendicular and parallel to the beam direction.  $E=0.5u^2$  is the electron energy and the cross section is plotted logarithmically. The ring-shaped pattern is the binary-encounter peak located at  $u=2v_p \cos\theta$ . It is clearly seen that the peak is inverted to a dip. Thus not only the cusp at  $u=0$  is inverted but the binary-encounter peak too. Physically this is easy to understand. Since the cross section at the location of the binary-encounter peak is proportional to the electron momentum density projected on the beam direction,<sup>7</sup> the usual peak gets a dip: The  $2p_0$  momentum distribution is vanishing everywhere perpendicular to the beam direction; thus the projection is zero. In contrast, the  $2p_1$  distribution peaks perpendicular to the beam direction. This yields an ordinary peak of the cross section at  $u=2v_p \cos\theta$ . It follows that the Fano-Macek alignment parameter  $A_0^{\text{col}}$  (Ref. 8) reaches its maximum possible positive value  $A_0^{\text{col}} = +0.5$  at the position of the binary-encounter peak. This holds for projectile velocities where the peak is really inverted, i.e., for  $v_p/v_{\text{or}} > 1$ . From the discussion above it is evident that every magnetic substate shows peak inversion if the momentum distribution has a nodal plane perpendicular to the beam direction. Thus any substate  $|nlm\rangle$  shows inversion if the sum  $(l+m)$  is odd. In conclusion, we note that peak inversion is a general property of certain magnetic substates and occurs in any first-order theory both for ionization and bound-state charge transfer.<sup>9</sup>

This work has been partly supported by the Bundesministerium für Forschung und Technologie and Gesellschaft für Schwerionenforschung, Darmstadt.

<sup>1</sup>J. Burgdörfer, Phys. Rev. Lett. **51**, 374 (1983).

<sup>2</sup>I. A. Sellin, M. Breinig, W. Brandt, and R. Laubert, Nucl. Instrum. Methods **194**, 395 (1982).

<sup>3</sup>M. H. Day, J. Phys. B **14**, 231 (1981).

<sup>4</sup>M. H. Day, J. Phys. B **13**, L65 (1980).

<sup>5</sup>J. S. Briggs and M. H. Day, J. Phys. B **13**, 4797 (1980).

<sup>6</sup>F. Drepper and J. S. Briggs, J. Phys. B **9**, 2063 (1976).

<sup>7</sup>F. Bell, H. Böckl, M. Z. Wu, and H.-D. Betz, J. Phys. B **16**, 187 (1983).

<sup>8</sup>U. Fano and J. Macek, Rev. Mod. Phys. **45**, 553 (1973).

<sup>9</sup>D. H. Jakubassa-Amundsen, J. Phys. B **14**, 2647 (1981).