

An
International
Journal
for
Reviews
and
Communications
in
Heterocyclic
Chemistry

HETEROCYCLES

Founded
by
Tetsuji Kametani

Volume 29

Number 1



The Japan Institute of Heterocyclic Chemistry



Elsevier Science Publishers

HETEROCYCLES

An International Journal for Reviews and Communications
in Heterocyclic Chemistry

CONTENTS

COMMUNICATIONS

- M. AVALOS GONZÁLEZ, R. BABIANO CABALLERO, P. CINTAS MORENO, J. FUENTES MOTA, J. L. JIMÉNEZ REQUEJO, and J. C. PALACIOS ALBARRÁN. Reaction of Glycosylisothiocyanates with 2-Chloroethylamine 1
- KENJI YAMAGATA, HIROSHI MARUOKA, YOSHICHIKA HASHIMOTO, and MOTOYOSHI YAMAZAKI. Ring Conversion of 2-Amino-4,5-dihydro-3-furancarbonitriles Catalyzed by Halide Ions 5
- ULF PINDUR, MANFRED EITEL, and ERFANIAN ABDOUSTHOUSHANG. Regioselectivity in the Diels-Alder Reactions of 2-Vinylindoles and Pyrano[3,4-*b*]indol-3-ones with CC-Dienophiles. 11
- KIYOSHI MATSUMOTO, KENSAKU HAMADA, TAKANE UCHIDA, and HIROSHI YOSHIDA. Formation of Novel 1 : 3 Adduct in the High Pressure Reaction of 2(1H)-Pyridones with Dimethyl Acetylenedicarboxylate : An X-Ray Crystal Structure Determination 21
- AKIRA MATSUDA, HITOMI OKAJIMA, and TOHRU UEDA. Synthesis of 2',3'-Dideoxy-3'-methylidenethymidine and 2',3'-Didehydro-2',3'-dideoxy-3'-methylthymidine : Deoxygenation of the Allylic Alcohol System in 3'-Deoxy-3'-methylidene-5-methyluridine 25
- KLAUS TH. WANNER and ILONA PRASCHAK. Asymmetric Electrophilic α -Amidoalkylation 6 : Syntheses of Tetrahydroisoquinolines of High Enantiomeric Purity. 29
- YONG-GYUN LEE, HIDEHARU IWASAKI, YOHSUKE YAMAMOTO, KATSUO OHKATA, and KIN-YA AKIBA. (4+2)Type Cycloaddition of Chromones with α,β -Unsaturated Ketones Mediated by *t*-Butyldimethylsilyl Triflate : One-pot Preparation of Xanthone Derivatives 35
- MASAMITSU OCHI, KENSUKE FUTATSUGI, YASUHIRO KUME, HIYOSHIZO KOTSUKI, KAZUHIKO ASAO, and KOZO SHIBATA. Litophytolides A and B, Two New Lipid Meta-

bolites of a Soft Coral <i>Litophyton</i> sp.	39
---	----

PAPERS

RONALD G. MICETICH, RAJESHWAR SINGH, CHEN C. SHAW, SAMARENDRA N. MAITI, MAYA P. SINGH, and PAUL SPEVAK. The Chemistry of the 6 β -Diacylaminopenicillins- . . .	43
ABDULLA J. HAMDAN and HAROLD W. MOORE. A New Syn- thetic Route to Heterocyclic Quinones	51
ANA M. CUADRO, JULIO ALVAREZ-BUILLA, and JUAN J. VAQUERO. Preparation of New Benzimidazole Derivatives from <i>N</i> -[(Methylthio)thiocarbonylmethyl]azinium Salts	57
PÉTER MÁTYUS, NÁNDOR MAKK, ENDRE KASZTREINER, and GYULA JERKOVICH. A Novel Method for the Synthesis of 2-Haloalkyl-3(2 <i>H</i>)-pyridazinones by <i>O</i> → <i>N</i> -Alkyl Rearrange- ment	67
JAN KEIJRSERS, BENNO HAMS, CHRIS KRUSE, and HANS SCHEEREN. High-Pressure Diels-Alder Reactions of 1- Methoxycarbonyl-3-methylthio- and -3-phenylthiopyrrole. An Efficient Entry to 7-Azabicyclo[2.2.1]heptanes	79
SALIMUZZAMAN SIDDIQUI, BINA SHAHEEN SIDDIQUI, TARIQ MAHMOOD, and SHAHEEN FAIZI. Tetranortriterpenoids from <i>Azadirachta Indica</i> A.Juss (Meliaceae)	87
FRANCESCO LUCCHESINI, NEVIO PICCI, MARCO POCCHI, AN- GELA DE MUNNO, and VINCENZO BERTINI. Synthesis of 4-Unsubstituted Isothiazoles	97
ANDRÉ MAQUESTIAU, ROBERT FLAMMANG, and FOUAD- BACHIR B. ABDELOUAHAB. Flash-Vacuum Pyrolysis of 1, 2,4-Triazolides : A New Synthesis of Functionalized Oxazoles .	103
MARÍA J. CLIMENT, HERMENEGILDO GARCÍA, SARA IBORRA, MIGUEL A. MIRANDA, and JAIME PRIMO. Photosensitized Dehydrogenation of Flavanones to Flavones using 2,4,6- Triphenylpyrylium Tetrafluoroborate (TPT)	115
TOKUHIRO WATANABE, KAZUHIKO HAYASHI, JUN SAKURA- DA, MICHIYO OHKI, NORIKO TAKAMATSU, HARUMI HIROHATA, KEIKO TAKEUCHI, KAYO YUASA, and AKIHIRO OHTA. Alkylation and Arylation of Pyrazines by Organotin Compounds.	123
EDDIE V. TAO, JOHN BRENNAN, JOHN K. SWARTZEN- DRUBER, and JACK B. DEETER. Synthesis of New Racemic Bicyclic γ -and δ -Lactams Based on Two-fold Intramolecular Cyclization	132
JEAN-MICHEL VIERFOND, LUCIEN LEGENDRE, JACQUELINE MAHUTEAU, and MARCEL MIOCQUE. Annelation of Quinoxaline by Sulfur Stabilized Carbanions	141


TATSUO NAGASAKA, HIROTO YAMAMOTO, HIDEKI HAYASHI, MARIKO WATANABE, and FUMIKO HAMAGUCHI. Wittig Reactions of 1-Alkoxy carbonyl-2-hydroxypyrrolidines and -piperidines : Syntheses of (±)-Hygrine and (±)-2-Epilasubine II	154
M. TERESA RAMOS, CARMEN AVENDAÑO, JOSÉ ELGUERO, M. LUISA JIMENO, JUANA BELLANATO, FELICIANA FLORENCIO, and JULIA SANZ-APARICIO. Tautomerism of Bis (2-benzothiazolyl) arylmethanes	165
JULIO A. SEIJAS, M. PILAR V. TATO, RAMÓN ESTÉVEZ, LUIS CASTEDO, and RICARDO RIGUERA. New Total Synthesis of Patulin	181
LING-LING TIEN, SHAW-TAO LIN, and HUI-JEAN CHIANG. Mass Spectrometric Fragmentation of the Protonated Sydnones under Chemical Ionization	185
JIN YUN SUN, YOSHIO HANO, and TARO NOMURA. On the Structure of Sanggenon Q, A New Diels-Alder Type Adduct from <i>Morus mongolica</i> Schneider.	195

ASYMMETRIC ELECTROPHILIC α -AMIDOALKYLATION 6¹:
 SYNTHESIS OF TETRAHYDROISOQUINOLINES OF HIGH ENANTIOMERIC PURITY.

Klaus Th. Wanner* and Ilona Praschak
 Institut für Pharmazie und Lebensmittelchemie der
 Universität München
 Sophienstr. 10, 8000 München 2

Abstract - Syntheses of 1-substituted tetrahydroisoquinolines of high enantiomeric purity are described. As key step a diastereoselective trapping reaction of a chiral acyliminium ion with silyl enol ethers is involved. The starting acyliminium ion can be prepared by hydride abstraction with triphenylcarbonium tetrafluoroborate.

There are at present several different methods for preparing 1-substituted 1,2,3,4-tetrahydroisoquinolines in enantiomerically pure form. They are based on resolution, enantioselective catalytic hydrogenation, diastereoselective alkylation of chiral formamide derived anions and on the incorporation of carbon fragments derived from the chiral pool. A survey of these methods has been provided in a recent paper². The interest in tetrahydroisoquinolines is due to their wide distribution among both alkaloids³ and biologically active compounds of synthetic origin⁴.




We have now found a novel method for the synthesis of α -substituted amines in optically pure form that we have termed asymmetric electrophilic α -amidoalkylation (AE α A). The trapping reaction of a chiral acyliminium ion ($R^*CO=$ chiral auxiliary in 1), which represents an asymmetric electrophilic α -amidoalkylating agent, with a nucleophile ("Nu-") was expected to proceed with stereoselective bond formation. As a result thereof one diastereomer (2 or 3) should predominate which after purification to a single diastereomer and subsequent removal of the chiral

auxiliary would yield an enantiomerically pure amine. This plan has been successfully implemented with pyrrolidines⁵ and piperidines⁶.

In this communication we wish to report the extension of this concept to the synthesis of tetrahydroisoquinolines.

First there was need for an efficient method for generating the acyliminium ion 6. Hydride abstraction⁷ from 4 seemed to be the most direct route. Upon treatment of amide 4 with triphenylcarbonium tetrafluoroborate 5 at room temperature (in CH₂Cl₂) the starting material was indeed completely consumed within 16 h and a yellow precipitate formed.



Addition of extra solvent resulted in a clear yellow solution to which enol ether 8a (1.2 eq. in CH₂Cl₂) was added at -78°C. After workup a mixture of diastereomers (S)-9a and (R)-10a was obtained. On SiO₂ no separation of (S)-9a from (R)-10a could be effected although various solvents were employed. The composition of the diastereomeric mixture was determined as 20.3/79.7 by HPLC using a chiral stationary phase⁸ (see Table 1). The reaction also proceeded smoothly at -90°C. Thereby the (S)-9a/(R)-10a ratio was improved to 17.6/82.4 and the mixture of diastereomers was obtained in a combined yield of 97.2%. An even more stereoselective reaction could be brought about by adding TiCl₄ (1 eq. at -78°C) prior to the addition of silyl enol ether 8a. In this case at -78°C and -93°C (temp. of enol ether addition) diastereomeric ratios of 12.4/87.6 and 11.2/88.8 (Table 1, entries 3 and 4), respectively, could be achieved. The results of the bond forming

reactions (at -78°C , without TiCl_4) with silyl enol ethers 8b and 8c can be taken from Table 1. In these cases the influence of lower temperatures and TiCl_4 has not been investigated yet.

Table 1.

en- try	<u>8</u>	R ¹	R ²	Reaction cond.		Yield [%]	Ratio		(R)- <u>12</u>	
				Add.	Temp.[$^{\circ}\text{C}$]		(S)- <u>9</u> /(R)- <u>10</u>	(S)- <u>9</u> /(R)- <u>10</u>	(S)- <u>11</u> /(R)- <u>12</u>	Yield [%] ^{a)} [α] ₅₇₈
1	a	H	H		-78	---	20.3/79.7	---	---	---
2	"	"	"		-90	97.2	17.6/82.4	16.7/83.3	59.1	-119.6 $^{\circ}$
3	"	"	"	1 eq. TiCl_4	-78	---	12.4/87.6	---	---	---
4	"	"	"	1 " "	-93	---	11.2/88.8	---	---	---
5	b	H	Cl		-78	98.9	14.8/85.2	9.9/90.1	5.3 ^{b)} 51.9 ^{c)}	-102.9 $^{\circ}$
6	c	OCH_3	OCH_3		-78	96.8	17.6/82.4	18.5/81.5	37.8	- 83.0 $^{\circ}$

a) Pure compound according to HPLC. b) Pure compound for analytical purposes only. c) Hard to separate mixture of (R)-12a and (R)-12b.

Compounds (S)-9/(R)-10 a-c have been employed in the synthesis of 1-substituted tetrahydroisoquinolines (R)-13 including the alkaloid (-)-homolaudosine ((R)-13d) as outlined below. By hydrogenation in acetic acid/1% trifluoroacetic acid using Pd on carbon (10% Pd) as catalyst the amides (S)-11/(R)-12 a-c were obtained. Said diastereomers (S)-11/(R)-12 proved to be separable on silica gel. The ratio (S)-11/(R)-12 was found to be almost the same as the (S)-9/(R)-10 ratio (see Table 1.) except for the compounds with a p-chlorophenyl substituent (Table 1. entry 5). This might be due to a difference in the rate of a side reaction which occurred during hydrogenation. Beside the carbonyl oxygen also the chlorine substituent had been removed to some extent (up to 36%).

The yields of the major diastereomers (R)-12 in pure form obtained after flash chromatography are also set forth in Table 1. The next step involved the cleavage of the amide bond and was first attempted by means of LiOH in dioxane/ H_2O (3/1, sealed tube, -150°C), but without any noticeable success.

Finally we found that amides (R)-12a - (R)-12c can be cleaved by treatment with LiAlH_4 in THF (2.5 - 5.0 mol.-eq., 2-3 h, 20°C). The pure amines (R)-13a - (R)-13c were thus obtained in fairly good yields (Table 2) after flash chromatography

(SiO₂, Et₂O/NEt₃=95/5). The lower yield for (R)-13b is due to the fact that for convenience a mixture of (R)-12b and (R)-12a (obtained from hydrogenolysis, vide supra) had been employed as starting material.




Table 2.

(R)- <u>13</u>					
	R ¹	R ²	R ³	Yield [%]	[α] ₅₇₈
a	H	H	H	64.7	+25.5°
b	Cl	H	H	16.1 ^{a)}	+12.4°
c	OCH ₃	OCH ₃	H	77.0	+15.1°
d	OCH ₃	OCH ₃	CH ₃	64.3	-13.5°

a) Based on starting material contaminated with 36% (R)-12a.

The R stereochemistry of (R)-13b has been established by comparing the optical rotation ([α]₅₇₈ = +12.4°, [α]₅₄₆ = +11.5°; c = 1.05; CH₃OH) with reported literature values⁹ ([α]_D = +15°; CH₃OH). The R configuration of (R)-13a could be deduced from the product distribution that was observed after hydrogenation of (S)-9b/(R)-10b. Thereby in addition to (S)-11b/(R)-12b a mixture of (S)-11a/(R)-12a was formed (vide supra) wherein the predominant isomer was the same as in the product obtained along the direct route from (S)-9a/(R)-10a (verified by HPLC).

The final step in the synthesis of (R)-homolaudanosine ((R)-13d) was the methylation of (R)-13c (CH₂O, 2.5 eq. NaCNBH₃) which afforded (R)-13d in 64.3% yield. The optical rotation of the obtained product ([α]₅₇₈ = -13.5°, c = 0.89, EtOH) compared to that of the natural enantiomer¹⁰ ((S)-13d: [α]_D = + 11°, c = 0.21, EtOH) indicates that the synthesized compounds and its precursors belong to the (R)-series. In order to verify the optical purity of the amines (R)-13 a sample of (R)-13a was treated with (-)-camphanic acid chloride. HPLC revealed (R)-12a as the prevailing compound, accompanied by trace amounts (< 0.4%) of a side product (possibly (S)-11a). Therefore the optical purity (ee) of (R)-13a - (R)-13d should at least be higher than 99%.

In conclusion, it has been shown that the chiral acyliminium ion (6) can be obtained by a simple hydride abstraction reaction. Subsequent trapping reactions

proceed with stereoselective bond formation and the resulting products are useful intermediates in the synthesis of 1-substituted tetrahydroisoquinolines of high enantiomeric purity.

ACKNOWLEDGEMENT

We are greatly indebted to Prof. F. Eiden for his generous support. Financial support of this work by Deutsche Forschungsgemeinschaft is gratefully acknowledged.

REFERENCES

1. Part 5: K.Th. Wanner, A. Kärtner, and E. Wadenstorfer, Heterocycles, in print.
2. J.M.P. Huber and D. Seebach, Helv. Chim. Acta, 70, 1944 (1987).
3. R.H.F. Manske: "Isoquinoline Alkaloids", In: The Alkaloids (Manske, ed.) Academic Press, London, 1960, Vol. 7, 423; J. Lundström: "Simple Isoquinoline Alkaloids". In: The Alkaloids (Brossi, ed.), Academic Press, London, 1983, Vol. 21, 255.
4. A. Brossi, H. Besendorf, L.A. Pirk, and A. Rheiner: "Isoquinoline Analgetics". In: Analgetics, Medicinal Chemistry (G. deStevens, ed.), Academic Press, London, 1965, Vol. 5, 281.
5. K.Th. Wanner and G. Höfner, Arch. Pharm. (Weinheim), in print.
6. K.Th. Wanner and A. Kärtner, Heterocycles, 26, 921 (1987); K.Th. Wanner and A. Kärtner, Arch. Pharm. (Weinheim), 320, 1253 (1987).
7. For a related oxidation with DDQ see: M. Onda, K. Yuasa, and J. Okada, Chem. Pharm. Bull., 22, 2365 (1974).
8. Covalently bound (R)-N-(3,5-dinitrobenzoyl)phenylglycine (J.T. Baker, Chemical Co.).
9. A. Brossi and F. Burkhardt, Helv. Chim. Acta, 44, 1558 (1961).
10. A.J. Aladesanmi, Ch. J. Kelley, and J.D. Ceasy, J. Nat. Prod., 46, 127 (1983).

Received, 8th September, 1988